

25

EPIDEMIOLOGY OF PRIMARY MALIGNANT MESOTHELIAL TUMORS IN CANADA*

A. D. McDONALD, A. HARPER, O. A. EL ATTAR, AND J. C. McDONALD

All fatal malignant mesothelial tumors known to pathologists in Canada between 1959 and mid-1968 were registered. They numbered 165 (1 per million population per annum): two thirds were in males. Occupational and residential histories were obtained "blind" from relatives and friends of 90% of the cases and 2 matched control series. An association with definite or probable occupational exposure to asbestos was clearly demonstrated, but only 20% of male cases and one female case had any such contact. Almost all the excess was in the manufacture and industrial application of asbestos rather than in mining or milling. No association was found with lesser degrees of occupational exposure or residence in asbestos-mining areas, but there was a small excess of possible domestic exposures. The smoking histories in the mesothelial tumor and main control groups were almost identical and unlike those for cases of primary lung cancer.

EVIDENCE FOR AN ASSOCIATION BETWEEN Malignant mesothelioma and asbestos exposure rests mainly on the remarkable concentration of cases reported by Wagner et al.⁵ in the crocidolite mining areas of South Africa, the case-control studies of Elmes et al.¹ in Belfast, and Newhouse and Thompson² in London, and the considerable excess mortality due to this disease observed by Selikoff et al.³ in American insulation workers. The survey described here was undertaken to obtain a more generally representative view of the problem in one of the 2 major chrysotile-producing countries of the world. Our aim has been to study all fatal cases known to have occurred in Canada since 1959.

METHOD

At the end of 1967 we wrote to the 425 members of either the Canadian Association

From the Department of Epidemiology and Health, McGill University, Montreal, Canada.

*A preliminary report on these results was presented at the International Conference on Pneumoconiosis at Johannesburg in April, 1969.

Supported by a grant from the Institute of Occupational and Environmental Health of the Quebec Asbestos Mining Association.

This survey was made possible by the great help given us by pathologists and members of provincial and other health departments throughout Canada. We are also much indebted to all those who did the field work and to the friends and relatives of the deceased for their cooperation.

Received for publication April 9, 1970.

of Pathologists or the Quebec Association of Laboratory Physicians inquiring whether they had seen any fatal cases of primary malignant tumors of the pleura or peritoneum, diagnosed by autopsy or biopsy since 1959. In due course we obtained a reply from them all, and, if this was in the affirmative, a personal visit was made by one of us to review the records and to exclude cases which did not meet these criteria or about which the pathologist had serious doubt. In all, 165 cases were found: 113 were pleural, 45 peritoneal, 3 pleural and peritoneal, and 4 pericardial. The diagnosis was supported by autopsy in two thirds of the cases and by biopsy in the remainder. Later, we plan to have material from as many as possible reviewed by a panel of pathologists.

During the visit to the pathologist, 2 fatal control cases—one primary and one secondary lung cancer—were selected from the autopsy or biopsy records, matched for sex and, as closely as possible, for age and date of death. In 2 instances, no adequately matching case of primary lung cancer could be found. Two control series were chosen in order to examine 2 types of question. Primarily, we wished to know whether persons who had died from malignant mesothelial tumors had differed in their past exposure to asbestos and certain other materials from comparable persons in the general population. Comparability demanded that the control series should comprise persons matched with the cases for

No. 4

age, sex, date and mortem diagnostic information obtained of equal quality. terminal illness secondary lung cancer criteria reasonably were advantage that were included, but second main question mesothelial tumor-smoking in the same, and, for this cases of this disease

A field survey by investigators (doctor and public health relatives and friend pleted a detailed questionnaire French relating to direct household exposure, and smoking was given to the companies were listed of the exact nature relative was then as involved contact with cement, copper, nickel a check list was used subject had ever been following: locomotive ing, overhauling, or large boilers or steam mestic heating equipment, fireproofing, fireproofing prefabricated ing buildings; man or clutch plates; mi

Year	
1960	
1961	
1962	
1963	
1964	
1965	
1966	
1967	
1968 (1st h	
Incidence/mil	
1960-1967	

T

ALD

be-
lion
den-
the
able
6 of
was
a in
upa-
mall
elial
for

Association of
g whether they
nary malignant
neum diagnosed
l. In due course
all, and, if this
sonal visit was
he records and
not meet these
pathologist had
es were found;
l. 3 pleural and
The diagnosis
wo thirds of the
mainder. Later,
om as many as
pathologists.
hologist, 2 fatal
nd one second-
d from the au-
ied for sex and
ge and date of
equately match-
ancer could be
re chosen in or-
tion. Primarily,
ersons who had
lial tumors had
to asbestos and
comparable per-
on. Comparabil-
ol series should
th the cases for

age, sex, date and place of death, and post-mortem diagnostic procedure. To insure that information obtained from relatives would be of equal quality, it was important that the terminal illness should also be similar. Secondary lung cancer seemed to meet these criteria reasonably well and had the additional advantage that several primary cancer sites were included, but none were in the lung. A second main question was whether primary mesothelial tumors were related to tobacco smoking in the same way as primary lung cancer, and, for this reason, a matched series of cases of this disease was also selected.

A field survey was arranged in which investigators (doctors, public health nurses, and public health inspectors) sought out the relatives and friends of the deceased and completed a detailed questionnaire in English or French relating to occupation, residence, indirect household exposure to dust, family history, and smoking habits. Special emphasis was given to the occupational history. So far as possible, all employers or employing companies were listed with dates and a note made of the exact nature of the work done. The relative was then asked whether each job involved contact with wood, rubber, asbestos, cement, copper, nickel, or fiber glass. Finally, a check list was used, to inquire whether the subject had ever been employed in any of the following: locomotive repair shop; shipbuilding, overhauling, or breaking; installation of large boilers or steam pipes; insulation of domestic heating equipment; building—soundproofing, fireproofing, heat retention, assembling prefabricated buildings, and demolishing buildings; manufacture of brake linings or clutch plates; mining industry; textile in-

dustry; cement industry; dock or transport industry.

No interviewer knew to which group the case under investigation belonged, and the information recorded in the questionnaire was also coded blind. Questionnaires were completed for 447 (91%) of the 493 cases and controls.

FINDINGS

The distribution of cases by province and year of death is shown in Table 1, and by age, sex, and site in Table 2. More cases were found in 1966 and 1967 than in previous years. Recent interest in these tumors may have contributed to this, and, as the records in some hospitals were poorer for the earlier years, memory may also have been a factor. With these points in mind, the true rate of increase is difficult to assess. During the 8 complete years, 1960–1967, the average annual incidence was 1.0 per million of population. The rate was appreciably higher in Quebec (1.5) than in Ontario (0.8) or in the rest of Canada (0.9), 1966 and 1967 again accounting for much of the difference. Three quarters of the cases were aged 50 or more at death, and males predominated by 2.4 to 1. Under 50, there was little difference in the sex distribution.

The case and 2 control series were very similar in age distribution and in the proportion of each for which the questionnaire was completed (Table 3). Table 4 shows the number of men aged 18 years or more who were occupationally exposed up to 10 years before death to asbestos and certain other materials in dust or vapor form. Whereas all 3

TABLE 1. Distribution of Cases by Year of Death and Province

Year	Ontario	Quebec	Other Provinces	Canada
1960	—	6	9	15
1961	1	6	3	10
1962	4	4	5	13
1963	9	3	7	19
1964	6	8	3	17
1965	7	7	5	19
1966	8	14	9	31
1967	6	13	6	25
1968 (1st half)	5	9	2	16
	46	70	49	165
Incidence/million/year 1960–1967	0.8	1.5	0.9	1.0

TABLE 2. Distribution of Mesothelial Tumors by Age, Sex, and Site

Age (years)	Pleura		Peritoneum		Pleura and peritoneum		Pericardium		Total	
	M	F	M	F	M	F	M	F	M	F
0-	—	—	—	1	—	—	—	—	—	1
10-	2	1	—	—	—	—	—	—	2	1
20-	—	4	—	—	—	—	—	—	—	4
30-	7	4	1	1	—	—	—	1	8	6
40-	9	7	4	1	—	—	—	1	13	9
50-	18	3	11	4	1	—	1	1	31	8
60-	26	5	2	5	—	2	—	—	28	12
70-	15	5	7	6	—	—	—	—	22	11
80-	3	4	1	1	—	—	—	—	4	5
All	80	33	26	19	1	2	1	3	108	57
TOTAL	113		45		3		4		165	

groups were equally exposed to wood dust, relatively more men in the mesothelial tumor group were exposed to all the other materials, the greatest excess being for asbestos. Men who were exposed to asbestos tended also to be exposed to other materials, especially cement and fiber glass (Table 4), but these were only responsible for part of the excess exposure to the other materials. The excess of "definite" or "probable" exposure to asbestos (Table 5) in the mesothelial tumor group compared with both control groups was substantial ($P < .001$), but there was virtually no difference in the frequency of jobs in which exposure was classified as "possible." In 60% of the cases, however, occupational exposure to asbestos was considered "unlikely." Only 2 women in all had any occupational exposure to asbestos; there was a definite exposure in the mesothelial tumor

group and a possible exposure in the secondary lung cancer group.

The occupations involving definite or probable exposure to asbestos are listed in Table 6. Men employed in asbestos textile manufacture, the installation of brake linings, or as insulators were responsible for most of the excess of cases over controls. Mine or mill workers made little contribution to the difference, and only one person, a hardware storeman, had worked with asbestos-cement products. The time between first exposure and death for the 21 cases in the definite or probable class ranged from 16 to 50 years; 4 men and one woman who worked on the job for less than 4 years had intervals of 18, 26, 26, 29, and 33 years.

Materials other than asbestos also deserve further consideration since several men had worked with more than one of them. Exclud-

TABLE 3. Distribution by Age and Sex of Cases and Controls

Age	Males			Females		
	Mesothelial tumors	Secondary lung cancer	Primary lung cancer	Mesothelial tumors	Secondary lung cancer	Primary lung cancer
0-		1 (1)	1 (1)	1 (1)	1 (1)	1 (1)
10-	2 (2)			1 (1)		
20-		2 (2)		4 (3)	2 (2)	
30-	8 (8)	6 (6)	2 (2)	6 (6)	6 (5)	3 (2)
40-	13 (10)	14 (11)	24 (20)	9 (8)	10 (10)	14 (13)
50-	31 (31)	26 (22)	24 (23)	8 (7)	7 (7)	8 (7)
60-	27 (25)	35 (32)	35 (33)	12 (11)	15 (14)	12 (7)
70-	23 (22)	20 (19)	17 (15)	11 (10)	13 (11)	15 (15)
80-	4 (3)	4 (4)	4 (4)	5 (4)	3 (3)	3 (2)
All	108 (101)	108 (97)	107 (95)	57 (51)	57 (53)	56 (47)
Mean	58.7	59.1	59.1	55.8	57.4	58.8
SD	13.9	13.3	12.5	19.3	17.3	15.5

No. of questionnaires completed shown in parentheses ().

* Definite number of 1

ing wood difference to cement ber were were in 1 to asbestos in 19 mer 2 control An analy Table 7, appeared tact with (9:1:1) ra seemed u bestos. Tl in the n copper: 1.

TABLE

Males

Expost

Males: 1
P < 0.001

Female

Expost

* One ch

TABLE 4. Occupational Exposure of Men to Certain Materials*

Total		Mesothelial tumors	Secondary lung cancer	Primary lung cancer
F		(100 men)	(96 men)	(97 men)
1	Asbestos	20	3	4
1	Cement	17 (4)	9 (0)	8 (0)
4	Copper	6 (2)	1 (0)	1 (0)
6	Fiber glass	6 (3)	0	0
9	Nickel	5 (2)	0	0
8	Rubber	3 (1)	0	0
12	Wood	18 (1)	19 (0)	16 (0)
11	Total exposures recorded	75 (13)	32 (0)	29 (0)
5				
57				

* Definite or probable exposure to the substances named in dust or vapor form in men aged 18 or more. The number of men who were also definitely or probably exposed to asbestos is given in parentheses.

165

in the second.

Definite or probable exposure to asbestos is listed in Table 4; textile manufacture, linings, or for most of the

Mine or mill work on the job for 10 years: 4 men on the job for 15, 18, 26, 26,

as also deserve consideration. Several men had no exposure to asbestos. Exclud-

ing wood, for which there was evidently no difference between the 3 groups, 37 exposures to cement, copper, fiber glass, nickel, or rubber were recorded for the cases, of which 12 were in men definitely or probably exposed to asbestos. The 25 remaining exposures were in 19 men, and the comparable figures for the 2 control groups were 10 and 9, respectively. An analysis of these exposures is shown in Table 7, where it may be seen that the excess appeared to be in occupations involving contact with copper, nickel, fiber glass, or rubber (9:1:1) rather than with cement only, and seemed unrelated to possible exposure to asbestos. The relevant occupations of the 9 men in the mesothelial group were as follows: *copper*: lathe operator (2), munition manu-

facture, brass foundry; *nickel*: lathe operator (2), mine worker; *rubber*: tire manufacture, plantation work; *fiber glass*: municipal maintenance work, wood-caulking (ship).

Residential histories indicated that 45% of male cases and 38% of female cases had spent their childhood entirely in a rural area compared with 40% of males and 38% of females with secondary lung cancer, but the primary lung cancers were less often reared in rural areas (males 31%, females 28%). No difference was found between cases and both control groups after the age of 15 years. Excluding 5 cases (3 mine or mill workers and 2 textile workers) and one mine and mill worker in each control group, who lived in mining towns, only 3 men and 2 women had

TABLE 5. Distribution of Cases and Controls According to Their Occupational Exposure to Asbestos

Males	Mesothelial tumors		Secondary lung cancer		Primary lung cancer	
Exposure to asbestos	No.	%	No.	%	No.	%
Definite	11	20.0	1	3.1	1	4.1
Probable	9		2		3	
Possible	20	20.0	22	22.9	16	16.5
Unlikely	60	60.0	71	74.0	77	79.4
All*	100	100.0	96	100.0	97	100.0

Males: Distribution of definite and probable groups compared with possible and unlikely: $\chi^2 = 21.5$, 2 d.f., $P < 0.001$

Females	Mesothelial tumors		Secondary lung cancer		Primary lung cancer	
Exposure to asbestos	No.	%	No.	%	No.	%
Definite	1	2.0	—	—	—	—
Probable	—	—	—	—	—	—
Possible	—	—	1	1.9	—	—
Unlikely	49	98.0	51	98.1	46	100.0
All*	50	100.0	52	100.0	46	100.0

* One child in each group dying under the age of 18 has been excluded.

3 (2)

14 (13)

8 (7)

12 (7)

15 (15)

3 (2)

56 (47)

58.8

15.5

TABLE 6. Distribution of Occupations Classified Under Definite or Probable Exposure to Asbestos

	Mesothelial tumors	Secondary lung cancer	Primary lung cancer
Asbestos mine or mill	3	1	1
Asbestos textile manufacture	4*	0	0
Brakelining installation	2	0	0
Insulation	5	0	0
Other occupations involving contact with insulation materials	5	2	3
Hardware store (handling asbestos sheet and pipe)	1	0	0
Filled gas masks with asbestos textile	1	0	0
TOTAL	21	3	4

* Includes one person who worked in brakelining manufacture.

ever lived within 20 miles of an asbestos mine. One of the 5 was a case. 3 had secondary lung cancer, and one had primary lung cancer.

The possibility of contact as a result of some other member of the family bringing home dusty clothes from occupations involving the use of asbestos was also examined. Excluding 7 persons (5 cases and one in each control group) who were occupationally exposed themselves (definitely or probably), there was one case in a female with definite home exposure and one male with probable home exposure. In addition to 5 persons (one case and 4 with primary lung cancer) who were possibly occupationally exposed to asbestos themselves, there were 26 persons with a possible home exposure: 14 cases, 6 with secondary lung cancer, and 6 with primary lung cancer. The category of possible home exposure was given in households where a person possibly exposed at work to asbestos brought home dusty clothes. Thus, excluding those persons with definite or probable occupational exposure, there were 15 persons with mesothelial tumors, 8 with secondary lung cancer, and 10 with primary lung cancer, who had had at least possible home exposure to asbestos dust.

The frequency of cancer reported in close relatives did not differ from that in the control groups. An analysis of cigarette smoking habits is shown in Table 8. In both males and females, persons in the primary lung cancer group had smoked considerably more than those in either the case or secondary lung cancer groups. The latter 2 groups had almost an identical distribution of smoking habits. The 20 men with definite or probable occupational exposure to asbestos showed no great difference in smoking habits from cases without occupational exposure.

DISCUSSION

Though our study is not directly comparable with those of Elmes et al.¹ in Belfast, and Newhouse and Thompson² in London, it seems fairly clear that the difference between cases and controls regarding asbestos exposure was less in our findings than in theirs. There are several possible explanations for this. The first is that our study was based on a national sample of cases derived from all parts of a very large country, whereas theirs were from 2 specific industrial areas, one a ship-building city and the other from a hospital close to a large asbestos factory. A second

TABLE 7. Occupational Exposure to Certain Materials of Men without Definite or Probable Exposure to Asbestos

	Mesothelial tumors	Secondary lung cancer	Primary lung cancer
Cement only	10 (5)	9 (6)	8 (2)
Copper, nickel, rubber, or fiber glass	9* (2)	1 (1)	1 (0)
	19 (7)	10 (7)	9 (2)

* Including three men also exposed to cement.

The number of men possibly exposed to asbestos is given in parentheses.

Cigarettes per day

Nil
10 or less
11-29
30 or more

TOTAL*

* Excluding

possibility of cases which third point sure that t were collec of the gro was not so possible in Thompson not compa gation was selected fo on a popul.

It was n bestos in tl Several stu exposure i risk of this our serie c to match and geogr considered less, it was worth not asbestos w lung cance in the cau cancers w

The fin dicate tha

1. Elmes, O. L.: Dis asbestos. Br

2. Newho lioma of ple asbestos in 22:261-269

3. Selikoff

TABLE 8. Average Daily Number of Cigarettes Smoked

Cigarettes per day	Male				Mesothelial tumor with asbestos exposure	Female		
	Mesothelial tumors	Secondary lung cancer	Primary lung cancer			Mesothelial tumors	Secondary lung cancer	Primary lung cancer
Nil	21 22	19 20	4 4		2 10	29 53	36 71	21 47
10 or less	17 18	21 22	10 11		4 20	9 18	5 10	9 29
11-29	49 41	37 39	47 50		13 65	10 20	7 14	12 27
30 or more	19 20	18 19	34 36		1 5	2 4	3 6	5 7
TOTAL*	97 100	95 100	95 100		20 100	50 100	51 100	45 100

* Excluding 6 persons under 15 years of age and 6 men and 2 women whose smoking history was not recorded

possibility is that our series was diluted by cases which they would not have accepted. A third point is that we took great care to insure that the histories of case and control were collected and coded without knowledge of the group to which they belonged. This was not so stated in the Belfast survey and not possible in the study of Newhouse and Thompson. The findings of Selikoff et al.³ are not comparable with ours since their investigation was based on an occupational group selected for its exposure to asbestos and not on a population sample.

It was not our aim to assess the role of asbestos in the etiology of primary lung cancer. Several studies have shown that occupational exposure may be associated with increased risk of this tumor, especially in smokers.⁴ Since our series of primary lung cancer was selected to match the mesothelial cases in age, sex, and geographical distribution, it cannot be considered representative in itself. Nevertheless, it was probably sufficiently so for it to be worth noting that the history of exposure to asbestos was very similar to that for secondary lung cancer. Any contribution from asbestos in the causation of this series of primary lung cancers was, therefore, not apparent.

CONCLUSION

The findings from this national survey indicate that primary malignant mesothelial tu-

mors are rare in Canada, and it is somewhat doubtful whether there has been a true increase in incidence since 1959. An association between these tumors and definite or probable occupational exposure to asbestos was clearly demonstrated, but only a minority of male cases (20%) and one female case had any such exposure. No difference was found between cases and the 2 control groups regarding lesser degrees of occupational exposure or residence in asbestos-mining areas, but, in 10% of the mesothelial tumors compared with 5% of the secondary lung cancer group, there was at least a possible domestic contact with asbestos dust.

Probably our most noteworthy finding was that almost all the excess in occupational exposure was in the manufacture and industrial application of asbestos rather than in mining or milling. Taken in conjunction with the results of other epidemiologic studies, this suggests either that chrysotile is less associated with mesothelial tumors than other forms of asbestos or that something in addition to asbestos is necessary. If there is an additional factor, our findings imply that it is not cigarette smoking.

Nine men with mesothelial tumors had worked in occupations unrelated to asbestos but involving contact with copper, nickel, rubber, or fiber glass compared with one in each of the control groups. The interpretation of this finding is doubtful.

REFERENCES

1. Elmes, P. C., McCaughey, W. T. E., and Wade, O. L.: Diffuse mesothelioma of the pleura and asbestos. *Brit. Med. J.* 1:350-353, 1965.
2. Newhouse, M. L., and Thompson, H.: Mesothelioma of pleura and peritoneum following exposure to asbestos in the London area. *Brit. J. Industr. Med.* 22:261-269, 1965.
3. Selikoff, I. J., Churg, J., and Hammond, E. C.: Relation between exposure to asbestos and mesothelioma. *New Eng. J. Med.* 272:560-565, 1965.
4. Selikoff, I. J., Hammond, E. C., and Churg, J.: Asbestos exposure, smoking and neoplasia. *JAMA* 204:106-112, 1968.
5. Wagner, J. C., Sleggs, C. A., and Marchand, P.: Diffuse pleural mesothelioma and asbestos exposure in the north western Cape Province. *Brit. J. Industr. Med.* 17:260-271, 1960.

orted in close
t in the con-
ette smoking
t both males
primary lung
lerably more
or secondary
? groups had
of smoking
or probable
s showed no
s from cases

ctly compa-
Belfast, and
London, it
nce between
os exposure
eirs. There
is for this.
ed on a na-
m all parts
theirs were
one a ship-
a hospital
A second