

Facts About Asbestos-Cement Pipe

CertainTeed

PLAINTIFF'S EXHIBIT

CT-1022

Asbestos—an irreplaceable material.

Fact:

A/C pipe is one of more than 3,000 products containing asbestos that serve people beneficially worldwide. In many products, such as A/C pipe, brake linings, fire protection equipment, and textiles, asbestos is irreplaceable.

Fact:

Although asbestos represents only about 15% of the material used in A/C pipe, it contributes its tensile strength, almost five times greater than that of steel. It gives A/C pipe flexibility, excellent chemical resistance, and provides the taxpayer with a very cost-effective pipe material.

A/C pipe is used worldwide.

Fact:

A/C water pipe is recommended and installed in virtually every country in the world and in all 50 of the United States. Over 1,600,000 miles are in service worldwide, with 250,000 miles in the U.S. where over a quarter-billion dollars worth is installed annually. One-third of all U.S. water pipe in use today is A/C.

A/C pipe is safe to use.

Fact:

There is no evidence that anyone in the general public has ever contracted any disease from naturally-occurring asbestos found in most water systems.

Fact:

The release of asbestos fibers from A/C pipe is infinitesimal and is far less than the amount of fiber found in many natural-source drinking waters. This has been confirmed by government, industry, and scientific studies in the U.S. and abroad.

Fact:

Despite more than 40 years of rapidly increasing A/C pipe usage in the U.S., a 1977 American Cancer Society report: "Cancer Facts and Figures," states that "gastrointestinal cancer rates for all sites (where A/C pipe was being used) for the period 1952-54 and 1972-74 declined 33 percent."

A 1977 study in Connecticut said there was "no correlation between the incidence of GI cancer and the use of A/C water pipe."

Knoxville, Tenn., has been using A/C pipe since 1938. Nashville, Tenn., has no A/C pipe in its system. A study by HEW, NCI, and the National Institute of Health: "U.S. Cancer Mortality by County," showed that over a 20-year period, Nashville, with metallic pipe, had a 29 percent higher death rate from GI cancer than Knoxville, with A/C pipe.

Fact:

Asbestos fibers in A/C pipe are securely bound into the chemically stable cement and silica materials from which the pipe is formed.

The hazards of aggressive water

Fact:

Certain types of "aggressive" water will attack all water system components, even including corrosion-resistant A/C pipe, as well as metallic pipe and pipe installed in homes. Aggressive (very soft, highly acidic) water costs the U.S. water-works industry a billion dollars annually in replacement and repair. No piping is suitable for use in highly aggressive water areas. The answer to aggressive water is to neutralize it at the source, regardless of the pipe materials used.

A/C pipe is safe to install

Fact:


Manufacturers and the water utility industry have made mass distribution of detailed work practices to contractors and installers of A/C water pipe. These present proven methods of cutting, beveling and tapping pipe to insure a safe work environment.

Fact:

Complete data on A/C pipe can be obtained from:

Mr. Wally Werner
CertainTeed Corporation
Government Relations Office
1627 K St. NW.,
Washington, D.C. 20006
Telephone: (202) 833-3670.

CTD033123




A/C Pipe and Health

CTD033124


Contents




3.
Perceptions
of Risk




17.
Asbestos
in Water...
The Risks




5.
Asbestos
and A/C Pipe




21.
A Commitment
to Quality




7.
Asbestos
and Health



9.
A/C Pipe
and Field
Operations



11.
A/C Pipe and
Waterborne
Asbestos




CTD033126

Perceptions of Risk

The world has never been . . . and never will be free from risk. It is inescapable. It is life itself . . . and since the dawn of time, man has accepted risk as a normal part of his daily activity.

The perception of risk varies . . . from situation to situation . . . from individual to individual. What one person considers safe, another may perceive as downright foolish. This simply illustrates that the perception of risk is subjective, emotional and often has little basis in fact.

All risks do have one thing in common: they can be measured . . . objectively, scientifically. There is a name for this measurement . . . it is called *probability*. Just as the meteorologist predicts the probability of rain, so can odds be calculated for driving an automobile and having a severe accident or lying on the beach and getting sunburned.




SAFE VS. ZERO RISK
Something is judged "safe" if its probability of risk is acceptable. However, to

say something is "safe" does not necessarily mean it is "free from risk". A seemingly "safe" 6-minute canoe ride, for example, carries the same risk of death as a 500 mph commuter flight from Washington, D.C., to New York City . . . traveling the same distance by automobile, the risk increases five-fold.¹

Very few things to which we are exposed are "zero risks". But like the automobile, electricity and X-rays, there are non-harmful ways to use potentially harmful


things . . . it's simply a matter of probabilities.

CANCER AND CARCINOGENS Just as cavemen had their risks, modern man has his. Certainly none is as well-publicized, personal and fraught with anxiety, as the risk of cancer. While the fear of cancer can and does alter our perception of risk, it cannot . . . does not . . . alter the *probability* of risk.




In the simplest sense, cancer is an unrestrained growth of cells. It is not a single disease, but a

collection of many diseases caused by agents medical science calls *carcinogens*. Asbestos is a carcinogen . . . it is not alone. It shares this dubious distinction with a growing list of common and exotic substances. As a matter of fact, nearly 300 organic and inorganic chemicals, many of them known or potential carcinogens, have been identified in drinking water *alone*. This fact has prompted a much closer examination of water pipe materials in common use to determine if any are above suspicion, since all contain toxic materials.



DOSE RESPONSE AND THRESHOLD
The critical question with asbestos, as with any toxic material, is

"at what exposure level does it represent a hazard?" Medical research determines this



CTD033128

Asbestos and A/C Pipe

level by a concept known as *dose-response*. The basic principle is rather straightforward: for every given dose, there is a proportionate biological response. Let's add another wrinkle to this concept . . . and it is a common sense wrinkle . . . *there is a dose small enough which will result in no measurable response*. This dose level is called a *threshold*. Threshold effects are common. We have thresholds for hearing, pain, heat and martinis.

Everything known about the potential hazards of asbestos points to the fact that it observes the rules of dose-response. Therefore, a practical threshold or safe level of exposure to asbestos must exist. If it did not, it would be difficult to reconcile the prevalence of asbestos in the environment with the absence of related diseases in the general population.

The probability of finding asbestos in the environment is very high. Asbestos-bearing rock formations are found in over 50 countries throughout the world. Geologists tell us these formations were created 470 million years ago by the action of hot, subterranean water on rock containing iron, magnesium and silica. These waters dissolved the surrounding rock layer producing "veins" of liquid minerals. As the earth cooled, these minerals crystallized into the fibrous ore we know today as asbestos.²


More than 2,000 years ago, pharaohs and kings used this unique product of nature which the Greeks named *asbesta*, meaning "unquenchable". Today, you don't have to be royalty to use asbestos . . . practically everyone does. There are over 3,000 products which use asbestos, usually in combination with other materials such as cement.

Micro-Reinforcement of Cement						
	Strength	Flex. Modulus	Chem. Resist.	Impact Resist.	Comp. Resist.	Age Resist.
Asbestos	++	++	++	++	++	++
Steel	++	++	++	++	++	++
Carbon	++	++	++	++	++	++
Glass	++	++	++	++	++	++
Organic Fibers	++	++	++	++	++	++

UNIQUE PROPERTIES
As the world's only naturally occurring mineral fiber, asbestos is truly unique.

Its soft and silky fibers exhibit incredible tensile strength . . . 4 to 5 times greater than steel. Its flexibility, heat and chemical resistance are unmatched by any other natural or man-made fiber.³ Asbestos is not only unique, it is *irreplaceable* for the micro-reinforcement of high performance products like asbestos-cement (A/C) pipe.

FREE VS. BOUND ASBESTOS Asbestos-cement pipe contains a blend of asbestos fibers: *chrysotile* for reinforcement and




CTD033130

Asbestos and Health

crocidolite or blue asbestos also for reinforcement as well as for facilitation of the pipe manufacturing process.

The total asbestos portion of A/C pipe is less than 20 percent. *More importantly, the asbestos fibers are not free, but encapsulated by or "locked-in" the cement binder.* Picture, if you will, the use of steel reinforcing rods in concrete.



The distinction between "free" and "bound" asbestos is extremely important. Why?

- (1) It determines the probability of fiber release, which in turn, directly relates to the probability of risk.
- (2) It explains why asbestos-cement pipe is a non-harmful use of asbestos.


The ability of asbestos to separate into microscopically small fibers is the key to its biological availability, in other words, its ability to become airborne or water-borne and thereby inhaled or ingested. The biological effects of asbestos are believed to be a function of fiber shape and size, rather than chemical composition. Generally, the longer and thinner the fiber, the greater the potential health risk.

For regulatory purposes, an asbestos fiber has been defined as "particulate material whose length is at least three times longer than its width."⁴ Fiber dimensions are measured in micrometers . . . that's about 39 millionths of an inch or about one-seventieth the diameter of a human hair. Asbestos fibers longer than 5 micrometers are of prime interest from a health standpoint.

SUSCEPTIBILITY Like all biological effects, asbestos health effects vary significantly from person to person. Physical activity, inherited immunities, the pre-existence of other diseases, and the efficiency of the body in clearing particles, all affect individual susceptibility.

Cigarette smoking, for example, increases by over 90 times, the risk of lung cancer in workers exposed to asbestos. Non-smoking asbestos workers, on the other hand, show no more than the normally expected rate of lung cancer in the general population.^{5,6}

DOSE AND ROUTES OF ENTRY While the relationship between asbestos fiber exposure and "probability of risk" is not a simple one, we do know this: *the body's response to asbestos is dependent on dose and route of entry*, that is, whether it is inhaled or ingested. Inhalation is the more



CTD033132

A/C Pipe and Field Operations

common means by which asbestos enters the body. Although natural cleansing mechanisms catch and clear as much as 98 percent of inhaled dust, prolonged or heavy exposure to airborne asbestos can lead to *asbestosis*, a disease which impairs breathing and in the case of smokers, high risk of *lung cancer*. To a much lesser degree, asbestos may cause *mesothelioma*, a rare cancer of the chest or abdominal cavity. Scientists believe that asbestosis and lung cancer are associated with fibers longer than 5 micrometers and mesothelioma with fibers longer than 20 micrometers.⁷ To date, all identified occurrences of these diseases attributed to asbestos, have involved exposures in or related to occupational settings.



In contrast to inhalation, a clear "cause and effect" relationship between ingested asbestos and gas-

trointestinal or GI cancer has not been established. In certain heavily exposed occupational groups, a somewhat higher than normal rate of GI cancer has been reported.⁸ It might therefore be concluded that ingested asbestos may be related to gastrointestinal cancer. Yet . . . everything from mice to baboons has been watered, force-fed or dined at will on asbestos . . . as much as 6 percent of their total food intake.^{9,10,11,12,13,14,15} Results? No cancers.

These then are the risks for exposure to asbestos: "high" for heavy or prolonged inhalation exposure; solely "presumptive", with no direct evidence of risk for ingestion.


Recognizing the risks of airborne asbestos, regulatory agencies internationally have set safe levels of exposure based on the number of fibers longer than 5 micrometers. In the United States, an *occupational exposure standard* is used for workers in asbestos product manufacturing industries. A *construction industry standard*, yet to be established, will apply to workers who may handle asbestos-containing products such as A/C pipe.

OSHA Asbestos Exposure Standards		
	Eight Hour Time Weighted Average	Ceiling Concentration
Occupational		
Existing	2.0 fibers/cc	10 fibers/cc
Proposed	0.5 fibers/cc	5 fibers/cc
Construction		
Existing*	2.0 fibers/cc	10 fibers/cc
Proposed**	—	—
*Identical with existing Occupational Standard		
**Pending		
Source: Code of Federal Regulations, 29 Part 1910.1001		

FIELD OPERATIONS Let's compare the existing and proposed occupational standards to fiber levels measured when cutting or machining A/C pipe by recommended work practices.

Loading, unloading and stringing out A/C pipe are basically dust-free operations. Even in the most confined working space, dust levels are barely measurable. A survey conducted by the Port of Tyne Authority in England measured dust levels in the hold of a ship loading A/C pipe.¹⁶ The highest dust measurement was 0.01 fibers/cc . . . one-fiftieth of the proposed OSHA Occupational Standard.


The adaptability of A/C pipe and the availability of a complete line of short lengths and standard fittings greatly minimize the need for field operations.



CTD033134

A/C Pipe and Waterborne Asbestos

Even when cutting, machining or finishing are necessary, independent industrial hygiene studies¹⁷ have confirmed that when recommended work practices are followed, fiber levels are not only far below the present 2 fiber standard, but also meet the proposed ½ fiber standard (see Appendix A).




Abrasive Disc Cutting

This should come as no surprise. Remember . . . the asbestos fiber is locked-in the cement binder. The infrequency and short duration of field operations further reduce even these minimal exposures. Conclusion: when recommended work practices are followed, the risk of working with A/C pipe is virtually non-existent.

What about waterborne asbestos? Does potable water conveyed through A/C pipe pick up asbestos fiber and cause cancer?

These questions about the risks of ingested asbestos are being raised because of the recognized hazards of excessive or prolonged exposure to airborne asbestos. To answer them, let's see how asbestos can enter a potable water system . . . the facts will speak for themselves.





SOURCE WATERS "The general prevalence of asbestos in soil results in its presence in most waters of

lake, river and well origin, and in distribution systems whether fabricated of asbestos-cement or other materials."¹⁸ Natural background levels have been confirmed by nation-wide surveys conducted by the United States Environmental Protection Agency.¹⁹ Although EPA found sites with asbestos levels below present limits of detection, natural and treated waters in Montana, Oregon, Pennsylvania, California, Connecticut, Massachusetts and Georgia had levels in the range of millions of fibers per liter.

Similar studies in Canada concluded: ". . . there is asbestos in all surface waters of all habited areas of Ontario, and perhaps there always has been some asbestos there."²⁰

The presence of asbestos in ice core samples dating back hundreds of years²¹ confirms that natural weathering and erosion have in the past and will for generations to come, release small amounts of



CTD033136

asbestos to surface watersheds and underground aquifers around the world.


SOILS Another potential source of waterborne asbestos can be asbestos-bearing soil deposited in the pipe during installation or repair. The chances of this occurring are the same whether the pipe is ductile iron, plastic or asbestos-cement. A thorough flushing of the distribution system is a simple enough precaution to eliminate any long term effects of soil contamination.

TAPPING A/C PIPE To provide house connections, water distribution mains are tapped approximately once every 140 feet. With A/C pipe, tapping can lead to a temporary increase in the asbestos fiber level of the water. *The industry recommends when tapping A/C pipe while under pressure, that equipment with a positive purge be used.* This ejects 99 percent of the asbestos-cement chips and downstream flushing takes care of the rest. Another alternative is the use of heavy-walled tapped couplings supplied directly from the factory.

AGGRESSIVE WATERS Now the critical question . . . *does A/C pipe contribute asbestos to drinking water conveyed through it?* A no-nonsense answer to this question is a simple "yes" . . . and "no." *It depends on the water . . . not the pipe. As long as A/C pipe's cement binder remains intact, the release, if any, of the locked-in fiber is infinitesimal.*

A/C pipe is remarkably corrosion resistant . . . much more so than metallic piping products. There are, however, certain types of corrosive or *aggressive* waters that will attack even corrosion resistant asbestos-cement. The most common of these are low

hardness or *soft* water and low pH or *acidic* water.




Imagine aggressive waters as being *hungry* waters . . . not only hungry for calcium and magnesium in

cement, but hungry for iron, lead, copper and cadmium in metallic pipe, hungry for organic residuals in plastic pipe, polyaromatic hydrocarbons in coal tar linings, hungry for almost all materials in waterworks use today.

Aggressive waters take a staggering toll from the U.S. waterworks industry every year . . . an estimated \$400 million²² in replacement costs and perhaps as much as \$600 million in protective devices and measures. But the heaviest toll of all may be its toll in human lives. In 1974, a "consistent and highly significant relationship" was confirmed between aggressive water and cardiovascular diseases such as heart attacks, still the number one killer in the United States. The study showed that the cities with the softest water had twice the death rate from coronary disease as cities where the water was hardest.²³ Economics alone argue strongly for national treatment standards for aggressive water . . . the loss of life punctuates the urgency.

AGGRESSIVE WATER GUIDELINES

Though the A/C pipe industry is not the first to spotlight the need for stabilization, it has been a leader in taking positive steps to include aggressive water guidelines in its national product specifications. Any potential loss of business the industry



CTD033138

could suffer as a result of these guidelines, is more than offset by the knowledge that where A/C pipe is used, both the utility and consumer are assured safe, trouble-free service. Based on a half-century of industry experience, *these guidelines do not recommend the use of A/C pipe with highly aggressive waters.*

Aggressive Water Guidelines		
Type of Water	Aggressive Index	A/C Pipe Recommended
Non-Aggressive	>12.0	Types I & II
Moderately Aggressive	10.0-11.9	Type II
Highly Aggressive	<10.0	

* The serviceability of pipe for such applications should be established by the purchaser in conjunction with the manufacturer.

Source: American Water Works Association Standards C-400-75 C-402-75
American Society for Testing & Materials Standard C 500-76

ASBESTOS IN A/C PIPE SYSTEMS
Measurements taken in the field confirm EPA laboratory results that for "moderately" and "non-aggressive" waters, A/C pipe does not contribute significant or in many cases, even measurable amounts of asbestos to drinking water conveyed through it.

Northeast Illinois . . . an analysis of 15 public water systems containing A/C pipe of various ages, lengths, and diameters. *Conclusion: No significant release of asbestos from A/C pipe was observed.*²⁴


California . . . an analysis of water at the end of long pipe runs in a distribution system with over 1,000 miles of A/C pipe. *Conclusion: All tests for asbestos negative.*²⁵

New Mexico . . . an analysis of asbestos fiber concentrations in water systems containing 548 miles of A/C pipe. *Conclusion: "No evidence of asbestos pick up from A/C pipe."*²⁶

Texas, Ohio and Connecticut . . . EPA monitorings of A/C pipe systems. *Conclusion: Fiber levels either "not statistically significant" or "below detectable limits."*



Common sense alone tells us that "real world" risk must relate to "real world" exposure. For that reason, these results are important . . . they measure actual exposures to water-borne asbestos in A/C pipe systems. *Clearly, the amount of asbestos, if any, contributed to drinking water supplied through A/C pipe, is infinitesimal . . . certainly far less than is found in many natural source waters.*




CTD033140

Asbestos in Water... The Risks

What about water supply systems which have measurable quantities of asbestos fiber . . . what's the risk of contracting cancer?

This time there is a straight answer to the questions . . . there is no evidence that anyone in the general public has ever contracted any disease from naturally occurring asbestos or the minute amounts of fiber released from A/C pipe. This is not a judgment of safety . . . it's a statement of fact.

A possible explanation may be the extremely small size of waterborne asbestos. You may recall that asbestos fibers less than 5 micrometers in length are not generally associated with adverse health effects. Commenting on asbestos in natural water supplies, EPA noted ". . . few samples show any evidence of fibers greater than 5 micrometers long and when these are present, they represent only a small percentage of the total fiber count."²⁷



**INGESTED
ASBESTOS
AND CANCER**
Whether it's
the fiber size
or the sub-
threshold
"dose", one

fact is irrefutable: if there is a "cause and effect" relationship between GI cancers, ingested asbestos or the use of A/C pipe, it does not show in state, national or world cancer statistics.

Let's start with the most publicized case, Duluth, Minnesota . . . millions of asbestos-like fibers in its drinking water, and Houston, Texas, with none. A court-ordered comparison of tissues from de-

ceased residents from both cities failed to disclose any asbestos fibers of waterborne origin.²⁸

Duluth again . . . this time a comparison with Minneapolis-St. Paul, which has no known asbestos in its water. No statistical difference in the incidence of gastrointestinal cancer.²⁹

Switzerland . . . the highest per capita use of A/C pipe in Western Europe. From 1951, more than 20 years after A/C pipe was introduced, to 1973, the death rate from GI cancer dropped a significant 29 percent.³⁰


The United States . . . despite over 40 years of rapidly increasing A/C pipe usage, gastrointestinal cancer rates for all sites for the period 1952-54 to 1972-74, declined 33 percent.³¹

Tennessee . . . Knoxville, A/C pipe in use since 1938 and Nashville, no A/C pipe. A twenty-year comparison shows that Nashville with metallic pipe, had a 29 percent higher death rate from GI cancer than Knoxville with A/C pipe.³²

Connecticut . . . Preliminary results from an Environmental Protection Agency study show no correlation between the incidence of GI cancer and the use of A/C water pipe.³³


This inability to relate a "real world" dose with a "real world" response again says in a common sense way that like the amount of asbestos contributed to drinking water by A/C pipe, the cancer risk from ingesting that asbestos must be infinitely small. An exact number? Well, we did say that all risks could be measured.

MEASURING THE RISKS Let's imagine a hypothetical human feeding experiment



CTD033142

based on magnified doses used in actual animal feeding studies which have failed to establish a cancer risk from ingested as-



bestos. Based on a diet containing 0.27 percent asbestos, this amounts to about 5 pounds of as-

bestos . . . roughly the amount of butter you eat in a year. Using the *highest* fiber levels found in drinking water, you would have to drink 3.8 million quarts of water per day to consume 5 pounds of asbestos in a year . . . or over 50,000 quarts per day for a lifetime.

This is, of course, preposterous. But it does quantify and place in perspective the probability of risk from a "real world" dose of ingested asbestos from A/C pipe . . . a dose which is not uncommonly a billion times less than asbestos levels proven to be non-carcinogenic in animal feeding studies.

One in a billion! That figure probably doesn't register with most people. In terms of quality, it's one bad apple in 2,000,000 barrels. In terms of volume, it's a very, very dry martini . . . 1 drop of vermouth in 15,750 gallons of gin.³⁴ In terms of risk . . . it's a thousand times safer than smoking 1½ cigarettes or drinking a half-bottle of your favorite wine.³⁵

NO RISK TO PUBLIC In spite of perceptions and publicity to the contrary, these incredibly low probabilities of risk tell us that A/C pipe is not . . . and never will be a risk to the public . . . a judgment that has been confirmed again and again by some

of the most respected health authorities in the world.


International Agency for Research on Cancer. The question: "Is there evidence of an increased risk of cancer resulting from asbestos fibers present in water, beverages, food or in the fluids used for the administration of drugs?" The answer: "Such evidence as there is does not indicate any risk."³⁶

Dr. Irving J. Selikoff, Director of the Environmental Sciences Laboratory, Mount Sinai School of Medicine. There is "no evidence at all at this time that the presumably very low level exposure by ingestion which might occur by intake of water supplied through asbestos-cement pipe has resulted in a recognized health hazard."³⁷

National Cancer Institute. "NCI scientists . . . have found no unusual cancer mortality patterns so far among persons residing where drinking water is contaminated by asbestos . . ."³⁸

Select Study Committee of the American Water Works Association Research Foundation . . . "No firm evidence shows that the proper use of asbestos-cement pipe poses a hazard to health by reason of ingestion of asbestos fibers. Calculations comparing the probable ingestion exposure in occupational groups to that likely to occur as a result of ingestion of potable water from asbestos-cement pipe systems suggests that the probability of risk to health from the use of such systems is small—approaching zero."³⁹

False accusations and "guilt by association" have created a perception of risk about A/C pipe. Scientific measurements and common sense show this distorted perception has no basis in fact.



CTD033144

A Commitment to Quality

Designed for value . . . engineered for performance . . . A/C pipe is the optimum balance between high performance, long life and cost effectiveness. Quality control second to none, assures high pressure reliability with sensible factors of safety . . . consistently, inexpensively . . . *and that's a fact!*

The A/C pipe industry has a commitment . . . a commitment to quality water. This commitment encompasses much more than the simple assurance of water quality at the consumer's tap. Regardless of how bountiful or clean water reserves might be . . . whimsical as nature can be . . . and short-sighted as man has been, quality water demands conservation in use, diligence in treatment, and economical reliability in distribution. Providing distribution pipe of integrity at a cost the user can and will support is the goal of the A/C pipe industry.

Appendices

Appendix A Ceiling Concentrations of Airborne Asbestos Fiber from A/C Pipe Field Operations

Field Operation	Type of Pipe	Operation Time (Min.)	Exposure (Fibers/cc)	
			Operator	Helper
Recommended Work Practices				
Manual Lathe	Pressure	12-13	0.51	0.22
Manual Lathe	Sewer	4-5	0.15	0.13
Power Lathe	Pressure	5-6	0.33	0.56
Power Lathe	Sewer	6-8	0.08	0.09
Power Hole Cutting	Pressure	1.5-2	0.44	0.23
Power Hole Cutting	Sewer	1	1.65	0.38
Cutting w/Chisel & Rasp	Pressure	10-11	1.99	0.90
Cutting w/Chisel & Rasp	Sewer	4-6	0.30	0.25
Chain-type Cutter	Pressure	2-3	0.85	0.87
Chain-type Cutter	Sewer	2-3	<0.06	<0.06
Pilot ^a -type Gutter	Sewer	2-3	0.09	—
Non-Recommended Work Practices				
Abrasive Disc Cutting*	Pressure	<1	20.30	59.70
Abrasive Disc Cutting*	Sewer	<1	35.50	64.00

*Workers should be provided with respiratory protection approved by the United States Department of the Interior, Bureau of Mines, Schedule 21C or Department of Health, Education and Welfare, National Institute of Occupational Safety and Health, under the provisions of the Code of Federal Regulations, 30, Part 11

Note: OSHA Standard for Occupational Exposure to Asbestos Dust (Ceiling Concentration) Ten (10) fibers per cubic centimeter of air

Appendix B
U.S. Trends in Age-Adjusted Cancer Death Rates
1952-54 to 1972-74 (per 100,000 population)

Source: "1977 Facts and Figures", American Cancer Society.

Site	Sex	1952-54	1972-74	Change (%)
Colon & Rectum	Male	19.3	18.9	- 2
	Female	18.4	14.9	- 19
Stomach	Male	16.4	7.4	- 54
	Female	8.6	3.6	- 58
Change:				- 33%

Appendix C
Nashville-Knoxville Cancer Mortality Rates
(per 100,000 population)-1950 to 1969

Source: National Cancer Institute Statistics.

Site	Sex*	Davidson County (Nashville)	Loudon County (Knoxville)
Stomach	White Males	9.8	7.8
	White Females	4.4	4.3
Large Intestine	White Males	14.1	7.5
	White Females	13.6	13.3
Rectum	White Males	4.7	3.4
	White Females	3.5	2.6
Totals		50.1	38.9

Difference 28.7%

*Complete statistics not available for non-white males and females. (Higher in Nashville)

CTD033147

Bibliography

- ¹Pochin, E. E., "The Acceptance of Risk," *British Medical Bulletin*, Volume 31, No. 3, September, 1975.
- ²Smith, M. F., "The Origin of Asbestos," *Asbestos Magazine*, July, 1940.
- ³Erik, H., "The Micro-Reinforcement of Cement," *A/C Underground*, Issue 04, 1976.
- ⁴United States Code of Federal Regulations, Chapter XVII, Part 1910.1001.
- ⁵Selikoff, I. J., Hammond, E. C. and Churg, J., "Asbestos Exposure, Smoking and Neoplasia," *Journal of the American Medical Association*, April 8, 1968.
- ⁶Kannerstein, M. and Churg, J., "Pathology of Carcinoma of the Lung Associated with Asbestos Exposure," *Cancer*, American Cancer Society, July, 1972.
- ⁷Smither, W. J., "Asbestosis—Some Comments on the Present Position of Medical Opinion," *International Conference of Asbestos Associations and Asbestos Information Bodies*, Paris, France, November, 1974.
- ⁸Selikoff, I. J., Hammond, E. C., and Churg, J., "Asbestos Exposure and Neoplasia," *Journal of the American Medical Association*, 188:22.
- ⁹Westlake, G. E., Spjut, H. J. and Smith, M. N., "Penetration of Colonic Mucosa by Asbestos Particles: An Electron Microscope Study in Rats Fed Asbestos Dust," *Laboratory Investigation*, 14:2029, 1965.
- ¹⁰Bonser, G. M. and Clayson, D. B., "Feeding of Blue Asbestos to Rats," *45th Annual Reports of the British Empire Cancer Campaign*, 1967.
- ¹¹Smith, W. E. L., Miller, L., Elasser, R. E. and Hubert, D. D., "Tests for Carcinogenicity of Asbestos," *Annals of the New York Academy of Science*, 132:456, 1965.
- ¹²Smith, W. E., "Talc and Nitrites in Relation to Gastric Cancer," *American Industrial Hygiene Association Journal*, 34:227, 1973.
- ¹³Webster, I., "The Ingestion of Asbestos Fibers," *Environmental Health Perspectives*, 9:199, 1974.
- ¹⁴Gross, P., Harley, R. A., Swinburne, L. M., Davis, J. M. G. and Greene, W. B., "Ingested Mineral Fibers," *Archives of Environmental Health*, 29:341, 1974.
- ¹⁵Bolton, R. E. and Davis, J. M. G., "The Short Term Effects of Chronic Asbestos Ingestion in Rats," *Annals of Occupational Hygiene*, 19:121, 1976.
- ¹⁶"News from TAC," reporting on a survey by the Occupational Hygiene Unit of the North of England Industrial Health Service, TAC Construction Materials Ltd., February, 1977.
- ¹⁷Equitable Environmental Health, Inc., Berkley, Ca., Contract P76S-036 for the A/C Pipe Producers Association.
- ¹⁸Kuschner, M., Lee, R., Robeck, G. G., Rossum, J. R., Schneiderman, M. A. and Taylor, E. W., "A Study of the Problem of Asbestos in Water," *American Water Works Association Journal*, 66:513.
- ¹⁹U.S. Environmental Protection Agency, "The Impact of Point and Non-Point Sources on Levels of Waterborne Asbestos," EPA Report No. EPA-560/6-76-018, October, 1976.
- ²⁰Kay, G. H., "Ontario Intensifies Search for Asbestos in Drinking Water," *Water and Pollution Control*, September, 1973.
- ²¹Johns-Manville Corporation, Personal Communications.
- ²²Hudson, H. E., and Gilreas, F. W., "Health and Economic Aspects of Water Hardness and Corrosiveness," *Journal of the American Water Works Association*, April, 1976.
- ²³Schroeder, H. A., and Kraemer, L. A., "Cardiovascular Mortality, Municipal Water and Corrosion," *Archives of Environmental Health*, 28:303, 1974.
- ²⁴Hallenbeck, W. H., Chen, E. H., Hesse, C. H., Kusum, P. M. and Wolff, A. H., "Is Chrysotile Asbestos Released from Asbestos-Cement Pipe Into Drinking Water?" (In preparation of submission to *Journal of American Water Works Association*).
- ²⁵Garretson, F., "Asbestos Absent From Water," *Oakland Tribune*, September 18, 1974.
- ²⁶Oliver, T. and Murr, L. E., "An Electron Microscope Study of Asbestiform Fiber Concentrations in Municipal and Private Water Supplies in the Rio Grande Valley of New Mexico: The Role of Asbestos-Cement Pipe," (In preparation of submission to *Journal of American Water Works Association*).
- ²⁷Op. Cit. (Environmental Protection Agency).
- ²⁸United States Court of Appeals for the Eighth Circuit, No. 73-1239 "Reserve Mining Co. vs. Ruckelshaus," filed March 14, 1975.
- ²⁹Levy, B. S., Sigurdson, E., Mandel, J., Laudon, E. and Pearson, J., "Investigation of Possible Effects of Asbestos in City Water: Surveillance of Gastrointestinal Cancer in Duluth, Minnesota," *American Journal of Epidemiology*, Volume 103, No. 4, April, 1976.
- ³⁰(a) Epidemiological and Vital Statistics Report, Volume 18, No. 6, World Health Organization.
(b) World Health Statistics Annual, 1973-1976, Volume 1, World Health Organization.
- ³¹American Cancer Society, 1977 *Cancer Facts and Figures*, 1976.
- ³²U. S. Department of Health, Education and Welfare, National Institutes of Health, National Cancer Institute, *U. S. Cancer Mortality by County—1950-1969*.
- ³³Craun, G. F., Millette, J. R., Woodhull, R. S. and Laiuppa, R., "Exposure to Asbestos Fibers in Water Distribution Systems," *97th Conference of American Water Works Association*, Anaheim, Ca., May, 1977.
- ³⁴Crummett, W. B., "One Part Per Trillion Is a Very Finely Split Hair," *Chemecology*, Manufacturing Chemists Association, December, 1976.
- ³⁵Op. Cit. (Pochin).
- ³⁶"The Biological Effects of Asbestos," Report of the Advisory Committee on Asbestos Cancers to the Director of the International Agency for Research on Cancer, *British Journal of Industrial Medicine*, 30:180, 1973.
- ³⁷Selikoff, I. J., Private Communications.
- ³⁸U.S. Department of Health, Education and Welfare, National Institutes of Health, National Cancer Institute, "Fact Sheet: Atlas of Cancer Mortality for U.S. Counties: 1950-1969," 1976.
- ³⁹Op Cit. (Kuschner et al).

A/C Pipe Producers Association

REGULAR MEMBERS

Asbestos de Mexico, S.A.
Tlalnepantla, Mexico

Atlas Asbestos Company
Montreal, Quebec, Canada

Canadian Johns-Manville Co., Ltd.
Mississauga, Ontario, Canada

Cement Asbestos Products Company
Birmingham, Alabama

Certain-teed Corporation
Valley Forge, Pennsylvania

Johns-Manville Sales Corporation
Denver, Colorado

Mexalit del Norte, S.A.
Mexico City, Mexico

ASSOCIATE MEMBERS

Amiantus Dienst Export
Hamburg, Germany

Asbestos Monterrey, S.A.
Mexico City, Mexico

Association de l'Industrie l'Asbeste-Ciment
Nyon, Switzerland

Eternit, B.V.
Amsterdam, Holland

Eventube
Neuilly-sur-Seine, France

James Hardie & Company Pty., Ltd.
Sydney, New South Wales, Australia

Hellenit, S.A.
Athens, Greece

Hyderabad Asbestos Cement Products Ltd.
Hyderabad, India

Paraisten Kalkki Oy
Pargas, Finland

Sociedad Industrial Pizarreno, S.A.
Santiago, Chile

TAC Construction Materials, Ltd.
Manchester, United Kingdom

The A/C Pipe Producers Association functions as a non-profit, non-partisan world-wide trade association on behalf of the A/C Pipe industry. Inquiries for information on specific products, their individual attributes and differences, should be directed to member companies.

The Statements contained in this document are those of the Association of Asbestos-Cement Pipe Producers and are not intended as a warranty.



Printed in U.S.A.

CTD033149



A/C Pipe Producers Association

1600 Wilson Boulevard, Suite 1308, Arlington, Virginia 22209

CTD033150