




Industrial Hygiene Digest

INDUSTRIAL HEALTH NEWS

LITERATURE ABSTRACTS



MEDICAL

ENGINEERING

CHEMICAL

TOXICOLOGICAL

LEGAL . . . decisions and trends

JULY, 1956
(Vol. 20, No. 7)

INDUSTRIAL HYGIENE FOUNDATION

MELLON INSTITUTE

4400 FIFTH AVENUE •• PITTSBURGH 13, PA.

PLAINTIFF'S
EXHIBIT

WV-00466

PLAINTIFF'S
EXHIBIT

1339.03



FOUNDATION FACTS

Foundation Facts is a monthly news-letter issued to industrial concerns holding membership in Industrial Hygiene Foundation. The Foundation is a nonprofit association of industries for the advancement of healthful working conditions.

Volume 18

July, 1956

No. 7

WELCOME TO MEMBERSHIP

Wald Industries, Inc. is the latest company to affiliate with Industrial Hygiene Foundation. We are pleased to add them to the membership roster.

INTERNATIONAL INTEREST IN FOUNDATION ACTIVITIES

The work of Industrial Hygiene Foundation is of interest not only to industries within the United States but to scientists and industrial health specialists in other countries as well. The Foundation is pleased to receive such visitors and to explain the American system and techniques for maintaining health in industry. Among the distinguished visitors recently was Dr. W. E. George, Medical Officer of the Joint Civil Control New South Wales. A visitor from Birmingham, England was Dr. J. G. Lawson, Medical Officer of Joseph Lucas, Ltd. Dr. Mario Battigelli, a surgeon from the Istituto Del Lavoro in Milan, Italy, visited the Foundation and plans to devote further time to studying I-H-F activities. Dr. R. S. Koo of the College of Medicine of Seoul National University in Korea also paid a visit to the Foundation.

SCHRENK APPOINTED TO POLLUTION ABATEMENT COMMITTEE

Dr. H. H. Schrenk, Research Director of Industrial Hygiene Foundation, has been appointed to the Pennsylvania State Chamber of Commerce Committee on Pollution for the 1956-57 period.

SCIENTIFIC EXHIBITS FOR THE 21st ANNUAL MEETING OF I-H-F

Each year at the annual meeting of the Foundation a limited number of exhibits are displayed illustrating the accomplishments of industrial hygiene. Companies desiring to describe through their exhibits the maintenance programs or some special industrial health activities are invited to reserve space for the November 14-15, 1956 meeting at the Madison Hotel. Please advise the Foundation as soon as possible.

PRODUCTS OF INDUSTRIAL HYGIENE FOUNDATION

A chart prepared by the Foundation, showing the functions of a biological laboratory as a trademark in industrial hygiene appears on the following page. It forms the basis for a paper, entitled "The Importance of Biological Research in Medicine" which was published in the June 18, 1956 issue of the Journal of the American Medical Association. The chart was prepared to show concisely and in a graphic manner the various functions of a biological laboratory. Reprints of the paper will be available at a later date.

It is evident from this chart that Industrial Hygiene Foundation's toxicology laboratory can begin its contribution even during the research development of new materials in helping to eliminate those substances which are obviously too toxic for the intended purpose. The extent of the biologic work can expand as the development phase of the material expands, and thus adequate biologic data parallels the development and commercial production and utilization of a new product.

evidence of previous infection. Coal-mine dust alone produced no fibrosis. The combination of bacilli and dust produced tuberculous cavities involving large areas of lung tissue, and killed the guinea pigs. -- Am. Rev. Tuberc.

- 910 Experimental Infective Tuberculosis. 4. Massive Pulmonary Fibrosis Produced by Coal-Mine Dust and Isoniazid-Resistant Tubercle Bacilli of Low Virulence. S. H. Zaidi, C. V. Harrison, E. J. King, and D. A. Mitchison. *Brit. J. Exptl. Pathol.* 36, 553-559 (Dec. 1955).

The effects of intratracheal injection of coal-mine dust, a catalase-negative isoniazid-resistant strain of tubercle bacilli of low virulence, and a combination of dust and bacilli were studied in the lungs of guinea pigs. The bacilli alone produced lesions which tended to disappear after 90 days, and the only evidence of tuberculosis at 300 days was a few calcified nodules in some of the lungs. While coal-mine dust itself produced no pulmonary fibrosis, the combined action of the dust and bacilli produced massive pulmonary fibrosis. In its early stages, the fibrosis was composed mainly of thick reticulin fibers and collagen, but later more collagen was formed. Serial cultures of the lungs injected with coal dust and bacilli yielded growth of tubercle bacilli up to, but not beyond, 120 days. The largest number of organisms was found at 30 days. In animals infected with bacilli alone, no growth could be obtained from lung cultures other than those from guinea pigs killed one day after injection. -- Am. Rev. Tuberc.

- 911 Costal Pleura and Pleural Adhesion Silicotic Pigmentation. D. Schwenkenbecher. *Tuberkulosearzt* 9, 592-595 (Oct. 1955). German.

A pneumothorax for cavernous tuberculous disease was induced in a patient who had worked as a bricklayer for fourteen years. During a pneumonolysis, a silicotic deposit was observed in the pleural adhesions as well as in the costal pleura. The suspicion of silicotuberculosis was verified by biopsy following eventual lobectomy. Silicotic deposits in the costal pleura are rare. They are usually localized in the neighborhood of the pleuro-pulmonary adhesions and are believed to come from the lymphatic system. In the present case the chest roentgenogram gave no indication of silicotuberculosis. -- Am. Rev. Tuberc.

- 912 Roentgenographic Complications of Pulmonary Asbestosis. G. Jacob and H. Bohlig. *Fortschr. Gebiete Rontgenstrahlen* 83, 515-525 (Oct. 1955). German.

Over a period of approximately twenty years, 343 cases of asbestosis, 130 in women and 213 in men, were observed in Dresden, a center for industries using asbestos. The most common complications were pleural thickening, compensatory emphysema, and nonspecific peribronchial, peribronchiectatic, and bronchopneumonic inflammations. Pleural changes were found in 49 per cent of the cases, 5 per cent of which had extensive, irregularly shaped calcifications. Active pulmonary tuberculosis was present in 4.4 per cent of the cases; bronchogenic cancer was not more frequent than in the general population but, when present, was found relatively often in the lower lobe. -- Am. Rev. Tuberc.

PHYSICAL ASPECTS OF THE ENVIRONMENT

- 913 Industrial Noise and How to Deal with It. H. A. Carter. *Arch. Phys. Med.* 37, 152-158 (Mar. 1956).

This paper deals briefly with noise sources, dangerous noise levels, noise measurement, effects on hearing, protection, noise reduction, and medico-legal aspects.