



INTERNAL CORRESPONDENCE



UNION CARBIDE CORPORATION
CHEMICALS AND PLASTICS

P. O. BOX 3370, BROWNSVILLE, TEXAS 78520

To (Name) A. L. Barefoot
Company
Location Brownsville Plant

Date June 25, 1982

Originating Dept.

Answering letter date

Subject Contractor Information
Safety and Health

Copy to J. L. Wyatt (w/o att)

Dear Ashton:

Attached is the information you requested relating to contract workers and required medical surveillance.

List of items included:

- ✓ 1. Summary of medical testing requirements.
- ✓ 2. Excerpts from OSHA on medical requirements on Asbestos, Lead and a summary by a medical testing company that will bring a unit into the plant and test the workers.
3. ~~The Corporation Occupational Health Contractor policy.~~
4. ~~The contractor guideline found in the Safety and Health Information Manual, CMD.~~

Call if there are any questions.

KGT:mfw
Attachment

K. Gran Townsend

HAZARDS	REQUIREMENTS						
	WORK EXPOSURE HISTORY AND MONITORING	PRE- EMPLOYMENT MEDICAL EXAM	PERSONAL HISTORY	BLOOD PRESSURE	LUNG FUNCTION	X-RAY	HERMATOLOGY
ASBESTOS	YES	YES	YES	YES	YES	YES	--
INORGANIC LEAD	YES	YES	YES	YES	--	--	YES
NOISE	YES	YES	YES	--	--	--	--
(1) RESPIRATOR USERS	YES	YES	YES	YES	YES	YES	--
SILICA	YES						
(2) AIR CONTAMINANTS	YES						
HOT ENVIRONMENTS	YES						
FORK LIFT DRIVERS							
WELDING FUMES	YES						

(1) Includes fire brigade members.

(2) These should include the major products and raw materials as a minimum (Industrial Hygienist to determine)

HAZARDS	REQUIREMENTS					
	URINALYSIS	BLOOD/ URINE TRACE METALS	AUDIOGRAM	ANNUAL MEDICAL EXAM	RECORD KEEPING	TRAINING PROGRAM
ASBESTOS	--	--	--	YES	YES	YES
INORGANIC LEAD	YES	YES	--	YES	YES	YES
NOISE	--	--	YES	YES	YES	YES
(1) RESPIRATOR USERS	--	--	--	YES	YES	YES
SILICA					YES	YES
(2) AIR CONTAMINANTS					YES	YES
HOT ENVIRONMENTS					YES	YES
FORK LIFT DRIVERS					YES	YES
WELDING FUMES					YES	YES

MEDICAL TESTING
REQUIREMENTS

HAZARDS

HAZARDS	MEDICAL TESTING REQUIREMENTS						
	WORK EXPOSURE HISTORY AND MONITORING	PRE- EMPLOYMENT MEDICAL EXAM	PERSONAL HISTORY	BLOOD PRESSURE	LUNG FUNCTION	X-RAY	HEMATOLOGY
ASBESTOS	YES	YES	YES	YES	YES	YES	--
INORGANIC LEAD	YES	YES	YES	YES	--	--	YES
NOISE	YES	YES	YES	--	--	--	--
(1) RESPIRATOR USERS	YES	YES	YES	YES	YES	YES	--
SILICA	YES						
(2) AIR CONTAMINANTS	YES						
HOT ENVIRONMENTS	YES						
FORK LIFT DRIVERS							
WELDING FUMES	YES						

(1) Includes fire brigade members.

(2) These should include the major products and raw materials as a minimum (Industrial Hygienist to determine)

MEDICAL TESTING
REQUIREMENTS

HAZARDS

	URINALYSIS	BLOOD/ URINE TRACE METALS	AUDIOGRAM	ANNUAL MEDICAL EXAM	RECORD KEEPING	TRAINING PROGRAM
ASBESTOS	--	--	--	YES	YES	YES
INORGANIC LEAD	YES	YES	--	YES	YES	YES
NOISE	--	--	YES	YES	YES	YES
(1) RESPIRATOR USERS	--	--	--	YES	YES	YES
SILICA					YES	YES
(2) AIR CONTAMINANTS					YES	YES
HOT ENVIRONMENTS					YES	YES
FORK LIFT DRIVERS					YES	YES
WELDING FUMES					YES	YES

Legend

Notation

Breathing Asbestos Dust. 14 point Gothic.
May Be Hazardous To
Your Health.

Spacing between lines shall be at least equal to the height of the upper of any two lines.

(2) **Caution labels**—(i) **Labeling.** Caution labels shall be affixed to all raw materials, mixtures, scrap, waste, debris, and other products containing asbestos fibers, or to their containers, except that no label is required where asbestos fibers have been modified by a bonding agent, coating, binder, or other material so that during any reasonably foreseeable use, handling, storage, disposal, processing, or transportation, no airborne concentrations of asbestos fibers in excess of the exposure limits prescribed in paragraph (b) of this section will be released.

(ii) **Label specifications.** The caution labels required by subdivision (i) of this subparagraph shall be printed in letters of sufficient size and contrast as to be readily visible and legible. The label shall state:

CAUTION

Contains Asbestos Fibers

Avoid Creating Dust

Breathing Asbestos Dust May Cause
Serious Bodily Harm

(h) **Housekeeping**—(1) **Cleaning.** All external surfaces in any place of employment shall be maintained free of accumulations of asbestos fibers if, with their dispersion, there would be an excessive concentration.

(2) **Waste disposal.** Asbestos waste, scrap, debris, bags, containers, equipment, and asbestos-contaminated clothing, consigned for disposal, which may produce in any reasonably foreseeable use, handling, storage, processing, disposal, or transportation airborne concentrations of asbestos fibers in excess of the exposure limits prescribed in paragraph (b) of this section shall be collected and disposed of in sealed impermeable bags, or other closed, impermeable containers.

(i) **Recordkeeping**—(1) **Exposure records.** Every employer shall maintain records of any personal or environmental monitoring required by this section. Records shall be maintained for a period of at least 20 years and shall be made available upon request to the Assistant Secretary of Labor for Occupational Safety and Health, the Director of the National Institute for Occupational Safety and Health, and to authorized representatives of either.

[1910.1001(i)(1) amended at 41 FR 11505, March 19, 1976]

(2) **Access.** Employee exposure records required by this paragraph shall be provided upon request to employees, designated representatives, and the Assistant Secretary in accordance with 29 CFR 1910.20 (a)-(e) and (g)-(i).

[Section 1910.1001(i)(2) amended at 45 FR 35212, May 23, 1980, effective August 21, 1980]

(3) **Employee notification.** Any employee found to have been exposed at any time to airborne concentrations of asbestos fibers in excess of the limits prescribed in paragraph (b) of this section shall be notified in writing of the expo-

sure as soon as practicable but not later than 5 days of the finding. The employee shall also be timely notified of the corrective action being taken.

(j) **Medical examinations**—(1) **General.** The employer shall provide or make available at his cost, medical examinations relative to exposure to asbestos required by this paragraph.

(2) **Preplacement.** The employer shall provide or make available to each of his employees, within 30 calendar days following his first employment in an occupation exposed to airborne concentrations of asbestos fibers, a comprehensive medical examination, which shall include, as a minimum, a chest roentgenogram (posterior-anterior 14 x 17 inches), a history to elicit symptomatology of respiratory disease, and pulmonary function tests to include forced vital capacity (FVC) and forced expiratory volume at 1 second (FEV_{1.0}).

(3) **Annual examinations.** On or before January 31, 1973, and at least annually thereafter, every employer shall provide, or make available, comprehensive medical examinations to each of his employees engaged in occupations exposed to airborne concentrations of asbestos fibers. Such annual examination shall include, as a minimum, a chest roentgenogram (posterior-anterior 14 x 17 inches), a history to elicit symptomatology of respiratory disease, and pulmonary function tests to include forced vital capacity (FVC) and forced expiratory volume at 1 second (FEV_{1.0}).

(4) **Termination of employment.** The employer shall provide, or make available, within 30 calendar days before or after the termination of employment of any employee engaged in an occupation exposed to airborne concentrations of asbestos fibers, a comprehensive medical examination which shall include, as a minimum, a chest roentgenogram (posterior-anterior 14 x 17 inches), a history to elicit symptomatology of respiratory disease, and pulmonary function tests to include forced vital capacity (FVC) and forced expiratory volume at 1 second (FEV_{1.0}).

(5) **Recent examinations.** No medical examination is required of any employee, if adequate records show that the employee has been examined in accordance with this paragraph within the past 1-year period.

(6) **Medical records**—(i) **Maintenance.** Employers of employees examined pursuant to this paragraph shall cause to be maintained complete and accurate records of all such medical examinations. Records shall be retained by employers for at least 20 years.

(ii) **Access.** Records of the medical examinations required by this paragraph shall be provided upon request to employees, designated representatives, and the Assistant Secretary in accordance with 29 CFR 1910.20(a)-(e) and (g)-(i). These records shall also be provided upon the request to the Director of NIOSH. Any physician who conducts a medical examination required by this paragraph shall furnish to the employer of the examined employee all the information specifically required by this paragraph, and any other

medical information related to occupational exposure to asbestos fibers.

[Section 1910.1001(j)(6)(ii) amended at 45 FR 35212, May 23, 1980, effective August 21, 1980; corrected by 45 FR 54333, August 15, 1980]

§ 1910.1002 Coal tar pitch volatiles; interpretation of term.

As used in Sec. 1910.1000 (Table Z-1), coal tar pitch volatiles include the fused polycyclic hydrocarbons which volatilize from the distillation residues of coal, petroleum, wood, and other organic matter.

§ 1910.1003 4-Nitrobiphenyl.

(a) **Scope and application.** (1) This section applies to any area in which 4-Nitrobiphenyl, Chemical Abstracts Service Registry Number 92933 is manufactured, processed, repackaged, released, handled, or stored, but shall not apply to transshipment in sealed containers, except for the labeling requirements under paragraphs (c) (2), (3), and (4) of this section.

(2) This section shall not apply to solid or liquid mixtures containing less than 0.1 percent by weight or volume of 4-Nitrobiphenyl.

(b) **Definitions.** For the purposes of this section: (1) "Absolute filter" is one capable of retaining 99.97 percent of a mono disperse aerosol of 0.3 μ m particles.

(2) "Authorized employee" means an employee whose duties require him to be in the regulated area and who has been specifically assigned by the employer.

(3) "Clean change room" means a room where employees put on clean clothing and/or protective equipment in an environment free of 4-Nitrobiphenyl. The clean change room shall be contiguous to and have an entry from a shower room, when the shower room facilities are otherwise required in this section.

(4) "Closed system" means an operation involving 4-Nitrobiphenyl where containment prevents the release of 4-Nitrobiphenyl into regulated areas, non-regulated areas, or the external environment.

(5) "Decontamination" means the inactivation of 4-Nitrobiphenyl or its safe disposal.

(6) "Director" means the Director, National Institute for Occupational Safety and Health, or any person directed by him or the Secretary of Health, Education, and Welfare to act for the Director.

(7) "Disposal" means the safe removal of 4-Nitrobiphenyl from the work environment.

(8) "Emergency" means an unforeseen circumstance or set of circumstances resulting in the release of 4-Nitrobiphenyl which may result in exposure to or contact with 4-Nitrobiphenyl.

(9) "External environment" means any environment external to regulated and nonregulated areas.

(10) "Isolated system" means a fully enclosed structure other than the vessel or containment of 4-Nitrobiphenyl.

LEAD

IT WOULD BE CATERED TO BE
OF THE LEAD PAINT - KGT
REFERENCE FILE

(2) *Cleaning floors.* (i) Floors and other surfaces where lead accumulates may not be cleaned by the use of compressed air.

(ii) Shoveling, dry or wet sweeping and brushing may be used only where vacuuming or other equally effective methods have been tried and found not to be effective.

(3) *Vacuuming.* Where vacuuming methods are selected, the vacuums shall be used and emptied in a manner which minimizes the reentry of lead into the workplace.

[Section 1910.1025(h)(2)(ii) corrected at 44 FR 5446, January 26, 1979, effective February 24, 1979]

[Section 1910.1025(h) effective date was March 31, 1979, to extent that employers must acquire vacuums to comply, according to action by U.S. Court of Appeals for District of Columbia Circuit. (44 FR 14554, March 13, 1979)]

(i) *Hygiene facilities and practices.*

(1) The employer shall assure that in areas where employees are exposed to lead above the PEL, without regard to the use of respirators, food or beverage is not present or consumed, tobacco products are not present or used, and cosmetics are not applied, except in change rooms, lunchrooms, and showers required under paragraphs (i)(2)-(i)(4) of this section.

(2) *Change rooms.* (i) The employer shall provide clean change rooms for employees who work in areas where their airborne exposure to lead is above the PEL, without regard to the use of respirators.

(ii) The employer shall assure that change rooms are equipped with separate storage facilities for protective work clothing and equipment and for street clothes which prevent cross-contamination.

(3) *Showers.* (i) The employer shall assure that employees who work in areas where their airborne exposure to lead is above the PEL, without regard to the use of respirators, shower at the end of the work shift.

(ii) The employer shall provide shower facilities in accordance with §1910.141(d)(3) of this Part.

(iii) The employer shall assure that employees who are required to shower pursuant to paragraph (i)(3)(i) do not leave the workplace wearing any clothing or equipment worn during the work shift.

(4) *Lunchrooms.* (i) The employer shall provide lunchroom facilities for employees who work in areas where their airborne exposure to lead is above the PEL, without regard to the use of respirators.

(ii) The employer shall assure that lunchroom facilities have a tempera-

ture controlled, positive pressure, filtered air supply, and are readily accessible to employees.

(iii) The employer shall assure that employees who work in areas where their airborne exposure to lead is above the PEL without regard to the use of a respirator wash their hands and face prior to eating, drinking, smoking or applying cosmetics.

(iv) The employer shall assure that employees do not enter lunchroom facilities with protective work clothing or equipment unless surface lead dust has been removed by vacuuming, downdraft booth, or other cleaning method.

(5) *Lavatories.* The employer shall provide an adequate number of lavatory facilities which comply with §1910.141(d)(1) and (2) of this Part.

[Sections 1910.1025(i)(1), (i)(2), (i)(3)(i) and (ii), and (i)(4)(i) and (iii) amended, and (i)(6) deleted, at 44 FR 5446, January 26, 1979, effective February 24, 1979]

(j) *Medical surveillance.* (1) *General.*

(i) The employer shall institute a medical surveillance program for all employees who are or may be exposed above the action level for more than 30 days per year.

(ii) The employer shall assure that all medical examinations and procedures are performed by or under the supervision of a licensed physician.

(iii) The employer shall provide the required medical surveillance including multiple physician review under paragraph (j)(3)(iii) without cost to employees and at a reasonable time and place.

(2) *Biological monitoring.* —(i) *Blood lead and ZPP level sampling and analysis.* The employer shall make available biological monitoring in the form of blood sampling and analysis for lead and zinc protoporphyrin levels to each employee covered under paragraph (j)(1)(i) of this section on the following schedule:

(A) At least every 6 months to each employee covered under paragraph (j)(1)(i) of this section;

(B) At least every two months for each employee whose last blood sampling and analysis indicated a blood lead level at or above 40 µg/100 g of whole blood. This frequency shall continue until two consecutive blood samples and analyses indicate a blood lead level below 40 µg/100 g of whole blood, and

(C) At least monthly during the removal period of each employee re-

moved from exposure to lead due to an elevated blood lead level.

(ii) *Follow-up blood sampling tests.* Whenever the results of a blood lead level test indicate that an employee's blood lead level exceeds the numerical criterion for medical removal under paragraph (k)(1)(i), the employer shall provide a second (follow-up) blood sampling test within two weeks after the employer receives the results of the first blood sampling test.

(iii) *Accuracy of blood lead level sampling and analysis.* Blood lead level sampling and analysis provided pursuant to this section shall have an accuracy (to a confidence level of 95 percent) within plus or minus 15 percent or 6 µg/100ml, whichever is greater, and shall be conducted by a laboratory licensed by the Center for Disease Control, United States Department of Health, Education and Welfare (CDC) or which has received a satisfactory grade in blood lead proficiency testing from CDC in the prior twelve months.

(iv) *Employee notification.* Within five working days after the receipt of biological monitoring results, the employer shall notify in writing each employee whose blood lead level exceeds 40 µg/100 g: (A) of that employee's blood lead level and (B) that the standard requires temporary medical removal with Medical Removal Protection benefits when an employee's blood lead level exceeds the numerical criterion for medical removal under paragraph (k)(1)(i) of this section.

(3) *Medical examinations and consultations.* —(i) *Frequency.* The employer shall make available medical examinations and consultations to each employee covered under paragraph (j)(1)(i) of this section on the following schedule:

(A) At least annually for each employee for whom a blood sampling test conducted at any time during the preceding 12 months indicated a blood lead level at or above 40 µg/100 g;

(B) Prior to assignment for each employee being assigned for the first time to an area in which airborne concentrations of lead are at or above the action level; -

(C) As soon as possible, upon notification by an employee either that the employee has developed signs or symptoms commonly associated with lead intoxication, that the employee desires medical advice concerning the effects of current or past exposure to lead on the employee's ability to procreate a healthy child, or that the employee has demonstrated difficulty in

breathing during a respirator fitting test or during use; and

(D) As medically appropriate for each employee either removed from exposure to lead due to a risk of sustaining material impairment to health, or otherwise limited pursuant to a final medical determination.

(ii) *Content.* Medical examinations made available pursuant to paragraph (j)(3)(i)(A)-(B) of this section shall include the following elements:

(A) A detailed work history and a medical history, with particular attention to past lead exposure (occupational and non-occupational), personal habits (smoking, hygiene), and past gastrointestinal, hematologic, renal, cardiovascular, reproductive and neurological problems;

(B) A thorough physical examination, with particular attention to teeth, gums, hematologic, gastrointestinal, renal, cardiovascular, and neurological systems. Pulmonary status should be evaluated if respiratory protection will be used;

(C) A blood pressure measurement;

(D) A blood sample and analysis which determines:

(1) Blood lead level;

(2) Hemoglobin and hematocrit determinations, red cell indices, and examination of peripheral smear morphology;

(3) Zinc protoporphyrin;

(4) Blood urea nitrogen; and,

(5) Serum creatinine;

(E) A routine urinalysis with microscopic examination; and

(F) Any laboratory or other test which the examining physician deems necessary by sound medical practice.

The content of medical examinations made available pursuant to paragraph (j)(3)(i)(C)-(D) of this section shall be determined by an examining physician and, if requested by an employee, shall include pregnancy testing or laboratory evaluation of male fertility.

(iii) *Multiple physician review mechanism.* (A) If the employer selects the initial physician who conducts any medical examination or consultation provided to an employee under this section, the employee may designate a second physician:

(1) To review any findings, determinations or recommendations of the initial physician; and

(2) To conduct such examinations, consultations, and laboratory tests as the second physician deems necessary to facilitate this review.

(B) The employer shall promptly notify an employee of the right to seek a second medical opinion after each occasion that an initial physician conducts a medical examination or consultation pursuant to this section. The employer may condition its participation in, and payment for, the multiple physician review mechanism upon the employee doing the following within fifteen (15) days after receipt of the foregoing notification, or receipt of the initial physician's written opinion, whichever is later:

(1) The employee informing the employer that he or she intends to seek a second medical opinion; and

(2) The employee initiating steps to make an appointment with a second physician.

(C) If the findings, determinations or recommendations of the second physician differ from those of the initial physician, then the employer and the employee shall assure that efforts are made for the two physicians to resolve any disagreement.

(D) If the two physicians have been unable to quickly resolve their disagreement, then the employer and the employee through their respective physicians shall designate a third physician:

(1) To review any findings, determinations or recommendations of the prior physicians; and

(2) To conduct such examinations, consultations, laboratory tests and discussions with the prior physicians as the third physician deems necessary to resolve the disagreement of the prior physicians.

(E) The employer shall act consistently with the findings, determinations and recommendations of the third physician, unless the employer and the employee reach an agreement which is otherwise consistent with the recommendations of at least one of the three physicians.

(iv) *Information provided to examining and consulting physicians.* (A) The employer shall provide an initial physician conducting a medical examination or consultation under this section with the following information:

(1) A copy of this regulation for lead including all Appendices;

(2) A description of the affected employee's duties as they relate to the employee's exposure;

(3) The employee's exposure level or anticipated exposure level to lead and

to any other toxic substance (if applicable);

(4) A description of any personal protective equipment used or to be used;

(5) Prior blood lead determinations; and

(6) All prior written medical opinions concerning the employee in the employer's possession or control.

(B) The employer shall provide the foregoing information to a second or third physician conducting a medical examination or consultation under this section upon request either by the second or third physician, or by the employee.

(v) *Written medical opinions.* (A) The employer shall obtain and furnish the employee with a copy of a written medical opinion from each examining or consulting physician which contains the following information:

(1) The physician's opinion as to whether the employee has any detected medical condition which would place the employee at increased risk of material impairment of the employee's health from exposure to lead;

(2) Any recommended special protective measures to be provided to the employee, or limitations to be placed upon the employee's exposure to lead;

(3) Any recommended limitation upon the employee's use of respirators, including a determination of whether the employee can wear a powered air purifying respirator if a physician determines that the employee cannot wear a negative pressure respirator; and

(4) The results of the blood lead determinations.

(B) The employer shall instruct each examining and consulting physician to:

(1) Not reveal either in the written opinion, or in any other means of communication with the employer, findings, including laboratory results, or diagnoses unrelated to an employee's occupational exposure to lead; and

(2) Advise the employee of any medical condition, occupational or nonoccupational, which dictates further medical examination or treatment.

(vi) *Alternate Physician Determination Mechanisms.* The employer and an employee or authorized employee representative may agree upon the use of any expeditious alternate physician determination mechanism in lieu of the multiple physician review mechanism provided by this paragraph so long as the alternate mechanism otherwise satisfies the requirements contained in this paragraph.

Appendix C to Section 1910.1025— Medical Surveillance Guidelines

Introduction

The primary purpose of the Occupational Safety and Health Act of 1970 is to assure, so far as possible, safe and healthful working conditions for every working man and woman. The occupational health standard for inorganic lead was promulgated to protect workers exposed to inorganic lead including metallic lead, all inorganic lead compounds and organic lead soaps.

Under this final standard in effect as of March 1, 1979, occupational exposure to inorganic lead* is to be limited to 50 $\mu\text{g}/\text{m}^3$ (micrograms per cubic meter) based on an 8 hour time-weighted average (TWA). This level of exposure eventually must be achieved through a combination of engineering, work practice and other administrative controls. Periods of time ranging from 1 to 10 years are provided for different industries to implement these controls. The schedule which is based on individual industry considerations is given in Table 1. Until these controls are in place, respirators must be used to meet the 50 $\mu\text{g}/\text{m}^3$ exposure limit.

The standard also provides for a program of biological monitoring and medical surveillance for all employees exposed to levels of inorganic lead above the action level of 30 $\mu\text{g}/\text{m}^3$ (TWA) for more than 30 days per year.

The purpose of this document is to outline the medical surveillance provisions of the standard for inorganic lead, and to provide further information to the physician regarding the examination and evaluation of workers exposed to inorganic lead.

[Section 1910.1025, Appendix C footnote added at 44 FR 68827, November 30, 1979]

*The term inorganic lead used throughout the medical surveillance appendices is meant to be synonymous with the definition of lead set forth in the standard.

Section 1 provides a detailed description of the monitoring procedure including the required frequency of blood testing for exposed workers, provisions for medical removal protection (MRP), the recommended right of the employee to a second medical opinion, and notification and recordkeeping requirements of the employer. A discussion of the requirements for respirator use and respirator monitoring and OSHA's position on prophylactic chelation therapy are also included in this section.

Section 2 discusses the toxic effects and clinical manifestations of lead poisoning and effects of lead intoxication on enzymatic pathways in heme synthesis. The adverse effects on both male and female reproductive capacity and on the fetus are also discussed.

Section 3 outlines the recommended medical evaluation of the worker exposed to inorganic lead including details of the medical history, physical examination, and recommended laboratory tests, which are based on the toxic effects of lead as discussed in Section 2.

Section 4 provides detailed information concerning the laboratory tests available for the monitoring of exposed workers. Included also is a discussion of the relative value of each test and the limitations and precautions which are necessary in the interpretation of the laboratory results.

[Section 1910.1025, Appendix C corrected at 44 FR 68827, November 30, 1979]

I. Medical surveillance and monitoring requirements for workers exposed to inorganic lead

Under the occupational health standard for inorganic lead, a program of biological monitoring and medical surveillance is to be made available to all employees exposed to lead above the

action level of 30 $\mu\text{g}/\text{m}^3$ TWA for more than 30 days each year. This program consists of periodic blood sampling and medical evaluation to be performed on schedule which is defined by previous laboratory results, worker complaints or concerns, and the clinical assessment of the examining physician.

Under this program, the blood lead level of all employees who are exposed to lead above the action level of 30 $\mu\text{g}/\text{m}^3$ is to be determined at least every six months. The frequency is increased to every two months for employees whose last blood lead level was between 40 $\mu\text{g}/100$ g whole blood and the level requiring employee medical removal to be discussed below. For employees who are removed from exposure to lead due to an elevated blood lead, a new blood lead level must be measured monthly. A zinc protoporphyrin (ZPP) measurement is strongly recommended on each occasion that a blood lead level measurement is made. ZPP monitoring is one of several requirements of the standard that has been temporarily stayed by pending litigation over the lead standard. Unless otherwise noted in this document, all provisions of the standard pertaining to medical surveillance and medical removal protection are currently in effect.

An annual medical examination and consultation performed under the guidelines discussed in Section 3 is to be made available to each employee for whom a blood test conducted at any time during the preceding 12 months indicated a blood lead level at or above 40 $\mu\text{g}/100$ g. Also, an examination is to be given to all employees prior to their assignment to an area in which airborne lead concentrations reach or exceed the action level. In addition, a medical examination must be provided as soon as possible after notification by an employee that the employee has developed signs or symptoms commonly associated with lead intoxication, that the employee desires medical advice regarding lead exposure and the ability to procreate a healthy child, or that the employee has demonstrated difficulty in breathing during a respirator fitting test or during respirator use. An examination is also to be made available to each employee removed from exposure to lead due to a risk of sustaining material impairment to health, or otherwise

Table 1

Permissible airborne lead levels by industry ($\mu\text{g}/\text{m}^3$)*	Effective date					
	Mar. 1, 1979	Mar. 1, 1980	Mar. 1, 1981	Mar. 1, 1982	Mar. 1, 1984	Mar. 1, 1989 (final)
1. Primary lead production.....	200	200	200	100	100	50
2. Secondary lead production.....	200	200	200	100	50	50
3. Lead-acid battery manufacturing.....	200	200	100	100	50	50
4. Nonferrous foundries.....	200	100	100	100	50	50
5. Lead pigment manufacturing.....	200	200	200	100	50	50
6. All other industries.....	200	50	50	50	50	50

*Airborne levels to be achieved without reliance on respirator protection through a combination of engineering, work practice and other administrative controls. While these controls are being implemented respirators must be used to meet the 50 $\mu\text{g}/\text{m}^3$ exposure limit.

Table 2

	Effective date				
	Mar. 1, 1979	Mar. 1, 1980	Mar. 1, 1981	Mar. 1, 1982	Mar. 1, 1983 (final)
A. Blood lead level requiring employee medical removal. (Level must be confirmed with second follow-up blood lead level within two weeks of first report.)	$\geq 80 \mu\text{g}/100 \text{ g}$	$\geq 70 \mu\text{g}/100 \text{ g}$	$\geq 60 \mu\text{g}/100 \text{ g}$	$\geq 60 \mu\text{g}/100 \text{ g}$	$\geq 60 \mu\text{g}/100 \text{ g}$ or average of last three blood samples or all blood samples over previous 6 months (whichever is over a longer time period) is $50 \mu\text{g}/100 \text{ g}$ or greater unless last blood sample is $40 \mu\text{g}/100 \text{ g}$ or less.
B. Frequency which employee exposed to action level of lead ($30 \mu\text{g}/\text{m}^3$ TWA) must have blood lead level checked (ZPP is also strongly recommended in each occasion that a blood lead is obtained):					
1. Last blood lead level less than $40 \mu\text{g}/100 \text{ g}$	Every 6 months.	Every 6 months.	Every 6 months.	Every 6 months.	Every 6 months.
2. Last blood lead level between $40 \mu\text{g}/100 \text{ g}$ and level requiring medical removal (see A above).	Every 2 months.	Every 2 months.	Every 2 months.	Every 2 months.	Every 2 months.
3. Employees removed from exposure to lead because of an elevated blood lead level.	Every 1 month.	Every 1 month.	Every 1 month.	Every 1 month.	Every 1 month.
C. Permissible airborne exposure limit for workers removed from work due to an elevated blood lead level (without regard to respirator protection).	$<100 \mu\text{g}/\text{m}^3$ 8 hr TWA.	$<50 \mu\text{g}/\text{m}^3$ 8 hr TWA.	$<30 \mu\text{g}/\text{m}^3$ 8 hr TWA.	$<30 \mu\text{g}/\text{m}^3$ 8 hr TWA.	$<30 \mu\text{g}/\text{m}^3$ 8 hr TWA.
D. Blood lead level confirmed with a second blood analysis, at which employee may return to work. Permissible exposure without regard to respirator protection is listed by industry in Table 1.	$\leq 60 \mu\text{g}/100 \text{ g}$	$\leq 50 \mu\text{g}/100 \text{ g}$	$\leq 40 \mu\text{g}/100 \text{ g}$	$\leq 40 \mu\text{g}/100 \text{ g}$	$\leq 40 \mu\text{g}/100 \text{ g}$

NOTE.—When medical opinion indicates that an employee is at risk of material impairment from exposure to lead, the physician can remove an employee from exposures exceeding the action level (or less) or recommend special protective measures as deemed appropriate and necessary. Medical monitoring during the medical removal period can be more stringent than noted in the table above if the physician so specifies. Return to work or removal of limitations and special protections is permitted when the physician indicates that the worker is no longer at risk of material impairment.

[Section 1910.1025, Appendix C, Table 2 corrected at 44 FR 68827, November 30, 1979]

limited or specially protected pursuant to medical recommendations.

Results of biological monitoring or the recommendations of an examining physician may necessitate removal of an employee from further lead exposure pursuant to the standard's medical removal protection (MRP) program. The object of the MRP program is to provide temporary medical removal to workers either with substantially elevated blood lead levels or otherwise at risk of sustaining material health impairment from continued substantial exposure to lead. The following guidelines which are summarized in Table 2 were created under the standard for the temporary removal of an exposed employee and his or her subsequent return to work in an exposure area.

Under the standard's ultimate worker removal criteria, a worker is to be removed from any work having any eight hour TWA exposure to lead of $30 \mu\text{g}/\text{m}^3$ or more whenever either of the following circumstances apply: (1) a blood lead level of $60 \mu\text{g}/100 \text{ g}$ or greater is obtained and confirmed by a second follow-up blood lead level performed within two weeks after the employer receives the results of the first blood sampling test, or (2) the average of the previous three blood lead determinations or the average of all blood lead determinations conducted during the previous six months,

whichever encompasses the longest time period, equals or exceeds $50 \mu\text{g}/100 \text{ g}$, unless the last blood sample indicates a blood lead level at or below $40 \mu\text{g}/100 \text{ g}$ in which case the employee need not be removed. Medical removal is to continue until two consecutive blood lead levels are $40 \mu\text{g}/100 \text{ g}$ or less.

During the first two years that the ultimate removal criteria are being phased in, the return criteria have been set to assure that a worker's blood lead level has substantially declined during the period of removal. From March 1, 1979 to March 1, 1980, the blood lead level requiring employee medical removal is $80 \mu\text{g}/100 \text{ g}$. Workers found to have a confirmed blood lead at this level or greater need only be removed from work having a daily 8 hour TWA exposure to lead at or above $100 \mu\text{g}/\text{m}^3$. Workers so removed are to be returned to work when their blood lead levels are at or below $60 \mu\text{g}/100 \text{ g}$ of whole blood. From March 1, 1980 to March 1, 1981, the blood lead level requiring medical removal is $70 \mu\text{g}/100 \text{ g}$. During this period workers need only be removed from jobs having a daily 8 hour TWA exposure to lead at or above $50 \mu\text{g}/\text{m}^3$ and are to be returned to work when a level of $50 \mu\text{g}/100 \text{ g}$ is achieved. Beginning March 1, 1981, return depends on a worker's blood lead level declining to $40 \mu\text{g}/100 \text{ g}$ of whole blood.

As part of the standard, the employer is required to notify in writing each

employee whose blood lead level exceeds $40 \mu\text{g}/100 \text{ g}$. In addition each such employee is to be informed that the standard requires medical removal with MRP benefits, discussed below, when an employee's blood lead level exceeds the above defined limits.

In addition to the above blood lead level criteria, temporary worker removal may also take place as a result of medical determinations and recommendations. Written medical opinions must be prepared after each examination pursuant to the standard. If the examining physician includes a medical finding, determination or opinion that the employee has a medical condition which places the employee at increased risk of material health impairment from exposure to lead, then the employee must be removed from exposure to lead at or above the action level. Alternatively, if the examining physician recommends special protective measures for an employee (e.g., use of a powered air purifying respirator) or recommends limitations on an employee's exposure to lead, then the employer must implement these recommendations. Recommendations may be more stringent than the specific provisions of the standard. The examining physician, therefore, is given broad flexibility to tailor special protective procedures to the needs of individual employees. This flexibility extends to the evaluation and

management of pregnant workers and male and female workers who are planning to raise children. Based on the history, physical examination, and laboratory studies, the physician might recommend special protective measures or medical removal for an employee who is pregnant or who is planning to conceive a child when, in the physician's judgment, continued exposure to lead at the current job would pose a significant risk. The return of the employee to his or her former job status, or the removal of special protections or limitations, depends upon the examining physician determining that the employee is no longer at increased risk of material impairment or that special measures are no longer needed.

During the period of any form of special protection or removal, the employer must maintain the worker's earnings, seniority, and other employment rights and benefits (as though the worker had not been removed) for a period of up to 18 months. This economic protection will maximize meaningful worker participation in the medical surveillance program, and is appropriate as part of the employer's overall obligation to provide a safe and healthful workplace. The provisions of MRP benefits during the employee's removal period may, however, be conditioned upon participation in medical surveillance.

On rare occasions, an employee's blood lead level may not acceptably decline within 18 months of removal. This situation will arise only in unusual circumstances, thus the standard relies on an individual medical examination to determine how to protect such an employee. This medical determination is to be based on both laboratory values, including lead levels, zinc protoporphyrin levels, blood counts, and other tests felt to be warranted, as well as the physician's judgment that any symptoms or findings on physical examination are a result of lead toxicity. The medical determination may be that the employee is incapable of ever safely returning to his or her former job status. The medical determination may provide additional removal time past 18 months for some employees or specify special protective measures to be implemented.

The lead standard provides for a multiple physician review in cases where the employee wishes a second

opinion concerning potential lead poisoning or toxicity. If an employee wishes a second opinion, he or she can make an appointment with a physician of his or her choice. This second physician will review the findings, recommendations or determinations of the first physician and conduct any examinations, consultations or tests deemed necessary in an attempt to make a final medical determination. If the first and second physicians do not agree in their assessment they must try to resolve their differences. If they cannot reach an agreement then they must designate a third physician to resolve the dispute. This multiple physician review mechanism has been temporarily stayed during the pending litigation, but OSHA recommends that it be used if disputes arise over medical determinations.

The employer must provide examining and consulting physicians with the following specific information: a copy of the lead regulations and all appendices, a description of the employee's duties as related to exposure, the exposure level to lead and any other toxic substances (if applicable), a description of personal protective equipment used, blood lead levels, and all prior written medical opinions regarding the employee in the employer's possession or control. The employer must also obtain from the physician and provide the employee with a written medical opinion containing blood lead levels, the physician's opinion as to whether the employee is at risk of material impairment to health, any recommended protective measures for the employee if further exposure is permitted, as well as any recommended limitations upon an employee's use of respirators.

Employers must instruct each physician not to reveal to the employer in writing or in any other way his or her findings, laboratory results, or diagnoses which are felt to be unrelated to occupational lead exposure. They must also instruct each physician to advise the employee of any occupationally or non-occupationally related medical condition requiring further treatment or evaluation.

The standard provides for the use of respirators where engineering and other primary controls have not been fully implemented. However, the use of respirator protection shall not be used in lieu of temporary medical removal due to elevated blood lead levels or findings

that an employee is at risk of material health impairment. This is based on the numerous inadequacies of respirators including skin rash where the facepiece makes contact with the skin, unacceptable stress to breathing in some workers with underlying cardiopulmonary impairment, difficulty in providing adequate fit, the tendency for respirators to create additional hazards by interfering with vision, hearing, and mobility, and the difficulties of assuring the maximum effectiveness of a complicated work practice program involving respirators. Respirators do, however, serve a useful function where engineering and work practice controls are inadequate by providing supplementary, interim, or short-term protection, provided they are properly selected for the environment in which the employee will be working, properly fitted to the employee, maintained and cleaned periodically, and worn by the employee when required.

In its final standard on occupational exposure to inorganic lead, OSHA has prohibited prophylactic chelation. Diagnostic and therapeutic chelation are permitted only under the supervision of a licensed physician with appropriate medical monitoring in an acceptable clinical setting. The decision to initiate chelation therapy must be made on an individual basis and take into account the severity of symptoms felt to be a result of lead toxicity along with blood lead levels, ZPP levels, and other laboratory tests as appropriate. EDTA and penicillamine which are the primary chelating agents used in the therapy of occupational lead poisoning have significant potential side effects and their use must be justified on the basis of expected benefits to the worker. Unless frank and severe symptoms are present, therapeutic chelation is not recommended given the opportunity to remove a worker from exposure and allow the body to naturally excrete accumulated lead. As a diagnostic aid, the chelation mobilization test using CA-EDTA has limited applicability. According to some investigators, the test can differentiate between lead-induced and other nephropathies. The test may also provide an estimation of the mobile fraction of the total body lead burden.

Employers are required to assure that accurate records are maintained on exposure monitoring, medical

surveillance, and medical removal for each employee. Exposure monitoring and medical surveillance records must be kept for 40 years or the duration of employment plus 20 years, whichever is longer, while medical removal records must be maintained for the duration of employment. All records required under the standard must be made available upon request to the Assistant Secretary of Labor for Occupational Safety and Health and the Director of the National Institute for Occupational Safety and Health. Employers must also make environmental and biological monitoring and medical removal records available to affected employees and to former employees or their authorized employee representatives. Employees or their specifically designated representatives have access to their entire medical surveillance records.

In addition, the standard requires that the employer inform all workers exposed to lead at or above the action level of the provisions of the standard and all its appendices, the purpose and description of medical surveillance and provisions for medical removal protection if temporary removal is required. An understanding of the potential health effects of lead exposure by all exposed employees along with full understanding of their rights under the lead standard is essential for an effective monitoring program.

[Section 1910.1025, Appendix C(I) corrected at 44 FR 68827, November 30, 1979]

II. Adverse health effects of inorganic lead

Although the toxicity of lead has been known for 2,000 years, the knowledge of the complex relationship between lead exposure and human response is still being refined. Significant research into the toxic properties of lead continues throughout the world, and it should be anticipated that our understanding of thresholds of effects and margins of safety will be improved in future years. The provisions of the lead standard are founded on two prime medical judgments: first, the prevention of adverse health effects from exposure to lead throughout a working lifetime requires that worker blood lead levels be maintained at or below 40 $\mu\text{g}/100\text{ g}$ and second, the blood lead levels of workers, male or female, who intend to parent in the near future should be maintained below 30 $\mu\text{g}/100\text{ g}$ to minimize adverse reproductive health

effects to the parents and developing fetus. The adverse effects of lead on reproduction are being actively researched and OSHA encourages the physician to remain abreast of recent developments in the area to best advise pregnant workers or workers planning to conceive children.

The spectrum of health effects caused by lead exposure can be subdivided into five developmental stages: normal, physiological changes of uncertain significance, pathophysiological changes, overt symptoms (morbidity), and mortality. Within this process there are no sharp distinctions, but rather a continuum of effects. Boundaries between categories overlap due to the wide variation of individual responses and exposures in the working population. OSHA's development of the lead standard focused on pathophysiological changes as well as later stages of disease.

1. Heme Synthesis Inhibition. The earliest demonstrated effect of lead involves its ability to inhibit at least two enzymes of the heme synthesis pathway at very low blood levels. Inhibition of delta aminolevulinic acid dehydrase (ALA-D) which catalyzes the conversion of delta-aminolevulinic acid (ALA) to protoporphyrin is observed at a blood lead level below 20 $\mu\text{g}/100\text{ g}$ whole blood. At a blood lead level of 40 $\mu\text{g}/100\text{ g}$, more than 20% of the population would have 70% inhibition of ALA-D. There is an exponential increase in ALA excretion at blood lead levels greater than 40 $\mu\text{g}/100\text{ g}$.

Another enzyme, ferrochelatase, is also inhibited at low blood lead levels. Inhibition of ferrochelatase leads to increased free erythrocyte protoporphyrin (FEP) in the blood which can then bind to zinc to yield zinc protoporphyrin. At a blood lead level of 50 $\mu\text{g}/100\text{ g}$ or greater, nearly 100% of the population will have an increase in FEP. There is also an exponential relationship between blood lead levels greater than 40 $\mu\text{g}/100\text{ g}$ and the associated ZPP level, which has led to the development of the ZPP screening test for lead exposure.

While the significance of these effects is subject to debate, it is OSHA's position that these enzyme disturbances are early stages of a disease process which may eventually result in the clinical symptoms of lead poisoning. Whether or not the effects do progress

to the later stages of clinical disease, disruption of these enzyme processes over a working lifetime is considered to be a material impairment of health.

One of the eventual results of lead-induced inhibition of enzymes in the heme synthesis pathway is anemia which can be asymptomatic if mild but associated with a wide array of symptoms including dizziness, fatigue, and tachycardia when more severe. Studies have indicated that lead levels as low as 50 $\mu\text{g}/100\text{ g}$ can be associated with a definite decreased hemoglobin, although most cases of lead-induced anemia, as well as shortened red-cell survival times, occur at lead levels exceeding 80 $\mu\text{g}/100\text{ g}$. Inhibited hemoglobin synthesis is more common in chronic cases whereas shortened erythrocyte life span is more common in acute cases.

In lead-induced anemias, there is usually a reticulocytosis along with the presence of basophilic stippling, and ringed sideroblasts, although none of the above are pathognomonic for lead-induced anemia.

2. Neurological Effects. Inorganic lead has been found to have toxic effects on both the central and peripheral nervous systems. The earliest stages of lead-induced central nervous system effects first manifest themselves in the form of behavioral disturbances and central nervous system symptoms including irritability, restlessness, insomnia and other sleep disturbances, fatigue, vertigo, headache, poor memory, tremor, depression, and apathy. With more severe exposure, symptoms can progress to drowsiness, stupor, hallucinations, delirium, convulsions and coma.

The most severe and acute form of lead poisoning which usually follows ingestion or inhalation of large amounts of lead is acute encephalopathy which may arise precipitously with the onset of intractable seizures, coma, cardiorespiratory arrest, and death within 48 hours.

While there is disagreement about what exposure levels are needed to produce the earliest symptoms, most experts agree that symptoms definitely can occur at blood lead levels of 60 $\mu\text{g}/100\text{ g}$ whole blood and therefore recommend a 40 $\mu\text{g}/100\text{ g}$ maximum. The central nervous system effects frequently are not reversible following discontinued exposure or chelation

therapy and when improvement does occur, it is almost always only partial.

The peripheral neuropathy resulting from lead exposure characteristically involves only motor function with minimal sensory damage and has a marked predilection for the extensor muscles of the most active extremity. The peripheral neuropathy can occur with varying degrees of severity. The earliest and mildest form which can be detected in workers with blood lead levels as low as 50 $\mu\text{g}/100\text{ g}$ is manifested by slowing of motor nerve conduction velocity often without clinical symptoms. With progression of the neuropathy there is development of painless extensor muscle weakness usually involving the extensor muscles of the fingers and hand in the most active upper extremity, followed in severe cases by wrist drop or, much less commonly, foot drop.

In addition to slowing of nerve conduction, electromyographical studies in patients with blood lead levels greater than 50 $\mu\text{g}/100\text{ g}$ have demonstrated a decrease in the number of acting motor unit potentials, an increase in the duration of motor unit potentials, and spontaneous pathological activity including fibrillations and fasciculations. Whether these effects occur at levels of 40 $\mu\text{g}/100\text{ g}$ is undetermined.

While the peripheral neuropathies can occasionally be reversed with therapy, again such recovery is not assured particularly in the more severe neuropathies and often improvement is only partial. The lack of reversibility is felt to be due in part to segmental demyelination.

3. Gastrointestinal. Lead may also affect the gastrointestinal system producing abdominal colic or diffuse abdominal pain, constipation, obstipation, diarrhea, anorexia, nausea and vomiting. Lead colic rarely develops at blood lead levels below 80 $\mu\text{g}/100\text{ g}$.

4. Renal. Renal toxicity represents one of the most serious health effects of lead poisoning. In the early stages of disease nuclear inclusion bodies can frequently be identified in proximal renal tubular cells. Renal function remains normal and the changes in this stage are probably reversible. With more advanced disease there is progressive interstitial fibrosis and impaired renal function. Eventually extensive interstitial fibrosis ensues with sclerotic

glomeruli and dilated and atrophied proximal tubules; all represent end stage kidney disease. Azotemia can be progressive, eventually resulting in frank uremia necessitating dialysis. There is occasionally associated hypertension and hyperuricemia with or without gout.

Early kidney disease is difficult to detect. The urinalysis is normal in early lead nephropathy and the blood urea nitrogen and serum creatinine increase only when two-thirds of kidney function is lost. Measurement of creatinine clearance can often detect earlier disease as can other methods of measurement of glomerular filtration rate. An abnormal Ca-EDTA mobilization test has been used to differentiate between lead-induced and other nephropathies, but this procedure is not widely accepted. A form of Fanconi syndrome with aminoaciduria, glycosuria, and hyperphosphaturia indicating severe injury to the proximal renal tubules is occasionally seen in children.

5. Reproductive effects. Exposure to lead can have serious effects on reproductive function in both males and females. In male workers exposed to lead there can be a decrease in sexual drive, impotence, decreased ability to produce healthy sperm, and sterility. Malformed sperm (teratospermia), decreased number of sperm (hypospermia), and sperm with decreased motility (asthenospermia) can all occur. Teratospermia has been noted at mean blood lead levels of 53 $\mu\text{g}/100\text{ g}$ and hypospermia and asthenospermia at 41 $\mu\text{g}/100\text{ g}$. Furthermore, there appears to be a dose-response relationship for teratospermia in lead exposed workers.

Women exposed to lead may experience menstrual disturbances including dysmenorrhea, menorrhagia and amenorrhea. Following exposure to lead, women have a higher frequency of sterility, premature births, spontaneous miscarriages, and stillbirths.

Germ cells can be affected by lead and cause genetic damage in the egg or sperm cells before conception and result in failure to implant, miscarriage, stillbirth, or birth defects.

Infants of mothers with lead poisoning have a higher mortality during the first year and suffer from lowered birth weights, slower growth, and nervous system disorders.

Lead can pass through the placental barrier and lead levels in the mother's

blood are comparable to concentrations of lead in the umbilical cord at birth. Transplacental passage becomes detectable at 12-14 weeks of gestation and increases until birth.

There is little direct data on damage to the fetus from exposure to lead but it is generally assumed that the fetus and newborn would be at least as susceptible to neurological damage as young children. Blood lead levels of 50-60 $\mu\text{g}/100\text{ g}$ in children can cause significant neurobehavioral impairments and there is evidence of hyperactivity at blood levels as low as 25 $\mu\text{g}/100\text{ g}$. Given the overall body of literature concerning the adverse health effects of lead in children, OSHA feels that the blood lead level in children should be maintained below 30 $\mu\text{g}/100\text{ g}$ with a population mean of 15 $\mu\text{g}/100\text{ g}$. Blood lead levels in the fetus and newborn likewise should not exceed 30 $\mu\text{g}/100\text{ g}$.

Because of lead's ability to pass through the placental barrier and also because of the demonstrated adverse effects of lead on reproductive function in both the male and female as well as the risk of genetic damage of lead on both the ovum and sperm, OSHA recommends a 30 $\mu\text{g}/100\text{ g}$ maximum permissible blood lead level in both males and females who wish to bear children.

6. Other toxic effects. Debate and research continue on the effects of lead on the human body. Hypertension has frequently been noted in occupationally exposed individuals although it is difficult to assess whether this is due to lead's adverse effects on the kidney or if some other mechanism is involved. Vascular and electrocardiographic changes have been detected but have not been well characterized. Lead is thought to impair thyroid function and interfere with the pituitary-adrenal axis, but again these effects have not been well defined.

III. Medical Evaluation

The most important principle in evaluating a worker for any occupational disease including lead poisoning is a high index of suspicion on the part of the examining physician. As discussed in Section 2, lead can affect numerous organ systems and produce a wide array of signs and symptoms, most of which are non-specific and subtle in nature at least in the early stages of disease. Unless serious concern for lead

toxicity is present, many of the early clues to diagnosis may easily be overlooked.

The crucial initial step in the medical evaluation is recognizing that a worker's employment can result in exposure to lead. The worker will frequently be able to define exposures to lead and lead containing materials but often will not volunteer this information unless specifically asked. In other situations the worker may not know of any exposures to lead but the suspicion might be raised on the part of the physician because of the industry or occupation of the worker. Potential occupational exposure to lead and its compounds occur in at least 120 occupations, including lead smelting, the manufacture of lead storage batteries, the manufacture of lead pigments and products containing pigments, solder manufacture, shipbuilding and ship repair, auto manufacturing, construction, and painting.

Once the possibility for lead exposure is raised, the focus can then be directed toward eliciting information from the medical history, physical exam, and finally from laboratory data to evaluate the worker for potential lead toxicity.

A complete and detailed work history is important in the initial evaluation. A listing of all previous employment with information on work processes, exposure to fumes or dust, known exposures to lead or other toxic substances, respiratory protection used, and previous medical surveillance should all be included in the worker's record. Where exposure to lead is suspected, information concerning on-the-job personal hygiene, smoking or eating habits in work areas, laundry procedures, and use of any protective clothing or respiratory protection equipment should be noted. A complete work history is essential in the medical evaluation of a worker with suspected lead toxicity, especially when long term effects such as neurotoxicity and nephrotoxicity are considered.

The medical history is also of fundamental importance and should include a listing of all past and current medical conditions, current medications including proprietary drug intake, previous surgeries and hospitalizations, allergies, smoking history, alcohol consumption, and also non-occupational lead exposures such as hobbies (hunting, riflery). Also known childhood

exposures should be elicited. Any previous history of hematological, neurological, gastrointestinal, renal, psychological, gynecological, genetic, or reproductive problems should be specifically noted.

A careful and complete review of systems must be performed to assess both recognized complaints and subtle or slowly acquired symptoms which the worker might not appreciate as being significant. The review of symptoms should include the following:

General—weight loss, fatigue, decreased appetite.

Head, Eyes, Ears, Nose, Throat (HEENT)—headaches, visual disturbances or decreased visual acuity, hearing deficits or tinnitus, pigmentation of the oral mucosa, or metallic taste in mouth.

Cardio-pulmonary—shortness of breath; cough, chest pains, palpitations, or orthopnea.

Gastrointestinal—nausea, vomiting, heartburn, abdominal pain, constipation or diarrhea.

Neurologic—irritability, insomnia, weakness (fatigue), dizziness, loss of memory, confusion, hallucinations, incoordination, ataxia, decreased strength in hands or feet, disturbances in gait, difficulty in climbing stairs, or seizures.

Hematologic—pallor, easy fatigability, abnormal blood loss, melena.

Reproductive (male and female and spouse where relevant)—history of infertility, impotence, loss of libido, abnormal menstrual periods, history of miscarriages, stillbirths, or children with birth defects.

Musculo-skeletal—muscle and joint pains.

The physical examination should emphasize the neurological, gastrointestinal, and cardiovascular systems. The worker's weight and blood pressure should be recorded and the oral mucosa checked for pigmentation characteristic of a possible Burtonian or lead line on the gingiva. It should be noted, however, that the lead line may not be present even in severe lead poisoning if good oral hygiene is practiced.

The presence of pallor on skin examination may indicate an anemia, which if severe might also be associated with a tachycardia. If an anemia is suspected, an active search for blood loss should be undertaken including

potential blood loss through the gastrointestinal tract.

A complete neurological examination should include an adequate mental status evaluation including a search for behavioral and psychological disturbances, memory testing, evaluation for irritability, insomnia, hallucinations, and mental clouding. Gait and coordination should be examined along with close observation for tremor. A detailed evaluation of peripheral nerve function including careful sensory and motor function testing is warranted. Strength testing particularly of extensor muscle groups of all extremities is of fundamental importance.

Cranial nerve evaluation should also be included in the routine examination.

The abdominal examination should include auscultation for bowel sounds and abdominal bruits and palpation for organomegaly, masses, and diffuse abdominal tenderness.

Cardiovascular examination should evaluate possible early signs of congestive heart failure. Pulmonary status should be addressed particularly if respirator protection is contemplated.

As part of the medical evaluation, the lead standard requires the following laboratory studies:

1. Blood lead level
2. Hemoglobin and hematocrit determinations, red cell indices, and examination of the peripheral blood smear to evaluate red blood cell morphology
3. Blood urea nitrogen
4. Serum creatinine
5. Routine urinalysis with microscopic examination.
6. A zinc protoporphyrin level (This requirement is currently not in effect due to the pending litigation, but is recommended nonetheless).

In addition to the above, the physician is authorized to order any further laboratory or other tests which he or she deems necessary in accordance with sound medical practice. The evaluation must also include pregnancy testing or laboratory evaluation of male fertility if requested by the employee.

Additional tests which are probably not warranted on a routine basis but may be appropriate when blood lead and ZPP levels are equivocal include delta aminolevulinic acid and coproporphyrin concentrations in the

urine, and dark-field illumination for detection of basophilic stippling in red blood cells.

If an anemia is detected further studies including a careful examination of the peripheral smear, reticulocyte count, stool for occult blood, serum iron, total iron binding capacity, bilirubin; and, if appropriate, vitamin B12 and folate may be of value in attempting to identify the cause of the anemia.

If a peripheral neuropathy is suspected, nerve conduction studies are warranted both for diagnosis and as a basis to monitor any therapy.

If renal disease is questioned, a 24 hour urine collection for creatinine clearance, protein, and electrolytes may be indicated. Elevated uric acid levels may result from lead-induced renal disease and a serum uric acid level might be performed.

An electrocardiogram and chest x-ray may be obtained as deemed appropriate.

Sophisticated and highly specialized testing should not be done routinely and where indicated should be under the direction of a specialist.

[Section 1910.1025, Appendix C(III) corrected at 44 FR 68827, November 30, 1979]

IV. Laboratory Evaluation

The blood lead level at present remains the single most important test to monitor lead exposure and is the test used in the medical surveillance program under the lead standard to guide employee medical removal. The ZPP which has several advantages over the blood lead level is, due to the pending litigation, not required under the standard. Because of its relatively recent development and the lack of extensive data concerning its interpretation, the ZPP currently remains an ancillary test.

This section will discuss the blood lead level and ZPP in detail and will outline their relative advantages and disadvantages. Other blood tests currently available to evaluate lead exposure will also be reviewed.

The blood lead level is a good index of current or recent lead absorption when there is no anemia present and when the worker has not taken any chelating agents. However, blood lead levels along with urinary lead levels do not necessarily indicate the total body burden of lead and are not adequate measures of past exposure. One reason for this is that lead has a high affinity for bone and up to 90% of the body's

total lead is deposited there. A very important component of the total lead body burden is lead in soft tissue (liver, kidney, and brain). This fraction of the lead body burden, the biologically active lead, is not entirely reflected by blood lead levels since it is a function of the dynamics of lead absorption, distribution, deposition in bone and excretion. Following discontinuation of exposure to lead, the excess body burden is only slowly mobilized from bone and other relatively stable body stores and excreted. Consequently, a high blood lead level may only represent recent heavy exposure to lead without a significant total body excess and likewise a low blood lead level does not exclude an elevated total body burden of lead.

Also due to its correlation with recent exposures, the blood lead level may vary considerably over short time intervals.

To minimize laboratory error and erroneous results due to contamination, blood specimens must be carefully collected after thorough cleaning of the skin with appropriate methods using lead-free blood containers and analyzed by a reliable laboratory. Under the standard, samples must be analyzed in laboratories which are approved by the Center for Disease Control (CDC) or which have received satisfactory grades in proficiency testing by the CDC in the previous year. Analysis is to be made using atomic absorption spectrophotometry, anodic stripping voltammetry, or any method which meets the accuracy requirements set forth by the standard.

The determination of lead in urine is generally considered a less reliable monitoring technique than analysis of whole blood primarily due to individual variability in urinary excretion capacity as well as the technical difficulty of obtaining accurate 24 hour urine collections. In addition, workers with renal insufficiency, whether due to lead or some other cause, may have decreased lead clearance and consequently urine lead levels may underestimate the true lead burden. Therefore, urine lead levels should not be used as a routine test.

The zinc protoporphyrin test, unlike the blood lead determination, measures an adverse metabolic effect of lead and as such is a better indicator of lead

toxicity than the level of blood lead itself. The level of ZPP reflects lead absorption over the preceding 3 to 4 months, and therefore is a better indicator of lead body burden. The ZPP requires more time than the blood lead to reach significantly elevated levels; the return to normal after discontinuing lead exposure is also slower. Furthermore, the ZPP test is simpler, faster, and less expensive to perform and no contamination is possible. Many investigators believe it is the most reliable means of monitoring chronic lead absorption.

Zinc protoporphyrin results from the inhibition of the enzyme ferrochelatase which catalyzes the insertion of an iron molecule into the protoporphyrin molecule, which then becomes heme. If iron is not inserted into the molecule then zinc, having a greater affinity for protoporphyrin, takes the place of the iron, forming ZPP.

An elevation in the level of circulating ZPP may occur at blood lead levels as low as 20-30 $\mu\text{g}/100\text{ g}$ in some workers. Once the blood lead level has reached 40 $\mu\text{g}/100\text{ g}$ there is more marked rise in the ZPP value from its normal range of less than 100 $\mu\text{g}/100\text{ ml}$. Increases in blood lead levels beyond 40 $\mu\text{g}/100\text{ g}$ are associated with exponential increases in ZPP.

Whereas blood lead levels fluctuate over short time spans, ZPP levels remain relatively stable. ZPP is measured directly in red blood cells and is present for the cell's entire 120 day life-span. Therefore, the ZPP level in blood reflects the average ZPP production over the previous 3-4 months and consequently the average lead exposure during that time interval.

It is recommended that a hematocrit be determined whenever a confirmed ZPP of 50 $\mu\text{g}/100\text{ ml}$ whole blood is obtained to rule out a significant underlying anemia. If the ZPP is in excess of 100 $\mu\text{g}/100\text{ ml}$ and not associated with abnormal elevations in blood lead levels, the laboratory should be checked to be sure that blood leads were determined using atomic absorption spectrophotometry, anodic stripping voltammetry, or any method which meets the accuracy requirements set forth by the standard by a CDC approved laboratory which is experienced in lead level determinations. Repeat periodic blood lead studies should be obtained in all individuals with elevated ZPP levels to be certain that an associated elevated blood lead level has not been missed.

[Sec. 1910.1025, Appendix C]

due to transient fluctuations in blood leads.

ZPP has a characteristic fluorescence spectrum with a peak at 594 nm which is detectable with a hematofluorimeter. The hematofluorimeter is accurate and portable and can provide on-site, instantaneous results for workers who can be frequently tested via a finger prick.

However, careful attention must be given to calibration and quality control procedures. Limited data on blood lead—ZPP correlations and the ZPP levels which are associated with the adverse health effects discussed in Section 2 are the major limitations of the test. Also it is difficult to correlate ZPP levels with environmental exposure and there is some variation of response with age and sex. Nevertheless, the ZPP promises to be an important diagnostic test for the early detection of lead toxicity and its value will increase as more data is collected regarding its relationship to other manifestations of lead poisoning.

Levels of delta-aminolevulinic acid (ALA) in the urine are also used as a measure of lead exposure. Increasing concentrations of ALA are believed to result from the inhibition of the enzyme delta-aminolevulinic acid dehydrase (ALA-D). Although the test is relatively easy to perform, inexpensive, and rapid,

the disadvantages include variability in results, the necessity to collect a complete 24 hour urine sample which has a specific gravity greater than 1.010, and also the fact that ALA decomposes in the presence of light.

The pattern of porphyrin excretion in the urine can also be helpful in identifying lead intoxication. With lead poisoning, the urine concentrations of coproporphyrins I and II, porphobilinogen and uroporphyrin I rise. The most important increase, however, is that of coproporphyrin III; levels may exceed 5,000 $\mu\text{g}/\text{l}$ in the urine in lead poisoned individuals, but its correlation with blood lead levels and ZPP are not as good as those of ALA. Increases in urinary porphyrins are not diagnostic of lead toxicity and may be seen in porphyria, some liver diseases, and in patients with high reticulocyte counts.

[Section 1910.1025, Appendix C(IV) corrected at 44 FR 68827, November 30, 1979]

Summary. The Occupational Safety and Health Administration's standard for inorganic lead places significant emphasis on the medical surveillance of all workers exposed to levels of inorganic lead above the action level of 30 $\mu\text{g}/\text{m}^3$ TWA. The physician has a fundamental role in this surveillance program, and in the operation of the medical removal protection program.

Even with adequate worker education on the adverse health effects of lead and appropriate training in work practices, personal hygiene and other control measures, the physician has a primary responsibility for evaluating potential lead toxicity in the worker. It is only through a careful and detailed medical and work history, a complete physical examination and appropriate laboratory testing that an accurate assessment can be made. Many of the adverse health effects of lead toxicity are either irreversible or only partially reversible and therefore early detection of disease is very important.

This document outlines the medical monitoring program as defined by the occupational safety and health standard for inorganic lead. It reviews the adverse health effects of lead poisoning and describes the important elements of the history and physical examinations as they relate to these adverse effects.

Finally, the appropriate laboratory testing for evaluating lead exposure and toxicity is presented.

It is hoped that this review and discussion will give the physician a better understanding of the OSHA standard with the ultimate goal of protecting the health and well-being of the worker exposed to lead under his or her care.

[The next page is 31:8451]

SUMMARY OF MEDICAL REQUIREMENTS

WORK EXPOSURE (REFERENCE AND DATE) BOLD TYPE FOR OSHA STANDARDS	A. PRE-PLACEMENT REQUIREMENT FOR THOSE WHO WILL BE OR ARE SUBJECT TO OCCUPATIONAL EXPOSURE (FOR REC- OMMENDATIONS), OR FOR THOSE WHO WILL BE OR ARE EMPLOYED IN REGULATED AREAS (FOR STANDARDS), UNLESS OTHER- WISE STATED.	B. PERIODIC REQUIREMENTS FOR THOSE SUBJECT TO OCCUPATIONAL EXPOSURE (FOR RECOMMENDATIONS), OR FOR THOSE WHO ARE AUTHORIZED EMPLOYEES (FOR STANDARDS), UNLESS OTHERWISE STATED.	C. OTHER REQUIREMENTS FOR THOSE SUBJECT TO OCCUPATIONAL EXPOSURE (FOR RECOMMENDATIONS), OR FOR THOSE WHO ARE AUTHORIZED EMPLOYEES (FOR STANDARDS), UNLESS OTHERWISE STATED.
Asbestos (1910.1001, 11/78)	- For those exposed to airborne concentrations. - Medical exam, with a minimum of: 14" x 17" post/ant chest X-ray; - Pulmonary function tests (FVC, FEV1.0). - Medical history, eliciting symptomatology of respiratory diseases.	At least annually - For those exposed to airborne concentrations. - Same as in A.	30 days prior to termination of employment Same as in A.
Asphalt fumes (NIOSH 78-106, 10/78)	- Comprehensive medical and work histories, emphasizing: - Eyes; - Skin; - Respiratory system. - Physical exam, with attn to abnormalities in: - Eyes; - Skin; - Respiratory system. - Ability to use positive or negative pressure respirator.	As DBRP Same as in A.1 and 2.	In the event of illness known or suspected to be due to asphalt fumes Physical examination.
Benzene (1910.1028, 4/79)	- For those who are or may be exposed to benzene at or above the action level and those subjected to an emergency. - Work history: - Past exposure to benzene and other hematologic toxins; - Exposure to marrow toxins outside current work situation, including volatile cleansing agents and insecticides.	At least semi-annually - Same as in A. - Interim work history: - New exposure to potential marrow toxins. - Interim personal history: - Change in drug/alcohol intake; - Change in appearance of physical symptoms relating to blood disorders.	In an emergency - Urinary phenol test. - If C-1 is 75 mg phenol/l of urine, then provide complete blood count as in A.5.a.i, ii, and iii. This should also be done one month later. - If any of the blood tests taken for either A or B indicate abnormal red cell, white cell, or thrombocyte counts or hemoglobin level, employee should be referred to a hematologist.

DBRP - DETERMINED BY RESPONSIBLE PHYSICIAN

WORK EXPOSURE (REFERENCE AND DATE) BOLD TYPE FOR OSHA STANDARDS	A. PRE-PLACEMENT REQUIREMENT FOR THOSE WHO WILL BE OR ARE SUBJECT TO OCCUPATIONAL EXPOSURE (FOR REC- OMMENDATIONS), OR FOR THOSE WHO WILL BE OR ARE EMPLOYED IN REGULATED AREAS (FOR STANDARDS), UNLESS OTHER- WISE STATED.	B. PERIODIC REQUIREMENTS FOR THOSE SUBJECT TO OCCUPATIONAL EXPOSURE (FOR RECOMMENDATIONS), OR FOR THOSE WHO ARE AUTHORIZED EMPLOYEES (FOR STANDARDS), UNLESS OTHERWISE STATED.	C. OTHER REQUIREMENTS FOR THOSE SUBJECT TO OCCUPATIONAL EXPOSURE (FOR RECOMMENDATIONS), OR FOR THOSE WHO ARE AUTHORIZED EMPLOYEES (FOR STANDARDS), UNLESS OTHERWISE STATED.
Crystalline silica (NIOSH 75-120, 11/74)	- Medical and work history: - Exposure to free silica; - Signs/symptoms of respiratory disease. 14" x 17" or 14" x 14" post/ant chest X-ray. - Baseline FVC and FEV1.0. - Ability to use negative or positive pressure respirator. - Body weight, height, and age.	At least every 3 yrs Same as in A, except 3.	
1,2-Dibromo-3-chloropropane (1910.1044, 7/78)	- Medical and work histories, including reproductive history. - Physical exam, including: - Genitourinary tract, testical size, body habitus, including sperm count; - Serum specimen: FSH, LH, total estrogen (for women), and others, as DBRP.	Annually - Same as in A.1 and 2. - Additional appropriate examinations, as DBRP.	In an emergency situation - Sperm count as soon as practicable, and 3 mos later. - If vasectomized, serum specimen as in A.2.b.
3,3-dichlorobenzidine (and its salts) (1910.1007, 11/78)	- Physical exam, with attn to: - Decreased immunocompetence; - Treatment with steroid or cytotoxic agent; - Pregnancy; - Cigarette smoking. - Personal, family, and occupational histories, including: - Genetic factors; - Environmental factors.	At least annually Same as in A.	

WORK EXPOSURE (REFERENCE AND DATE) BOLD TYPE FOR OSHA STANDARDS	A. PRE-PLACEMENT REQUIREMENT FOR THOSE WHO WILL BE OR ARE SUBJECT TO OCCUPATIONAL EXPOSURE (FOR REC- OMMENDATIONS), OR FOR THOSE WHO WILL BE OR ARE EMPLOYED IN REGULATED AREAS (FOR STANDARDS), UNLESS OTHER- WISE STATED.	B. PERIODIC REQUIREMENTS FOR THOSE SUBJECT TO OCCUPATIONAL EXPOSURE (FOR RECOMMENDATIONS), OR FOR THOSE WHO ARE AUTHORIZED EMPLOYEES (FOR STANDARDS), UNLESS OTHERWISE STATED.	C. OTHER REQUIREMENTS FOR THOSE SUBJECT TO OCCUPATIONAL EXPOSURE (FOR RECOMMENDATIONS), OR FOR THOSE WHO ARE AUTHORIZED EMPLOYEES (FOR STANDARDS), UNLESS OTHERWISE STATED.
Dinitro-ortho-cresol (continued) Dioxane (NIOSH 77-226, 9/77) Emergency egress from elevated work stations (NIOSH 76-128, 1976)	b. Complete blood count; c. Hematocrit; d. Liver function; e. Kidney function. 4. Ability to use positive and negative pressure respirators. 1. Comprehensive medical and work histories, emphasizing disorders of: a. Upper respiratory system; b. Hepatic function; c. Renal function. 2. Physical exam, with attn to: a. Nares; b. Hepatic function; c. Renal function. 3. Clinical tests, at least: a. Liver function test, e.g., SGOT, SGPT; b. Kidney function test, e.g., complete urinalysis. 4. Ability to use positive pressure respirator. 1. Medical and work histories. 2. If work at <80 ft but >15 ft: a. Depth perception; b. Fields of vision; c. Reaction time; d. Manual dexterity; e. Coordination; f. Abnormal tendency toward dizziness.	Annually 1. Same as in A.1, 2, and 3. Annually, or as DBRP 1. Same as in A.	

WORK EXPOSURE (REFERENCE AND DATE) BOLD TYPE FOR OSHA STANDARDS	A. PRE-PLACEMENT REQUIREMENT FOR THOSE WHO WILL BE OR ARE SUBJECT TO OCCUPATIONAL EXPOSURE (FOR REC- OMMENDATIONS), OR FOR THOSE WHO WILL BE OR ARE EMPLOYED IN REGULATED AREAS (FOR STANDARDS), UNLESS OTHER- WISE STATED.	B. PERIODIC REQUIREMENTS FOR THOSE SUBJECT TO OCCUPATIONAL EXPOSURE (FOR RECOMMENDATIONS), OR FOR THOSE WHO ARE AUTHORIZED EMPLOYEES (FOR STANDARDS), UNLESS OTHERWISE STATED.	C. OTHER REQUIREMENTS FOR THOSE SUBJECT TO OCCUPATIONAL EXPOSURE (FOR RECOMMENDATIONS), OR FOR THOSE WHO ARE AUTHORIZED EMPLOYEES (FOR STANDARDS), UNLESS OTHERWISE STATED.
Emergency egress from elevated work stations (continued)	3. If work at >80 ft: a. Visual acuity to ensure at least 20/30 Snellen in one eye and 20/50 in the other (correct with lenses if necessary); b. Color perception; c. Hearing tests, with or w/o aids (to ensure at least 15/20 in one ear; or a hearing loss of ≤ 20 db in normally tested frequencies); d. Strength; e. Endurance; f. Agility; g. Coordination; h. Speed of reactions; i. Beware of: i. Cardiovascular diseases; ii. Epilepsy; iii. Diabetes; iv. Emotional instability.		
Epichlorohydrin (NIOSH 76-206, 9/76)	1. Comprehensive medical and work histories. 2. Complete physical exam, with attn to: a. Kidneys; b. Liver; c. Respiratory tract; d. Hematopoietic system; e. Complaints and evidence of eye, mucous membrane, and skin irritations. 3. As DBRP: a. SGPT and SGOT; b. Lactic dehydrogenase. 4. Ability of workers to use respirators.	Annually 1. Same as in A.1, 2, and 3.	Additional 1. For overexposure. 2. First aid.

WORK EXPOSURE (REFERENCE AND DATE) DOLD TYPE FOR OSHA STANDARDS	A. PRE-PLACEMENT REQUIREMENT FOR THOSE WHO WILL BE OR ARE SUBJECT TO OCCUPATIONAL EXPOSURE (FOR REC- OMMENDATIONS), OR FOR THOSE WHO WILL BE OR ARE EMPLOYED IN REGULATED AREAS (FOR STANDARDS), UNLESS OTHER- WISE STATED.	B. PERIODIC REQUIREMENTS FOR THOSE SUBJECT TO OCCUPATIONAL EXPOSURE (FOR RECOMMENDATIONS), OR FOR THOSE WHO ARE AUTHORIZED EMPLOYEES (FOR STANDARDS), UNLESS OTHERWISE STATED.	C. OTHER REQUIREMENTS FOR THOSE SUBJECT TO OCCUPATIONAL EXPOSURE (FOR RECOMMENDATIONS), OR FOR THOSE WHO ARE AUTHORIZED EMPLOYEES (FOR STANDARDS), UNLESS OTHERWISE STATED.
Glycidyl ethers (continued)	c. Pulmonary function tests; d. Manual dexterity tests; e. Visual-motor coupling tests.		
Hot environments (NIOSH 72-10269, 6/72)	1. Medical examination, with attn to: a. Cardiovascular system; b. Renal system; c. Hepatic system; d. Endocrine system; e. Respiratory system; f. Skin 2. History of previous heat-related disorders and illnesses.	Biennially 1. For those ≤ 45 yrs old. 2. Same as in A. Annually 3. For those > 45 yrs old. 4. Same as in A.	
Hydrazines (NIOSH 78-172, 6/78)	1. Comprehensive medical and work histories, with special attn to: a. Dermal contact. 2. Comprehensive physical exam, with special attn to: a. Dermal contact. 3. Clinical tests, including: a. Complete and differential blood count; b. Liver function test (SGOT, SGPT included); c. Urinalysis, including specific gravity, glucose, protein, microscopic exam; d. 14" x 17" post/ant chest X-ray. 4. Urobilinogen and serum bilirubin tests, as DBRP. 5. Ability to use positive pressure respirator.	At least annually 1. Same as A. 1, 2, 3, and 4.	Additional 1. For those with illnesses due to exposure. 2. Appropriate medical services. In an emergency 3. Immediate medical attention and appropriate follow-up medical care.

WORK EXPOSURE (REFERENCE AND DATE) BOLD TYPE FOR OSHA STANDARDS	A. PRE-PLACEMENT REQUIREMENT FOR THOSE WHO WILL BE OR ARE SUBJECT TO OCCUPATIONAL EXPOSURE (FOR REC- OMMENDATIONS), OR FOR THOSE WHO WILL BE OR ARE EMPLOYED IN REGULATED AREAS (FOR STANDARDS), UNLESS OTHER- WISE STATED.	B. PERIODIC REQUIREMENTS FOR THOSE SUBJECT TO OCCUPATIONAL EXPOSURE (FOR RECOMMENDATIONS), OR FOR THOSE WHO ARE AUTHORIZED EMPLOYEES (FOR STANDARDS), UNLESS OTHERWISE STATED.	C. OTHER REQUIREMENTS FOR THOSE SUBJECT TO OCCUPATIONAL EXPOSURE (FOR RECOMMENDATIONS), OR FOR THOSE WHO ARE AUTHORIZED EMPLOYEES (FOR STANDARDS), UNLESS OTHERWISE STATED.
Ketone (NIOSH 78-173, 10/78)	- Comprehensive medical and work histories: - Pre-existing neuropathy; - Disorders of: - Nervous system; - Respiratory system; - Eyes. - Physical exam, with attn to: - Central and peripheral nervous system; - Respiratory system; - Eyes. - Urinalysis, as indicator of kidney function. - Appropriate liver tests, as DBRP. - An electrodiagnostic exam (for those exposed to methyl n-butylketone), including electromyography and nerve conduction velocity measurement. - Ability to use negative and positive pressure respirators.	Annually, and as DBRP Same as in A, except 6.	Additional - For those with adverse effects from exposure. - Appropriate medical management.
Lead (inorganic) (1910.1025, 4/79) <i>SEE OSHA STANDARD</i>	- For those who are or may be exposed above the action level for more than 30 days/yr. - Work history, including past lead exposure at and outside work. - Personal or medical history, with attn to: - Smoking; - Hygiene and other habits; - Past gastrointestinal problems; - Past reproductive problems; - Past cardiovascular problems; - Past neurologic problems.	At least annually - For those in A.1 who have a blood lead level $\geq 40 \mu\text{g}/100 \text{ g}$ in the past yr. - Same as in A.	Additional - For those developing signs/symptoms associated with lead intoxication. - Same as in A. Additional - For those removed from area due to risk of impairment to health, or for those restricted due to a final medical exam. - Appropriate medical management.

WORK EXPOSURE (REFERENCE AND DATE) BOLD TYPE FOR OSHA STANDARDS	A. PRE-PLACEMENT REQUIREMENT FOR THOSE WHO WILL BE OR ARE SUBJECT TO OCCUPATIONAL EXPOSURE (FOR REC- OMMENDATIONS), OR FOR THOSE WHO WILL BE OR ARE EMPLOYED IN REGULATED AREAS (FOR STANDARDS), UNLESS OTHER- WISE STATED.	B. PERIODIC REQUIREMENTS FOR THOSE SUBJECT TO OCCUPATIONAL EXPOSURE (FOR RECOMMENDATIONS), OR FOR THOSE WHO ARE AUTHORIZED EMPLOYEES (FOR STANDARDS), UNLESS OTHERWISE STATED.	C. OTHER REQUIREMENTS FOR THOSE SUBJECT TO OCCUPATIONAL EXPOSURE (FOR RECOMMENDATIONS), OR FOR THOSE WHO ARE AUTHORIZED EMPLOYEES (FOR STANDARDS), UNLESS OTHERWISE STATED.
Lead (inorganic) (continued)	4. Physical exam, with attn to: a. Teeth and gums; b. Hematologic system; c. Gastrointestinal system; d. Renal system; e. Neurologic system; f. Pulmonary status (to be evaluated if respiratory protection will be used). 5. Lab tests: a. Blood pressure; b. Blood sample and analysis: i. Blood lead levels; ii. Hemoglobin; iii. Hematocrit; iv. Red cell indices; v. Peripheral smear morphology; vi. Zinc protoporphyrin; vii. Blood urea nitrogen; viii. Serum creatinine; c. Routine urinalysis, with microscopic exam; d. Any other tests, as DBRP.		
Logging (from felling to first haul) (NIOSH 76-188, 7/76)	1. Medical examination for previously existing and new conditions that would endanger worker's safety: a. Cardiovascular system; b. Musculoskeletal system; c. Central nervous system; d. Integrity of extremities and digits; e. Integrity of hearing; f. Visual acuity; g. Peripheral vasculature (Reynaud's syndrome), for chain saw operators. 2. Complete series of tetanus toxoid inoculations.	Biennially 1. For those <40 yrs old. 2. Same as in A.1. Annually 3. For those ≥40 yrs old. 4. Same as in A.1. Every 10 yrs, unless injury intervenes 5. Booster doses of tetanus toxoid inoculations.	After absence for injury of 5 or more days 1. Medical exam to determine fitness. In an emergency 2. First aid, with emergency assistance and medical services, as necessary.

WORK EXPOSURE (REFERENCE AND DATE) BOLD TYPE FOR OSHA STANDARDS	A. PRE-PLACEMENT REQUIREMENT FOR THOSE WHO WILL BE OR ARE SUBJECT TO OCCUPATIONAL EXPOSURE (FOR REC- OMMENDATIONS), OR FOR THOSE WHO WILL BE OR ARE EMPLOYED IN REGULATED AREAS (FOR STANDARDS), UNLESS OTHER- WISE STATED.	B. PERIODIC REQUIREMENTS FOR THOSE SUBJECT TO OCCUPATIONAL EXPOSURE (FOR RECOMMENDATIONS), OR FOR THOSE WHO ARE AUTHORIZED EMPLOYEES (FOR STANDARDS), UNLESS OTHERWISE STATED.	C. OTHER REQUIREMENTS FOR THOSE SUBJECT TO OCCUPATIONAL EXPOSURE (FOR RECOMMENDATIONS), OR FOR THOSE WHO ARE AUTHORIZED EMPLOYEES (FOR STANDARDS), UNLESS OTHERWISE STATED.
N-Nitrosodimethylamine (continued) Noise (NIOSH 73-11001, 8/72) Organotin compounds (NIOSH 77-115, 11/76)	2. Personal, family, and occupational histories, including: a. Genetic factors; b. Environmental factors. 1. Baseline audiogram, which includes: a. Aural medical history; b. Last exposure to high level noise; c. Number of hours since exposure; d. Type of exposure; e. Noise level, if known. 1. Comprehensive medical and work histories. 2. Complete physical exam, with attn to: a. 14" x 17" post/ant chest X-ray; b. FVC and FEV1.0; c. SGOT, SGPT, and any other tests of hepatic function, as DBRP; d. Eye exam, including tests for: i. Visual acuity; ii. Color vision; iii. Pupillary reactions; iv. Glaucoma; v. Possibility of choked disc; e. ECG (for ≥ 40 yrs old); f. Prior history of increased intracranial pressure; g. If spinal fluid is measured, then perform Quekenstedt maneuver; h. Urinalysis. 3. Ability to use positive or negative pressure respirator.	Within 30 days of assignment, and every 6 yrs 1. Same as in A. Biennially 1. See pp 1-6 and 7 of Reference for specific levels for testing. Annually, or as DBRP 1. Same as in A. 1 and 2, with the exception of neurologic exam, as DBRP.	In an emergency 1. Examination to determine need for treatment, with attn to: a. Eyes; b. Lungs. Additional 2. For those with adverse health effects. 3. Appropriate medical services.

