

Common External Marks of Occupation or Occupational Diseases or Other Diseases (continued)

502

Mark or sign	Probable cause	Significance	Comment
Lips, Mouth, Teeth, Tongue, Neck, continued			
"Beefy" tongue.	Vitamin B deficiency.	One manifestation of vitamin B lack. R. M. I.	Enlarged tongue, roughened, rugose and red; somewhat resembles raw beef.
Herpes (labial)—"cold sores."	Action of a virus, allergy.	Virus probably always present but only becomes active under unfavorable conditions such as infection.	Not to be confused with labial chancre.
Occupational fungal disease around mouth.	Condition limited to Mediterranean countries. Involves chiefly quail force-feeders.	Organism transferred by quail pecking feeders' faces.	Condition not known in United States.
Discolored gums.	Various metal or mineral deposits, tobacco stain, etc.	The purple gum line of lead absorption, the similar line from bismuth, the blue line following trinitrotoluene exposure or poisoning, purple line from copper, bluish line from mercury or zinc, purple gums from scurvy. R. M. I.	Lines or discoloration of gums are not conclusive proof of any occupational disease.
Burton line (lead line).	Bright purple line at gum margin. Common after exposure to lead.	Not diagnostic of lead poisoning.	Not readily visible as a sign of abnormality. R. M. I.
Acute multiple dental caries.	Specific form attributed to work with citric acid, chiefly bakers and confectioners.	Well recognized in European countries. Uncommon in exaggerated form in United States.	All sugars as used in bakeries and confectionaries may produce some excess of dental caries.
Dental erosions.	Erosion from chemicals, such as from acids or mechanical erosion such as from glass blowers' pipes; numerous variations; specific variations for many trades, such as battery makers from acid, sewers from thread biting, wind instrument players, "tack spitters," brush makers who trim bristles by biting, pencil chewers in offices.	Profoundly eroded teeth of acid workers seldom decayed and are not painful.	Apart from dental erosion constituting a sign of occupational damage in many trades, the premature loss of teeth may possibly be related to work as a contributory cause.
"Mulberry molars."	Congenital syphilis.	Dome-shaped, stunted first molars due to overgrowth of enamel.	Various other dental defects suggest congenital syphilis.
Hutchinson's teeth.	Congenital syphilis.	Notched incisors with narrow edges and peg topped.	May be associated with impaired hearing and baked cornea.

CAREY P. MCCORD