

ently sensitive
In our opinion,
ssi

Mutation Research, 41 (1976) 131-142
© Elsevier/North-Holland Biomedical Press

o Mrs. J. Novot-

CARCINOGENIC, MUTAGENIC AND TERATOGENIC RISKS ASSOCIATED WITH VINYL CHLORIDE

PETER F. INFANTE, JOSEPH K. WAGONER and RICHARD J. WAXWEILER

Division of Surveillance, Hazard Evaluations and Field Studies, National Institute for Occupational Safety and Health, Main Post Office Building, Cincinnati, Ohio 45202 (U.S.A.)

(Received May 13th, 1976)

Summary

The data presented demonstrate clearly that vinyl chloride (VC) is related to a significant excess of mortality from cancer of the liver, lung and brain among workers occupationally exposed to VC. The risk of dying from cancer of the lymphatic and hematopoietic system also appears to increase with an increase in latency. These cancer sites could have been predicted by the animal bioassay conducted by Maltoni. With regard to the liver, even the histopathologic type of cancer (angiosarcoma) was observed first in experimental animals. A study of cancer mortality among populations residing proximate to VC polymerization facilities also demonstrated an increased risk of dying from CNS and lymphatic cancer. These latter findings raise cause for concern about out-plant emissions of VC, but without further study these cancers obviously cannot be interpreted as being related to out-plant exposure to VC.

Various test systems now have elicited a positive mutagenic response to VC. Thus, our observations of a significant excess of fetal mortality among the wives of males, who were occupationally exposed to VC, raise public health concern that VC may be mutagenic in humans.

With regard to the teratogenicity of VC, observations of a significant excess of children born with birth defects were reported among populations residing proximate to VC polymerization facilities. Additional epidemiologic study is needed to determine whether a repeated pattern of excessive numbers of children born with birth defects can be observed in other communities with VC polymerization facilities.

Introduction

In 1930, the first adverse health effects of vinyl chloride (VC) were reported [22]. Since then, numerous investigators have reported the toxic effects of VC on the central nervous system, the liver, the bones of the fingers and the lungs

tion production, III,

ties, Vol. 3. Plenum

some change during

nosome aberrations,

n. Humangenetik, 6

t Imuran (Azathiop-

s. Osterman-Golther,

i with Azathioprine,

the preparation of

(1965) 333-338,

ing treatment with

ne of the frequen-

d ical agents,

orbs drin. I. Testing

(1976) 271-278.

osomal aberrations

on Res., 34 (1976)

vinylchloride workers,

is, 3/2 (1973) 86-

on. Health

Arch. Toxicol., 28

it in vitro for the

" Springer-Verlag,

the occupational

f cytovan-induced

980 002045

[2,5,9-12,18,20,28]. More recently, the study of VC toxicity has broadened to assess the spectrum of carcinogenesis [2,15,16,21,26,29,30], mutagenesis [1,3,4,6,13,14,23,24], and teratogenesis [8]. These efforts were stimulated by the work of Viola et al. [29] who, in 1971, reported the induction of tumors of the skin, lungs and bones in rats exposed by inhalation to 30,000 ppm of VC over a period of twelve months. Widespread concern for the carcinogenic activity of VC, however, did not occur until early 1974, when Creech and Johnson [2] reported four deaths from angiosarcoma of the liver among workers employed in the manufacture of polyvinyl chloride (PVC) resins. It was later learned that Maltoni had previously demonstrated VC-induced hepatic angiosarcomas, lung adenomas, brain neuroblastomas, lymphomas and various other tumors in mice, rats and hamsters [15,16].

With regard to mutagenicity, several investigators have induced mutations via microbial test systems [1,14,24]. Also, VC metabolites have induced mutations in mammalian cells [6]. In addition, reports from several countries have demonstrated significant excesses of chromosomal aberrations among workers exposed to VC as contrasted with those not exposed [3,4,13,23]. Because of these observations and of the widespread exposure to VC among both workers and the general population, the National Institute for Occupational Safety and Health (NIOSH) in the U.S.A. undertook an epidemiologic program to evaluate the magnitude and spectrum of VC toxicity to humans. This program was 4-fold in nature and sought to evaluate: (1) The site-specific risk of cancer among workers exposed to VC [30]. (2) The risk of some cancers among populations residing proximate to VC polymerization facilities [8]. (3) The site specific risk of congenital anomalies among populations residing proximate to VC polymerization facilities [8]. (4) The risk of fetal wastage among the wives of workers occupationally exposed to VC [7].

Cancer risk among workers exposed to VC

To assess the neoplastic risk of workers exposed to VC, a population consisting of employees from four VC polymerizing facilities was selected for cohort mortality study [30]. Since occupationally induced cancers often take many years to become manifest, the study cohort was restricted to workers who had achieved five or more years of employment and for whom at least 10 years had lapsed since initial employment. Thus, the exposure period was five or more years and the latency period was 10 or more years. Follow-up for study cohort members was greater than 99%.

Table I shows the total mortality experience among the study cohort. The expected numbers are based on United States mortality data applied to NIOSH's modified life-table method. For all malignant neoplasms, there were 35 observed vs 23.5 expected, the SMR was 149. This excess was significant at the $P < 0.05$ level of confidence. For total mortality, 136 deaths were observed vs 126.3 expected. Selecting a cohort with at least five years work experience and 10 years latency eliminated the "healthy worker effect" [19] and so there were more total deaths than expected and this was mostly at the expense of cancer mortality.

Table II shows cancer mortality experience by greater than 10 and 15 year latency periods. At the greater than 10 year latency period, only biliary and

TABLE
MORT

Cause

All mal
Heart
Non-m
Cirrho
All oth

a ICD
b SMR
c Sign

liver
with
cate;
canc
how
Thes
occu

TABLE
CANC
TO V

Site
All m

Brain

Respi

Bihar

Lym

All o

a Ob
b Ex
c SM
d Su
e Su

was broadened
mutagenesis
stimulated
induction of
for 30,000
concern for the
1974, when
of the liver
chloride (PVC)
strated VC-in-
s, lymphomas

mutations via
ed mutations
ountries have
ong workers
l. Because of
both workers
al Safety and
n to evaluate
program was
sk of cancer
ncers among
(3) The site
proximate to
ng the wives

a con-
selected for
ncers often
restricted to
for whom at
posure period
s. Follow-up

cohort. The
applied to
, there were
significant at
re observed
experience
and so there
expense of

and 15 year
biliary and

TABLE I
MORTALITY EXPERIENCE AMONG COHORT WORKERS EXPOSED TO VINYL CHLORIDE

Cause of death	ICD code ^a	Observed	Expected	SMR ^b
All malignant neoplasms	(140-205)	35	23.5	149 ^c
Heart	(400-443)	57	54.7	104
Non-malignant respiratory diseases	(470-527)	6	3.4	176
Cirrhosis	(581)	2	4.0	50
All other causes		36	40.7	88

^a ICD, International Classification of Diseases, 7th revision.

^b SMR, standardized mortality ratio.

^c Significant at $P < 0.05$.

liver cancer deaths were significantly in excess; however, when the sub-cohort with 15 years of latency since initial employment was used, deaths from three categories of cancer were significantly in excess. The excess in mortality from cancer of the lymphatic and hematopoietic systems was not significant; however, the SMR increased from 159 to 176 with an increase in latency. These comparisons show the importance of latency when looking for occupationally-induced cancers.

TABLE II
CANCER MORTALITY BY INTERVAL SINCE INITIAL EXPOSURE AMONG WORKERS EXPOSED TO VINYL CHLORIDE

Site of malignancy		10+ years		15+ years
All malignant neoplasms	Obs. ^a	35		31
	Exp. ^b	23.5		16.9
	SMR ^c	149 ^d	<	184 ^e
Brain and CNS cancer	Obs.	3		3
	Exp.	0.9		0.6
	SMR	329	<	498 ^d
Respiratory system cancer	Obs.	12		11
	Exp.	7.7		5.7
	SMR	156	<	194 ^d
Biliary and liver cancer	Obs.	7		7
	Exp.	0.6		0.4
	SMR	1153 ^c	<	1606 ^e
Lymphatic and hematopoietic system cancer	Obs.	4		3
	Exp.	2.5		1.7
	SMR	159	<	176
All other malignant neoplasms	Obs.	9		7
	Exp.	11.7		8.4
	SMR	77	<	83

^a Obs., observed.

^b Exp., expected.

^c SMR, standardized mortality ratio.

^d Significant at $P < 0.05$.

^e Significant at $P < 0.01$.

BBG 002047

In laboratory studies, Maltoni has induced hepatic angiosarcomas, lung adenomas, brain neuroblastomas and lymphomas [16]. Therefore, the predictive value of animal bioassay studies can be demonstrated, not only by the observation of a significant excess of angiosarcoma of the liver among VC workers, but also by significant excesses of lung and brain tumors.

Cancer risks in populations residing near VC polymerization facilities

The risk of some cancers among adult populations residing in the only three Ohio communities with VC polymerization facilities was assessed [8]. All three communities had at least one facility in operation by 1954 and Painesville had a second plant in operation by 1967. The community population sizes varied from 12,000–24,000. Between 1960–70 the population had remained stable in Painesville and Ashtabula, but had increased by 30% in Avon Lake. As reported previously, for the three communities taken as a whole, there were no apparent differences in racial origin or family income as compared to the average for the State [8].

As a result of previous findings [26], cancers of the central nervous system, lymphatic and hematopoietic systems, were selected for study. Because of possible error in death certificate data in terms of metastases from other primary sites, data for lung and liver cancers were not analyzed.

Table III shows data for observed versus expected CNS cancer deaths in the white population by sex for the period 1958–73. North Ridgeville is the only community shown which does not have a PVC polymerization facility. It was included in the analyses of the cancer data because it is located contiguous to Avon Lake and because it had a high incidence of children born with birth defects [8]. If North Ridgeville had not been included in the analyses of cancer mortality, little difference in the results would have been observed. Expected values are based on the occurrence in the balance of the counties over the same period of time. As shown in Table III, there was a significant excess of CNS cancer deaths in males. The excess was greatest in Painesville and North Ridgeville. With all communities combined, there were 27 CNS cancer deaths observed in males versus 14.1 expected; the SMR was 191. The difference was significant at $P < 0.01$.

In females, with all groups combined, there was only a slight excess. With sex groups combined, the excess was significant at $P < 0.01$. It may be noteworthy that one father-daughter combination for CNS tumor deaths occurred in Painesville. The daughter died of a papillary ependymoma in 1963 at the age of 16. The father died two years later of a glioblastoma multiforme at the age of 58. It would be difficult to determine whether this observation is the result of genetic or environmental factors.

Table III also shows deaths from lymphomas. Although the differences between observed and expected deaths in males were not significant, there was a consistent excess in each community. In females, there were excess lymphoma deaths in two of the four communities. With sex groups combined, a significant excess of lymphoma deaths was observed in Ashtabula. With sex groups and communities combined, the number of observed deaths was 61 versus 48.7 expected. The SMR was 125. This was significant at

TABLE III
OBSERVED AND EXPECTED
YEARS AND OLDER
FACILITIES, 1958–73

Expected numbers of
the counties in which

	M	F
Ashtabula	1	1
Painesville	1	1
Avon Lake	1	1
N. Ridgeville	1	1
Communities Combined	2	2

a International Class
b $P < 0.05$.
c $P < 0.01$.
d $P < 0.002$.

0.05 < $P < 0.1$
ty. As shown i
identical.

Birth defects a

The occurrence
communities v
North Ridgevil
in the initial a
North Ridgevil
subsequent an
subsequent an
tions among c

GGC 002048

TABLE III

OBSERVED AND EXPECTED DEATHS FOR THREE TYPES OF CANCER FOR RESIDENTS 45 YEARS AND OLDER IN THE OHIO COMMUNITIES WITH VINYL CHLORIDE POLYMERIZATION FACILITIES, 1958-73

Expected numbers of cancer deaths based on occurrence over the same period of time in the balance of the counties in which the communities are located.

	Males		Females		Sexes combined	
	Obs./Exp.	SMR	Obs./Exp.	SMR	Obs./Exp.	SMR
CNS cancer (191-192) ^a						
Ashtabula	7/ 6.1	115	6/ 3.8	158	13/ 9.9	131
Painesville	12/ 3.8	316 ^d	2/ 2.8	71	14/ 6.6	212 ^b
Avon Lake	2/ 2.3	87	1/ 1.8	56	3/ 4.1	73
N. Ridgeville	6/ 1.9	316 ^b	2/ 1.5	133	8/ 3.4	235 ^b
Communities Combined	27/14.1	191 ^c	11/ 9.9	111	38/24.0	158 ^c
Leukemia and aleukemia (204-207) ^a						
Ashtabula	13/14.7	88	11/ 8.6	128	24/23.3	103
Painesville	8/ 7.1	113	7/ 6.8	103	15/13.9	108
Avon Lake	1/ 3.6	28	5/ 2.2	227	6/ 5.8	103
N. Ridgeville	2/ 3.2	63	0/ 1.9	0	2/ 5.1	39
Communities Combined	24/28.6	84	23/19.5	118	47/48.1	98
Lymphomas (200-203) ^a						
Ashtabula	18/12.6	143	14/ 5.8	241 ^c	32/18.4	174 ^c
Painesville	12/10.4	115	4/ 8.3	48	16/18.7	86
Avon Lake	4/ 3.3	121	5/ 2.9	172	9/ 6.2	145
N. Ridgeville	3/ 2.8	107	1/ 2.6	38	4/ 5.4	74
Communities Combined	37/29.1	127	24/19.6	122	61/48.7	125

^a International Classification of Diseases, 8th revision codes.

^b $P < 0.05$.

^c $P < 0.01$.

^d $P < 0.002$.

$0.05 < P < 0.10$. Data were then analyzed for leukemia and aleukemia mortality. As shown in Table III, the observed versus expected mortality was virtually identical.

Birth defects among populations residing near VC polymerization facilities

The occurrence of birth defects was studied [8] for the same three Ohio communities with VC polymerization facilities. Birth data for residents of North Ridgeville were not combined with data for the three index communities in the initial analyses (Table IV) because the high incidence of birth defects in North Ridgeville, which lies contiguous to Avon Lake, was identified in subsequent analyses [8]. Data for North Ridgeville, however, were included in subsequent analyses (Table V) for specific birth malformations. The observations among community residents were compared to both the occurrence in

360 002949

TABLE IV

RESIDENT BIRTHS, MALFORMATION RATE PER 1000 LIVE BIRTHS IN OHIO AND IN THREE SELECTED COMMUNITIES AND OBSERVED VERSUS EXPECTED NUMBERS OF MALFORMATIONS IN EACH CITY, YEARS COMBINED, 1970-73

Malformations are based on codes 740-759 of the International Classification of Diseases, 8th revision, 1968.

Area	Births	Malformations			χ^2
		Rate/10 ³	Number observed	Number expected ^a	
Entire state of Ohio	719,287	10.1	7295	—	—
Ashtabula city	1900	17.4	33	19.3	9.78 ^b
Painesville city	1381	18.1	25	14.0	10.29 ^b
Avon Lake city	738	20.3	15	7.5	7.56 ^b
All three communities combined	4019	18.2	73	40.8	27.13 ^c

^a Expected numbers are based on state rate per 1000 live births.

^b $P < 0.01$.

^c $P < 0.001$.

the balance of the counties in which the communities are located and to the occurrence in the State for the period 1970-73. Between 1970-73, the rate for birth defects changed from 9.3 to 11.3 per 1000 live births for the balance of the counties, from 9.6 to 10.8 for the entire state, whereas, the rate for the three index cities combined changed from 17.0 to 22.5. Thus, the incidence of birth defects for children born in the index cities was almost twice as great as the incidence in the balance of the counties or in the entire State, and the differences appear to be increasing with time.

Table IV shows data for malformation rates in each index community versus the rate for the entire State. These differences were all highly significant. The rates ranged from 17.47 in Ashtabula to 20.33 in Avon Lake, as compared to 10.14 for the entire State. When the community experience was compared to the occurrence in the balance of the counties in which the communities were

TABLE V

OBSERVED, EXPECTED AND RELATIVE RISK FOR SPECIFIC CONGENITAL ANOMALIES IN INDEX AREAS INCLUDING N. RIDGEVILLE, 1970-73^a

Defect category	Number of defects		RR ^b
	Observed	Expected	
All defects (740-756, 758, 759) ^c	109	56.0	1.95
Central nervous system (740-749)	17	5.6	3.02
Cleft palate and lip (749)	10	6.5	1.53
Genital organs (752)	16	8.4	1.90
Clubfoot (754)	23	8.2	2.79
All other defects	43	27.2	1.58

^a Excludes skin, hair and nails (757).

^b RR, relative risk (observed/expected).

^c International Classification of Disease codes, 8th revision, are shown in parentheses.

located,
the occu
the Stat
congenit

Table
total an
was obs
the CNS
have be

Fetal m

Since
systems
chromo
exposed
germin
ascerta
born a
polym
include
selecte
matche
with w
indirec
percen
contra
("cont
respec

The
accord
fetal
respec
16.5%

As
husba
30 ye
prima
husba
(5.3%
differ
of fet
of a
in jo
thesis
paren
and i
adju

ND IN THREE MALFORMA-

use revision,

0.78 b
0.29 b
7.56 b
7.13 c

d and to the
-73, the rate
the balance
rate for the
incidence of
e as great as
ate, and the

unity versus
ficant. The
b. red to
is compared
unities were

ANOMALIES IN

located, the differences remained significant. Therefore, whether you compare the occurrence in the communities to the expected, based on the average for the State, or for the balance of the counties, the excess of children with congenital anomalies in the communities appears to be significant.

Table V shows observed versus expected numbers and the relative risk for total and selected malformations. Although an increase in most organ systems was observed, the greatest excess of severe defects included malformations of the CNS, cleft lip and palate, club foot and genital organs. These observations have been reported previously in more detail [8].

Fetal mortality among wives of workers exposed to VC

Since VC had elicited a positive mutagenic response via microbial test systems [1,6,14,24] and also had been associated with significant excesses of chromosomal aberrations in the lymphocytes of workers occupationally exposed to VC [3,4,13,23], concern was expressed that VC may induce germinal mutations. Thus, a questionnaire-interview survey was conducted to ascertain the incidence of fetal loss (defined as any product of conception not born alive) among wives of workers exposed to VC [7a]. All current VC polymerization and polyvinyl chloride (PVC) fabrication workers were included for study together with a similar number of current rubber workers selected from work areas relatively free from known toxic materials and matched as a group to the VC workers by age. No interviews were conducted with workers' wives and no data were obtained concerning maternal age, except indirectly through paternal age. Group participation rates ranged from 62-77 percent. Data for the wives of VC polymerization workers (study group) were contrasted with data for the wives of PVC fabrication and rubber workers ("controls"), who were known to have had very low or no VC exposure, respectively.

The data in Table VI show the paternal age distribution for fetal deaths according to the husband's exposure. Prior to husband's exposure, the crude fetal death rates for the control and study group were 6.9 and 10.1%, respectively. Subsequent to husband's exposure, the crude rates were 8.8 and 16.5%, respectively.

As can be seen in Table VI, the excess in fetal mortality subsequent to husband's exposure, was associated with younger-aged husbands. For husbands 30 years of age and older, the rates for the control group 17/142 (12.0%) and primary VC exposure group 9/69 (13%) were about the same; whereas, for husbands less than 30 years of age, the rates for the control group 7/131 (5.3%) and primary VC exposure group 14/70 (20.0%) were significantly different, $P < 0.001$, $\chi^2 = 10.52$, with one degree of freedom (df). The excess of fetal mortality among wives of younger-aged husbands may be a reflection of a practice of placing newly hired personnel because of little or no seniority, in jobs where occupational exposures to VC may have been worse. This hypothesis, however, needs further assessment in other working populations. Since parental age was positively correlated with fetal mortality in this population and in a previous study [25], fetal mortality rates for the study group were adjusted to the age-distribution of the control group.

680 002051

TABLE VI

PATERAL AGE DISTRIBUTION FOR FETAL DEATHS ACCORDING TO HUSBAND'S VC EXPOSURE

Paternal age group (years)	"Controls"			Primary VC Exposure		
	Pregnancies	Fetal deaths		Pregnancies	Fetal deaths	
	(N)	(N)	(%)	(N)	(N)	(%)
Prior to husband's exposure						
<20	31	2	6.5	7	0	0.0
20-24	80	4	5.0	44	2	4.5
25-29	38	4	10.5	56	7	12.5
30-34	6	1	16.7	27	5	18.5
≥35	4	0	0.0	14	1	7.1
All ages crude rate	159	11	6.9	148	15	10.1
Mean paternal age at conception (year)	23.0			26.4		
Age-adjusted ^a rate			(6.9)			(6.1)
Subsequent to husband's exposure						
>20	1	0	0.0	0	0	0.0
20-24	45	4	9.3	22	3	13.6
25-29	87	3	3.4	48	11	22.9
30-34	87	7	8.0	36	3	8.3
≥35	55	10	18.2	33	6	18.2
All ages crude rate	273	24	8.8	139	23	16.5
Mean paternal age at conception (year)	30.4			30.2		
Age-adjusted ^a rate			(8.8)			(15.8)

^a Fetal mortality rates for primary VC exposure group are direct age-adjusted to the paternal age-distribution of the pregnancies in the control group (shown in parentheses).

TABLE VII

MEAN PATERAL, NUMBER OF PREGNANCIES AND AGE-ADJUSTED FETAL DEATH RATES ACCORDING TO HUSBAND'S VC EXPOSURE

	"Controls" ^a	Primary VC exposure ^b
Prior to husband's exposure		
Number of families	95	70
Mean paternal age at conception (years)	23.0	26.4
Number of fetal deaths among wives	11	15
Number of pregnancies	159	148
Age-adjusted fetal deaths/100 preg. ^c	6.9	6.1
Subsequent to husband's exposure		
Number of families	113	62
Mean paternal age at conception (years)	30.4	30.2
Number of fetal deaths among wives	24	23
Number of pregnancies	273	139
Age-adjusted fetal deaths/100 preg. ^c	8.8	15.8 ^d

^a Rubber and PVC fabrication workers.

^b VC polymerization workers.

^c Rates age-adjusted to "control" group paternal age distribution.

^d Subsequent to husbands' exposure, the frequency of fetal deaths among wives was significantly greater in the primary VC exposure group as compared to the "controls" ($P < 0.05$) or to the frequency in the study group prior to husband's exposure ($P < 0.02$) by age-adjusted Chi-square testing.

The data wives of mal exposures, t respectively. 8.8% for th was signific [17] ($\chi^2 = 4$ indicated ch to a change after husba ($P < 0.025$, before expo whereas, af greater redu a difference be seen in study group group, whe groups wer situations n To deter might have primary VC three or fo then recall carriage ra maintained

TABLE VIII

NUMBER OF BAND'S VC DEATHS

Rates for the

Before husba
After husban

Before husba
After husban

Before husba
After husban

VC EXPO.

Fetal deaths

(N) (%)

0	0.0
2	4.5
7	12.5
5	18.5
1	7.1
13	10.1

(6.1)

0	0.0
1	13.6
1	22.9
1	8.3
1	18.2
1	16.5

(15.8)

re-distrib-

TH RES

posure b

its greater
nev in the

The data in Table VII show fetal death rates per 100 pregnancies for the wives of male workers in the control and study groups. Prior to their husband's exposures, the rates were 6.9% and 6.1% for the control and study groups, respectively. Subsequent to husband's exposure, however, the rates were 8.8% for the control group versus 15.8% for the study group. This difference was significant at the $P < 0.05$ level by Mantel-Haenszel Chi-square testing [17] ($\chi^2 = 4.81$, $df = 1$). Further, the before and after exposure comparisons indicated changes in rates from 6.9 to 8.8% for the control group as compared to a change from 6.1 to 15.8% for the study group. The rates for before and after husband's exposure in the study group were significantly different ($P < 0.025$, $\chi^2 = 5.78$, $df = 1$). For the study group, it may be noted that before exposure age-adjustment reduced the crude rate from 10.1 to 6.1%; whereas, after exposure the crude rate was reduced from 16.5 to 15.8%. The greater reduction in rates for the before exposure age-adjustment resulted from a difference in the age distribution of the husbands in the two groups. As can be seen in Table VII, prior to husband's exposure, the mean paternal age in the study group was 26.4 years as compared to only 23.0 years for the control group, whereas, subsequent to husband's exposure, the mean ages of the two groups were virtually the same, i.e., 30.4 years versus 30.2 years. In both situations mean age was a good measure of central tendency.

To determine whether women who had chronically experienced abortions might have weighted the results in favor of a higher fetal mortality rate in the primary VC exposure group, data for the pregnancies of women who had two, three or four or more spontaneous abortions were eliminated. The data were then recalculated to determine whether or not the trend of a greater miscarriage rate could be maintained. As shown in Table VIII, the trend was maintained for each analysis.

TABLE VIII

NUMBER OF PREGNANCIES AND AGE-ADJUSTED FETAL DEATH RATES ACCORDING TO HUSBAND'S VC EXPOSURE EXCLUDING PREGNANCIES IN WOMEN WITH ≥ 2 , 3 OR 4 FETAL DEATHS

Rates for the primary VC exposure group are age-adjusted to the Control group

	Controls		Primary VC exposure	
	Number of pregnancies	Fetal death rate	Number of pregnancies	Fetal death rate
≥ 2 Fetal deaths excluded				
Before husband's exposure	155	5.8%	126	1.7%
After husband's exposure	255	4.7%	111	6.2%
≥ 3 Fetal deaths excluded				
Before husband's exposure	159	6.9%	141	3.1%
After husband's exposure	265	6.8%	120	10.8%
≥ 4 Fetal deaths excluded				
Before husband's exposure	159	6.9%	142	5.8%
After husband's exposure	265	6.8%	127	11.8%

500-000053

As reported previously [7a], additional analyses suggested that the significant excess in fetal mortality after the husband's exposure would not seem to be the result of bias from interviewers nor from respondents. Because of the highly volatile nature of vinyl chloride [27], carry-home exposure to the wife would seem unlikely. Therefore, the leading possibility for the mechanism involved would seem to be gerin-cell damage in the male through direct VC exposure.

References

- 1 Bartsch, H., C. Malavielle and R. Montesano, Human, rat and mouse liver-mediated mutagenicity of vinyl chloride in *S. typhimurium* strains, *Int. J. Cancer*, 15 (1975) 429-437.
- 2 Creech, J.L., Jr. and M.N. Johnson, Angiosarcoma of liver in the manufacture of polyvinyl chloride, *J. Occup. Med.*, 16 (1974) 150-151.
- 3 Ducatman, A., K. Hirschhorn and L.J. Selikoff, Vinyl chloride exposure and human chromosome aberrations, *Mutation Res.*, 31 (1975) 163-168.
- 4 F. Cravioto, F., B. Lambert and J. Lindsten, Chromosome aberrations in workers exposed to vinyl chloride, *Lancet* 1 (1975) 459.
- 5 Gedrigk, P., R. Muller and H. Bechtelsheimer, Morphology of liver damage among polyvinyl chloride workers, *Ann. NY Acad. Sci.*, 246 (1975) 278-285.
- 6 Huberman, E., H. Bartsch and L. Sachs, Mutation induction in Chinese hamster V79 cells by two vinyl chloride metabolites, chloroethylene oxide and 2-chloroacetaldehyde, *Int. J. Cancer*, 16 (1975) 639-644.
- 7a Infante, P.F., J.K. Wagoner and A.J. McMichael, Genetic risks of vinyl chloride, *Lancet*, 1 (1976) 734-735.
- 7b Infante, P.F., J.K. Wagoner, A.J. McMichael, R.J. Waxweiler and H. Falk, Genetic risks of vinyl chloride, *Lancet*, 1 (1976) 1289-1290.
- 8 Infante, P.F., Oncogenic and mutagenic risks in communities with polyvinyl chloride production facilities, *Ann. NY Acad. Sci.*, 271 (1976) 49-57.
- 9 Kramer, C.G. and J.E. Mutchler, The correlation of clinical and environmental measurements for workers exposed to vinyl chloride, *J. Amer. Ind. Hyg. Assoc.*, 33 (1972) 19-30.
- 10 Lange, C.E., S. Juhe, G. Stein and G. Veltman, Further results in polyvinyl chloride production workers, *Ann. NY Acad. Sci.*, 246 (1975) 18-21.
- 11 Lester, D., L.A. Greenberg and W.R. Adams, Effects of single and repeated exposures of humans and rats to vinyl chloride, *J. Amer. Ind. Hyg. Assoc.*, 24 (1963) 265-275.
- 12 Lilis, R., H. Anderson, W.J. Nicholson, S. Daum, A.S. Fischheim and L.J. Selikoff, Prevalence of disease among vinyl chloride and polyvinyl chloride workers, *Ann. NY Acad. Sci.*, 246 (1975) 22-41.
- 13 Loken, E. and E. Thils-Evensen, Preliminary report on the medical examination of 288 employees at the PVC plant, Norsk Hydro a.s., Porsgrunn Fabrikk, Porsgrunn, Norway, unpublished.
- 14 Loprieno, N., R. Barale and S. Baroncelli, Evaluation of the genetic effects induced by vinyl chloride monomer (VCN) under mammalian metabolic activation: studies in vitro and in vivo, *Mutation Res.*, 40 (1976) 85-95.
- 15 Maltoni, C., Preliminary report on the carcinogenicity bioassays of vinyl chloride, U.S. Dept. Labor Informal Fact-Finding Hearing on Possible Hazards of Vinyl Chloride Manufacture and Use, Washington, D.C., February 1974.
- 16 Maltoni, C. and G. Lefemine, Carcinogenicity bioassays of vinyl chloride, Current results, *Ann. NY Acad. Sci.*, 246 (1975) 195-218.
- 17 Mantel, N. and W. Haenszel, Statistical aspects of the analysis of data from retrospective studies of disease, *J. Natl. Cancer Inst.*, 22 (1959) 719-748.
- 18 Marsteller, H.J., W.K. Lebach, R. Muller and P. Gedrigk, Unusual splenomegalic liver disease as evidenced by peritoneoscopy and guided liver biopsy among polyvinyl chloride production workers, *Ann. NY Acad. Sci.*, 246 (1975) 95-134.
- 19 McMichael, A.J., S.G. Haynes and H.A. Tyroler, Observations on the evaluation of occupational mortality data, *J. Occup. Med.*, 17 (1975) 128-131.
- 20 Miller, A., A.S. Teirstein, M. Chuang, L.J. Selikoff and R. Warshaw, Changes in pulmonary function in workers exposed to vinyl chloride and polyvinyl chloride, *Ann. NY Acad. Sci.*, 246 (1975) 12-52.
- 21 Nicholson, W.J., E.C. Hammond, H. Seidman and L.J. Selikoff, Mortality experience of a cohort of vinyl chloride-polyvinyl chloride workers, *Ann. NY Acad. Sci.*, 246 (1975) 225-230.
- 22 Patty, F.A., W.P. Yant and C.P. Waite, Acute response of guinea pigs to vapors of some new commercial organic compounds, *Public Health Reports*, 45 (1930) 1963-1971.
- 23 Purchase, I.F.H., C.R. Richardson and D. Anderson, Chromosomal and dominant lethal effects of vinyl chloride, *Lancet*, II (1975) 410-411.

- 24 Rannug, U., metabolic act
- 25 Shapiro, S., loss, Mullbani
- 26 Tabershaw, I and its polyn
- 27 United State Task III, vin
- 28 Veltman, G vinyl chlorid
- 29 Viola, P.L., ride, Cancer
- 30 Waxweiler, workers exp

the significant
them to be the
of the highly
would
ism involved
exposure.

mutagenicity of
vinyl chloride, J.
can chromosome
exposed to vinyl
polyvinyl chloride
79 cells by two
cancer, 16 (1975)
Lancet, 1 (1976)
of vinyl chloride,
ride production
measurements for
ride production
of humans and
ence of
(1975) 22-41.
88 employees at
ed.
by vinyl chloride
Mutation Res.,
U.S. Dept. Labor
and Use, Wash-
results, Ann. NY
ective studies of
r disease as evi-
n workers, Ann.
of occupational
ary function in
(1975) 42-52.
of a cohort of
some new com-
thal effects of

- 24 Rannung, U., A. Johansson, C. Ramel and C.A. Wachtmeister, The mutagenicity of vinyl chloride after metabolic activation, *Ambio*, 3 (1974) 194-197.
- 25 Shapiro, S., E.W. Jones and P.M. Densen, A life table of pregnancy terminations and correlates of fetal loss, *Millbank Quarterly*, 40 (1962) 7-15.
- 26 Tabershaw, I.R., and W.R. Gaffey, Mortality study of workers in the manufacture of vinyl chloride and its polymers, *J. Occup. Med.* 16 (1974) 509-518.
- 27 United States Environmental Protection Agency, Sampling and analysis of select toxic substances, Task III, vinyl chloride, Contract No. 68-01-2646, Jan 20, 1976.
- 28 Veltman, G., C.E. Lange, S. Juhe, G. Stem and U. Bachner, Clinical manifestations and course of vinyl chloride disease, *Ann. NY Acad. Sci.*, 246 (1975) 6-17.
- 29 Viola, P.L., A. Bigotti and A. Caputo, Oncogenic response of rat skin, lungs and bones to vinyl chloride, *Cancer Res.* 31 (1971) 516-519.
- 30 Waxweiler, R.J., W. Stringer, J.K. Wakoner, J. Jones, H. Falk and C. Carter, Neoplastic risk among workers exposed to vinyl chloride, *Ann. NY Acad. Sci.*, 271 (1976) 39-48.

200 000055

strains, *Mutation Res.*, 19 (1973)
regency of xeroderma pigmen-
ry, 238 (1972) 80-83.
plementation group in xeroderma
plementation data regarding

Mutation Research, 37 (1976) 313-316
© Elsevier/North-Holland Biomedical Press

Short communication

HIGH RATE OF CHROMOSOMAL ABERRATION IN PVC WORKERS

I. SZENTESI, É. HORNYÁK, G. UNGVÁRY, A. CZEIZEL, Z. BOGNÁR and
M. TIMAR

*Laboratory of Human Genetics, National Institute of Hygiene; Department of
Experimental Pathology, State Institute of Occupational Health, H-1966 Budapest,
Gyáli ut 2-6 (Hungary)*

(Received March 26th, 1976)

(Revision received June 15th, 1976)

(Accepted June 23rd, 1976)

Occupational diseases, such as acro-osteolysis, Raynaud's phenomenon and chronic liver lesion, were first diagnosed among PVC workers in the fifties [3, 15, 17]. Experimental studies and case reports published in the seventies demonstrated a causal relationship between exposure to PVC and angiosarcoma of the liver [2, 5, 8, 11-13, 16]. The mutagenic properties of cancerogenic compounds are of great interest, since mutagenicity can be proved earlier and by simpler methods. Besides investigations on experimental mutagenesis [1, 9, 10], the results of four studies of chromosome aberrations in 7, 11, 13 and 56 workers exposed to VC were published in 1975 [4, 6, 7, 14]. Exposure time was in the range of 9-29 and 4-28 years in two reports [4, 6] but was not known in the other two [7, 14]. The occurrence of chromosomal aberrations was invariably higher in the exposed groups than in the controls.

This paper reports on the chromosome examination of 45 PVC workers exposed to the compound VC for 0.5 to 12 years.

45 PVC workers, 44 industrial controls (workers engaged in other chemical plants, not exposed to PVC, and only indirectly exposed to other chemicals) and 49 normal controls (no occupational exposure to chemicals) were examined. The chromosome analyses were performed in parallel and simultaneously in 48-h cultures of peripheral blood. The chromosome aberrations (Table I) were blind scored by one microscopist. Table I also indicates the distribution of age and sex.

The following conclusions were drawn. The rate of numerical chromosome aberrations did not differ significantly between PVC workers and controls. The frequency of chromatid-type aberrations was higher in PVC workers than in the two control groups ($P < 0.001$). Unstable chromosome-type aberrations were also significantly higher in PVC workers ($P < 0.001$).

The exposure-time of PVC workers is shown in Fig. 1. Values above the confidence limit of the normal control group were regarded as "pathological"

VC Health Info
313

their sanitation in the production and utilization of vinyl chloride plastics, *Gig. Sanit.*, 10 (1949) 38.

16 Viola, P.L., A. Bigotti and A. Caputo, Oncogenic response of rat skin, lungs and bones to vinyl chloride, *Cancer Res.*, 31 (1971) 516.

17 Wilson, R.H., W.E. McCormick, C.F. Tatum and J.L. Creech, Occupational acroosteolysis, *J. Amer. Med. Ass.*, 201 (1967) 572.

TABLE I
OCCURRENCE OF CHROMOSOME ABERRATIONS

Group	Number of persons	Number of males	Age	S.D.	Number of mitoses		Non-modal cells		Chromatid-type aberrations		Unstable chromosome-type aberrations	
					$\bar{x}$	S.D.	Num-ber	%	Num-ber	%	Num-ber	%
I. Controls	49	34	29.1	8.44	2523		55	2.18	149	5.90	9	0.35
II. Industrial controls	44	19	43.9	10.30	2988		98	3.27	199	6.67	17	0.56
III. PVC workers	45	41	27.3	8.31	1600		52	3.25	165	10.31	25	1.56




Fig. 1. Distribution of exposure-time and number of "pathological" cases in PVC workers group. Normal: (□) male, (○) female. Pathological: (■) male, (●) female.

which means that 7 or 8 and more chromatid-type aberrations were found in 44–55 or 55–66 mitoses and 2 or more unstable chromosome-type aberrations in 15–101 mitoses. In contrast with earlier studies [6,7], our "pathological" PVC workers had been exposed to the compound for longer periods. These persons are being given a clinical check-up and thereafter it will be decided whether they can continue their present work.

References

- 1 Bartsch, H., C. Malaveille, R. Montesano and L. Tomatis, Tissue-mediated mutagenicity of vinylidene chloride and 2-chlorobutadiene in *Salmonella typhimurium*, *Nature*, 255 (1975) 641–643.
- 2 Creech, J.L. and M.N. Johnson, Angiosarcoma of the liver in the manufactures of PVC, *J. Occup. Med.*, 16 (1974) 150–151.
- 3 Dinman, B.D., W.A. Cook, W.H. Whitehouse, H.J. Magnusson and T. Ditchek, Occupational acroosteolysis. I. An epidemiological study, *Arch. Environ. Health*, 22 (1971) 61–73.
- 4 Ducatman, A., K. Hirschhorn and J. Selikoff, Vinyl chloride exposure and human chromosome aberrations, *Mutation Res.*, 31 (1975) 163–168.
- 5 Editorial, Angiosarcoma of the liver in vinyl chloride, polyvinyl chloride workers, *J. Occup. Med.*, 16 (1974) 809.
- 6 Funes-Cravioto, F., B. Lambert, J. Lindstein, L. Ehrenberg, A.T. Natarajan and S. Osterman-Golkar, Chromosome aberrations in workers exposed to vinyl chloride, *Lancet*, 22 (1975) 459.
- 7 Hansteen, I.L., L. Hillestad and E. Thils-Evenson, Chromosome studies in workers exposed to vinyl chloride, *Mutation Res.*, 38 (1976) 112 (abstract).
- 8 Lange, C.E., S. Jühe and G. Veltman, Liver angiosarcoma in two workers in a PVC-producing factory, *Dtsch. Med. Wochenschr.*, 99 (1974) 1598–1599.
- 9 Loprieno, N., A. Abbondandolo, R. Barale, S. Baroncelli, Mutagenicity of industrial compounds: vinyl chloride, styrene and their possible metabolites, *Mutation Res.*, 38 (1976) 114–115 (abstract).
- 10 Magnussen, J. and C. Ramel, Mutagenic effects of vinyl chloride in *Drosophila melanogaster*, *Mutation Res.*, 38 (1976) 115 (abstract).
- 11 Makk, L., J.L. Creech, J.G. Whelan and M.N. Johnson, Liver damage and angiosarcoma in vinyl chloride workers, *J. Amer. Med. Ass.*, 230 (1974) 64–68.
- 12 Maltoni, C. and G. Lefemine, Carcinogenicity bioassays of vinyl chloride. I. Research plan and early results, *Environ. Res.*, 7 (1974) 387–405.
- 13 Maltoni, C. and G. Lefemine, Carcinogenicity bioassays of vinyl chloride: current results, *Ann. New York Acad. Sci.*, 246 (1975) 195–218.
- 14 Purchio, J.F.H., C.R. Richardson and D. Anderson, Chromosomal and dominant lethal effects of vinyl chloride, *Lancet*, 30 (1975) 410–411.
- 15 Tikhomirova, S.U., N.P. Tikhomirova, S.U. Levina and L.A. Kozlov, Working conditions and m... s for