Dermatology 1964; 89:793-797.

44. Dugois P, Amblard P, Almard M, and Deshors G. Acne chlorique collective et accidentell d'un type nouveau. Bull Soc Fran Derm Syph 1968; 75:260-261.

45. Mocarelli P, Marocchi A, and Brambilla P. Clinical laboratory manifestations of exposure to dioxin in children. A six-year study of the effects of an environmental disaster near Seveso, Italy. JAMA 1986; 256:2687-2695.

46. Doss M, Sauer H, von Tipermann R, and Colombi AM. Development of chronic hepatic porphyria (porphyria cutanea tarda) with inherited uroporphyrinogen decarboxylase deficiency under exposure to dioxin. Int J Biochem 1984; 16:369-373.

47. Centen AHJ, Strik JJTWA, and Colombi A. Coproporphyrinuria and chronic hepatic porphyria type A found in people from Seveso (Italy) exposed to 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD). Chemical Porphyria in Man, Strik JJTWA and Koeman JH, eds. Elsevier/North-Holland Biomedical Press 1979; pp 75-81.

48. Bauer H et al. Occupational intoxication during the production of chorphenol compounds. Arch Gewerbepath Gewerbehyg 1961; 18:538-555.

49. Oliver RM. Toxic effects of 2,3,7,8 tetrachorodibenzo 1,4 dioxin in laboratory workers. Br J Indus Med 1975; 32:49-53.

50. Lathrop GD et al. An epidemiologic investigation of health effects in Air Force personnel following exposure to herbicides: baseline morbidity results. National Technical Information Service No. AD A 138 340, 1984.

51. Hoffman RE, Stehr-Green PA, Webb KB, et al. Health effects of long-term exposure to 2,3,7,8-tetrachlorodibenzo-p-dioxin. JAMA 1986; 255:2031-2038.

52. Walker AE and Martin JV. Lipid profiles in dioxin-exposed workers. Lancet 1979 1:446-447.

53. Martin JV. Lipid abnormalities in workers exposed to dioxin. Br J Indust Med 1984; 41:254-256.