

Washington Post, Sept. 13, 1976

SPECIAL TO THE WASHINGTON POST

Small Mineral Pigments Corporation factory in Beltsville—which makes pigments for coloring paint, ink, plastic and cement—complained to state authorities about what he thought was lead poisoning from dust in the plant.

Twice that year, Maryland health inspectors went to the factory to investigate. But each time, according to state records, company officials shut down potentially dusty operations and the inspectors were unable to measure for lead contamination.

This year, Clarence Wolfe, a pale and gaunt-looking worker at Mineral Pigments, which now has a different owner, also complained of symptoms of lead poisoning. "I'm always tired, I'm run down. I don't have any energy or anything," the 43-year-old worker said.

There's times out there, it's terrible," Wolfe said of the dust in an interview earlier this summer. "I've seen it like a fog around here."

100 cubic centimeters of whole blood, considerably above the normal range level most health experts consider a danger line for lead absorption. "Above 80 micrograms," explained Dr. Julian Chisholm, associate professor of pediatrics at Johns Hopkins University Medical School and a nationally recognized expert on lead disease, "the risk of serious symptoms of lead poisoning increases dramatically."

During the last few years, state and federal officials have been aware of a problem at Mineral Pigments. Yet deadlines for correcting it have been set and reset repeatedly.

Once again, authorities are investigating Mineral Pigments. The history is tangled and, to some government safety officials, troubling.

"It's not a good story," said Morton Corn, assistant secretary of Labor for the Occupational Safety and Health Administration (OSHA). He termed

See MINERAL, A1, Col. 1

Industrial Chemical PCB Is Linked to Liver Cancer

By Morton Hiltz

Washington Post Staff Writer

COLD SPRING HARBOR, N.Y., Sept. 13—A toxic industrial chemical that recently was found in the milk of 93 of 97 tested American nursing mothers was linked to liver cancer in humans today by a leading Japanese scientist.

The chemical, polychlorinated biphenyls (PCBs), is believed to be present in the fatty tissue of nearly every person in the United States, Japan and possibly other industrialized countries.

The association of PCBs with human liver cancer—which is always fatal—generally within a year—developed from an episode in which about 1,000 persons in Yusho, Japan, over a period of a few weeks in 1968, ate rice oil accidentally contaminated with a total of 0.5 to 6.0 grams of PCBs.

Varying but apparently extremely high amounts of PCBs persisted in some of the victims' bodies. Many still report fatigue, headaches, fever, coughs, numbness in the limbs, severe

skin eruptions, and digestive and menstrual disorders. Some infants born to victims had PCBs in their tissues at birth, and also had severe skin discolorations.

PCBs, extremely stable chemicals, are used in electrical equipment, paints, printing ink, adhesives and other products. They have been used widely for 45 years but were not regarded as environmental contaminants until 1968.

Dr. Titechi Hirayama, a specialist in tracing the causes of human disease at the National Cancer Center Research Institute in Tokyo, told an international conference on cancer at Cold Spring Harbor Laboratory that five of the Yusho victims died of liver cancer within five years after eating the contaminated cooking oil. This figure was out to a rate of 500 per 100,000, compared with the expected rate of 31 per 100,000.

Hirayama, in an interview later, said that the association between PCBs and liver cancer was definite.

See PCB, A6, Col. 2

Study Links PCB To Liver Cancer

PCB, From A1

two cases and possible or probable in the remaining three. The two definite cases alone translate into a rate six times higher than normal, while all five cases translate into a 16-fold higher incidence, he said.

Kirayama's presentation followed a report by pathologist James R. Allen of the University of Wisconsin, who warned that "any level" of human exposure to PCBs "may be injurious to human health."

The Yusho rates—reinforced by data on brain changes in monkeys and other adverse effects in animals cited by Allen—suggest a "high possibility" that PCBs caused the liver cancer, Kirayama said.

As for the hundreds of millions of persons who have ingested much smaller amounts of PCBs, mainly in their food, Kirayama said, "I am concerned about the implications for people everywhere. . . . There is no question the problem is serious."

"Although Kirayama previously had reported the liver cancer risk calculations in Japan, they came, as a surprise to several attending scientists who specialize in causes of environmental cancer."

Kirayama's disclosure "is one of the most worrisome sets of data that we have recently seen," said one of those scientists, Dr. Irving J. Selikoff of the Mount Sinai School of Medicine in New York City.

The Environmental Protection Agency recently reported that PCBs had been found in tests of milk of mothers who live in 10 states. EPA and Department of Health, Education and Welfare officials have set Sept. 23 for a meeting with obstetricians and pediatricians about how to advise mothers about breast feeding.

The concentration of PCBs in the mothers' milk fat was highest, 16.6 parts per million, in a Michigan woman, while it averaged 1.4 ppm in 18 Virginia women. Overall, the average was 1.7 ppm.

The temporary maximum allowed by the Food and Drug Administration is 62 ppm for infant and junior food, but 1.5 ppm for cow's milk on grocer's shelves.

In recent studies reported by pathologist Allen, female monkeys that consumed PCBs at a rate of 2.5 ppm had much higher concentrations—up to 16 ppm—in their milk fat. Exposed to this "severe toxicity," Allen said, their breast-fed infants developed skin discoloration, scum, loss of eyelashes, and swelling and congestion of the eyelids.

The infants improved when weaned from milk containing PCBs, but, Allen said, a 1976 study showed that "behavioral and learning deficits" had persisted for at least a year and may still continue.

Allen recalled that the first indication that PCBs may be linked to cancer came from a Japanese study of mice in 1972. In a study of rats reported in 1973, 98 per cent of PCB-fed rats developed apparently malignant liver growths.

Not yet knowing of Kirayama's calculations of the human liver cancer increase at Yusho, Allen based his concern that PCBs may be carcinogens of animal data, saying, "There are only suggestions that a higher incidence of cancer occurs in persons exposed to PCB."

Magruder's			
10 YEARS OF QUALITY MEATS			
WAGERS BEEF 50¢ WAGERS BEEF 50¢	OUR FAMOUS SIRLOIN STEAK 1.50		BACON CHILI 1.39
FOLLY WHITE ROUND STEAK 1.19		BONELESS ROUND STEAK 1.49	
FRESH GROUND BEEF 69¢		PORTERHOUSE GROUND BONE STEAK 2.49	
LOCAL NEARBY FARMS FRESH VEAL SALE!		COUNTRY HILL ROAST 1.99	
VEAL CUTLET 3.29	ROUND BONE CHICK 1.19	BONELESS STEAK 1.19	
ROYAL DAIRY COTTAGE CHEESE 59¢	KRAZY AMERICAN SINGLES 89¢		
SUGAR 59¢		THOMAS ENGLISH MUFFINS 89¢	
DOLE PINEAPPLES 39¢	THOMSON SEEDLESS GRAPES 39¢	EQUUS WIRE CALIFORNIA 49¢	
GRAND GRAND 50¢	VIENNA HOT 50¢	ROCKVILLE HOT 50¢	