

TABLE 2.—Increase in Individual Incapacity Rating after Physical Medicine Treatment—Continued

Incapacity of Hand Before and After Treatment, per Cent									
Patient	Group	Case	Before		After		Difference		Weeks Before Treatment
			M*	PM*	M	PM	M	PM	
53	13	516	5	40	25	40	20	0	15
54		590	0	6	8	6	8	0	3
55		663	6	6	9	6	3	0	4
..	14		..	..	..	..	..	..	..
..	15		..	..	..	..	..	..	..
56	16	126	1.5	12	2.5	10	1	- 2	3
57		320	0	5	15	2.5	15	- 2.5	6
58		439	0	0	1	0	1	0	10
59		532	0	0	8	0	8	0	2.5
60		573	5	5	20	0	15	- 5	14
61		710	20	20	30	20	10	0	12
62		736	0	9	3	3	3	- 6	4
63	16	915	0	35	50	35	50	0	12
64		1267	30	60	67	60	37	0	6
65		1644	60	60	67	60	7	0	4
66		425	0	15	1	5	1	-10	4
67		1274	4.5	9	9	5.5	4.5	- 3.5	4
68	17	84	50	70	70	70	20	0	8
69		121	1	1	2.5	1.5	1.5	+ 0.5	3.5
70		113	10	10	12	10	2	0	4.5
71		255	0	1	1	1	1	0	6.5
72		349	0	10	7.5	10	7.5	0	3
73		377	10	8	25	8	15	0	5
74		497	4.5	4.5	10	4.5	5.5	0	5.5
75		476	0	3	5	3	5	0	16
76		524	2.5	2.5	4.5	0	2	- 2.5	11
77		547	0	10	0.5	0	0.5	-10	1
78		569	0	9	3.5	2	3.5	7	5
79		588	0	25	5	3	5	-22	4
80		620	0	15	15	0	15	-15	2
81		726	0	12	3.5	2	3.5	-10	1.5
82		430	5	25	15	0	10	-25	4
83		507	40	40	60	40	20	0	19
84		1030	10	10	15	5	5	- 5	20
85		1247	4.5	9	9	5.5	4.5	- 3.5	4
86		1392	0	6	8	2	8	- 4	20
87		1542	9	9	12	5	3	- 4	5
88		1552	7	7	12	3	5	- 4	4
89		1604	5	5	7	5	2	0	8
90		1921	30	67	60	67	30	0	9
91	18	548	10	15	14	15	4	0	20
92		1265	60	60	90	60	30	0	6
93		1577	7	40	25	20	18	-20	8
94		1693	20	75	85	75	65	0	14
95		1793	70	70	75	70	5	0	3
96		702	10	60	25	60	15	0	24
97		1522	5	25	50	15	45	-10	3
98		1399	60	85	70	75	10	-10	4
99	19	98	20	20	33	5	13	-15	8.5
100		645	0	30	10	10	10	-20	4
101		897	30	30	40	40	10	+10	3
		Av.	16.7	29.0	28.6	22.8	11.9	- 6.2	6.5

* M stands for incapacity determined by the Medical Division; PM, for incapacity determined by the Department of Physical Medicine and Rehabilitation.

Twenty-six patients had no provisional disability rating by the Medical Division on admission to the Department. Yet, 24 of these patients did have incapacities, ranging from 5 to 50 per cent of the hand, as measured in the Center, before starting rehabilitation measures.

Fifty-three patients did not improve, according to the figures of the Department of Physical Medicine and Rehabilitation. Twenty-five patients showed a decrease of 10 per cent or less; 8 additional patients (33 in all), of 20 per cent or less, and 12 (45 in all), of less than 35 per cent.

Only 3 workmen showed an increase in disability rating, as confirmed by the Center; i. e., + 3 per cent, + 0.5 per cent and + 10 per cent.