

Plasmid Purification Your Lab Can Afford.



Making Manual Plasmid Mini-Preps a Thing of the Past.

High Purity

Sufficient for automated,
fluorescent and manual
sequencing

- Direct Loading of Culture
- Up to 24 Preps in 60 Minutes
- Easy Operation
- No centrifugation, no organic
solvent or extractions

Step into the future of automated

1-800-366-7469

DEFENDANT'S
EXHIBIT

-164

Harvey Mudd: Top Ranked

In her News article "Some small schools are big on manufacturing scientists" (Innovations on Campus, 4 Nov., p. 850), Anne Simon Moffat recognizes what we at Harvey Mudd College have known all along: small student-faculty ratios combined with a demanding curriculum that stresses hands-on laboratory experience provide undergraduates in the sciences with the education and tools to succeed at the graduate level.

However, Moffat's article may leave the impression that only the schools she lists produce high percentages of graduates who go on to earn Ph.D.'s. This is not the case. In a study prepared by the National Research Council and the U.S. Department of Education that reviewed the number and percentage of alumni from 1413 colleges and universities who graduated with bachelor's degrees between 1971 and 1980 and earned Ph.D.'s between 1979 and 1988, Harvey Mudd College ranked first in the nation in the percentage of graduates who went on to earn Ph.D.'s in chemistry and mathematics and second in physics and astronomy. The Higher Education Data Sharing Consortium study she cites does not include the Carnegie Classification—specialty engineering schools—to which Harvey Mudd College belongs; most schools in this category have outstanding science departments, as is the case at Harvey Mudd College.

Henry E. Riggs
President,

Harvey Mudd College,
301 East Twelfth Street,
Claremont, CA 91711-5990, USA

Chrysotile, Tremolite, and Mesothelioma

Five years ago (Letters, 24 Aug. 1990, p. 844), we discussed the possibility that the deaths of Quebec chrysotile miners and millers by mesothelioma might have been caused by contamination with amphibole fibers in the tremolite series. Clear evidence is now available to address this question. We recently reported that in our cohort of almost 11,000 men born between 1891 and 1920 (followed continuously since 1966), there had been 33 cases of mesothelioma among some 7300 deaths from all causes by the end of 1988 (1). Analysis at that stage suggested that the risk of mesothelioma was appreciably higher among miners and millers in Thetford Mines than at Asbestos, and higher still among workers in a small asbestos products factory in Asbestos. It was known

that the level of fibrous tremolite contamination was several-fold higher in Thetford than in Asbestos and that crocidolite had been used in the factory.

Since then the follow-up has continued, and by the end of 1992, 37 mesothelioma deaths had been identified among 8000 deaths from all causes. Twenty-four of the 37 were from Thetford Mines and 13 from Asbestos, including five from the factory. The 24 mesothelioma cases in Thetford have now been the subject of detailed study.

First, 10 controls were sought for each subject from men in the cohort with the same year of birth who survived. In one case, only six controls could be found and, in another, only one; these two subjects and their controls were excluded from further study. From detailed employment records, the number of years worked by subjects and controls in each of 15 mines in the Thetford region were calculated. Periods of service within 20 years of death of a subject were excluded, as in the etiology of mesothelioma such exposure would probably be unimportant. It was immediately obvious that man-years of employment of the cases was concentrated in a localized area of five mines (Area A), where the ratio of man-years for controls to cases was 5.3 (1881/354), compared with 21.6 (1275/59) in the 10 mines located peripherally (Area B)—a fourfold difference. Data for the two excluded cases and their available controls had the same pattern.

The possibility that this distribution might be related to the concentration of fibrous tremolite in the two areas was then tested with data on asbestos fiber concentrations in lung tissue from 83 cohort members from Thetford Mines who had died from causes other than mesothelioma and had been examined by electron microscopy in 1988 (2). The number of lungs examined was 58 from Area A and 25 from Area B; the groups were similar in duration of employment (36 and 37 years) and time from termination to death (8 years in both), but estimated accumulative dust exposure was about 30% higher in group B. The geometric mean concentrations of fibers equal to or greater than 5 micrometers in length per microgram of dried lung were as follows: chrysotile, Area A, 7; Area B, 13 (not significant); tremolite, Area A, 32; Area B, 7 ($P = 0.0002$).

The more than fourfold observed difference in tremolite concentration between the two areas suggests that the relatively rare cases of mesothelioma mainly or perhaps wholly resulted from amphibole fiber contamination. These findings have important implications, but it should be emphasized that they represent the result of lo

heavy exposures 20 to 70 years ago. The geological distribution of tremolite within Quebec chrysotile ore body may well vary in time and place and, at present levels of environmental control, any mesothelioma risk from exposure in either area A or B would be far below the limits of epidemiological detection.

J. Corbett McDonald
Alison D. McDonald

National Heart and Lung Institute,
London University,
Royal Brompton Hospital,
London SW3 6LY, United Kingdom

References

1. J. C. McDonald, F. D. K. Liddell, A. Dufresne, A. D. McDonald, *Br. J. Ind. Med.* 50, 1073 (1993).
2. P. S. Sébastien, J. C. McDonald, A. D. McDonald, B. Case, R. Harley, *ibid.* 46, 180 (1989).


Corrections and Clarifications

In the report "GDNF: A potent survival factor for motoneurons present in peripheral nerve and muscle" by C. E. Henderson *et al.* (11 Nov., p. 1062), the seventh author's name should have been "Laura Simmons." The affiliation for authors H. S. Phillips, M. Armanini, and Arnon Rosenthal should have been "Department of Neuroscience, Genentech,

South San Francisco, CA 94080-4990, USA." The affiliation for B. Moffet and R. A. Vandlen should have been "Department of Protein Chemistry, Genentech," and that for Laura Simmons should have been "Department of Cell Genetics, Genentech." Alun M. Davies is at the University of St. Andrews, St. Andrews, Fife, Scotland KY16 9AJ, United Kingdom, and Vassilis E. Koliatsos is in the Departments of Pathology, Neurology, and Neuroscience and the Neuropathology Laboratory, Johns Hopkins University School of Medicine, Baltimore, MD 21208, USA. Beginning on line 19 of the first column on page 1064, the text should have read: "The average volume of the surviving motoneurons in the facial nuclei that received GDNF was $5186 \pm 308 \mu^3$, close to that in the control nuclei ($6081 \pm 105 \mu^3$) ($n = 3$). In contrast, although BDNF and NT-4/5 both enhanced motoneuron survival to a similar degree as did GDNF (2, 3), neither of them prevented the axotomy-induced shrinkage of facial motoneurons. In BDNF and NT-4/5-treated nuclei, the volume of the surviving motoneurons was only $65.5 \pm 4\%$ and $63.3 \pm 2.6\%$ of the control nuclei, respectively ($n = 3$) (14)."

Throughout the Research Article "Autoproteolysis in hedgehog protein biogenesis" by John J. Lee *et al.* (2 Dec., p. 1528) the word "disc" was incorrectly spelled "disk." In figure 4C (p. 1531), lane 9 should have been labeled "flu227," not "flu408."

We'll Help You Build A New Lab



Let the free Science product information service put you in touch with the vendors whose products you will need.

Simply write us a letter stating the specifics about your proposed lab and the instruments and supplies you need. We will do the rest. Write to:

SCIENCE Magazine
New Lab Service Department
1333 H Street, N.W., Rm 830
Washington, DC 20005

SCIENCE
The Official Journal of Research

NEW! EndNote Plus 2.0

EndLink 2

EndNote 2

Bibliographies made easy

More than 75,000 researchers use EndNote to create bibliographies and organize references in word. Now version 2 makes this even easier with great new features such as term lists, new search options, and 240 journal styles! And to make it even easier, EndLink lets

you import references

from more than 100

sources without typing

any references in word.

Niles & Associates, Inc.

800 Jones Street

Berkeley, CA 94710

Voice: 510-541-3416

FAX: 510-541-3883

email: endnote@aol.com

Mail to Niles & Associates, Inc., 800 Jones St., Berkeley, CA 94710

☐ Yes! send me information about EndNote for...

☐ Mac

Version 2.0

☐ Windows

Under Development

☐ DOS

Version 1.3.3
2.0 not available

Name _____

Address _____

City/State/Zip _____