Analytical Chemistry
Special Study 71-9
Job No. 13840

# DETERMINATION OF POLYCHLORINATED BIPHENYL RESIDUES IN BEAGLE DOG TISSUES FROM A TWO YEAR ORAL CHRONIC TOXICITY STUDY

## INTRODUCTION

This report covers the results of the analysis of tissues from IBT # C7299, "Two Year Chronic Toxicity Study with Aroclor 1242, Aroclor 1254, and Aroclor 1260 in Beagle Dogs", for PCB residues.

The specific objectives of this study were to establish the relationship between tissue storage levels and exposure period, exposure level, degree of chlorination and any observed toxic effects as well as determining the homolog distribution of the retained PCB residues. Enough animals were also included in this study to determine the rate at which the retained residues washed out once oral exposure to the PCB products was ceased.

## SUMMARY

The PCB residues found in the dogs were an order of magnitude less than the PCB residues found in a similar study with rats.(1) Material balance calculations indicate that in all cases less than 1% of the total test material ingested was stored in any of the animal tissues.

Significant alterations of the homolog distributions of all products were observed. The amount of alteration was greatest for Aroclor 1242 and least for Aroclor 1260.

The estimated biological half lives of the residues retained from each product were: Aroclor 1242 - 47 days; Aroclor 1254 - 108 days; and Aroclor 1260 - 301 days. These half lives were found to be independent of both tissue type and exposure level.

## DETAILS

The samples were analyzed as outlined in Analytical Chemistry Method 71-35. The method involves isolation of the PCB's by solvent extraction, followed by chemical and alumina column clean-up of the extract, and detection of the type and level of PCB's retained in the tissues by electron capture gas chromatography (EC/GC).

Standard solutions of the Aroclor fed were used to prepare the calibration curves for quantitation in the cases of Aroclor 1254 and Aroclor 1260. For Aroclor 1242, the Aroclor 1254 standard solutions were used to prepare the calibration curves, since the principal isomers found in the tissue extracts were those that are dominant in production Aroclor 1254.

The PCB residue levels found in each of the tissue samples are shown in Tables I and II. The values in Table I were calculated using the wet weight of the tissue, those in Table II were calculated on the basis of the hexane soluble lipid weight.

After the 24 month sacrifice samples had been analyzed, the large number of samples (180 total) and hence time and money needed for analysis of the complete set of samples prompted a reevaluation. It was decided that little information would be lost by analyzing only a balanced subset of the recovery samples and using multiple regression statistics to predict the remaining points.(2) The open boxes in Tables I and II represent the samples which were not analyzed. The regression equation obtained indicated that, relative to the rat study completed earlier, there was more scatter in the dog data. This increased variability is attributable to the lower storage levels observed and to the fact that the tissue samples from the recovery sacrifices were from individual animals rather than group composite samples analyzed in the rat study. However, an acceptable regression equation was obtained by eliminating results where the difference between the predicted and actual values exceeded two standard deviations. The predicted PCB residue levels (lipid weight basis) are shown in Table III.

Figures I thru IX are semi-logarithmic plots relating average PCB residue level to exposure period for each Aroclor and for each tissue (muscle, liver, and fat) at three exposure levels (1, 10 and 100 ppm). On these plots solid lines were used for actual data while dashed lines were used for predicted data. These figures illustrate the decrease of PCB residue levels with time in those animals placed on a recovery diet. The figures also show that the recovery was much slower for those animals fed the higher chlorinated products such as Aroclor 1254 and Aroclor 1260. There is little evidence, at least under the conditions employed in this study, that exposure level or tissue differences influenced the residue washout rate.

The significantly higher PCB residue levels found in liver seem to indicate this tissue is a primary storage area and/or site of metabolism.

Chromatograms I, II and III compare the typical homolog distributions of the retained residue to that of the Aroclor fed. Isomeric alterations were observed with each Aroclor at all feed levels. Aroclor 1242 was altered to a greater extent than either Aroclor 1254 or Aroclor 1260.

The chromatographic conditions employed while suitable for quantitation of the PCB residues in the tissues are not ideal for the identification of the specific isomers which are retained. Nevertheless with these limitations, the chlorine number(s) of the principal isomer(s) are shown above each peak on the chromatogram. An inspection

of the chromatograms reveals that the less chlorinated isomers are metabolized and/or excreted rapidly. Further inspection of the chromatograms from the recovery samples shows that the ratios of the peak heights continue to change which along with the decrease in PCB residue levels, indicates that the higher chlorinated isomers can also be metabolized and/or excreted, albeit more slowly. Chromatogram IV is a comparison of the extracts from the animals after 60 days on a recovery diet free from PCB's. It is evident, that the isomers observed are the same irrespective of the material fed. This again illustrates the much lower wash out rate encountered with the higher chlorinated isomers.

## TOXICITY

No untoward behavorial reactions or significant abnormalities were observed during the two year study.(1)

## REFERENCES

1. "Two Year Chronic Oral Toxicity Study With Aroclor 1242, Aroclor 1254, and Aroclor 1260 in Beagle Dogs", Industrial Bio-Test Laboratories Report (IBT # C7299).

2. "PCB Residues in Beagle Dogs", Statistics Special Study 71-25.

db

Monsanto Industrial Chemicals Company
Applied Sciences Section
St. Louis, Missouri

12/72 - W. M. Mees, E. S. Tucker, W. J. Litschgi, J. Cowell

TABLE I

# IBT #C7299 Two-Year Chronic Oral Toxicity Study of Aroclor in Beagle Dogs

## ppm PCB in Dog Tissues (Wet Weight Basis)

| Material | Tissue | 1 ppm | | | | | | 10 ppm | | | | | | 100 ppm | | | | | | Exposure Level |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | 24 mo. | | 30 d. | | 60 d. | | 24 mo. | | 30 d. | | 60 d. | | 24 mo. | | 30 d. | | 60 d. | | Sacrifice |
| | | ♂ | ♀ | ♂ | ♀ | ♂ | ♀ | ♂ | ♀ | ♂ | ♀ | ♂ | ♀ | ♂ | ♀ | ♂ | ♀ | ♂ | ♀ | Sex |
| Aroclor 1242 | Muscle | 0.07 | 0.08 | 0.09 | 0.06 | | | 0.3 | 0.2 | | 0.2 | | 0.04 | 0.6 | 0.5 | 0.3 | 0.4 | 0.5 | | |
| | Liver | | 0.08 | 0.3 | 0.04 | | 0.04 | 0.2 | 0.1 | | 0.1 | 0.05 | | 0.6 | | 0.4 | 0.3 | | | |
| | Fat | 2 | 2 | 1 | 0.5 | 0.5 | | 1 | 1 | | 2 | 1 | 2 | 4 | 4 | 9 | 2 | | | |
| Aroclor 1254 | Muscle | 0.3 | 0.08 | 0.2 | | 0.1 | 0.1 | 0.7 | 0.3 | | 0.7 | | | 4 | 3 | | | | 3 | |
| | Liver | 0.1 | 0.2 | | 0.2 | | | 0.4 | 0.7 | | | 0.5 | | 2 | 2 | 1 | | | | |
| | Fat | 4 | 3 | 2 | | | 2 | 5 | 4 | | 18 | | | 95 | 73 | | | | 34 | |
| Aroclor 1260 | Muscle | 0.2 | 0.4 | | 1 | | 0.2 | 1 | 0.4 | | 6 | | 3 | 26 | 10 | | 34 | 26 | | |
| | Liver | 0.2 | 0.1 | 0.1 | | | 0.3 | 2 | 0.7 | 2 | 1 | 0.9 | 1 | 11 | 4 | | 2 | 9 | | |
| | Fat | 11 | 14 | | 14 | | 4 | 101 | 31 | 40 | 47 | | 62 | 93 | 221 | | 352 | 173 | | |

MONS 067522

TABLE II

# IBT #C7299 Two-Year Chronic Oral Toxicity Study of Aroclor in Beagle Dogs

## ppm PCB in Dog Tissues (Lipid Weight Basis)

| Material | Tissue | 1 ppm | | | | | | 10 ppm | | | | | | 100 ppm | | | | | | Exposure Level |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | 24 mo. | | 30 d. | | 60 d. | | 24 mo. | | 30 d. | | 60 d. | | 24 mo. | | 30 d. | | 60 d. | | Sacrifice |
| | | ♂ | ♀ | ♂ | ♀ | ♂ | ♀ | ♂ | ♀ | ♂ | ♀ | ♂ | ♀ | ♂ | ♀ | ♂ | ♀ | ♂ | ♀ | Sex |
| Aroclor 1242 | Muscle | 2 | 2 | 1 | 2 | | | 6 | 3 | | 2 | | 1 | 19 | 20 | 10 | 10 | 6 | | |
| | Liver | | 4 | 11 | 2 | | 2 | 8 | 8 | | 9 | 4 | | 29 | | 23 | 18 | | | |
| | Fat | 2 | 3 | 1 | 1 | 0.6 | | 1 | 2 | | 2 | 2 | 2 | 6 | 7 | 13 | 3 | | | |
| Aroclor 1254 | Muscle | 10 | 4 | 4 | | 2 | 2 | 18 | 18 | | 9 | | | 61 | 95 | | | | 32 | |
| | Liver | 8 | 6 | | 9 | | | 23 | 40 | | | 43 | | 110 | 130 | 111 | | | | |
| | Fat | 5 | 3 | 3 | | | 2 | 6 | 10 | | 22 | | | 128 | 132 | | | | 42 | |
| Aroclor 1260 | Muscle | 9 | 7 | | 8 | | 3 | 78 | 20 | | 66 | | 37 | 716 | 100 | | 491 | 301 | | |
| | Liver | 17 | 13 | 11 | | | 21 | 175 | 43 | 112 | 135 | 55 | 120 | 762 | 269 | | 81 | 659 | | |
| | Fat | 14 | 18 | | 20 | | 5 | 159 | 39 | 54 | 72 | | 78 | 1261 | 287 | | 439 | 233 | | |

MCNS 067523

**TABLE III**


IBT #C7299 Two-Year Chronic Oral Toxicity Study of Aroclor in Beagle Dogs


Predicted ppm PCB in Dog Tissues (Lipid Weight Basis)


| Material | Tissue | 1 ppm | | | | | | 10 ppm | | | | | | 100 ppm | | | | | | Exposure Level |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | 24 mo | | 30 d. | | 60 d. | | 24 mo. | | 30 d. | | 60 d. | | 24 mo. | | 30 d. | | 60 d. | | Sacrifice |
| | | ♂ | ♀ | ♂ | ♀ | ♂ | ♀ | ♂ | ♀ | ♂ | ♀ | ♂ | ♀ | ♂ | ♀ | ♂ | ♀ | ♂ | ♀ | Sex |
| Aroclor 1242 | Muscle | 2 | 2 | 1 | 1 | 0.7 | 0.7 | 5 | 4 | 4 | 3 | 2 | 2 | 20 | 10 | 9 | 6 | 6 | 4 | |
| | Liver | 3 | 3 | 2 | 2 | 1 | 1 | 10 | 8 | 7 | 5 | 5 | 4 | 28 | 18 | 18 | 12 | 11 | 8 | |
| | Fat | 2 | 2 | 1 | 1 | 0.7 | 0.7 | 5 | 4 | 4 | 3 | 2 | 2 | 20 | 10 | 9 | 6 | 6 | 4 | |
| Aroclor 1254 | Muscle | 5 | 5 | 4 | 4 | 3 | 3 | 24 | 20 | 20 | 16 | 16 | 13 | 121 | 80 | 100 | 65 | 80 | 54 | |
| | Liver | 10 | 10 | 8 | 8 | 6 | 6 | 47 | 38 | 39 | 31 | 33 | 26 | 238 | 156 | 195 | 128 | 157 | 103 | |
| | Fat | 5 | 5 | 4 | 4 | 3 | 3 | 24 | 20 | 20 | 16 | 16 | 13 | 121 | 80 | 100 | 65 | 80 | 54 | |
| Aroclor 1260 | Muscle | 8 | 8 | 8 | 8 | 7 | 7 | 54 | 44 | 50 | 40 | 47 | 38 | 355 | 233 | 333 | 217 | 308 | 202 | |
| | Liver | 16 | 16 | 14 | 14 | 14 | 14 | 106 | 84 | 98 | 80 | 91 | 74 | 698 | 457 | 654 | 426 | 605 | 397 | |
| | Fat | 8 | 8 | 8 | 8 | 7 | 7 | 54 | 44 | 50 | 40 | 47 | 38 | 355 | 233 | 333 | 217 | 308 | 202 | |


Regression Equation – log ppm PCB = −1.3450 − 0.0190 (day recovery) − 0.3888 (log ppm fed) + 0.2929 (if liver) + 0.0376 (%$Cl_2$) + 0.0202 (log ppm fed × %$Cl_2$) − 0.0917 (log ppm fed − if fat) + 0.0003 (day recovery × if fat)


MONS 067524



Figure I
Aroclor 1242
Muscle
Average PCB Residue Level (ppm)
500
200
100
50
20
10
5
2
100 ppm
10 ppm
1 ppm
24 Mo
30 day
60 day
Time →
MONS 067525



Figure II
Aroclor 1242
Liver
500
200
100
50
20
10
5
2
Average PCB Residue Level (ppm)
100 ppm
10 ppm
1 ppm
24 mo.
30 day
60 day
Time
MONS 067526



Figure III
Aroclor 1242
Fat
500
200
100
50
20
10
5
2
Average PCB Residue Level (ppm)
100 ppm
10 ppm
1 ppm
24 mo
30 day
60 day
Time
MONS 067527



Figure IV
Aroclor 1254
Muscle
500
200
100
50
20
10
5
2
Average PCB Residue Level (ppm)
100 ppm
10 ppm
1 ppm
24 mo.
30 day
60 day
Time
MONS 067528



Figure II
Aroclor 1254
Liver
500
200
100
50
20
10
5
2
1
Average PCB Residue Level (ppm)
100 ppm
10 ppm
1 ppm
24 mo
30 day
60 day
Time
MONS 067529


Figure VI
Aroclor 1254
Fat
500
200
100
50
20
10
5
2
Average PCB Residue Level (ppm)
100 ppm
10 ppm
1 ppm
24 mo
30 day
60 day
Time
MONS 067530



Figure III
Aroclor 1260
Muscle
100 ppm
10 ppm
1 ppm
Average PCB Residue Level (ppm)
1000
500
200
100
50
20
10
5
2
1
24 mo
30 day
60 day
Time
MCNS 067531


Figure VIII
Aroclor 1260
Liver
100 ppm
10 ppm
1 ppm
Average PCB Residue Level (ppm)
1000
500
200
100
50
20
10
5
2
24 mo
30 day
60 day
Time
MONS 067532



Figure IX
Aroclor 1260
Fat
100 ppm
10 ppm
1 ppm
Average PCB Residue Level (ppm)
1000
500
200
100
50
20
10
5
2
24 mo
30 day
60 day
Time
MONS 067533


Chromatogram I
Aroclor 1242 Std.
Aroclor 1254 Std.
Aroclor 1242 Fed
24 mo.
Aroclor 1242 Fed
30 day
Aroclor 1242 Fed
60 day
6+7
7+8
8


MONS 067534


Chromatogram II
Aroclor 1242 Std
Aroclor 1254 Std
Aroclor 1254 Fed
24 mo
Aroclor 1254 Fed
30 day
Aroclor 1254 Fed
60 day
MONS 067535

Chromatogram III





Aroclor 1260 Std





Aroclor 1260 Fed

24 mo




Aroclor 1260 Fed.


30 day




Aroclor 1260 Fed

60 day

MONS 067536


Chromatogram IV
Aroclor 1242 Fed
60 day
6+7
7+8
8
6
Aroclor 1254 Fed
60 day
Aroclor 1260 Fed
60 day
Aroclor 1260 Std
MONS 067537

Copyright, 1973

Vol. 13, No. 3, December 1973

# Analabs, Inc. Research Notes

A subsidiary of NEN New England Nuclear

## Polycyclic Aromatic Hydrocarbons

O. HUTZINGER AND S. SAFE

*Atlantic Regional Laboratory, National Research Council of Canada, Halifax, Nova Scotia*

AND

M. ZANDER

*Rutgerswerke, Castrop-Rauxel, Germany*

### INTRODUCTION

In recent years there has been increasing concern about the possible harmful effects of organic compounds in the environment. An important class of pollutants are the polycyclic aromatic hydrocarbons (PAH) which have been shown to occur in air, water and food (see Table 1) *via* automobile and engine exhausts, the burning of fossil fuels, oil spills, food processing, tobacco smoking, and from natural sources. Moreover several PAH have been shown to possess carcinogenic (1-4) and mutagenic (5, 6) properties. This review is designed as an introduction to the general literature on PAH with particular emphasis on environmental analysis of these compounds.

### NOMENCLATURE

Although several systems of nomenclature for polycyclic aromatic hydrocarbons have coexisted in the past (7, 8) the IUPAC system (9) is becoming generally accepted now.

Common names for the basic ring systems are retained (see Fig. 1 on page 2). For the construction of more highly condensed aromatic hydrocarbons the basic system is lettered (a, b, c, etc., (see Fig. 2 on page 3) and the "added" rings numbered in the fashion shown in Fig. 2.

IN THIS ISSUE

| | |
|---|---|
| Article, Polycyclic Aromatic Hydrocarbons | 1-11 |
| Polynuclear Hydrocarbon Standards | 12-13 |
| Article, The Solubilities of Twenty-one Chlorobiphenyls in Water | 14-16 |
| Article, Progress Reports on PCB Isomers | 16-19 |
| PCB Isomers, 99% Pure | 20 |
| Polyurethan Foam Plugs PCB and Pesticide Extraction | 21 |
| Perchlorination Kits for Analysis of Polychlorinated Aromatics | 21 |
| Polychlorinated Biphenyl Mixtures, Dibenzo-P-Dioxins and DiBenzoFurans | 22 |
| Pesticide Standards | 28-30 |
| Precoated TLC Plates | 30 |
| Pesticide Solvents | 31 |
| Pollution Standards, Catalog | 32 |

### REACTIONS AND REACTIVITY


The standard reference books on the chemistry of polycyclic hydrocarbons are Clar's two volumes (10) and chemical-technical aspects of diphenyl and the terphenyls are discussed in an encyclopedia article (11). The analysis of PAH has been reviewed (12).

Most polycyclic hydrocarbons undergo the typical substitution reactions which are known from the parent compound, benzene.

In addition, many of the polycyclic hydrocarbons show a tendency for addition reactions (*cf.* Fig. 3 on page 3) ("fixed double bonds").

MONS 067538

Fig. 1. Common names structure and numbering of basic ring systems.




The reactivity of polycyclic hydrocarbons towards addition reactions can be explained and predicted qualitatively by Clar's π -sextett theory (13, 14). A quantum mechanical explanation of Clar's theory has been given (15, 16).

The stability of polycyclic aromatic hydrocarbons depends on the structure as well as on the number of rings. Hexabenzocoronene (Fig. 4) for example is an extremely



MONS 067539

## TABLE 1
## Polycyclic hydrocarbons in man's environment*

| | |
|---|---|
| Benzo [a] pyrene in city air (mcg/1000 m³) | |
| Los Angeles | 5 |
| London | 68 |
| Bonn | 133 |
| Benzo [a] pyrene in food (mcg/kg) | |
| Vegetables | 10 |
| Smoked foods | 1-10 |
| Carcinogenic Polycyclic Hydrocarbons in water (mcg/m³) | |
| Groundwater | 1-10 |
| Waste water | 100,000 |
| Benzo [a] pyrene in cigarette smoke | |
| 1 mcg in the smoke of 40 cigarettes | |

*Adapted from M. Zander, "Analysis of Polycyclic Aromatics" presented at the "Third Annual Symposium on Recent Advances in the Analytical Chemistry of Pollutants," Athens, Ga., May 1973.





Fig. 2. Nomenclature for more highly condensed ring systems.





Fig. 3. Examples for addition reactions occurring with PAH.



Fig. 4. Hexabenzocoronene ring system.




Fig. 5. Heptacene ring system.

stable hydrocarbon; even at 700°C neither decomposition nor melting takes place. Heptacene (Fig. 5) on the other hand, is extremely unstable with a reactivity that surpasses many free radicals.

The photochemical stability of a number of PAH has been investigated (17); the literature on the photodecomposition of the carcinogen benzo [a] pyrene is discussed in reference (18).



MONS 067540

## OCCURRENCE OF PAH

PAH occur naturally in petroleum and are also synthesized by living organisms such as higher plants, algae and microorganisms (*cf.* 18, 19).

PAH are formed as pyrolysis products at high temperatures (20-22) and can, therefore, be found in coal tar, automobile exhaust, in emission from burning fossil fuels and tobacco smoke.

A panel, studying particulate polycyclic organic matter (POM) concluded recently that the nation-wide emmission of benzo [a] pyrene in the US amount to 500 tons from heat and power generation, 600 tons from refuse burning, 200 tons from coke production and 20 tons from motor vehicles (23).

The distribution of PAH is almost universal and numerous analyses show their presence (5, 18, 24-29) in air (particularly soot samples), water and soil. Polycyclic hydrocarbons and hetero-aromatic compounds, known to occur in coal tar and pitch are listed and described in references (30-32).

Because of the direct intake in human nutrition, much work has been carried out to develop analytical methods and determine levels of polycyclic aromatic hydrocarbons in foods (25, 33-43). Levels are highest in smoked foods (occasionally values > 10 ppm total polycyclic hydrocarbon are observed), whereas levels in unsmoked food products are generally lower than 1 ppm.

A large number of polycyclic aromatic (including benzo [a] pyrene) and heteroaromatic hydrocarbons as well as related compounds have been identified in tobacco smoke (44-46).

Polycyclic hydrocarbons are also released into the environment from general oil pollution and the ever-increasing oil spills (47, 48). Analytical methods for the identification of hydrocarbon pollutants on seas and beaches are reviewed in reference (49). The uptake and effect of oil (including the polycyclic aromatic hydrocarbons) on marine life is of increasing concern (50-52).

## ANALYSIS

The extraction, separation (clean-up) and analysis of PAH have been extensively investigated (53-56) and a review article on chromatographic procedures gives a useful comparison of the various methods (57).

As is general for the analysis of any trace component, the need is felt (97) for an extended collection of reference standards of polynuclear hydrocarbons and their alkyl and hydro derivatives. The later type of derivatives is of great importance since it is not ordinarily discovered by the conventional UV or fluorescence methods and is best identified and measured by MS. The alkyl derivatives are in many instances more carcinogenic than the parent compound.

The carcinogenic PAH benzo [a] pyrene has understandably been studied in some detail and a thorough comparison of the analytical methods used in the determination of this compound from air particulates has been reported by Sawicki (58). Analysis of benzo [a] pyrene in cigarette smoke (59), air (60), bitumen and plants (61) and water and other commodities (62) has recently been described.

*Thin layer and paper chromatography* – TLC and paper chromatography has been used successfully for the separation of polycyclic hydrocarbons (57, 63). Cellulose, acetylated cellulose and aluminum oxide (64) and silica are the preferred layers. A methanol-ether-water (4:4:1) mixture (65) has been recommended for acetylated cellulose layers. A typical chromatogram employing this system is shown in Fig. 6. (66) Paper chromatography on acetylated paper has been used for the chromatography of PAH with solvents similar to the TLC separation on acetylated cellulose powder (67).

Electron acceptor reagents can be used to visualize PAH on paper and thin layer chromatograms. The hydrocarbons can be recovered unchanged from the characteristically colored charge transfer complexes (see section on mass spectrometry). The native fluorescence or phosphorescence of PAH, however, offers a much more sensitive method of detection (see section on luminescence spectroscopy).



MONS 067541





Fig. 6. Two dimensional thin layer chromatogram. Layer: acetylated cellulose – aluminum oxide, 1:2. Solvents: System I, n-hexane, n-pentane, toluene (90:5:5). System II, methanol, ether, water 4:4:1. Compounds 1, dibenzo [a, e] pyrene; 2, dibenz [a, h] anthracene; 3, dibenz [a, c] anthracene; 4, perylene; 7, benzo [e] pyrene; 8, pyrene; 9, anthracene.

*Liquid chromatography* – The behavior of 18 polycyclic hydrocarbons on two column materials (Durapak OPN and 40% cellulose acetate) were investigated using an ultraviolet detector (68). Although neither system could resolve all components of the mixture, the method was shown to be of potential utility. A number of pulynuclear aza-heterocyclic hydrocarbons were separated by high pressure liquid chromatography using Zipax packing impregnated with silver ions (69) and a chemically bonded phase (70). Polycyclic aromatic hydrocarbons were also separated by liquid chromatography using Corasil I impregnated with the complexing reagent 2, 4, 7-trinitrofluorenone (71). A number of column packings have been compared in the separation of polynuclear hydrocarbons in engine oils (72).

*Gas chromatography* – Ultraviolet absorption and luminescence analysis have dominated the analysis of polycyclic hydrocarbons for some time, more recently gas chromatography has played an important role in the separation, detection and identification of these compounds (57, 73-77).

The use of capillary gas chromatography with its great resolution potential has markedly increased the analytical potential of this technique. Application of capillary G. C. have been reported in the analysis of polycyclic aromatic hydrocarbons and related compounds in anthracene oil (78), cigarette smoke (45, 79), air and car exhaust (80, 81), soot samples (75) as well as for quantitative determinations (82). Gouw *et al.* (83) have shown that PAH can be separated effectively on short capillary columns (10 m). A typical capillary column chromatogram is shown in Fig. 7 (see Table 2).

*Ultraviolet spectroscopy* – U.V. spectra of PAH show in the spectral range accessible by commercial instruments three electronic transitions which are differentiated by oscillator-strength polarisation and temperature dependency. The most intense bands in the U. V.-spectra of PAH are the so-called $\beta$ – or $^1$Bb-bands which show a molar extinction coefficient of *ca.* ($10^5$). The extinction coefficient for the para – or $^1$La-bands and the $\gamma$ – or $^1$Lb-bands are *ca.* $10^4$ - $10^5$ and ($10^2$ - $10^3$) respectively. For analytical purposes, the para-bands are the most suitable.

U. V. spectra are usually recorded in liquid solution at room temperature. It is sometimes advantageous, to measure the U. V. spectra in a solid matrix at low temperatures, frequently the bands are more intense and very narrow (84).



MONS 067542

## TABLE 2
### Compounds separated on capillary column

| No. | Compounds | No. | Compounds |
|---|---|---|---|
| 1 | Naphthalene | 25a | Chrysene |
| 2 | Azulene | 25b | Triphenylene |
| 3 | Diphenyl | 26 | Naphthacene/Tetracene |
| 4 | Acetylnaphthalene | 27 | 7-Methyl-benz [a] anthracene |
| 5 | Acenaphtene | 28a | Benzo [a] fluoranthene |
| 6 | Fluorene | 28b | Benzo [b] fluoranthene |
| 7 | 2-Methylfluorene | 29 | Benzo [k] fluoranthene |
| 8 | 1-Methylfluorene | 30 | 7, 12-Dimethyl-benz [a] anthracene |
| 9 | Trans-stilbene | 31 | Benzo [e] pyrene |
| 10 | Phenanthrene | 32 | Benzo [a] pyrene |
| 11 | Anthracene | 33 | Perylene |
| 12 | 2-Methylanthracene | 34a | 3-Methylcholanthrene |
| 13 | 4H-Cyclopenta [def] phenanthrene | 34b | 20-Methylcholanthrene |
| 14 | 1-Methylphenanthrene | 35 | Dibenz [a, j] anthracene |
| 15 | 9-Methylanthracene | 36a | Dibenz [a, c] anthracene |
| 16 | Fluoranthene | 36b | Dibenz [a, h] anthracene |
| 17 | Pyrene | 37 | Benzotetraphene |
| 18 | 11H-Benzo [a] fluorene | 38a | Benzo (g, h, i) perylene |
| 19a | 11H-Benzo [b] fluorene | 38b | Picene |
| 19b | 7H-Benzo [c] fluorene | 39 | Anthanthrene |
| 20 | 1-Methylpyrene | 40 | Dibenzo [a, e] pyrene |
| 21 | 3-Methylpyrene | 41 | Coronene |
| 22 | Benzo [g, h, i] fluoranthene | 42 | Dibenzo [a,e] pyrene |
| 23 | Benzo [c] phenanthrene | 43 | Dibenzo [a, i] pyrene |
| 24 | Benzo [a] anthracene | 44 | Dibenzo [a, h] pyrene |

Quantitative analysis of PAH by U. V. spectroscopy is possible after separation (*e. g.* by TLC). For the analysis in mixtures the qualitative composition must be known and a large number of key bands be mensured. Mathematical treatment of data is usually done by computer (85). A collection of U. V. spectra including a large number of polycyclic compounds is available (86).





Fig. 7. Chromatogram of a standard mixture of hydrocarbons. Glass capillary column coated with SE-52. Linear programmed temperature 100-300°, rate 1.8/min. Numbers correspond to compounds in Table 2.

MONS 067543





Fig. 8. Electronic spectra of phenanthrene. A. absorption (UV) spectrum (cyclohexane); F, fluorescence spectrum (benzene); P, phosphorescence spectrum (ethanol, isopentane, ether, 2:5:5; 77°K).

*Luminescence spectroscopy* – The fluorescence quantum yield for most PAH is relatively high to make fluorimetry a very sensitive and therefore widely used analytical method for these compounds. (For spectra of polycyclic hydrocarbons see (87). Fluorescence of individual members of a mixture can be observed when their absorption spectra differ and selective excitation wavelengths are used (88).

Fluorescence spectra are usually recorded in solution at room temperature. Spectra obtained at low temperature in a suitable matrix (n -alkanes, n -pentane to n -decane; cyclohexane) frequently give very characteristic spectra showing many bands (Shpol'skiispectra).

An interesting new technique is the so-called "quenchofluorimetry" (89) of multicomponent mixtures. This method is based on the fact that the fluorescence of some compounds is quenched more than of others if a "quencher" is added to the mixture. PAH with the fluoranthene ring system (89; quencher: nitromethane) and perylene (90; quencher; methyl iodide) for instance can be determined in complex PAH mixtures by this method.

Phosphorescence spectra which are measured in solid solution at low temperatures usually show a higher selectivity than fluorescence spectra. Fig. 8 shows the U.V.–fluorescence and phosphorescence spectrum of phenanthrene. A comparison of U. V.–fluorescence and phosphorescence spectroscopy is given in Table 3. For the analysis of PAH in the environment separation prior to analysis by U. V. or luminescence spectroscopy is necessary. The most commonly used combination is TLC fluorescence. Samples for quantitative analysis can either be eluted from the TLC plates or more elegantly determined by *in situ* measurements (91).

*Mass spectrometry*–Mass spectrometric analysis plays an important role in the identification of polycyclic aromatic hydrocarbons (92-96) and this technique is particularly useful in combination with gas chromatography. The combined GC-MS

## TABLE 3

## Comparison* of U.V. fluorescence and phosphorescence spectroscopy

| Method | Selectivity | Sensitivity | Reproducibility |
|---|---|---|---|
| U. V. | + | + | ++ |
| Fluorescence | ++ | +++ | ++ |
| Phosphorescence | +++ | +++ | ++ |

+++ excellent
++ good
+ average

*Adapted from M. Zander, "Analysis of Polycyclic Aromatics" presented at the "third Annual Symposium on Recent Advances in the Analytical Chemistry of Pollutants," Athens, Ga., May 1973

MONS 067544




Fig. 9. 70 eV mass spectrum of benzo [a] pyrene 9-cyanomethylene-2, 4, 7-trinitrofluorene complex eluted from TLC at different sample temperatures (as indicated); (a), benzo [a] pyrene (b). CNTNF.

technique (97) has been used in studies in which the identity of the observed gas chromatographic peaks has been confirmed by the characteristic mass spectra of the components.

PAH may be located on TLC plates by the characteristic color they form with electron acceptor reagents. The complexes may be transferred directly to a mass spectrometer sample tube and individual spectra of the compounds be obtained by thermal dissociation and fractional sublimation in the ion source of a mass spectrometer (98). Mass spectra of benzo [a] pyrene and the complexing agent 9-dicyanomethylene-2, 4, 7-trinitrofluorene obtained at different sample temperatures from the complex are shown in Fig. 9.

The mass spectra of most polycyclic aromatic hydrocarbons exhibit several common fragmentation pathways; *i.e.*:

1) The molecular ion is usually the most abundant ionic species and this greatly facilitates the determination of the molecular formula of the hydrocarbon.
2) In addition, the $M^{+}$-1, $M^{+}$-2 and $M^{+}$-3 species are also relatively intense.
3) Methyl substituted aromatic hydrocarbons exhibit $M^{+}$-15 ions.
4) Doubly-charged ions are observed for most of the aforementioned ionic species.

These characteristics indicate that mass spectrometry does not readily distinguish between isomeric polycycolic aromatic hydrocarbons since very few fragment ions are produced and the molecular ion tends to dominate the spectra. This characteristic can, however, be an advantage with the GC-MS technique since the separation of many isomeric hydrocarbons can readily be effected by gas chromatography and the resulting mass spectra gives confirmation of the GC assignments.

*Infrared spectra* – IR spectra are useful for identifying alkyl derivatives of naphthalene. IR spectra, do not, however, have any advantages over U.V. spectra in the identification of unsubstituted PAH, but are very useful in the structure determination of new compounds (12).

*Nuclear Magnetic Resonance Spectra*

Proton magnetic resonance spectroscopy can be used for the identification of PAH and their alkyl derivatives (12, 14). Experimental difficulties exist with the highly condensed systems because of their low solubility.

220 MHz Proton magnetic resonance spectrum of benzo [a] pyrene has recently been published (99) and C-13 NMR spectra of several simple condensed systems have been recorded (100).

## BIOLOGICAL ACTIVITY AND METABOLISM

It is now generally accepted (101) that to be carcinogenic, polycyclic aromatic hydrocarbons undergo metabolic activation. Epoxides (102, 103) may be involved and reaction with nucleic acid components (99, 103, 104) have been suggested. The



MONS 067545

molecular properties and chemical reactivity of carcinogenic hydrocarbons with respect to their biological activity has been intensively studied (for references see 99; see also 105-107).

Metabolism of polycyclic aromatic hydrocarbons (45, 108, 109) in mammals and microsomal enzyme preparations yield oxygenated derivatives. Hydroxy compounds are usually formed and epoxides (102, 109, 110) have been shown to be intermediates. The synthesis, properties and metabolism of PAH-epoxides has recently been extensively studied (111-114).

## REFERENCES


1. Badger, G. M., "The Chemical Basis of Carcinogenic Activity," C. C. Thomas Publ., Springfield, 1962.
2. Schoental, R., "Carcinogenesis by Polycyclic Aromatic Hydrocarbons and by Certain Other Carcinogens" in E. Clar, "Polycyclic Hydrocarbons," Vol. 1, Academic Press, London, 1964, p. 133.
3. Shubik, P., Proc. Nat. Acad. Sci. USA, 69, 1052 (1972).
4. Gerarde, H. W., "Toxicology and Biochemistry of Aromatic Hydrocarbons," Elsevier, Amsterdam, 1960.
5. Fishbein, L., W. G. Flamm and H. L. Falk, "Chemical Mutagens," Academic Press, New York, 1970, p. 275.
6. Miller, E. C. and J. A. Miller, in "Chemical Mutagens" (A. Hollaender, Editor) Vol. 1, Plenum Press, New York, 1971, p. 105.
7. Stelzner, R. and Kuh, Lit. Reg. Org. Chem. 3, 21 (1921).
8. Patterson, A. M., J. Amer. Chem. Soc. 47, 543 (1925).
9. IUPAC – Rules 1957.
10. Clar, E., "Polycyclic Hydrocarbons," Vol. 1 and 2, Academic Press, London (1964).
11. Poffenberger, N. and H. L. Hubbard, "Diphenyl and Terphenyls" in Encyclopedia of Chemical Technology" Vol. 7, p. 191.
12. Zander, M. "Isolierung und Identifizierung von aromatischen Kohlenwasserstoffen." in, "Methoden der organischen Chemie," Thieme, Stuttgart, in press.
13. Zander, M., Angew. Chem. 72, 513 (1960).
14. Clar, E., "The Aromatic Sextett," J. Wiley, London, 1972.
15. Polansky, O. E. and G. Derflinger, Int. J. Quantum Chem. 1, 379 (1967).
16. Zander, M., Ann. Chem. 723, 27 (1969).
17. Kuratsune, M., and T. Hirohata, in "Analysis of Carcinogenic Air Pollutants," (E. Sawicki, Editor), National Cancer Institute Monograph No. 9, Washington, 1962, p. 117.
18. Andelman, J. B. and M. J. Suess, "The Photodecomposition of 3, 4-Benzpyrene sorbed on Calcium Carbonate" in, "Organic compounds in Aquatic Environments" (S. D. Faust and J. V. Hunter, Editors), Marcel Dekker, New York, 1971, p. 439.
19. Steubing, L., Angew. Bot. 45, 1 (1971).
20. Badger, G. M., Proc. Roy. Soc. 87 (1956).
21. Badger, G. M., Symposium on the Analysis of Carcinogenic Air Pollutants, Cincinnati, Ohio, August, 1961.
22. Lang, K. F., H. Buffleb and M. Zander, Erdoel Kohle 16, 944 (1963).
23. Anon., Chem. Eng. News Sept. 18, 1972, p. 17.
24. Hettche, H. O., Schriftenr. der Landesanstalt fur Immissions – und Bodennutzungsschutz des Landes Nordrhein-Westfalen 12, 92 (1968).
25. Sawicki, E., Arch. Environ. Health, 14, 40 (1967).
26. Anon., Chem. Eng. News Sept. 18, 1972, p. 17.
    J. Air Wat. Poll. 9, 515 (1965).
27. Chalot, G., M. Castegnaro, J. L. Roche and R. Fontanges, Chromatographia, 3, 507 (1970).




MONS 067546

28. Borneff, J. Gas – und Wasserfach. 108, 1072 (1967).
30. The Coal Tar Research Association, "The Coal Tar Data Book," 1953.
31. Kruber, D., A. Raeithel and S. Grigoleit, Erdol, Kohle 8, 637 (1955).
32. Lang, K. F. and I. Eigen, Fortschr. Chem. Forsch. 8, 91 (1967).
33. Grimmer, C., Erdol Kohle 19, 578 (1966).
34. Gunther, F. A. and F. Buzzetti, Residue Rev. 9, 90 (1965).
35. Howard, J. W., R. T. Teague Jr., R. H. White and B. E. Frey, Jr., J. Assoc. Off. Anal. Chem. 49, 595 (1966).
36. Howard, J. W., R. H. White, B. E. Frey Jr. and E. W. Turicchi. J. Assoc. Off. Anal. Chem. 49, 611 (1966).
37. Borneff, J. and B. Fabian, Arch. Hyp. Bakteriol. 150, 485 (1967).
38. Malanoski, A. J., C. L. Greenfield, C. J. Barnes, J. M. Worthington and F. L. Joe, Jr., J. Assoc. Off. Anal. Chem. 51, 114 (1968).
39. Haenni, E. O., Residue Rev. 24, 41 (1968).
40. Howard, J. H. and T. Fazio, J. Agr. Food Chem. 17, 527 (1969).
41. Tilgner, D. J. and H. Daun, Residue Rev. 27, 19 (1969).
42. Grimmer, G. and A. Hildebrandt, J. Assoc. Off. Anal. Chem. 55, 631 (1972).
43. Fazio, T., R. H. White and J. W. Howard, J. Assoc. Off. Anal. Chem. 56, 68 (1973).
44. Wynder, E. L. and D. Hoffmann, "Tobacco and Tobacco Smoke," Academe Press, New York, 1967.
45. Carugno, N. and S. Rossi, J. Gas Chromatog. 5, 103 (1967).
46. Neurath, G. B., Planta Medica, 22, 267 (1972).
47. Boyland, D. B. and B. W. Tripp, Nature 230, 44 (1971).
48. Blumer, M. and J. Sass, Science, 176, 1120 (1972).
49. Adlard, E. R., J. Inst. Petroleum, 58, 63 (1972).
50. Nelson-Smith, A., Advan. Marine Biol. 8, 215 (1970).
51. Lee, R. F., R. Sauerheber and A. A. Benson, Science, 177, 344 (1972).
52. Lee, R. F., R. Sauerheber and G. H. Dobbs, Marine Biol. 17, 201 (1972)
53. Sawicki, E., Chemist-Analyst, 53, 24, 56 and 88 (1964).
54. Sawicki, E., CRC Critical Rev. Anal. Chem. 275 (1970).
55. Leithe, W., "The Analysis of Air Pollutants," Ann Arbor-Humphrey Science Publishers, London, 1970, p. 253.
56. Dubois, L., Corkery, A. and Monkman, J. L., Int. J. Air Poll. 2, 236 (1960).
57. Schaad, R. E., Chromatog. Rev. 13, 61 (1970).
58. Sawicki, E., T. W. Stanley, W. C. Elbert, J. Mecker and S. McPherson, Atmosph. Environ. 1, 131 (1967).
59. Davis, H. J., Anal. Chem. 40, 1583 (1968).
60. Stromberg, L. E. and G. Widmark, J. Chromatog. 49, 340, (1970).
61. Schamp, N. and F. van Wassenhove, J. Chromatog. 69, 421 (1972).
62. Siddiqi, I. and K. H. Wagner, Chemosphere, 83, (1972).
63. Sawicki, C. R. and E. Sawicki, "Thin Layer Chromatography in Air Pollution Research" in Progess in Thin Layer Chromatography and Related Methods, A. Niederwieser and G. Pataki (Editors) Vol. III, Ann Arbor Science Publishers, Ann Arbor, 1972, p. 233.
64. Sawicki, E., T. W. Stanley, W. C. Elbert, and J. D. Pfaff, Anal. Chem. 36, 497 (1964).
65. Wieland, Th., G. Luben, and H. Determann, Experientia 18, 432 (1962).
66. Kohler, M., Golder, H. and Schiesser, R., Z Anal. Chem. 206, 430 (1964).
67. Wieland, Th., and W. Kracht, Angew. Chem. 69, 172 (1957).
68. Ives, N. F. and L. Guiffrida, J. Ass. Offic. Anal. Chem. 55, 757 (1972).
69. Vivilecchia, R., M. Thiebaud and R. W. Frei, J. Chromatog. Sci. 10, 411 (1972).
70. Ray, S. and R. W. Frei, J. Chromatog. 71. 451 (1972).
71. Karger, B. L., M. Martin, J. Loheac and J. Guiochon, Anal. Chem. 45, 496 (1973).



MONS 067547

72. Vaughan, C. G., Wheals, B. B. and Whitehouse, M. J., J. Chromatog. 78, 203 (1973).
73. Cantuti, V., C. P. Cartoni, A. Liberti and A. G. Torri, J. Chromatog. 17, 60 (1965).
74. DeMaio, L. and M. Corn, Anal. Chem. 38, 131 (1966).
75. Chakraborty, B. B. and R. Long, Environ. Sci. Technol. 1, 829 (1967).
76. Bhatia, K., Anal. Chem. 43, 609 (1971).
77. Frycka, J., J. Chromatog. 65, 432 (1972).
78. Proksch, E., Osterr. Chemiker Ztg., 67, 105 (1966).
79. Grob, K., Chem. Ind. (London), 248 (1973).
80. Grimmer, G. and H. Bohnke, Z. Anal. Chem. 261, 310 (1972).
81. Grimmer, G., A. Hildebrandt and H. Bohnke, Erdol, Kohle 25, 442, 531 (1972).
82. Grimmer, G., Erdol Kohle, 25, 339 (1972).
83. T. H. Gouw, J. M. Whittemore and R. E. Jentoft, Anal. Chem. 42, 1394 (1970).
84. Clar, E., Spectrochim. Acta 4, 116 (1950).
85. Schmidt, H., Erdol-Kohle 19, 275 (1966).
86. Friedel, R. A. and Orchin, M., "Ultraviolet Spectra of Aromatic Compounds," Wiley, New York, (1951).
87. Berlman, I. B., "Fluorescence Spectra of Aromatic Molecules" Academic Press, New York, (1965).
88. Sawicki, E., T. R. Hauser, u. T. W. Stanley, Inter. J. Air Pollution 2, 253 (1960).
89. Sawicki, E., T. W. Stanley, u. W. C. Elbert, Talanta 11, 1433 (1964).
90. Zander, M., Z. anal. Chem. 263, 10 (1973).
91. Sawicki, E., and J. D. Pfaff, Anal. Chim. Acta 32, 521 (1965).
92. Majer, J. R., Perry, R. and Reade, M. J., J. Chrom., 48, 328 (1970).
93. Lao, R. C., Thomas, R. S., Monkman, J. L. and Pottie, R. F., Symposium on the Identification and Measurement of Environmental Pollutants, National Research Council, Ottawa, Canada, 144 (1971).
94. Perry, R., Symposium on the Identification and Measurement of Environmental Pollutants, National Research Council, Ottawa, Canada, 130 (1971).
95. Lao, R. C., Thomas, R. S. and Monkman, J. L., Intern. J. Environ. Anal. Chem., 1, 187 (1972).
96. Safe, S., and O. Hutzinger, "Mass Spectrometry of Pesticides and Pollutants" CRC press, Cleveland, in press.
97. Lao, R. C., Thomas, R. S., Oja, H. and Dubois, L., Anal. Chem. 45, 908 (1973).
98. Hutzinger, O., and S. Safe, unpublished data.
99. Cavalieri, E. and M. Calvin, Proc. Nat. Acad. Sci. USA, 68, 1251 (1971).
100. Stothers, J. B., "Carbon-13 NMR Spectroscopy," Academic Press, New York, 1972, p. 90.
101. Miller, J. A., Cancer Res. 30, 559 (1970).
102. Selkirk, J. K., E. Huberman and C. Hiedelberger, Biochem. Biophys. Res. Commun. 43, 1010 (1971).
103. Grover, P. L. and P. Sims, Biochem. J. 129, 41P, (1972).
104. Cavalieri, E. and M. Calvin, Photochem. Photobiol. 14, 641 (1971).
105. Popp, F. A., Z. Naturforsch. 27b, 731 and 850 (1972).
106. Feldmeister, A., D. Tsai and N. D. Weiner, J. Pharm. Pharmacol. 61, 1065 (1972).
107. Johnson, M.D. and Calvin, M., Nature, 241, 271 (1973).
108. Williams, R. T., "Detoxication Medianisms" Chapman & Hall, London, (1959).
109. Daly, J. W., D. M. Jerina and B. Witkop, Experientia, 28, 1129 (1972).
110. Keysell, G. R., J. Booth, P. Sims P. L. Grover and A. Hewer, Biochem. J., 129, 41P, (1972).
111. Waterfall, J. F. and P. Sims, Biochem. J. 128, 265-277 (1972).
112. Sims, P., Xenobiotica, 2, 469 (1972).
113. Sims, P., Biochem. J. 130, 27 (1972).
114. Sims, P., Biochem. J. 131, 405 (1973).



MONS 067548

# Polynuclear Hydrocarbon Standards

| Catalog # | Product | Synonym | Min. Purity | Price |
|---|---|---|---|---|
| RNH001 | Acenaphthene | | 99% | 7.00/100mg |
| RNH002 | Acridine | | 99% | 14.00/100mg |
| RNH003 | Anthanthrene | | 99% | 45.00/10mg |
| RNH004 | Anthracene | | 99% | 7.00/100mg |
| RNH005 | Azulene | | 99% | 10.00/10mg |
| RNH006 | 1, 2-Benzanthracene | Tetraphene | 99% | 7.00/10mg |
| RHN007 | Benz [a] anthracene-7, 12-Dione | Benzanthraquinone | 80% | 14.00/100mg |
| ——— | 2, 3-Benzanthracene | See Tetracene | | |
| ——— | Benzanthraquinone | See Benz [a] anthrancene-7, 12-dione | | |
| ——— | 9, 10-o-Benzeno-9, 10-dihydroanthracene | See Triptycene | | |
| RNH026 | 2, 3 Benzocarbazole 5H-Benzo [b] carbazole | 5H-Benzo [b] carbazole See 2, 3-Benzocarbazole | 98% | 7.00/100mg |
| RNH027 | 1, 2-Benzocoronene | | 95% | 14.00/10mg |
| RNH028 | 1, 2-Benzodiphenylene-sulfide | | 99% | 12.00/100mg |
| RNH029 | Benzo [b] fluoranthene | | 95% | 13.00/100mg |
| RNH030 | Benzo [g, h, i] fluoranthene | | 95% | 17.00/100mg |
| RNH006 | 1, 2-Benzofluorene | | 99% | 10.00/10mg |
| RNH031 | 2, 3-Benzofluorene | | 99% | 10.00/10mg |
| RNH032 | Benzo g,h,i, (gi), perylene | 1, 12-Benzoperylene | 99% | 10.00/10mg |
| ——— | 1, 12-Benzoperylene | See Benzo [g, h, i] Perylene | | |
| ——— | 9, 10-Benzophenanthrene | See Triphenylene | | |
| RNH033 | Benzo (a) pyrene | 3, 4-Benzopyrene | 98% | 17.00/100mg |
| ——— | 3, 4-Benzopyrene | See Benzo [a] pyrene | | |
| RNH009 | 5, 6-Benzoquinoline | Benzo [f] quinoline or $\beta$ -Naphthoquinoline | 99% | 10.00/100mg |
| | Benzo [f] quinoline | See 5, 6-Benzoquinoline | | |
| RNH010 | 1, 1'-Binaphthyl | | 95% | 10.00/100mg |
| RCS001 | Biphenyl | | 99% | 7.00/500mg |
| RNH011 | Carbazole | | 99% | 7.00/100mg |
| PRH012 | Chrysene | | 99% | 14.00/100mg |
| RNH013 | Coronene | | 98% | 18.00/10mg |
| RNH016 | 1, 2, 3, 4-Dibenzanthracene | Dibenz [a, c] Anthracene | 99% | 10.00/10mg |
| RNH017 | 1, 2, 5, 6-Dibenzanthracene | Dibenz [a, h] Anthracene | 99% | 7.00/10mg |
| RNH018 | 1, 2, 7, 8-Dibenzanthracene | Dibenz [a, j] anthracene | 95% | 12.00/10mg |
| ——— | Dibenz [a, c] anthracene | See 1, 2, 3, 4-Dibenzanthracene | | |
| ——— | Dibenz [a, h] anthracene | See 1, 2, 5, 6-Dibenzanthracene | | |
| ——— | Dibenz [a, j] anthracene | See 1, 2, 7, 8-Dibenzanthracene | | |
| RNH019 | 2, 3, 5, 6-Dibenzfluoranthene | | 98% | 14.00/10mg |
| RNH020 | 3, 4, 5, 6-Dibenzocarbazole | | 95% | 8.00/10mg |
| RCS067 | Dibenzo-p-dioxin | | 99% | 15.00/50mg |
| RCS070 | Dibenzofuran | | 99% | 7.00/500mg |
| RNH021 | 1, 2, 4, 5-Dibenzopyrene | | 98% | 17.00/100mg |
| RNH014 | Dibenzothiophene | Diphenylenesulfide | 99% | 7.00/100mg |
| RNH015 | 1, 2, 8, 9-Dibenzpentacene | | 99% | 14.00/10mg |
| RNH022 | 9, 10-Dihydroanthracene | | 99% | 17.00/100mg |
| ——— | 10, 15-Dihydro-5H-Diindeno [1, 2a-1', 2'-c] Fluorene | See Truxene | | |

MONS 067549

# Polynuclear Hydrocarbon Standards

| Catalog # | Product | Synonym | Min. Purity | Price |
|---|---|---|---|---|
| RNH023 | Diindeno [1, 2, 3-cd-1', 2', 3'-lm] Perylene | Periflanthene | 98% | 9.00/100mg |
| RNH024 | 3, 6-Dimethylphenanthrene | | 99% | 7.00/10mg |
| RNH025 | 9, 10-Diphenylanthracene | | 99% | 7.00/100mg |
| ——— | Diphenylenesulfide | See Dibenzothiophene | | |
| RNH034 | Dodecahydrotriphenylene | | 99% | 7.00/100mg |
| RNH034 | Durene | 1, 2, 4, 5-Tetramethylbenzene | 99.9% | 7.00/100mg |
| RNH036 | Fluoranthene | | 99% | 7.00/100mg |
| RNH037 | Fluorene | | 99% | 14.00/100mg |
| RNH038 | Hexamethylbenzene | | 99% | 7.00/100mg |
| | Indeno [1, 2, 3-cd] pyrene | See o-Phenylenepyrene | | |
| RNH039 | 2-Methylanthracene | | 99% | 8.00/100mg |
| RNH040 | 9-Methylanthracene | | 99.99% | 14.00/100mg |
| RNH041 | 4-Methyl Biphenyl | 4-Phenyltoluene | 98% | 10.00/100mg |
| RNH042 | 4, 5-Methylenephenanthrene | | 99% | 19.00/100mg |
| RNH043 | 2-Methylfluorene | | 95% | 7.00/100mg |
| RNH044 | 1-Methylphenanthrene | | 99% | 7.00/100mg |
| RNH045 | 2-Methyl phenanthrene | | 99% | 7.00/100mg |
| RNH046 | 3-Methyl phenanthrene | | 95% | 7.00/100mg |
| ——— | Naphthacene | See Tetracene | | |
| RCS072 | Naphthalene | | 99% | 7.00/500mg |
| | $\beta$ -Naphthoquinoline | See 5, 6-Benzoquinoline | | |
| RNH047 | Octadecahydrochrysene | | 95% | 14.00/100mg |
| RNH048 | Pentacene | | 98% | 18.00/10mg |
| | Periflanthene | See Diindeno [1, 2, 3-cd-1', 2', 3'-lm] perylene | | |
| RNH049 | Perylene | | 99% | 14.00/100mg |
| RNH050 | Phenanthrene | | 99% | 7.00/100mg |
| RNH051 | Phenanthrenequinone | | 95% | 14.00/100mg |
| RNH052 | 9-Phenylanthracene | | 98% | 14.00/100mg. |
| RNH053 | o-Phenylenepyrene | Indeno [1, 2, 3-cd] pyrene | 99% | 7.00/10mg |
| RNH054 | 2-Phenylnaphthalene | | 99% | 19.00/100mg |
| RNH055 | 4-Phenyltoluene | 4-Methyl biphenyl | 99% | |
| RNH056 | Picene | | | 12.00/100mg |
| RNH057 | Pyrene | | 99% | 7.00/100mg |
| RNH058 | p-Quaterphenyl | | 98% | 14.00/10mg |
| RNH059 | m-Quinquephenyl | | 95% | 32.00/10mg |
| RCS059 | o-Terphenyl | | 99% | 7.00/100mg |
| RCS060 | m-Terpenyl | | 99% | 7.00/100mg |
| RCS061 | p-Terphenyl | | 99% | 7.00/100mg |
| RNH060 | Tetracene | Naphtacene | 99% | 14.00/10mg |
| RNH061 | Tetradecahydrophenanthrene | | 95% | 18.00/100mg |
| RNH062 | 1, 2, 3, 4-Tetrahydrofluoranthene | | 90% | 18.00/100mg |
| ——— | 1, 2, 4, 5-Tetramethylbenzene | See Durene | | |
| ——— | Tetraphene | See 1, 2-Benzanthracene | | |
| RNH063 | 1, 2, 3, 4-Tetraphenylnaphthalene | | 97% | 7.00/100mg |
| RNH064 | Triphenylene | 9, 10-Benzophenanthrene | 99% | 10.00/10mg |
| RNH065 | Triptycene | 9, 10-o-Benzeno-9, 10-dihydroanthracene | 98% | 18.00/10mg |
| RNH066 | Truxene | 10, 15-Dihydro-5H-diindeno [1, 2-a 1', 2',-c] fluorene | 95% | 14.00/100mg |



MONS 067550

# THE SOLUBILITIES OF TWENTY-ONE CHLOROBIPHENYLS IN WATER

P. R. Wallnofer, M. Koniger
Bayerische Landesanstalt fur Dodenkultur and Pflanzenbau, Munchen, Germany

and O. Hutzinger
Atlantic Regional Laboratory, National Research Council of Canada, Halifax, Nova Scotia, Canada

Commercial PCB preparations consist of complicated mixtures of chlorobiphenyls with different chlorine content and their isomers. The composition of these technical products has only recently been established [1-5].

A knowledge of the solubilities of PCB components is necessary for experiments involving individual isomers and may also be useful for the evaluation of PCB distribution in natural waters.

Considerable differences in the solubilities of chlorobiphenyls with different numbers of chlorination as well as positional isomers were indicated by change in the GC peak pattern of Aroclor 1254 re-extracted from aqueous solutions with hexane[1] Solubilities of five mono-and dichlorobiphenyls also vary widely[6].

We wish to present the results of our study on the solubilities of twenty-one chlorobiphenyls in water. Most of the compounds investigated are major components of commercial PCB preparations. Since water soluble humic substances were recently reported to act as carriers for lipophylic compounds including pollutants[9, 10] the solubility of 4, 4'-dichlorobiphenyl in aqueous extracts of humic acid was also measured. The solubility of this compound was also determined in the presence of different concentrations of Tween 80. Aroclor preparations have an increased solubility in water in the presence of solubilizers[11] and the presence of the biologically relatively inert Tween 80 (e.g. 2% Tween 80 in nutrient solution showed no measurable growth inhibition in a number of microorganisms [12]) may allow the use of higher concentrations of chlorobiphenyl in many experiments.

## EXPERIMENTAL

*Instrumentation.* Gas chromatography (GC) was carried out on a Varian 1800 instrument using a glass column (5 ft. x 2 mm i.d.) packed with 1.5% XE-60 on Anakrom Q operated at temperatures of 160-100°C depending on the compound. Carrier gas was nitrogen at a flow rate of 30 ml/min. Injector and electron capture detector were kept at 220°C.

*Chemicals.* Chlorobiphenyls (99% pure) were either Syntherized or obtained from Analabs Inc. (North Haven, Conn., U.S.A.). Humic acid was purchased from Aldrich Chem. Co. (Milwaukee, Wisc., U.S.A.).

*Methods.* An example for the determination of the solubility of one chlorobiphenyl in water is given below.

A hexane solution of the chlorobiphenyl (1 mg) was pipetted into a 300 ml Erlenmeyer flask and the solvent evaporated to give a thin film of the compound. Water (100 ml) was added, the flask sealed with parafilm and the mixture shaken at 30°C for 10 days. After this time the aqueous suspension was allowed to equilibrate at 22-24°C for 3 hr. Contact of aqueous chlorobiphenyl solution with glass was kept to a minimum after this step because of the known adsorption effects[13]. An aliquot (10 ml) was drawn using a syringe carrying a small filter compartment and immediately



MONS 067551

extracted with hexane (2 x 10 ml). The hexane extract was concentrated to 10 ml and 2 µl samples were injected into the gas chromatograph. Quantitative evaluation was carried out by measuring the peak height using calibration curve of known amounts of chlorobiphenyl standard in hexane. Sampling and analyses were carried out in duplicate.

## RESULTS AND DISCUSSION

In view of the problems with adsorption and low solubility, the experiments were not designed to give solubility data as absolute physical constants but were intended to yield comparable values of reasonable accuracy for biological experiments.

The data for twenty-one chlorobiphenyls in water are shown in Table 1.

TABLE 1

Solubility of chlorobiphenyls in water

| Compound | Solubility (mg) | (ppm) |
|---|---|---|
| **Monochlorobiphenyls** | | |
| 2- | 6.0 | 5.8 |
| 3- | 3.35 | 3.6 |
| 4- | 1.20 | 1.17 |
| **Dichlorobiphenyls** | | |
| 2, 4- | 1.40 | 1.40 |
| 2, 2'- | 1.56 | 1.44 |
| 2, 4'- | 1.90 | 1.85 |
| 4, 4'- | 0.08 | 0.07 |
| **Trichlorobiphenyls** | | |
| 2, 4, 4'- | 0.085 | 0.085 |
| 3, 4, 2'- | 0.075 | 0.080 |
| **Tetrachlorobiphenyls** | | |
| 2, 2', 5, 5'- | 0.046 | 0.046 |
| 2, 3, 2', 3'- | 0.037 | 0.030 |
| 2, 3, 2', 5'- | 0.170 | 0.170 |
| 2, 4, 2', 4'- | 0.070 | 0.065 |
| 2, 4, 3', 4'- | 0.055 | 0.060 |
| 2, 5, 3', 4'- | 0.041 | 0.041 |
| 3, 4, 3', 4'- | 0.175 | 0.175 |
| **Pentachlorobiphenyls** | | |
| 2, 3, 4, 2', 5'- | 0.021 | 0.023 |
| 2, 4, 5, 2', 5'- | 0.035 | 0.028 |
| **Hexachlorobiphenyl** | | |
| 2, 2', 4, 4', 5, 5'- | 0.0088 | 0.0088 |
| **Octachlorobiphenyl** | | |
| 2, 3, 4, 5, 2', 3', 4', 5'- | 0.0068 | 0.0072 |
| **Decachlorobiphenyl** | 0.015 | 0.015 |
| 4, 4'-Dichlorobiphenyl | | |
| + Tween 10 0.1% | 6.2 | 5.5 |
| + Tween 80 1% | 10.0 | 10.0 |
| + Humic acid extract | 0.075 | 0.06 |



MONS 067552

## ACKNOWLEDGMENTS

We thank Dr. Freed for supplying data prior to publication. P.R.W. expresses his gratitude to the "Deutsche Forschungsgeminschaft" for financial support. Part of this work was carried out while O. H. was on leave at the Bayerische Landesanstalt. This author would like to thank the "Deutsche Akademische Austauschdients" for the invitation to Germany and financial aid.

## REFERENCES

1. D. Sissons and D. Welti, J. Chromatogr. 60, 15 (1971).
2. R. C. Webb and A. C. McCall, J. Ass. Offic. Anal. Chem. 55, 746 (1972).
3. S. Salki, A. Tsutsui, K. Oguri, H. Yoshimura and M. Hamana, Fukuoka Igaku Zasshi 62, 21 (1971): Chem. Abstr. 74, 146294 (1971).
4. A. C. Tas and R. H. deVos, Environ. Sci. Technol. 5, 1216 (1971).
5. A. C. Tas and R. J. C. Kleinpool, Bull. Environ. Contam. Toxicol. 8, 32 (1972).
6. D. E. Willis and R. F. Addison, J. Fish. Res. Bd. Canada 29, 592 (1972).
7. V. H. Freed, personal communication (Some physical chemical properties of the polychlorinated biphenyls-Aroclor 1254; manuscript).
8. R. C. Webb, PCB Newsletter (1970).
9. G. Ogner and M. Schnitzer, Science 170, 317 (1970).
10. K. Matsuda and M. Schnitzer, Bull. Environ. Contam. Toxicol. 6, 200 (1971).
11. V. Zitko, Fish. Res. Bd. Canada, Manuscript Report Series No. 1083 (1970).
12. H. J. Rehm, S. Laufer-Heydenreich and P. Wallnofer, Z. Lebensm. Unters. Forchung 135, 117 (1966).
13. M. L. Hattula, Dansk. Tidskr. Farm. 45, 259 (1971).

# PROGRESS REPORTS ON PCB ISOMERS

By: Michael Bolgar

Given the toxicity and persistence of the widely dispersed polychlorinated biphenyls (PCB's) it seems logical that the U.S. Interdepartmental Taskforce on PCB's in their 181-page May 1972 Report (Com-$7_2$-10419) would include as part of their recommendation (page 18) the following additional studies, (italics are ours):

1. *Toxicological evaluation of a select number of representative, purified PCB isomers as well as purified trace contaminants such as the chlorinated dibenzofurans.*

2. Definitive mammalian elaboration of the kinetics, absorption, distribution, metabolism, and excretion of the technical PCBs as well as *a number of key isomers and the chlorinated dibenzofurans.*

3. Elaboration of the subcellular and intracellular actions of the technical PCBs as well as *a number of representative isomers and chlorinated dibenzofurans.*

4. More definitive epidemiological studies of the PCBs with both more representative population sampling and standardization on the basis of lipid content of the tissues.

In our list of biphenyls and 50 isomers of the 209 possible chlorobiphenyls we included a great number of the major components present in the commercial mixtures. For



MONS 067553

instance, Willis and Addison[1] report that the major components of Aroclor 1221 accounting for 92.3 weight per cent are biphenyl, 2-Cl, 4-Cl, 2, 2'-$Cl_2$, 2, 4'-$Cl_2$ (all available from Analabs stock). From Sissons and Welti's[2] list of the major components of the more important Aroclor 1254 (we list more than half). Methods have been developed[3] to recover PCB's present in waste and river water and to increase detectability by perchlorinating the samples sensitizing them to electron-capture detection by gas chromatography[4].

We have made the tools available (polyurethane foam plugs and perchlorination kits).

Assembled below is data of melting points taken of our pure isomers and taken from the literature. This list reveals interesting features of configuration/physical property corelation.

## MELTING POINTS OF POLYCHLORINATED BIPHENYLS

| *Chlorine Position* | *Melting Points* |
|---|---|
| *Mono* | |
| 2-Cl | 32.0-32.5°C |
| 3-Cl | Oil |
| 4-Cl | 75.5-77.0°C |
| *Di* | |
| 2, 3-$Cl_2$ | Oil (Lit. 27.7-28.2°C) |
| 2, 4-$Cl_2$ | Oil (Lit. 24-25°C) |
| 2, 5-$Cl_2$ | Oil (Lit. 22-23°C) |
| 2, 6-$Cl_2$ | 34-35°C |
| 3, 4-$Cl_2$ | 47-48°C |
| 3, 5-$Cl_2$ | Oil (Lit. 31-32°C) |
| 2, 2'-$Cl_2$ | 60-61°C |
| 3, 3'-$Cl_2$ | 25-26°C |
| 4, 4'-$Cl_2$ | 148-149°C |
| 2, 3'-$Cl_2$ | Lit. Oil |
| 2, 4'-$Cl_2$ | 42.5-43.5°C |
| 3, 4'-$Cl_2$ | Lit. Oil |
| *Tri* | |
| 2, 3, 4-$Cl_3$ | 101-102°C |
| 2, 3, 5-$Cl_3$ | Lit. 41°C |
| 2, 3, 6-$Cl_3$ | 58-60°C |
| 2, 4, 5-$Cl_3$ | 76-78°C |
| 2, 4, 6-$Cl_3$ | 62.0-63.5°C |
| 2, 3, 2'-$Cl_3$ | Lit. 28-29°C |
| 2, 3, 4'-$Cl_3$ | Lit. 73°C |
| 2, 4, 4'-$Cl_3$ | 57.0-57.5°C |
| 2, 5, 2'$Cl_3$ | 43-44°C |
| 2, 5, 3'-$Cl_3$ | 40.0-40.5°C |
| 2, 5, 4'-$Cl_3$ | 63.5-64.5°C |
| 3, 4, 2'-$Cl_3$ | 60.0-60.5°C |
| 3, 4, 4'-$Cl_3$ | Lit. 87-88°C |
| 3, 5, 2'-$Cl_3$ | Lit. 58°C |
| 3, 5, 4'-$Cl_3$ | Lit. 88°C |



MONS 067554

# MELTING POINTS OF POLYCHLORINATED BIPHENYLS

| *Chlorine Position* | *Melting Points* |
|---|---|
| ***Tetra*** | |
| 2, 3, 4, 5-$Cl_4$ | 91.5-92.5°C |
| 2, 3, 5, 6-$Cl_4$ | 77.0-77.5°C |
| 2, 3, 4, 4'-$Cl_4$ | Lit. 141-142°C |
| 2, 4, 5, 4'-$Cl_4$ | Lit. 125°C |
| 2, 3, 2' 3'-$Cl_4$ | 119.5-121.5°C |
| 2, 3, 2' 5'-$Cl_4$ | 46.5-47.0°C |
| 2, 4, 2' 4'-$Cl_4$ | 41-42°C |
| 2, 4, 2' 5'-$Cl_4$ | 64-66°C |
| 2, 4, 3' 4'-$Cl_4$ | 127-128°C |
| 2, 5, 2' 5'-$Cl_4$ | 86-87°C |
| 2, 5, 3' 4'-$Cl_4$ | 104-107°C |
| 2, 6, 2', 6'-$Cl_4$ | 190-200°C (sublimes) |
| 3, 4, 3', 4'-$Cl_4$ | 182-184°C |
| 3, 5, 3', 5'-$Cl_4$ | Lit. 164°C |
| ***Penta*** | |
| 2, 3, 4, 5, 6,-$Cl_5$ | 123-125°C |
| 2, 3, 4, 5, 4'-$Cl_5$ | Lit. 98-99°C |
| 2, 3, 4, 2', 5'-$Cl_5$ | 111.5-113.0°C |
| 2, 3, 4, 3' 4'-$Cl_5$ | Lit. 101-105°C |
| 2, 3, 6, 2' 5'-$Cl_5$ | 97-98°C |
| 2, 4, 5, 2' 3'-$Cl_5$ | 81-82°C |
| 2, 4, 5, 2' 5'-$Cl_5$ | 76.5-77.5°C |
| 2, 4, 5, 3' 4'-$Cl_5$ | Lit. 105-107°C |
| ***Hexa*** | |
| 2, 3, 4, 5, 6, 2'-$Cl_6$ | Lit. 134-137°C |
| 2, 3, 4, 5, 6, 3'-$Cl_6$ | Lit. 97-100°C |
| 2, 3, 4, 5, 6, 4'-$Cl_6$ | Lit. 160-165°C |
| 2, 3, 4, 5, 2' 4'-$Cl_6$ | Lit. 77-78°C |
| 2, 3, 4, 5, 2' 5'-$Cl_6$ | 89-91°C |
| 2, 3, 5, 6, 2' 5'-$Cl_6$ | 100-101°C |
| 2, 3, 4, 2' 3' 4'-$Cl_6$ | 155-8°C |
| 2, 3, 4, 2' 4' 5'-$Cl_6$ | 78.5-80.0°C |
| 2, 3, 5, 2' 3' 5'-$Cl_6$ | Lit. 128-129°C |
| 2, 3, 6, 2' 3' 6'-$Cl_6$ | 115-117°C |
| 2, 3, 6, 2' 4' 5'-$Cl_6$ | Lit. Oil |
| 2, 4, 5, 2' 4' 5'-$Cl_6$ | 102-104°C |
| 2, 4, 5, 2' 4' 6'-$Cl_6$ | Lit. Oil |
| 2, 4, 6, 2' 4' 6'-$Cl_6$ | 113.5-114.5°C |
| 3, 4, 5, 3', 4' 5'-$Cl_6$ | Lit. 198 |

MONS 067555

# MELTING POINTS OF POLYCHLORINATED BIPHENYLS

| *Chlorine Position* | *Melting Points* |
|---|---|
| *Hepta* | |
| 2, 3, 4, 5, 6, 2' 5'-$Cl_7$ | 147-150°C |
| 2, 3, 4, 5, 6, 3', 4'-$Cl_7$ | Lit. 116-118°C |
| 2, 3, 4, 5, 2', 3', 4'-$Cl_7$ | Lit. 134.5-135.5°C |
| 2, 3, 4, 5, 2' 3' 6'-$Cl_7$ | Lit. 130.5-130.7°C |
| 2, 3, 4, 5, 2' 4' 5'-$Cl_7$ | Lit. 109-110°C |
| 2, 3, 4, 5, 2' 4' 6'-$Cl_7$ | Lit. 152-153°C |
| 2, 3, 4, 5, 3' 4' 5'-$Cl_7$ | Lit. 162-163°C |
| *Octa* | |
| 2, 3, 4, 5, 2' 3' 4' 5'-$Cl_8$ | 158-159°C |
| 2, 3, 4, 6, 2' 3' 4' 6'-$Cl_8$ | Lit. 132°C |
| 2, 3, 5, 6, 2' 3' 5' 6'-$Cl_8$ | 161-162°C |
| *Nona* | |
| 2, 3, 4, 5, 6, 2' 3' 4' 5'-$Cl_9$ | 204.5-206.5°C |
| *Deca* | |
| 2, 3, 4, 5, 6, 2' 3' 4' 5' 6'-$Cl_{10}$ | 300-310°C (sublimes) |

All melting Points taken on a Fischer-Johns melting point stage.

## REFERENCES


1. D. E. Willis and R. F. Addison, J. Fish. Res. Board Canada, 29, No. 4. 592-595 (1972)
2. D. Sessons and D. Welti, J. Chromatography, 60, 15-32 (1971)
3. H. D. Gesser, A. Chow, F. C. Davis, J. F. Uthe and J. Reinke, Anal. Lett., 4, 883-6 (1971)
4. V. Zitko, O. Hutzinger and S. Safe, Bull. Environ. Contam. Toxicol. 6, 160 (1971)




MONS 067556

# 99% Pure PCB Isomers

For clarity we have kept the position numbering separate for each ring. For Chemical Abstracts nomenclature transpose numbers listing lowest number first.

| | | |
|---|---|---|
| RCS-001 | Biphenyl | $7.00/500mg |
| RCS-002 | 2-Chlorobiphenyl | 7.00/100mg |
| RCS-003 | 3-Chlorobiphenyl | 7.00/100mg |
| RCS-004 | 4-Chlorobiphenyl | 7.00/100mg |
| RCS-005 | 2, 2'-Dichlorobiphenyl | 18.00/100mg |
| RCS-006 | 3, 3'-Dichlorobiphenyl | 18.00/100mg |
| RCS-007 | 4, 4'-Dichlorobiphenyl | 18.00/100mg |
| RCS-008 | 2, 4'-Dichlorobiphenyl | 32.00/50mg |
| RCS-009 | 2, 3-Dichlorobiphenyl | 14.00/100mg |
| RCS-010 | 2, 4-Dichlorobiphenyl | 14.00/100mg |
| RCS-011 | 2, 5-Dichlorobiphenyl | 14.00/100mg |
| RCS-012 | 2, 6-Dichlorobiphenyl | 14.00/100mg |
| RCS-013 | 3, 4-Dichlorobiphenyl | 14.00/100mg |
| RCS-014 | 3, 5-Dichlorobiphenyl | 14.00/100mg |
| RCS-015 | 2, 4, 4'-Trichlorobiphenyl | 26.00/20mg |
| RCS-052 | 2, 4, 3', 4'-Tetrachlorobiphenyl | 32.00/20mg |
| RCS-016 | 2, 5, 2'-Trichlorobiphenyl | 28.00/50mg |
| RCS-017 | 2, 5, 3'-Trichlorobiphenyl | 28.00/50mg |
| RCS-018 | 2, 5, 4'-Trichlorobiphenyl | 28.00/50mg |
| RCS-019 | 3, 4, 2'-Trichlorobiphenyl | 28.00/50mg |
| RCS-020 | 2, 3, 4-Trichlorobiphenyl | 26.00/50mg |
| RCS-021 | 2, 3, 6-Trichlorobiphenyl | 30.00/50mg |
| RCS-022 | 2, 4, 5-Trichlorobiphenyl | 22.00/100mg |
| RCS-023 | 2, 4, 6-Trichlorobiphenyl | 26.00/100mg |
| RCS-024 | 2, 3, 2', 3'-Tetrachlorobiphenyl | 28.00/100mg |
| RCS-025 | 2, 4, 2', 4'-Tetrachlorobiphenyl | 28.00/100mg |
| RCS-026 | 2, 5, 2', 5'-Tetrachlorobiphenyl | 28.00/100mg |
| RCS-027 | 2, 6, 2', 6'-Tetrachlorobiphenyl | 28.00/100mg |
| RCS-028 | 3, 4, 3', 4'-Tetrachlorobiphenyl | 28.00/100mg |
| RCS-029 | 2, 3, 2', 5'-Tetrachlorobiphenyl | 35.00/50mg |
| RCS-030 | 2, 4, 2', 5'-Tetrachlorobiphemyl | 22.00/20mg |
| RCS-031 | 2, 5, 3', 4'-Tetrachlorobiphenyl | 35.00/50mg |
| RCS-032 | 2, 3, 4, 5,-Tetrachlorobiphenyl | 20.00/50mg |
| RCS-033 | 2, 3, 5, 6-Tetrachlorobiphenyl | 20.00/50mg |
| RCS-034 | 2, 3, 4, 2', 5'-Pentachlorobiphenyl | 25.00/20mg |
| RCS-035 | 2, 3, 6, 2', 5'-Pentachlorobiphenyl | 22.00/10mg |
| RCS-036 | 2, 4, 5, 2', 3'-Pentachlorobiphenyl | 22.00/10mg |
| RCS-038 | 2, 4, 5, 2', 5'-Pentachlorobiphenyl | 35.00/50mg |
| RCS-039 | 2, 3, 4, 5, 6-Pentachlorobiphenyl | 28.00/100mg |
| RCS-040 | 2, 3, 4, 2', 3', 4'-Hexachlorobiphenyl | 32.00/20mg |
| RCS-041 | 2, 3, 4, 2', 4', 5'-Hexachlorobiphenyl | 28.00/10mg |
| RCS-042 | 2, 3, 6, 2', 3', 6'-Hexachlorobiphenyl | 36.00/20mg |
| RCS-043 | 2, 4, 5, 2', 4', 5'-Hexachlorobiphenyl | 36.00/100mg |
| RCS-044 | 2, 4, 6, 2', 4', 6-Hexachlorobiphenyl | 32.00/50mg |
| RCS-045 | 2, 3, 4, 5, 2', 5'-Hexachlorobiphenyl | 28.00/10mg |
| RCS-046 | 2, 3, 5, 6, 2', 5'-Hexachlorobiphenyl | 28.00/10mg |
| RCS-047 | 2, 3, 4, 5, 6, 2', 5'-Heptachlorobiphenyl | 36.00/10mg |
| RCS-048 | 2, 3, 4, 5, 2', 3', 4', 5'-Octachlorobiphenyl | 26.00/20mg |
| RCS-049 | 2, 3, 5, 6, 1', 3', 5', 6'-Octachlrobiphenyl | 26.00/20mg |
| RCS-050 | 2, 3, 4, 5, 6, 2', 3', 4', 5'-Nonachlorobiphenyl | 36.00/10mg |
| RCS-051 | 2, 3, 4, 5, 6, 2', 3', 4', 5', 6'-Decachlorobiphenyl | 7.00/100mg |

Discount on single order of any 5 PCB's-10%  10 PCB's-15%
Bibliography on PCB Literature available on request.



MONS 067557

## POLYURETHANE-FOAM PLUGS

For complete extraction of pesticides and PCB's from environmental water samples (for instance – river water).
References:

1. The extraction and recovery of polychlorinated biphenyls (PCB) using porous polyurethane foam. H. D. Gesser, A. Chow, F. C. Davis, J. F. Uthe, and J. Reinke. Anal. Lett., 4, 883-6 (1971).
2. Extraction of organochlorine pesticides from water by porous polyurethane. J. F. Uthe, J. Reinke and H. Gesser, Environmental Letters, 3, 117-135 (1972).

Written procedure for extraction of pesticides and PCB's from water samples will be supplied with each order or on request.

Polyurethane-foam plugs (22mm x 38mm)

| | | |
|---|---|---|
| RCS-083 | Untreated | $10.00/50 |
| RCS-084 | Washed | 12.00/12 |
| RCS-085 | Treated with 0.5% DC-200 | 12.00/12 |

Chromatographic Tubes (20mm ID, 400mm long)

| | | |
|---|---|---|
| RCS-086 | Tube without fritted disc | $13.00/2 tubes |
| RCS-087 | Tube with fritted disc | 11.00/each |

## Perchlorination Kits for Analysis of Polychlorinated Aromatics

References:

1. Analysis of chlorinated aromatic hydrocarbons by exhaustive chlorination: Qualitative and structural aspects of the perchloro derivatives of biphenyl, naphthalene, terphenyl, dibenzofuran, dibenzodioxin and DDE. O. Hutzinger, S. Safe, and V. Zitko.
   International Journal of Environmental Analytical Chemistry, 2, 95 (1972).
2. Column chromatographic separation of polychlorinated biphenyls from chlorinated hydrocarbon pesticides, and subsequent gas chromatographic quantitation in terms of derivatives. O. W. Berg, P. L. Diosady and G. A. V. Rees Bulletin of Environmental Contamination and Toxicology, 7, #6, 338-347, (1972).
3. Screening method for the detection of chlorodibenzo-p-dioxins in the presence of chlorobiphenyls, chloronaphthalenes, and chlorodibenzofurans. D. T. Williams and B. J. Blanchfield, JAOAC. 55, 93-5 (1972).

| | | |
|---|---|---|
| RCS079 | Perchlorination reagents $A_1$ and $A_2$<br>For PCB's, PCDD's, PCF's, and p,p'-DDE (in equal volumes). | $10.00 |
| RCS080 | Perchlorination reagent B<br>For PCB, PCT and p,p'-DDE | 6.00 |
| RCS081 | Perchlorination reagent C<br>For PCN's | 6.00 |
| RCS082 | Perchlorination kit ($A_1$, $A_2$, B, C) | 16.00 |

Detailed perchlorination procedures will be supplied on request and with each order.



MONS 067558

## POLYCHLORINATED BIPHENYL MIXTURES

RCS066 ..... PCB-Kit (1 gm each of Aroclor 1232, 1242, 1248, 1254, 1260 and 1268) ..... 9.00/kit

## 99% POLYCHLORINATED NAPHTHALENES

RCS072 ..... Naphthalene ..... $ 7.00/500 mg
RCS073 ..... 1-Chloronaphthalene ..... 7.00/100 mg
RCS074 ..... 2-Chloronaphthalene ..... 7.00/100 mg
RCS075 ..... 1,2,3,4,5,6,7,8-Octachloronaphthalene ..... 15.00/50 mg
RCS076 ..... Halowax kit (1 gm each of 1031,1000, 1001, 1099, 1013, 1014, 1051) ..... 10.00/kit

| Halowax Number ..... | 1031 | 1000 | 1001 | 1099 | 1013 | 1014 | 1051 |
|---|---|---|---|---|---|---|---|
| Approx. % Chlorine ..... | 22 | 26 | 50 | 52 | 56 | 62 | 70 |

| Number of Chlorines ..... | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
|---|---|---|---|---|---|---|---|---|
| % of Chlorine ..... | 21.85 | 36.40 | 46.00 | 53.38 | 59.07 | 63.6 | 67.2 | 70.3 |

Reference: Behavior of chlorinated naphthalenes in analytical methods for organochlorinated pesticides and polychlorinated biphenyls. J. A. Armour and J. A. Burke, JAOAC, 54, 175-7 (1971).

## 99% POLYCHLORINATED DIBENZO-P-DIOXINS

RCS053 ... 1-Chlorodibenzo-p-dioxin ..... 20/50 mg
RCS054 ... 2-Chlorodibenzo-p-dioxin ..... 20/50 mg
RCS067 ... Dibenzo-p-dioxin ..... $15.00/50 mg
RCS055 ... 2, 3-Dichlorodibenzo-p-dioxin ..... 25/50 mg
RCS068 ... 2, 7-Dichlorodibenzo-p-dioxin ..... 25.00/50 mg
RCS069 ... 1, 2, 3, 4, 6, 7, 8, 9-Octachlorodibenzo-p-dioxin ..... 20.00/50 mg
RCS056 ... 1, 2, 4-Trichlorodibenzo-p-dioxin ..... 25/50 mg
RCS057 ... 1, 2, 3, 4-Tetrachlorodibenzo-p-dioxin ..... 25/50 mg

References:

1. Thin-layer chromatographic separation of two chlorodibenzo-p-dioxins from some polychlorinated biphenyls and organochlorine pesticides. D. T. Williams and B. J. Blanchfield, JAOAC, 54, 1429-31 (1971).
2. Determination of polychlorodibenzo-p-dioxins in chlorophenols and lipids. D. Firestone, Paper presented at the 162nd National ACS Meeting, Pesticide Div., Sept. 12-17, 1971.

## 99% POLYCHLORINATED DIBENZOFURANS

RCS070 ..... Dibenzofuran ..... $ 7.00/500 mg
RCS058 ..... 3, 6-Dichlorodibenzofuran ..... 36/20 mg
RCS071 ..... 1,2,3,4,5,6,7,8-Octachlorodibenzofuran ..... 15.00/50 mg


Reference: Identification and toxicological evaluation of chlorinated dibenzofuran and chlorinated naphthalene in two commercial polychlorinated biphenyls. J. G. Vos, J. H. Koeman, H. L. van der Maas, M. C. Ten Noever de Brauw and R. H. de Vos, Food Cosmet. Toxicol., 8, 625-633 (1970).

### OTHER POLLUTION STANDARDS

RCS077 ..... Hexachlorophene [2,2'-Methylenebis-(3,4,6-trichlorophenol)] ..... $ 5.00/1 mg
RCS078 ..... Perchlorinated p,p'-DDE ..... 20.00/50 mg



MONS 067559



# PESTICIDE STANDARDS

The list of pesticide standards that follows is possibly the most extensive compilation offered. For your convenience we have also listed the synonyms and a cross reference for different trade names. If the pesticide you need is not listed, please inquire. (*Available also in solution concentrations of .1 to 1000 PPM in powers of 10).

| Catalog # | Product | Synonym | Active Ingredient | % A. I. | Price |
|---|---|---|---|---|---|
| ——— | AAT | See Parathion | | | |
| ——— | Acquinite | See Trichloronitromethane | | | |
| ——— | Acricide | See Binapacryl | | | |
| ——— | Acti-Dione | See Cycloheximide | | | |
| RSP050 | Alanap | NPA | N-1-Naphthylphthalamic Acid | Tech | 10.00/g |
| RSP051 * | Aldrin | Octalene | 1, 2, 3, 4, 10, 10 Hexachloro-1, 4, 4, 5, 8, 8a-hexahydro-1, 4-endoexo-5, 8-dimethano naphthalene | 90% | 10.00/g |
| RSP057 | Aldrin, tech. | | 1, 2, 3, 4, 10, 10-Hexachloro-1, 4, 4, 5, 8, 8a-hexahydro-1, 4-endoexo-5, 8-dimethano naphthalene | 90% | 8.00/g |
| ——— | Allron | See Parathion | | | |
| RSP089 | Allyl Alcohol | | Allyl Alcohol | 97% | 8.00/g |
| RSP058 | Ametryne | | 2-Methyl mercapto-4-ethylamino-6-isopropylamino-S-triazine | 98.5% | 10.00/g |
| RSP059 | Ametryne, tech. | | 2-Methyl mercapto-4-ethylamino-6-isopropylamino-S-triazine | Tech | 8.00/g |
| ——— | Aminocarb | See Matacil | | | |
| RSP081 | Amitrole | | 3-Amino-1, 2, 4,-triazole | 90% | 20.00/100mg |
| RSP082 | Amitrole, tech. | | 3-Amino-1, 2, 4,-triazole | | 8.00/g |
| RSP083 * | Amobam | | Diammonium ethylene bisdithiocarbamate | 42% | 12.00/g |
| ——— | Apracarb | See Baygon | | | |
| RSP084 | Aramite | | 2-(p-tert-Butylphenoxy)-isopropyl-2'-chloroethyl sulfite | 93% | 20.00/100mg |
| ——— | Arathane | See Karathane | | | |
| RSP085 * | Atrazine | Gesaprim | 2-chloro-4-ethylamino-6-isopropylamino-s-triazine | 90% | 10.00/mg |
| RSP085 | Atrazine, tech. | | 2-chloro-4-ethylamino-6-isopropylamino-s-triazine | | 8.00/g |
| RSP088 | Avadex | Di-allate | S-2, 3-Dichloroallyl-N, N-diisopropyl-thiol carbonate | Tech | 12.00/g |
| RSP087 | Azinphosmethyl | Guthion Gothion Methyl guthion | O, O-Dimethyl-S-[4-oxo-1, 2, 3-benzotriazin-3 (4H)-yl methyl] phosphorodithioate | 96% | 15.00/g |
| RSP088 | Azinphosmethyl, tech. | | O, O-Dimethyl-S-[4-oxo-1, 2, 3-benzotriazin-3 (4H)-yl methyl] phosphorodithioate | | 6.00/g |
| RSP089 | Azobenzene | | Diphenyldiimide | 99% | 15.00/100mg |
| RSP070 | Bandane | | Polychlorodicyclopentadiene isomers | Tech | 12.00/100mg |
| RSP071 | Banvel D | Dicamba | 3, 6-Dichloro-o-anisic acid | 99% | 20.00/100mg |
| ——— | Baron | See Erbon | | | |
| ——— | Basudin | See Diazinon | | | |
| RSP078 | Baygon | Apracarb Unden | o-Isopropoxyphenyl methylcarbamate | Tech | 10.00/g |
| RSP073 | Benzyl Benzoate | | Benzyl Benzoate | 99% | 10.00/g |
| RSP063 * | BHC-mix | | Hexachlorocyclohexane isomer mix | Tech | 6.00/g |
| RSP049 | $\alpha$ -BHC | | Alpha-1, 2, 3, 4, 5, 6-Hexachlorocyclohexane | 99% | 20.00/100mg |
| RSP074 | $\beta$ -BHC | | Beta-1, 2, 3, 4, 5, 6-Hexachlorocyclohexane | 99% | 20.00/100mg |
| ——— | $\gamma$ -BHC | See Lindane | | | |
| RSP044 | $\delta$ -BHC | | Delta-1, 2, 3, 4, 5, 6 Hexachlorocyclohexane | 99% | 20.00/100mg |
| RSP075 | Butacrycl | Acricide Miticide | 2-sec-Butyl-4, 6-dinitrophenyl-3, 3-dimethylacrylate | 98% | 10.00/g |
| RSP076 | Butran | | 2, 6-Dichloro-4-nitroaniline | Tech | 10.00/g |
| ——— | Bromenil | See Bromoxynil | | | |
| RSP077 | Bromoxynil | Bromenil Buctryl | 3, 5-Dibromo-4-hydroxybenzonitrile | 98% | 10.00/100mg |
| ——— | Buctryl | See Bromoxynil | | | |
| RSP078 | Butan | | 2-Nitro-1, 1-bis (p-chlorophenyl) butane | 98% | 10.00/100mg |
| ——— | Butoxide | See Piperonyl Butoxide | | | |
| ——— | Caparol | See Prometryne | | | |



MONS 067560

# PESTICIDE STANDARDS

| Catalog # | Product | Synonym | Active Ingredient | % A. I. | Price |
|---|---|---|---|---|---|
| RSP079 | Captan | Orthocide | N-Trichloromethylthio-4-cyclohexene 1, 2-dicarboximide | 99% | 10.00/g |
| ——— | Carbaryl | See Sevin | | | |
| ——— | Carbophenothion | See Trithion | | | |
| ——— | Carbophos | See Malathion | | | |
| ——— | Cardibe | See Ethyl Hexane Diol | | | |
| RSP080 | Chlorbenside | Sulfide<br>Chlorparacide<br>Chlorsulphacide<br>Mitox | p-chlorobenzyl-p-chlorophenyl | 99% | 12.00/100mg |
| RSP004 • | Chlordane | Octachlor | 1, 2, 3, 4, 5, 6, 7, 8, 8-Octachloro-3a, 4, 7, 7a-tetrahydro-4, 7-methano indene | Tech. | 14.00/g |
| RSP081 | Chloroacetic acid | Monochloroacetic Acid | Chloroacetic Acid | 99% | 8.00/g |
| RSP082 | Chlorobenzilate | | Ethyl-4, 4'-dichlorobenrilate | 99% | 10.00/g |
| RSP083 | Chlorobenzilate, tech. | | Ethyl-4, 4'-dichlorobenzilate | 45% | 4.00/g |
| ——— | Chlorophos | See Trichlorfon | | | |
| ——— | Chloropicrin | See Trichloronitromethane | | | |
| ——— | Chlorsulphacide | See Chlorbenside | | | |
| ——— | Chlorparacide | See Chlorbenside | | | |
| RSP084 | Cinchrin | | $\alpha$ -Methyl benzyl-3-(dimethoxyphosphinyloxy)-cis-crotonate | 85% | 10.00/g |
| RSP088 • | Co-Ral | Coumaphos | O, O-Diethyl-O-(3-chloro-4-methyl-2-oxo-2H-1-benzopyran-7-yl phosphoro thioate | 99% | 2.100/100mg |
| RSP085 | Co-Ral, tech | | O, O-Diethyl-O-(3-chloro-4-methyl-2-oxo-2H-1-benzopyran-7-yl phosphoro thioate | | 6.00/g |
| ——— | Coumafene | See Warfarin | | | |
| ——— | Coumaphos | See Co-Ral | | | |
| RSP086 | Cyclohexamide | Acti-dione | 3-[2-(3, 5 Dimethyl-2-oxycyclohexyl) -2-hydroxyethyl] glutarimide | Tech. | 12.00/g |
| ——— | Cygon | See Dimethoate | | | |
| ——— | Cymen | See Ziraphos | | | |
| RSP087 | Cypova | Dodine | n-dodecylguanidine acetate | 95% | 20.00/100mg |
| RSP088 | Cypova, tech. | | n-dodecylguanidine acetate | | 6.00/g |
| RSP009 • | 2, 4-D Acid | | 2-(2, 4-Dichlorophenoxy) acetic acid | 99% | 12.00/100mg |
| RSP007 • | 2, 4-D Acid-Methyl Ester | | 2-(2, 4-Dichlorophenoxy) acetic acid methyl ester | 99% | 10.00/100mg |
| RSP089 | Dacthal | DCPA | Dimethyl-2, 3, 5, 6-tetrachloroterephthalate | 99% | 10.00/g |
| RSP090 | Dalapon | | 2, 2-Dichloropropionic Acid | 90% | 20.00/100mg |
| ——— | Dall | See Methyl Parathion | | | |
| RSP091 | Dasanet | Mylone | 3, 5-Dimethyl-1, 3, 5, 3H-tetrahydrothiadiazin-2-thone | 98% | 12.00/g |
| ——— | DCPA | See Dacthal | | | |
| RSP008 | DDD mix | TDE mix | 1, 1-Dichloro-2, 2-(chlorophenyl) ethane | Tech. (3 isomer mix) | 3.00/g |
| RSP098 • | , p-DDD | o, p-TDE | 1, 1-Dichloro-2-(o-chlorophenyl)-2-(p-chlorophenyl) ethane | 99% | 12.00/100mg |
| RSP009 • | p, p'-DDD | p, p'-TDE | 1, 1-Dichloro-2, 2-bis (p-chlorophenyl) ethane | 99% | 20.00/100mg |
| RSP047 | o, p-DDE | o, p-DDX | 1, 1-Dichloro-2-(o-chlorophenyl)-2-(p-chlorophenyl) ethylene | 99% | 20.00/100mg |
| RSP010 • | p, p'-DDE | p, p'-DDX | 1, 1-Dichloro-2, 2'-bis (p-chlorophenyl) ethylene | 99% | 25.00/g |
| RSP011 • | DDT mix | | 1, 1, 1-Trichloro-2, 2-bis (chlorophenyl) ethylene | Tech (3 isomer mix) | 2.00/g |
| RSP018 • | o, p-DDT | | 1, 1, 1-Trichloro-2-(o-chlorophenyl) -2-(p-chlorophenyl) ethane | 99% | 25.00/g |
| RSP013 • | p, p'-DDT | | 1, 1, 1-Trichloro-2, 2-bis (p-chlorophenyl) ethane | 99% | 12.00/g |
| ——— • | DDVP | See Dichlorvos | | | |
| ——— | o, p-DDX | See o, p-DDE | | | |
| ——— | p, p'-DDX | See p, p'-DDE | | | |
| RSP093 | DEET | Delphene<br>Detamide | N, N-Diethyl-m-toluamide | 95% | 12.00/g |



MONS 067561

# PESTICIDE STANDARDS

| Catalog # | Product | Synonym | Active Ingredient | % A. I. | Price |
|---|---|---|---|---|---|
| RSP015 • | DEF | | S, S, S-Tributyl phosphorotrithioate | 94% | 22.00/100mg |
| ——— | Delphene | See DEET | | | |
| RSP018 • | Demeton | Systox | Mix of O, O-Diethyl-O-2-(ethyl thio) ethyl phosphorothioate) and O, O-Diethyl S-2-(ethylthio) ethyl phosphorthioate) | Tech. | 20.00/100mg |
| ——— | Detsuide | See DEET | | | |
| RSP094 | Deuon | | p-Dimethylaminobenzenediazo sodium sulfonate | 84% | 12.00/100mg |
| ——— | DFA | See Diphenylamine | | | |
| ——— | Di-Allate | See Avadex | | | |
| RSP017 • | Diazinon | Basudin | O, O-Diethyl-O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate | 99% | 12.00/100mg |
| RSP086 | Diazinon, tech. | | O, O-Diethyl-O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate | 60% | 6.00/g |
| RSP018 • | Dibrom | Naled | 1, 2-Dibromo-2, 2-dichloroethyl dimethyl phosphate | 93% | 12.00/100mg |
| RSP088 | Dibutylphthalate | | Dibutylphthalate | 99% | 12.00/g |
| ——— | Dicamba | See Banvel D | | | |
| ——— | Dicapton | See Dicaphon | | | |
| RSP087 | Dicaphon | Dicapton | O-(2-Chloro-4-nitrophenyl)-O, O-dimethyl phosphorothioate | 99.8% | 12.00/100mg |
| ——— | Dichlone | See Phygon | | | |
| RSP088 | o-Dichlorobenzene | | o-Dichlorobenzene | 99% | 6.00/g |
| RSP089 | p-Dichlorobenzene | | p-Dichlorobenzene | 99% | 6.00/g |
| RSP180 | 2, 4-Dichlorophenol | | 2, 4-Dichlorophenol | 99% | 6.00/g |
| ——— | Dichlorprop | See 2, 4-DP-Acid | | | |
| RSP014 | Dichlorvos | DDVP<br>Herkol<br>Vapona | O, O-Dimethyl-O-(2, 2-dichlorovinyl) phosphate | 99% | 12.00/100mg |
| ——— | Dicofol | See Kelthane | | | |
| RSP019 • | Dieldrin | Hexol | 1, 2, 3, 4, 10, 10-Hexachloro-6, 7-epoxy-1, 4, 4a, 5, 6, 7, 8, 8a-octahydro-1, 4-endoexo-5, 8-dimethanonaphthalene | 85% | 10.00/g |
| RSP101 | Dikolutan | | N-(1, 1, 2, 2-Tetrachloroethylsulfenyl) cis-Δ-4-cyclohexene-1, 2-dicarboximide | 99% | 12.00/100mg |
| ——— | Dimecron | See Phosphamidon | | | |
| RSP102 | Dimethoate | Rogor<br>Cygon<br>Fostion | O, O-Dimethyl-S-(N-Methyl carbamoyl-methyl) phosphorodithioate | 99% | 22.00/g |
| RSP103 | Dimethyl Phthalate | | Dimethyl Phthalate | 99% | 12.00/g |
| ——— | Dinocap | See Karathane | | | |
| RSP104 | Dinosb | DNBP | 2, 4-Dinitro-6-sec-butyl phenol | 99% | 12.00/100mg |
| RSP105 | Diphacin | Diphacinone | 2-Diphenyl-acetyl-1, 3-indandione | Tech | 12.00/100mg |
| ——— | Diphenenone | See Diphacin | | | |
| RSP106 | Diphenamid | Dymid<br>Enid | N, N-Dimethyl 2, 2-diphenylacetamide | Tech | 18.00/g |
| RSP107 | Diphenyl-Amine | DFA | Diphenylamine | 99% | 18.00/g |
| ——— | Dipterex | See Trichlorfon | | | |
| ——— | Diptrine | See Trichlorfon | | | |
| ——— | Disulfoton | See Di-Syston | | | |
| RSP108 | Di-Syston | Disulfoton | O, O-Diethyl-S-2-(ethylthio) ethyl phosphorodithioate | 97% | 16.00/g |
| ——— | Dithane-Manganese | See Maneb | | | |
| ——— | Dithane-Zinc | See Zineb | | | |
| ——— | DMPA | See Zytron | | | |
| ——— | DNBP | See Dinoseb | | | |
| ——— | DNTP | See Parathion | | | |
| ——— | Dodine | See Cyprex | | | |
| ——— | Dowco 138 | See Ruelene | | | |
| RSP109 | 2, 4-DP-Acid | Dichlorprop | 2-(2, 4-Dichlorophenoxy) propionic acid | 98% | 12.00/100mg |
| ——— | DTMC | See Kelthane | | | |
| ——— | Duphor | See Tedion | | | |
| RSP110 | Durshan | | O, O-Diethyl-O-(3, 5, 6-trichloro-2-pyridyl) phosphorothioate | 99% | 12.00/100mg |

MONS 067562

# PESTICIDE STANDARDS

| Catalog # | Product | Synonym | Active Ingredient | % A.I. | Price |
|---|---|---|---|---|---|
| ------ | Dylox | See Trichlorfon | | | |
| ------ | Dyrmil | See Diphenamid | | | |
| ------ | Endosulfan | See Thiodan | | | |
| RSP020 • | Endrin | | 1, 2, 3, 4, 10, 10-Hexachloro-6, 7-epoxy-1, 4, 4a, 5, 6, 7, 8, 8a-octahydro-1, 4-endo, endo-5,8-dimethanonaphthalene | 99% | 10.00/g |
| RSP111 | Endrin, tech. | | 1, 2, 3, 4, 10, 10-Hexachloro-6, 7-epoxy-1, 4, 4a, 5, 6, 7, 8, 8a-octahydro-1, 4-endo, endo-5,8-dimethanonaphthalene | 97% | 8.00/g |
| ------ | Enid | See Diphenamid | | | |
| RSP021 • | EPN | | O-Ethyl-O-p-nitrophenyl phenylphosphonothioate | 97% | 12.00/100mg |
| RSP112 | Ethion | Rhodia | 2-(2, 4, 5-Trichlorophenoxy) ethyl-2, 2-dichloropropionate | Tech. | 8.00/g |
| ------ | Ethomate | See Oven | | | |
| RSP088 • | Ethion | Nialate | O, O, O', O'-Tetraethyl-S, S'-methylene bisphosphorodithioate | 95% | 22.00/100mg |
| RSP113 | Ethyl Hexanediol | Cardite | 3-Ethyl-1, 3 Hexanediol | Tech. | 12.00/g |
| ------ | Ethon | See Parathion | | | |
| RSP115 | Eugenol | | Eugenol | 99% | 12.00/g |
| RSP124 | Fenitrothion | Folithion<br>Sumithion | O, O-Dimethyl-O-(4-nitro-m-tolyl) phosphorothioate | 98% | 20.00/100mg |
| RSP033 | Fenitrothion, tech. | | O, O-Dimethyl-O-(4-nitro-m-tolyl) phosphorothioate | 95% | 12.00/g |
| RSP110 | Fenthion | | O, O-Dimethyl-O-(4-(methylthio) -m-tolyl) phosphorothioate | Tech. | 10.00/100mg |
| ------ | Fenuron | See Silvex | | | |
| ------ | Fenuron | See Parathion | | | |
| ------ | Folithion | See Fenitrothion | | | |
| ------ | Folpet | See Phaltan | | | |
| ------ | Fonofos | See Stauffer | | | |
| ------ | Fonofos | See Dyanthonate | | | |
| RSP117 | Fumazone | | 1, 2-Dibromo-3-chloropropane | Tech. | 8.00/g |
| ------ | Genafume | See Prometone | | | |
| ------ | Gemanil | See Propazine | | | |
| ------ | Geogrim | See Atrazine | | | |
| ------ | Geogard | See Prometryne | | | |
| ------ | Geotop | See Simazine | | | |
| ------ | Gonthion | See Azinphosmethyl | | | |
| ------ | Guthion | See Azinphosmethyl | | | |
| ------ | HCB | See Hexachlorobenzene | | | |
| ------ | HEOD | See Dieldrin | | | |
| RSP024 • | Heptachlor | Methanoindene | 1, 4, 5, 6, 7, 8, 8-Heptachloro-3a, 4, 7, 7a-tetrahydro-4, 7-methanoindene | 97.4% | 10.00/g |
| RSP025 • | Heptachlor | | 1, 4, 5, 6, 7, 8, 8-Heptachloro-2, 3-epoxy-3a, 4, 7, 7a-tetrahydro-4, 7-methanoindan | 99% | 35.00/100mg |
| ------ | Herbol | See Dichlorvos | | | |
| RSP051 | Hexachlorobenzene | HCB | Hexachlorobenzene | 99% | 10.00/g |
| RSP118 | Imidan | | O, O-Dimethyl-S-phthalimidomethyl phosphorodithioate | 95% | 10.00/g |
| RSP119 | IPC | | Isopropyl N-phenylcarbamate | Tech. | 6.00/g |
| ------ | Isocarbane | See Karathane | | | |
| RSP120 | Karathane | Arathane<br>Dinocap<br>Isocrotane<br>Mildex | 2 (1-Methyl-n-heptyl)-4, 6-dinitrophenyl crotonate | 85% | 20.00/g |
| ------ | Karbofos | See Malathion | | | |
| RSP026 • | Kelthane | Dicofol<br>DTMC | 1, 1-bis (4-Chlorophenyl)-2, 2, 2-trichloroethanol | 99% | 17.00/g |
| RSP121 | Kelthane, tech. | | 1, 1-bis (4-Chlorophenyl)-2, 2, 2-trichloroethanol | | 8.00/g |
| RSP045 | Lannate | Methomyl | S-Methyl-N-((Methylcarbamoyl) oxy) thioacetimidate | 90% | 10.00/g |
| RSP122 | Lethane 384 | | 2-(2-Butoxyethoxy) ethyl thiocyanate | 53% | 12.00/g |

MONS 067563

# PESTICIDE STANDARDS

| Catalog # | Product | Synonym | Active Ingredient | % A. I. | Price |
|---|---|---|---|---|---|
| RSP027 • | Lindane | Gamma-BHC (γ-BHC) | Gamma-1, 2, 3, 4, 5, 6-Hexachlorocyclohexane | 99% | 20.00/g |
| RSP028 • | Malathion | Carbophos Karbofos | S-(1, 2-bis (Ethoxycarbonyl) ethyl)-O, O-dimethyl phosphorodithioate | 99% | 20.00/g |
| RSP183 | Malathion, tech. | | S-(1, 2-bis (Ethoxycarbonyl) ethyl)-O, O-dimethyl phosphorodithioate | 95% | 8.00/g |
| ——— | Malix | See Thiodan | | | |
| RSP124 | Maneb | Dithane-manganese Manzate | Ethylenebisdithiocarbamate manganese | 80% | 12.00/100mg |
| ——— | Manzate | See Maneb | | | |
| RSP125 | Matacil | Aminocarb | 4-Dimethylamino-3-tolyl N-methyl carbamate | 98% | 12.00/100mg |
| RSP126 | Mesmin | | (N-Methylmercuri)-1, 4, 5, 6, 7, 7-hexachlorobicyclo (2.2.1) hept-5-ene-2, 3-dicarboximide | 99.7% | 20.00/100mg |
| RSP127 | Metasystox-R | Oxydemeton-methyl | O, O-Dimethyl-S-(2-(ethylsulfinyl) ethyl) phosphorothioate | 50% | 12.00/g |
| ——— | Methomyl | See Lannate | | | |
| RSP029 • | Methoxychlor | Methoxy DDT DMDT | 1, 1, 1-Trichloro-2, 2-bis (p-methoxyphenyl) ethane | 98% | 17.00/g |
| RSP128 | Methoxychlor, tech. | | 1, 1, 1-Trichloro-2, 2-bis (p-methoxyphenyl) ethane | 80.5% | 6.00/g |
| ——— | Methoxy Propazine | See Prometone | | | |
| ——— | Methyl Guthion | See Azinphosmethyl | | | |
| RSP129 | Methyl Parathion | | O, O-Dimethyl-O-p-nitrophenyl phosphate | 98% | 12.00/g |
| RSP068 | Methyl Parathion | Dalf Nitrox | O, O-Dimethyl-O-p-nitrophenyl phosphorothioate | 98% | 12.00/100mg |
| RSP130 | Methyl Parathion, tech. | | O, O-Dimethyl-O-p-nitrophenyl phosphorothioate | 85% | 8.00/g |
| ——— | Mevinphos | See Phosdrin | | | |
| ——— | Mildex | See Karathane | | | |
| ——— | Mintacol | See Para-zeon | | | |
| RSP050 | Mirex | | Dodecachloroctahydro-1, 3, 4-metheno-2H-cyclobuta (cd) pentalene | 99% | 16.00/g |
| ——— | Mitox | See Chlorbenside | | | |
| ——— | Monochloroacetic Acid | See Chloroacetic Acid | | | |
| RSP131 | Morestan, tech. | Fortan | 6-Methyl-2-oxo-1, 3-dithio (4, 5-b) quinoxaline | 87% | 12.00/g |
| ——— | Morocide | See Binapacryl | | | |
| ——— | Naled | See Dibrom | | | |
| ——— | Negaweon | See Trichlorfon | | | |
| ——— | Nemafos | See Zinophos | | | |
| ——— | Nialate | See Ethion | | | |
| ——— | Niran | See Parathion | | | |
| ——— | Nitrostigmine | See Parathion | | | |
| ——— | Nitox | See Methyl Parathion | | | |
| ——— | NPA | See Alanap | | | |
| ——— | Octachlor | See Chlordane | | | |
| ——— | Octalene | See Aldrin | | | |
| ——— | OMPA | See Schradan | | | |
| ——— | Orthocide | See Captan | | | |
| RSP132 | Ovex | Estonate Ovochlor Ovotran | p-Chlorophenyl p-chlorobenzenesulfonate | 98% | 12.00/g |
| RSP133 | Ovex, tech. | | p-Chlorophenyl p-chlorobenzenesulfonate | | 4.00/g |
| ——— | Ovochlor | See Ovex | | | |
| ——— | Ovotran | See Ovex | | | |
| ——— | Oxydemetonmethyl | See Metasystox-R | | | |
| RSP134 | Panogen | | Methylmercuric dicyanodiamide | min. 90% | 12.00/g |
| RSP030 • | Para-zeon | Mintacol | O, O-Diethyl-o-p-nitrophenyl phosphate (oxygen analog of parathion) | 99% | 12.00/100mg |
| RSP135 | Paraquat Dichloride | | 1, 1'-Dimethyl-4, 4'-dipyridinium dihydrochloride | 99% | 12.00/100mg |




MONS 067564

# PESTICIDE STANDARDS

| Catalog # | Product | Synonym | Active Ingredient | % A. I. | Price |
|---|---|---|---|---|---|
| RSP031 | Parathion | Folidol<br>Niran<br>Ethion<br>Rhodiatox<br>Thiophos<br>Nitrostigmine<br>Alkron<br>AAT; SNP;<br>DNTP; | O, O-Diethyl-o-p-nitrophenyl phosphorothioate | 99.7% | 15.00/100mg |
| RSP130 | Parathion, tech. | | O, O-Diethyl-o-p-nitrophenyl phosphorothioate | Tech. | 12.00/g |
| RSP137 | Paris Green | | Copper acetoarsenite | 91.7% | 6.00/g |
| RSP138 | Parmon | | Bis (p-chlorophenyl)-3-pyridine methanol | Tech. | 12.00/100mg |
| ------ | Parnate | See Zineb | | | |
| RSP139 | PCNB | | Pentachloronitrobenzene | 99% | 8.00/g |
| ------ | PCP | See Pentachlorophenol | | | |
| RSP140 | Pentachlorophenol | PCP<br>Santobrite | Pentachlorophenol | 99% | 20.00/g |
| RSP141 | Perthane | | 1, 1-dichloro-2, 2-bis (p-ethylphenyl) ethane | 95% | 12.00/g |
| ------ | Peyan | See Schradan | | | |
| RSP148 | Phaltan | Folpet<br>Thiophol | N-Trichloromethylthiophthalimide | 99% | 12.00/100mg |
| RSP143 | Phaltan, tech. | | N-Trichloromethylthiophthalimide | 93.3% | 8.00/g |
| ------ | Pharmacide | See Tenaphene | | | |
| ------ | Phermitin | See Tenaphene | | | |
| RSP038 * | Phenothiazine | | Phenothiazine | Pract. | 12.00/g |
| ------ | Phorate | See Thimet | | | |
| RSP033 * | Phosdrin | Mevinphos | 2-Carbomethoxy-1-propen-2yl dimethyl phosphate | 60% | 18.00/g |
| RSP144 | Phosphamidon | Dimecron | 1-chlorodiethyl carbamoyl-1-propen-2-yl-dimethyl phosphate | 99% | 15.00/g |
| RSP145 | Phosphamidon | | 1-chlorodiethyl carbamoyl-1-propen-2-yl-dimethyl phosphate | 87.8% | 6.00/g |
| RSP146 | Phygon | Dichlone | 2, 3-Dichloro-1, 4-naphthoquinone | Tech. | 12.00/g |
| RSP147 | Piperonyl Butoxide | Butacide | 12-(2-n-Butoxyethoxy)-ethoxy)-4, 5-methylene dioxy-2-propyl butkene | 90% | 10.00/g |
| RSP148 | Plantvax | | 2, 3-Dihydro-5-carboxanilide-6-methyl-1, 4-oxathiin-4, 4-dioxide | 97% | 22.00/g |
| RSP149 | Pramex | | N, N-Di-n-butyl-p-chlorobenzene sulfonamide | Tech. | 8.00/g |
| ------ | Prevenol I | See 2, 4, 5-Trichlorophenol | | | |
| RSP150 | Prolan | | 2-Nitro-1, 1-bis (p-chlorophenyl) propane | 99% | 16.00/100mg |
| RSP155 | Prometone | Methoxy-propazine<br>Gesafram<br>Caparol<br>Genagard | 2, 4-Bis-(isopropylamino)-6-methoxy-S-triazine | 99% | 14.00/g |
| RSP034 * | Prometryne | | 2-Methyl-mercapto-4, 6-bis (isopropyl amino)-S-triazine | 99% | 15.00/100mg |
| RSP035 * | Propazine | Gesamil | 2-Chloro-4, 6-bis (isopropyl amino) -S-triazine | 98.7% | 14.00/100mg |
| RSP151 | Propyliasate | | Di-n-propyl-3-methyl-6, 7-methylene-hexy-1, 2, 3, 4-tetrahydronaphthalene-1, 2-dicarboxylate | Tech. | 10.00/g |
| RSP152 | Randox | CDAA | 2-Chloro-N, N-diallyl acetamide | Tech. | 6.00/g |
| ------ | Rhodiatox | See Parathion | | | |
| ------ | Rogor | See Dimethoate | | | |
| RSP036 * | Ronnel | | Dimethyl 2, 4, 5-trichlorophenyl phosphorothioate | 99% | 14.00/g |
| RSP152 | Ronnel, tech. | | Dimethyl 2, 4, 5-trichlorophenyl phosphorothioate | Tech. | 6.00/g |
| RSP034 | Roshow | Dowco 128 | 4-tert-Butyl-2-chlorophenyl methylethyl phosphoramidite | 99% | 12.00/100mg |
| RSP154 | Roshow, tech. | | 4-tert-Butyl-2-chlorophenyl methylethyl phosphoramidite | | 8.00/g |
| ------ | Santobrite | See Pentachlorophenol | | | |



MONS 067565



# PESTICIDE STANDARD

| Catalog # | | Product | Synonym | Active Ingredient | % A.I. | Price |
|---|---|---|---|---|---|---|
| RSP038 | • | Schradan | OMPA<br>Pestan<br>hytess | Octamethylpyrophosphoramide | 99.8% | 14.00/g |
| RSP173 | | Schradan, tech. | | Octamethylpyrophosphoramide | 95% | 8.00/g |
| RSP170 | • | Sevin | Carbaryl | 1-Naphthyl N-methyl carbamate | 99% | 21.00/100mg |
| RSP171 | | Silvex | Fenoprop | 2-(2, 4, 5-Trichlorophenoxy) propionic Acid | 99% | 10.00/100mg |
| RSP172 | | Silvex, tech. | | 2-(2, 4, 5-Trichlorophenoxy) propionic Acid | | 8.00/g |
| RSP039 | • | Simazine | Gesatop | 2-chloro-4, 6-bis (ethylamino)-S-triazine | 99.3% | 11.00/100mg |
| ------ | | SNP | See Parathion | | | |
| RSP174 | | Spergon | Chloranil | Tetrachloro-p-benzoquinone | Tech | 6.00/g |
| RSP175 | | Strobane | | Terpene polychlorinates | 95%<br>Chlorene | 10.00/g |
| ------ | | Sulfo-Clde | See Sulfoxide | | | |
| RSP176 | | Sulfoxide | Sulfonyl<br>Sulfo-cide | 1, 2-Methylmedioxy-4-12-(octyl sulfinyl) propyl] benzene | 88% | 12.00/g |
| ------ | | Sulfonyl | See Sulfoxide | | | |
| ------ | | Sumithion | See Fenitrothion | | | |
| ------ | | Systox | See Demeton | | | |
| ------ | | Sytam | See Schradan | | | |
| RSP177 | | 2, 4, 5 T-acid | | 2, 4, 5-Trichlorophenoxyacetic Acid | 98% | 10.00/100mg |
| RSP178 | | 2, 4, 5 T-acid | | 2, 4, 5-Trichlorophenoxyacetic Acid | Tech. | 8.00/g |
| ------ | | TCTS | See Tedion | | | |
| ------ | | TDE mix | See DDD-mix | | | |
| ------ | | o, p-TDE | See o, p-DDD | | | |
| ------ | | p, p' TDE | See p, p'-DDD | | | |
| RSP158 | | Tedion | Duphar<br>Tetradifon<br>TCTS | p-Chlorophenyl 2, 4, 5-trichlorophenyl sulfone | 98% | 10.00/g |
| RSP187 | | 2, 3, 4, 6 Tetrachlorophenol | | 2, 3, 4, 6 Tetrachlorophenol | 98% | 12.00/100mg |
| ------ | | Tetradifon | See Tedion | | | |
| RSP156 | | Thimet | Phorate | O, O Diethyl S-(ethylthio) methyl phosphorodithioate | 98.8% | 15.00/g |
| RSP159 | | Thimet, tech. | | O, O Diethyl S-(ethylthio) methyl phosphorodithioate | 91% | 8.00/g |
| RSP040 | • | Thiodan | Endosulfan<br>Malix | 6, 7, 8, 9, 10, 10-Hexachloro-1, 5, 5a, 6, 9, 9a-hexahydro-6, 9-methano 2, 4, 3-benzodioxathiepin 3-oxide | 98% | 23.00/100mg |
| ------ | | Thiometon | See Zinephos | | | |
| ------ | | Thiophal | See Phaltan | | | |
| ------ | | Thiophos | See Parathion | | | |
| RSP041 | • | Tenaphene | Phenacide<br>Phenator | Polychlorinated camphene | Tech. | 17.00/g |
| RSP042 | • | Trichlorfon | Tugon<br>Chlorophos<br>Dipterex<br>Dylox<br>Neguvon<br>Dipterex | O, O Dimethyl (1-hydroxy-2, 2, 2-trichloroethyl) phosphonate | 98% | 12.00/g |
| RSP180 | | Trichloronitromethane | Chloropicrin<br>Acquisite | Trichloronitromethane | 95% | 12.00/g |
| RSP181 | | 2, 4, 5 Trichlorophenol | Preweted I | 2, 4, 5 Trichlorophenol | 99% | 10.00/100mg |
| RSP182 | | 2, 4, 6 Trichlorophenol | | 2, 4, 6 Trichlorophenol | 99% | 10.00/100mg |
| RSP055 | | Trithion | Carbaphenothion | S-[[ p-Chlorophenyl) thiol methyl] O, O diethyl phosphorodithioate | 95% | 10.00/g |
| ------ | | Tugon | See Trichlorfon | | | |
| ------ | | Undex | See Baygon | | | |
| ------ | | Vapona | See Dichlorvos | | | |
| RSP183 | | Vorlex | | Methylisothiocyanate | 20% | 12.00/g |
| RSP043 | • | Warfarin | Coumafene | 3-(α'-Acetonylbenzyl)-4-hydroxy coumarin | Tech | 10.00/100mg |

MONS 067566

## PESTICIDE STANDARDS

| Catalog # | Product | Synonym | Active Ingredient | % A.I. | Price |
|---|---|---|---|---|---|
| RSP104 | Zectran | | 4-dimethyl amino-3, 5 xylyl-N-Methyl carbamate | 99% | 10.00/g |
| RSP105 | Zectran, tech. | | 4-dimethyl amino-3, 5 xylyl-N-Methyl carbamate | 95% | 6.00/g |
| RSP106 | Zineb | Dithane-zinc Famate | Zinc ethylene bisdithiocarbamate | 75% | 20.00/g |
| RSP107 | Ziosphos | Thermorin Cyren Nemafos | O, O-Diethyl-O-2-pyrazinyl phosphorothioate | 98% | 10.00/100mg |
| RSP108 | Zytron | DMPA | O-(2, 4-Dichlorophenyl)-O-methyl isopropyl phosphoramidothioate | 98% | 12.00/100mg |
| RSP109 | Zytron, tech. | | O-(2, 4-Dichlorophenyl)-O-methyl isopropyl phosphoramidothioate | Tech. | 8.00/g |

# PRECOATED TLC PLATES from ANALABS

Analabs TLC plates are of precision quality. They are precoated using procedures developed by leading chromatographic investigators.

The prescored 20 x 20 cm plate can be snapped in sections for two 10 x 20 cm, or four 5 x 20 cm plates. Scoring on the uncoated side makes snapping the plates easy. A clear edge on the plate allows the plate to be grasped without disturbing the coating. Separation information can be marked in this area.

ORDERING INFORMATION
20 x 20 cm PRECOATED PLATES 250 $\mu$ thick. 25 PLATES/BOX

| *NUMBER CATALOG* | *TLC PLATE TYPE* | *PRESCORED PRICE* | | *UNSCORED PRICE* |
|---|---|---|---|---|
| | | 5 x 20 cm | 10 x 20 cm | 20 x 20 cm |
| TLP-001 | Anasil G | $30 | $26 | $22 |
| TLP-002 | Anasil GF (254) | $30 | $26 | $22 |
| TLP-003 | Anasil H | $30 | $26 | $22 |
| TLP-004 | Anasil HF (254) | $30 | $26 | $22 |
| TLP-005 | Silica Gel G | $30 | $26 | $22 |
| TLP-006 | Silica Gel GF (254) | $30 | $26 | $22 |
| TLP-007 | Microcrystalline Cellulose (Avicel) | $31 | $27 | $23 |
| TLP-008 | Microcrystalline Cellulose, F (254) (Avicel) | $31 | $27 | $23 |

Analabs precoated plates are guaranteed to arrive at your laboratory in condition for immediate use. Any plates that arrive damaged will be replaced at no cost to you and no troublesome paperwork.



MONS 067567

# PESTICIDEGRADE SOLVENTS

Pesticidegrade Solvents are certified to contain no interfering substance greater than that equivalent to 10 parts per trillion of a chlorinated pesticide.

| *Catalog #* | *Solvent* | | *Unit* | *Price* |
|---|---|---|---|---|
| SSP001 | Acetone, Presticidgrade | | | |
| | Water | 0.05% max. | 1 Liter | $14.00 |
| SSP002 | Acetonitrile, Pesticidegrade | | | |
| | Water | 0.05% max. | 1 Liter | 15.00 |
| SSP003 | Benzene, Pesticidegrade | | | |
| | Water | 0.03% max. | 1 Liter | 12.00 |
| SSP004 | 2-Butanone, Pesticidegrade | | | |
| | Water | 0.05% max. | 1 Liter | 13.00 |
| SSP005 | Butyl Alcohol, Pesticidegrade | | | |
| | Water | 0.05% max. | 1 Liter | 12.00 |
| SSP006 | iso-Butyl Alcohol, Pesticidegrade | | | |
| | Water | 0.05% max. | 1 Liter | 15.00 |
| SSP007 | Carbon Tetrachloride, Pesticidegrade | | | |
| | Water | 0.02% max. | 1 Liter | 16.00 |
| SSP008 | Chloroform, Pesticidegrade | | | |
| | Water | 0.02% max. | 1 Liter | 17.00 |
| SSP009 | Cyelohexane, Pesticidegrade | | | |
| | Water | 0.02% max. | 1 Liter | 16.00 |
| SSP010 | Dichloromethane, Pesticidegrade | | | |
| | Water | 0.02% max. | 1 Liter | 16.00 |
| SSP011 | N, N-Dimethylformamide, Pesticidegrade | | | |
| | Water | 0.05% max. | 1 Liter | 19.00 |
| SSP012 | p-Dioxane, Pesticidegrade | | | |
| | Water | 0.05% max. | 1 Liter | 17.00 |
| SSP013 | Ether, Pesticidegrade | | | |
| | Water | 0.05% max. | 1 Liter | 12.00 |
| SSP014 | Ethyl Acetate, Pesticidegrade | | | |
| | Water | 0.05% max. | 1 Liter | 13.00 |
| SSP015 | Ethyl Alcohol, Pesticidegrade Completely Denatured | | | |
| | Water | 0.05% max. | 1 Liter | 12.00 |
| SSP016 | Heptane, Pesticidegrade | | | |
| | Water | 0.03% max. | 1 Liter | 13.00 |
| SSP017 | Hexane, Pesticidegrade | | | |
| | Water | 0.02% max. | 1 Liter | 12.00 |
| SSP018 | Methanol, Pesticidegrade | | | |
| | Water | 0.05% max. | 1 Liter | 11.00 |
| SSP019 | Methylsulfoxide, Pesticidegrade | | | |
| | Water | 0.05% max. | 1 Liter | 21.00 |
| SSP020 | Pentane, Pesticidegrade | | | |
| | Water | 0.03% max. | 1 Liter | 15.00 |
| SSP021 | Petroleum Ether, Pesticidegrade | | | |
| | Water | 0.03% max. | 1 Liter | 12.00 |
| SSP022 | iso-propyl Alcohol, Pesticidegrade | | | |
| | Water | 0.05% max. | 1 Liter | 15.00 |
| SSP023 | Toluene, Pesticidegrade | | | |
| | Water | 0.02% max. | 1 Liter | 11.00 |
| SSP024 | 2, 2, 4-Trimethylpentane, Pesticidegrade | | | |
| | Water | 0.02% max. | 1 Liter | 16.00 |



MONS 067568

# Analabs ANNOUNCES for 1974
# A POLLUTION CHEMICAL STANDARD CATALOG

**Contains:**

**PCT's**

**PCB's**

**PCN's**

**Dioxins**

**Pesticides**

**Dibenzo Furans**

**Polynuclear Hydrocarbons**




Send in your name and address for a copy of our Pollution Control Catalog. This catalog will only be sent on request. Available February, 1974.

***It's That Time Again*** . . . Time to update our mailing list. If there are any changes to be made to your name and address please return the name tag to us, with any changes. If you are not interested in this publication, please circle the return requested notice in the lower left hand corner and we will pay the postage.

**Analabs, Inc.** A subsidiary of NEN New England Nuclear

80 Republic Drive, North Haven, Connecticut 06473
Telephone: (203) 288-8463 Telex: 96-3318

BULK RATE
U. S. POSTAGE
PAID
North Haven, Conn.
Permit No. 20

T I B

R E KELLER G LDR -
MONSANTO
[redacted]
ST LOUIS MO

63104

RETURN REQUESTED

MONS 067569