

PFOS:  
A DIETARY LC50 STUDY WITH THE NORTHERN BOBWHITE

WILDLIFE INTERNATIONAL LTD. PROJECT NO.: 454-103  
3M LAB REQUEST NO.: U2723

U.S. Environmental Protection Agency  
Series 850 – Ecological Effects Test Guidelines (draft)  
OPPTS Number 850.2200

FIFRA Subdivision E, Section 71-2

OECD Guideline 205

AUTHORS: Sean P. Gallagher  
Courtney S. Casey  
Joann B. Beavers  
Raymond L. Van Hoven

STUDY INITIATION: April 21, 1999

STUDY COMPLETION: April 26, 2000

SUBMITTED TO

3M Corporation  
Environmental Laboratory  
935 Bush Avenue  
St. Paul, Minnesota 55144

***Wildlife International, Ltd.***

8598 Commerce Drive  
Easton, Maryland 21601  
(410) 822-8600

Page 1 of 62

**Exhibit  
2800**

State of Minnesota v. 3M Co.,  
Court File No. 27-CV-10-28862

3MA00233956

2800.0001

- 2 -

## GOOD LABORATORY PRACTICE COMPLIANCE STATEMENT

SPONSOR: 3M Corporation

TITLE: PFOS: A Dietary LC50 Study with the Northern Bobwhite

WILDLIFE INTERNATIONAL LTD. PROJECT NO.: 454-103

3M LAB REQUEST NO.: U2723

STUDY COMPLETION: April 26, 2000

This study was conducted in compliance with Good Laboratory Practice Standards as published by the U.S. Environmental Protection Agency, 40 CFR Part 160 and 792, 17 August 1989; OECD Principles of Good Laboratory Practice, (OCDE/GD(92) 32, Environment Monograph No. 45, Paris, 1992); and Japan MAFF, 59 NohSan, Notification No. 3850, Agricultural Production Bureau, 10 August 1984 with the following exception:

The test substance was not characterized in accordance with full GLP compliance; however, the characterization was performed according to 3M Standard Operating Procedures and Methods, and all raw data are being maintained in the 3M archives. The test substance is being recharacterized in accordance with GLP.

The stability of the test substance and reference standard under conditions of storage at the test site was not determined in accordance with Good Laboratory Practice Standards.

STUDY DIRECTOR:

Sean P. Gallagher  
Sean P. Gallagher  
Senior Biologist

DATE

4/26/00SPONSOR'S REPRESENTATIVE

Susan A. Beach  
Ms. Susan A. Beach

DATE

4/27/00

- 3 -

## QUALITY ASSURANCE STATEMENT

This study was examined for compliance with Good Laboratory Practice Standards as published by the U.S. Environmental Protection Agency, 40 CFR Part 160 and 792, 17 August 1989; OECD Principles of Good Laboratory Practice, (OCDE/GD (92) 32, Environment Monograph No. 45, Paris, 1992); and Japan MAFF, 59 NohSan, Notification No. 3850, Agricultural Production Bureau, 10 August 1984. The dates of all audits and inspections and the dates that any findings were reported to the Study Director and Laboratory Management were as follows:

| ACTIVITY                                      | DATE CONDUCTED              | DATE REPORTED TO: |                    |
|-----------------------------------------------|-----------------------------|-------------------|--------------------|
|                                               |                             | STUDY DIRECTOR    | MANAGEMENT         |
| Test Substance Prep.<br>& Analytical Sampling | April 22, 1999              | April 22, 1999    | April 23, 1999     |
| Matrix Fortification                          | April 22, 1999              | April 22, 1999    | April 23, 1999     |
| Feed Consumption<br>& Analytical Sampling     | April 27, 1999              | April 27, 1999    | May 4, 1999        |
| Analytical Data and<br>Draft Report           | July 7, 8, 9, 1999          | July 9, 1999      | July 16, 1999      |
| Biology Data and<br>Draft Report              | August 26, 27, 30, 31, 1999 | August 31, 1999   | September 13, 1999 |
| Final Report                                  | April 17-18, 2000           | April 18, 2000    | April 19, 2000     |

Susan L. Coleman  
Susan L. Coleman  
Senior Quality Assurance Representative

DATE 4-19-00

- 4 -

## REPORT APPROVAL


SPONSOR: 3M Corporation


TITLE: PFOS: A Dietary LC50 Study with the Northern Bobwhite

WILDLIFE INTERNATIONAL LTD. PROJECT NO.: 454-103

3M LAB REQUEST NO.: U2723


STUDY DIRECTOR:

  
Sean P. Gallagher  
Senior Biologist, Avian Toxicology


  
Date


CHEMISTRY PRINCIPAL INVESTIGATOR:

  
Raymond L. Van Hoven, Ph.D  
Scientist, Analytical Chemistry

  
Date

REPORT APPROVED BY:

  
Joann B. Beavers  
Director, Avian Toxicology

  
Date

  
Willard B. Nixon, Ph.D.  
Manager, Analytical Chemistry

  
Date

## TABLE OF CONTENTS

|                                                     |         |
|-----------------------------------------------------|---------|
| TITLE PAGE.....                                     | Page 1  |
| GOOD LABORATORY PRACTICE COMPLIANCE STATEMENT ..... | Page 2  |
| QUALITY ASSURANCE STATEMENT .....                   | Page 3  |
| REPORT APPROVAL.....                                | Page 4  |
| TABLE OF CONTENTS .....                             | Page 5  |
| Tables and Appendices .....                         | Page 6  |
| SUMMARY .....                                       | Page 7  |
| INTRODUCTION.....                                   | Page 8  |
| OBJECTIVE .....                                     | Page 8  |
| MATERIALS AND METHODS .....                         | Page 8  |
| Test Substance.....                                 | Page 8  |
| Treatment Groups.....                               | Page 9  |
| Duration of the Test.....                           | Page 9  |
| Test Birds .....                                    | Page 9  |
| Animal Diet.....                                    | Page 10 |
| Diet Preparation.....                               | Page 10 |
| Diet Sampling.....                                  | Page 10 |
| Analytical Method .....                             | Page 11 |
| Housing and Environmental Conditions .....          | Page 13 |
| Observations.....                                   | Page 14 |
| Animal Body Weights/Feed Consumption.....           | Page 14 |
| Gross Necropsy .....                                | Page 14 |
| Statistical Analyses .....                          | Page 14 |
| RESULTS .....                                       | Page 15 |
| Diet Analysis .....                                 | Page 15 |
| Mortalities and Clinical Observations.....          | Page 15 |
| Body Weight and Feed Consumption.....               | Page 17 |
| Gross Necropsy .....                                | Page 17 |
| CONCLUSION.....                                     | Page 18 |
| REFERENCES .....                                    | Page 19 |

## TABLE OF CONTENTS

- continued -

## TABLES AND APPENDICES

|                |                                                                                                                              |         |
|----------------|------------------------------------------------------------------------------------------------------------------------------|---------|
| TABLE 1:       | Cumulative Mortality from a Northern Bobwhite Acute Dietary Toxicity Study with PFOS.....                                    | Page 20 |
| TABLE 2:       | Mean Body Weight (g) from a Northern Bobwhite Acute Dietary Toxicity Study with PFOS.....                                    | Page 21 |
| TABLE 3:       | Mean Feed Consumption (g/bird/day) from a Northern Bobwhite Acute Dietary Toxicity Study with PFOS.....                      | Page 23 |
| TABLE 4:       | Group Gross Pathological Observations from a Northern Bobwhite Acute Dietary Toxicity Study with PFOS.....                   | Page 25 |
| TABLE 5:       | Cumulative Mortality (Estimated Cumulative Dose, mg/kg) from a Northern Bobwhite Acute Dietary Toxicity Study with PFOS..... | Page 26 |
| APPENDIX I:    | Certificate of Analysis.....                                                                                                 | Page 27 |
| APPENDIX II:   | Diet Formulation.....                                                                                                        | Page 28 |
| APPENDIX III:  | Analytical Methods and Results.....                                                                                          | Page 29 |
| APPENDIX IV:   | Diet Preparation.....                                                                                                        | Page 43 |
| APPENDIX V:    | Cumulative Mortality by Pen from a Northern Bobwhite Acute Dietary Toxicity Study with PFOS .....                            | Page 44 |
| APPENDIX VI:   | Individual Body Weights (g) from a Northern Bobwhite Acute Dietary Toxicity Study with PFOS .....                            | Page 45 |
| APPENDIX VII:  | Feed Consumption (g/bird/day) by Pen from a Northern Bobwhite Acute Dietary Toxicity Study with PFOS .....                   | Page 58 |
| APPENDIX VIII: | Changes to Protocol .....                                                                                                    | Page 61 |
| APPENDIX IX:   | Personnel Involved in the Study .....                                                                                        | Page 62 |

- 7 -

## SUMMARY

SPONSOR: 3M Corporation

TEST SUBSTANCE: PFOS

WILDLIFE INTERNATIONAL LTD. PROJECT NO.: 454-103

STUDY: PFOS: A Dietary LC50 Study with the Northern Bobwhite

## RESULTS:

The dietary LC50 value for northern bobwhite exposed to PFOS was determined to be 220 ppm a.i. with a 95% confidence interval of 164 ppm a.i. to 289 ppm a.i. The slope of the concentration-response curve was 7.005 and the chi-square value was 0.023. The no mortality concentration was 73.2 ppm a.i. Based upon treatment related mortality, signs of toxicity and effects upon body weight gain at the 146 ppm a.i. test concentration, the no-observed-effect concentration was 73.2 ppm a.i.

## TEST DATES:

Hatch - April 12, 1999

Acclimation - April 12-22, 1999

Experimental Start - April 22, 1999

Experimental Termination - May 14, 1999

## NOMINAL TEST

CONCENTRATIONS: 0, 18.3, 36.6, 73.2, 146, 293, 586 and 1171 ppm a.i.

## TEST ANIMALS:

Northern Bobwhite (*Colinus virginianus*)

AGE TEST ANIMALS: 10 days of age at test initiation

SOURCE TEST ANIMALS: Wildlife International Ltd. Production Flock  
8598 Commerce Drive  
Easton, Maryland 21601

STUDY COMPLETION: April 26, 2000

- 8 -

### INTRODUCTION

This study was conducted by Wildlife International Ltd. for 3M Corporation at the Wildlife International Ltd. avian toxicology facility in Easton, Maryland. The in-life portion of the test was conducted from April 22, 1999 to May 14, 1999. Raw data generated at Wildlife International Ltd. and a copy of the final report are filed under Project Number 454-103 in archives located on the Wildlife International Ltd. site.

### OBJECTIVE

The objective of this study was to evaluate the toxicity of a test substance to the Northern Bobwhite (*Colinus virginianus*) administered through the diet for five days. An LC50 value will be calculated, if possible.

### MATERIALS AND METHODS

The methods used in conducting this study are based upon procedures specified in the U.S. Environmental Protection Agency Series 850 - Ecological Effects Test Guidelines OPPTS Number 850.2200 (1), Section 71-2 of the Environmental Protection Agency's Registration Guidelines, Pesticide Assessment Guidelines, FIFRA Subdivision E, Hazard Evaluation: Wildlife and Aquatic Organisms (2); OECD Guideline 205, Guideline for Testing of Chemicals, Avian Dietary Toxicity Test (3); and upon ASTM Standard E857-87, "Standard Practice for Conducting Subacute Dietary Toxicity Tests with Avian Species" (4).

#### Test Substance

The test substance was received from 3M Corporation on October 29, 1998 and was assigned Wildlife International Ltd. Identification Number 4675 upon receipt. The test substance was white powder identified as: FC-95; Lot No.:217. The reported purity of the test substance was 98.9%, with an expiration date of 2008. Following test termination, the test material was reanalyzed. The results of reanalysis indicate a test substance purity of 90.49%. All test concentrations have been adjusted to reflect the purity reported on the new Certificate of Analysis (Appendix I). The test substance was stored under ambient conditions.

The internal standard was received from 3M Corporation on July 2, 1998 and was assigned Wildlife International Ltd. identification number 4526 upon receipt. The internal standard, a granular



material, was identified as: 1H, 1H, 2H, 2H Perfluorooctane Sulfonic Acid, Chemical Abstract Number: 27619-97-2. The standard was stored under ambient conditions.

#### Treatment Groups

The test consisted of a geometric series of seven test concentrations and a control group. Thirty northern bobwhite chicks were assigned to the control group and ten northern bobwhite chicks were assigned to each of the treatment groups. The birds were sorted by weight, then chosen indiscriminately from within each represented weight class for placement into control and treatment groups. The birds were housed in brooding pens containing five chicks each. Nominal dietary concentrations used in this study were 0, 18.3, 36.6, 73.2, 146, 293, 586 and 1171 parts per million active ingredient (ppm a.i.) of PFOS. The dietary concentrations were established based upon known toxicity data and information supplied by the Sponsor.

Each group was fed the appropriate test or control diet for five days. During the exposure period the control group received untreated feed. Following the five-day exposure period all groups were given untreated basal diet for three days. On Day 8, half of the surviving treatment and control birds were euthanized and liver tissue, blood, and bile samples were collected for analysis. The remaining birds were fed basal ration until Day 22. On Day 22, these birds were euthanized and also sampled for liver weight, blood, and bile.

#### Duration of the Test

The primary phases of this test and their durations were:

1. Acclimation - 10 days.
2. Exposure - 5 days.
3. Post-exposure observation - 3 or 17 days

#### Test Birds

All northern bobwhite (*Colinus virginianus*) were 10 days of age and appeared to be in good health at initiation of the test. The birds were obtained from Wildlife International Ltd. Production Flock, Easton, MD and were hatched on April 12, 1999. Birds ranged in weight from 18 to 23 grams at test initiation. The birds used in this study were immature and could not be differentiated by sex.

All birds were from the same hatch, pen-reared and phenotypically indistinguishable from wild birds. All birds were acclimated to the caging and facilities from the day of hatch until initiation of the test.

#### Animal Diet

Throughout acclimation and testing all test birds were fed a game bird ration formulated to Wildlife International Ltd.'s specifications (Appendix II). The chicks were given a vitamin supplement in their water from the day they were hatched until the initiation of the test. Water from the town of Easton public water supply, and feed were provided *ad libitum* during acclimation and testing. The birds received no form of antibiotic medication during acclimation or testing.

#### Diet Preparation

The test substance was mixed directly into the ration. Mixing was done with a Hobart mixer (Model Number AS200T). All dietary test concentrations were adjusted to 100% PFOS based upon the reported purity of the test substance. All dietary concentrations and the LC50 value are reported as ppm a.i. in the diet. Nominal dietary test concentrations used in this study were 18.3, 36.6, 73.2, 146, 293, 586, 1171 ppm a.i. (Appendix IV).

#### Diet Sampling

Samples of the test diets were collected to verify the test concentrations administered and to confirm the stability and homogeneity of the test substance in the diets. Homogeneity of the test substance in the diet was evaluated by collecting six samples from the 18.3 ppm a.i. test diet and six samples from the 1171 ppm a.i. test diet at preparation on Day 0. Homogeneity samples were collected from the top, middle and bottom of the left and right sections of the mixing vessel. The homogeneity samples also served as verification samples. One verification sample was collected from the control diet and two verification samples were collected from each remaining treatment group at preparation on Day 0. At the end of the exposure period (Day 5), one sample was collected from the control and two samples were collected from each treatment group to determine stability of the test substance in the diet under test conditions. The stability samples were collected from feed remaining in the feeders after being at ambient test pen conditions for five days. Samples were transferred immediately to Wildlife International Ltd. analytical chemistry.

### Analytical Method

The method used for the analysis of the avian diet samples was based upon methodology developed at Wildlife International Ltd. and entitled "Method Outline for the Determination of PFOS in Avian Feed".

Avian diet samples were extracted with methanol. Methanol was added to a requisite quantity of feed contained in a French-square glass bottle. Bottles were capped and shaken on a shaker table. Samples were vacuum filtered using qualitative filter paper. The retained feed was rinsed three times with methanol into the filtrate. The filtrate was transferred to a volumetric flask and brought to volume with methanol. As appropriate, samples were further diluted with methanol. Each sample then was diluted with a 50% methanol : 50% NANOpure® water solution containing 0.100 mg 4H PFOS (internal standard)/L and 0.05% formic acid (v/v) so that they fell within the calibration range of the PFOS methodology. A method flowchart is provided in Appendix III, Figure 1.

Concentrations of PFOS in the standards and samples were determined by reversed-phase high performance liquid chromatography using a Hewlett-Packard Model 1100 High Performance Liquid Chromatograph (HPLC) with a Perkin-Elmer API 100LC Mass Spectrometer equipped with a Perkin-Elmer TurboIonSpray ion source. HPLC separations were achieved using a Keystone Betasil C<sub>18</sub> analytical column (100 mm x 2 mm I.D., 3 µm particle size). The instrument parameters are summarized in Appendix III, Table 1.

Calibration standards of PFOS prepared in a 50% methanol : 50% NANOpure® water solution containing 0.100 mg 4H PFOS (internal standard)/L and 0.05% formic acid (v/v), ranging in concentration from 0.00229 to 0.0457 mg a.i./L were analyzed with the samples. The same and most prominent peak response for PFOS was utilized to monitor PFOS in all calibration, quality control, and study samples. No attempt was made to quantify PFOS on the basis of individual isomeric components. Linear regression equations were generated using peak area response ratios (PFOS : internal standard) versus the respective concentration ratios (PFOS : internal standard) of the calibration standards. A typical calibration curve is presented in Appendix III, Figure 2. The concentration of PFOS in the samples was determined by substituting the peak area response ratios into the applicable linear regression equation. Representative ion chromatograms of low and high calibration standards are presented in Appendix III, Figures 3 and 4, respectively.

The method limit of quantitation (LOQ) for these analyses was set at 1.15 ppm a.i. calculated as the product of the lowest calibration standard analyzed (0.00229 mg a.i./L) and the dilution factor of the matrix blank samples (500).

Two matrix blank samples were analyzed to determine possible interferences. No interferences were observed at or above the LOQ during sample analyses (Appendix III, Table 2). An interference in the feed appeared at approximately the same retention time as the peak of interest but it was well below the LOQ. A representative chromatogram of a matrix blank is presented in Appendix III, Figure 5.

Avian diet was fortified at 4.57, 183 and 1830 ppm a.i. and analyzed concurrently with the samples to determine the mean procedural recovery (Appendix III, Table 3). Sample concentrations were not corrected for the mean procedural recovery of 94.7%. A representative chromatogram of a matrix fortification is presented in Appendix III, Figure 6.

An example calculation is presented for sample number 454-103-2, nominal concentration of 18.3 ppm a.i. in avian diet.

Initial Weight: 10.0 g

Final Volume: 200 mL

Dilution Factor: 100 (intermediate dilution factor  $\times$  final dilution factor)

PFOS Peak Area: 113568

Internal Standard Peak Area: 413160

Peak Area Ratio: 0.2749

Calibration curve equation.

Slope: 2.77397

Intercept: 0.01894

Curve is weighted (1/x).

$$\text{PFOS (mg a.i./L) at instrument} = \frac{(\text{Peak area ratio} - (\text{Y-intercept})) \times \text{I.S. Concentration}}{\text{Slope}}$$

- 13 -

$$= \frac{(0.2749 - 0.01894) \times 0.100 \text{ mg/L}}{2.77397}$$

$$= 0.00923 \text{ mg a.i./L}$$

Note: I.S. = internal standard.

$$\text{PFOS (ppm a.i.) in sample} = \frac{\text{PFOS (mg a.i./L) at instrument} \times \text{Final Volume (L)} \times \text{Dilution Factor}}{\text{Initial Weight (Kg)}}$$

$$= \frac{0.00923 \times 0.200 \times 100}{0.01}$$

$$= 18.5 \text{ ppm a.i.}$$

$$\text{Percent of Nominal Concentration} = \frac{\text{PFOS (ppm a.i.) in sample}}{\text{PFOS (ppm a.i.) nominal}} \times 100$$

$$= \frac{18.5}{18.3} \times 100 = 101\%$$

#### Housing and Environmental Conditions

During acclimation and testing, all birds were housed indoors in batteries of thermostatically controlled brooding pens manufactured by Beacon Steel Products Co. (Model No. B735Q). Each pen had floor space that measured approximately 72 X 90 cm. Ceiling height was approximately 23 cm. External walls, ceilings and floors were constructed of galvanized steel wire and sheeting. Birds were sorted by weight, then chosen indiscriminately from within each represented weight class for assignment to pens. Each group of birds was identified by pen number and test concentration. Individual birds were identified by leg bands.

During the test the average temperature in the brooding compartment of the pens was  $38^{\circ}\text{C} \pm 2^{\circ}\text{C}$  (SD). Average ambient room temperature for this study was  $27.3^{\circ}\text{C} \pm 1.2^{\circ}\text{C}$  (SD) with an average relative humidity of  $31\% \pm 14\%$  (SD). The photoperiod (maintained by a time clock) was sixteen hours of light per day during acclimation and throughout the test. The light source was fluorescent lights which closely approximate noon-day sunlight. The birds were exposed to an average of

- 14 -

approximately 139 lux of illumination. Housing and husbandry practices were based on guidelines established by the National Research Council (5).

#### Observations

During acclimation all birds were observed daily. Birds exhibiting abnormal behavior or physical injury were not used. Following test initiation and continuing until termination, all birds were typically observed at least twice daily. A record was maintained of all mortality, signs of toxicity, and abnormal behavior.

#### Animal Body Weights/Feed Consumption

Individual body weights were measured at the initiation of the test, on Day 5, Day 8, and on Days 15 and Day 22 for all remaining birds. Average feed consumption values during the exposure period (Days 0-5) and the post-exposure observation period (Days 6-8) were determined by pen for each treatment group and the control group. Additionally, feed consumption was determined for Days 8-15 and 15-22 for the remaining treatment and control birds. Feed consumption was determined by measuring the change in the weight of the feed presented to the birds over a given period of time. The accuracy of feed consumption values may have been affected by the unavoidable wastage of feed by the birds.

#### Gross Necropsy

All test birds that died during the course of the test and all birds remaining at the termination were subjected to a gross necropsy. Additionally, livers were weighed and liver tissue, blood, and bile were collected from birds euthanized on Day 8 and 22, and when possible from those that died during the course of the study.

#### Statistical Analyses

Mortality data were analyzed using the computer program of C.E. Stephan (6). The program was designed to calculate the LC50 value and the 95% confidence interval by probit analysis, moving average method or the binomial probability method (7,8,9). In this study, the LC50 value was determined using the probit method. The slope of the concentration-response curve and results of the goodness of fit test are reported. Body weight data were compared by Dunnett's test using TOXSTAT software (10,11). No statistical analyses were applied to feed consumption data.

## RESULTS

### Diet Analysis

Avian diet samples were collected from the 18.3 and 1171 ppm a.i. test concentrations and analyzed to evaluate homogeneity of the test substance in the avian diet. The analysis of these samples also served as verification of test substance concentrations. Resulting mean measured concentrations, standard deviations and coefficients of variation (CV) for these test concentrations were  $19.5 \pm 2.13$  ppm a.i. (CV = 10.9%) and  $1196 \pm 70.2$  ppm a.i. (CV = 5.87%), respectively (Appendix III, Table 4). Control avian diet samples collected during the test showed no interferences above the LOQ. Samples collected during the test to verify the 36.6, 73.2, 146, 293 and 586 ppm a.i. test substance concentrations had mean measured concentrations of 40.2, 74.5, 174, 291 and 537 ppm a.i., respectively. These values represented 110, 102, 119, 99.3 and 91.6% of the nominal concentrations, respectively (Appendix III, Table 5). Analysis of avian diet samples collected from feeders after being held at ambient temperature for five days averaged 101, 122, 104, 101, 109, 114 and 102% of the Day 0 values for the 18.3, 36.6, 73.2, 146, 293, 586 and 1171 ppm a.i. test substance concentrations, respectively (Appendix III, Table 6). A representative chromatogram of a test sample is shown in Appendix III, Figure 7.

### Mortalities and Clinical Observations

One incidental mortality occurred in the control group during the course of the study (Table 1 and Appendix V). On the morning of Day 5, one bird was noted with a broken leg and was subsequently euthanized on Day 6. Additionally, two birds in the control group were intermittently noted with foot lesions associated with cage mate aggression. Otherwise, all control birds were normal in appearance and behavior throughout the test.

No treatment related mortalities or overt signs of toxicity were observed in the 18.3, 36.6, or 73.2 ppm a.i. treatment groups. One bird in the 18.3 ppm a.i. treatment group was noted as lame from Day 6 through Day 8, and blood of undetermined origin was noted on the underside of one bird from the 73.2 ppm a.i. treatment group on Day 22. Otherwise, all birds in the 18.3, 36.6 and 73.2 ppm a.i. treatment groups were normal in appearance and behavior throughout the test period.

- 16 -

There was 11% (1 of 9) mortality in the 146 ppm a.i. treatment group, 80% (8 of 10) mortality in the 293 ppm a.i. treatment group and 100% (10 of 10) mortality in the 586 and 1171 ppm a.i. treatment groups. In the 146 ppm a.i. treatment group, one bird was euthanized on Day 3 after sustaining a broken leg. This incidental mortality was not used in the calculation of the LC50 value. Additionally, there was one treatment-related mortality in the 146 ppm a.i. treatment group, a bird found dead on the morning of Day 7. Clinical signs of toxicity were observed in this treatment group on Day 5, when two birds displayed wing droop. All other birds at this test concentration were normal in appearance and behavior for the duration of the test.

In the 293 ppm a.i. treatment group there were eight treatment-related mortalities, occurring on Days 5, 6 and 7. Signs of toxicity were first observed on the morning of Day 4 and continued to be exhibited through the morning of Day 8 for the single bird euthanized on Day 8, and through the afternoon of Day 8 for the single bird surviving until Day 22. Signs of toxicity included a ruffled appearance, reduced reaction to stimuli (sound and motion), lethargy, wing droop, loss of coordination, lower limb weakness and convulsions. The single remaining bird appeared to have recovered and was normal in appearance and behavior from the afternoon of Day 9 until test termination.

In the 586 ppm a.i. treatment group mortalities were first noted on Day 3 and continued to be observed through Day 7, at which point all birds had died. Overt signs of toxicity were first observed on the afternoon of Day 2 and continued through the morning of Day 7, when the final birds were found dead. Signs of toxicity observed among birds in the 586 ppm a.i. treatment group included a ruffled appearance, reduced reaction to stimuli (sound and motion), lethargy, depression, wing droop, loss of coordination, lower limb weakness, lower limb rigidity, prostrate posture, and convulsions.

In the 1171 ppm a.i. treatment group 100% mortality had occurred by the morning of Day 4. Signs of toxicity in the 1171 ppm a.i. treatment group were first observed on the afternoon of Day 2, with the first mortalities noted on the morning of Day 3. Signs of toxicity observed prior to death included a ruffled appearance, reduced reaction to stimuli (sound and motion), lethargy, depression, loss of coordination, wing droop, and lower limb weakness and rigidity.



#### Body Weight and Feed Consumption

When compared to the control group, there were no apparent treatment related effects upon body weight among birds in the 18.3, 36.6 or 73.2 ppm a.i. treatment groups. However, there was a concentration responsive reduction in body weight gain or body weight loss in the 146, 293 and 586 ppm a.i. treatment groups during the exposure period (Days 0-5) (Table 2 and Appendix VI). Differences from the control group were statistically significant at  $p < 0.05$  for the 146 ppm a.i. level and at  $p < 0.01$  for the 293 and 586 ppm a.i. levels. A statistically significant ( $p < 0.01$ ) reduction in body weight gain continued to be observed at the 146 ppm a.i. test concentration through Day 8 of the study. At the 293 ppm a.i. concentration a statistically significant ( $p < 0.01$ ) mean weight loss continued through Day 8 of the study and a marked reduction in weight gain was noted through Day 15 of the study. Due to total mortality, body weight effects could not be determined for the 1171 ppm a.i. level during the exposure period or for the 586 and 1171 ppm a.i. treatment groups for post-exposure period.

There were no apparent treatment related effects upon feed consumption at the 18.3, 36.6, 73.2 or 146 ppm a.i. test concentrations (Table 3 and Appendix VII). However, a reduction in feed consumption was noted at the 293, 586 and 1171 ppm a.i. treatment groups during the exposure period (Days 0-5). There were no treatment-related effects on feed consumption in any of the surviving treatment groups during the Day 6-8 post-exposure period. In the 293 ppm a.i. treatment group only one bird survived to Day 22. The reduction in feed consumption observed at the 293 ppm a.i. test concentration during both the Day 8-15 and 15-22 post-exposure periods was the result of having only one bird in the pen, and was not considered to be treatment related.

#### Gross Necropsy

During the course of the test, all birds that died were subjected to a gross necropsy. Necropsy results for birds found dead were similar. Common observations included thin condition, loss of muscle mass, altered spleen color, autolysis of tissues and pale organs. Details of the necropsy findings are presented in Table 4.

Half of the surviving birds were subjected to gross necropsy on Day 8 and the remaining birds were necropsied on Day 22, following test termination. On Day 8, one bird in the 73.2 ppm a.i. treatment group was noted with a slightly pale liver. Due to the isolated nature of this finding, it was

- 18 -

not considered to be related to treatment. The single bird euthanized from the 293 ppm a.i. treatment group was observed at necropsy to have with a lack of muscle mass and general thinness. Since these findings correlated with an impact upon body weight noted at this concentration, the findings were considered to be treatment related. Necropsy results were unremarkable for all other birds euthanized on Day 8. Similarly, Day 22 necropsy findings were unremarkable for all birds.

#### CONCLUSION

The dietary LC50 value for northern bobwhite exposed to PFOS was determined to be 220 ppm a.i. with a 95% confidence interval of 164 ppm a.i. to 289 ppm a.i. The slope of the concentration-response curve was 7.005 and the chi-square value was 0.023. The no mortality concentration was 73.2 ppm a.i. Based upon treatment related mortality, signs of toxicity and effects upon body weight gain at the 146 ppm a.i. a.i test concentration, the no-observed-effect concentration was 73.2 ppm a.i..

REFERENCES

1. U.S. Environmental Protection Agency, 1996. Series 850 – Ecological Effects Test Guidelines (draft), OPPTS Number 850.2200: Avian Dietary Toxicity Test.
2. U.S. Environmental Protection Agency. 1982. *Pesticide Assessment Guidelines, FIFRA Subdivision E, Hazard Evaluation: Wildlife and Aquatic Organisms*, subsection 71-2, Environmental Protection Agency, Office of Pesticide Programs. Washington, D.C.
3. Organization for Economic Cooperation and Development. 1984. *Avian Dietary Toxicity Test*. OECD Guideline for Testing of Chemicals. Guideline 205. Paris.
4. American Society for Testing and Materials. 1987. Standard Practice for Conducting Subacute Dietary Toxicity Tests with Avian Species. ASTM Standard E857-87. Annual Book of ASTM Standards, Vol. 11.04. Philadelphia, PA.
5. National Research Council. 1996. *Guide for the Care and Use of Laboratory Animals*. Washington, D.C. National Academy Press. 125 pp.
6. Stephan, C.E. 1978. U.S. EPA, Environmental Research Laboratory, Duluth, MN. Personal Communication
7. Stephan, C.E. 1977. Methods for Calculating an LC50. Pages 65-84 *In Aquatic Toxicology and Hazard Evaluations*, American Society for Testing and Materials. Pub. No. STP 634. Philadelphia, PA.
8. Finney, D.J. 1971. *Statistical Methods in Biological Assay*, 2nd ed., Griffin Press, London.
9. Thompson, W.R. 1947. *Bacteriological Reviews*, Vol 2, No.2: 115-145.
10. Dunnett, C.W. 1955. A Multiple Comparison's Procedure for Comparing Several Treatments with a Control. *Jour. Amer. Statis. Assoc.* 50:1096-1121.
11. Gulley, D.D. 1990. TOXSTAT Release 3.2. The University of Wyoming.

- 20 -

TABLE 1

Cumulative Mortality from a Northern Bobwhite Acute Dietary Toxicity Study with PFOS

| Experimental Group<br>(ppm a.i.) | No. Dead Per No. Exposed<br>Exposure Period |       |       |                   |       |       | No. Dead Per No. Exposed<br>Post-Exposure Period |       |                    |
|----------------------------------|---------------------------------------------|-------|-------|-------------------|-------|-------|--------------------------------------------------|-------|--------------------|
|                                  | Day 0                                       | Day 1 | Day 2 | Day 3             | Day 4 | Day 5 | Day 6                                            | Day 7 | Day 8 <sup>1</sup> |
| <b>Control</b>                   |                                             |       |       |                   |       |       |                                                  |       |                    |
| 0                                | 0/30                                        | 0/30  | 0/30  | 0/30              | 0/30  | 0/30  | 1/30                                             | 1/30  | 1/30               |
| <b>Treatment</b>                 |                                             |       |       |                   |       |       |                                                  |       |                    |
| 18.3                             | 0/10                                        | 0/10  | 0/10  | 0/10              | 0/10  | 0/10  | 0/10                                             | 0/10  | 0/10               |
| 36.6                             | 0/10                                        | 0/10  | 0/10  | 0/10              | 0/10  | 0/10  | 0/10                                             | 0/10  | 0/10               |
| 73.2                             | 0/10                                        | 0/10  | 0/10  | 0/10              | 0/10  | 0/10  | 0/10                                             | 0/10  | 0/10               |
| 146                              | 0/10                                        | 0/10  | 0/10  | 1/10 <sup>2</sup> | 0/9   | 0/9   | 0/9                                              | 1/9   | 1/9                |
| 293                              | 0/10                                        | 0/10  | 0/10  | 0/10              | 0/10  | 2/10  | 4/10                                             | 8/10  | 8/10               |
| 586                              | 0/10                                        | 0/10  | 0/10  | 1/10              | 2/10  | 5/10  | 8/10                                             | 10/10 | 10/10              |
| 1171                             | 0/10                                        | 0/10  | 0/10  | 3/10              | 10/10 | 10/10 | 10/10                                            | 10/10 | 10/10              |

The LC50 value was calculated to be 220 ppm a.i. with a 95% confidence interval of 164 ppm a.i. to 289 ppm a.i..

<sup>1</sup> - No mortalities occurred in any of the control or treatment groups from Day 8 to Day 22.

<sup>2</sup> - Bird euthanized on day 3 after sustaining a broken leg.

- 21 -

TABLE 2  
Page 1

Mean Body Weight (g) from a Northern Bobwhite Acute Dietary Toxicity Study with PFOS

| Experimental Group<br>(ppm a.i.) |      | Exposure Period |       |                                | Post-Exposure Period |                                | Total Change <sup>1</sup> |
|----------------------------------|------|-----------------|-------|--------------------------------|----------------------|--------------------------------|---------------------------|
|                                  |      | Day 0           | Day 5 | Change <sup>1</sup><br>Day 0-5 | Day 8                | Change <sup>1</sup><br>Day 5-8 |                           |
| <b>Control</b>                   |      |                 |       |                                |                      |                                |                           |
| 0                                | Mean | 20              | 30    | 10                             | 38                   | 8                              | 18                        |
|                                  | SD   | 1               | 4     | 3                              | 5                    | 2                              | 4                         |
| <b>Treatment</b>                 |      |                 |       |                                |                      |                                |                           |
| 18.3                             | Mean | 21              | 31    | 11                             | 40                   | 9                              | 20                        |
|                                  | SD   | 1               | 4     | 3                              | 5                    | 2                              | 3                         |
| 36.6                             | Mean | 20              | 31    | 11                             | 39                   | 8                              | 19                        |
|                                  | SD   | 2               | 3     | 2                              | 3                    | 1                              | 2                         |
| 73.2                             | Mean | 20              | 30    | 9                              | 37                   | 7                              | 16                        |
|                                  | SD   | 1               | 2     | 1                              | 3                    | 1                              | 2                         |
| 146                              | Mean | 20              | 27*   | 7*                             | 33*                  | 6**                            | 13**                      |
|                                  | SD   | 2               | 3     | 3                              | 3                    | 2                              | 4                         |
| 293                              | Mean | 20              | 18**  | -2**                           | 18**                 | -2**                           | -1**                      |
|                                  | SD   | 1               | 2     | 2                              | 4                    | 4                              | 5                         |
| 586                              | Mean | 20              | 16**  | -4**                           | -                    | -                              | -                         |
|                                  | SD   | 1               | 2     | 2                              | -                    | -                              | -                         |
| 1171                             | Mean | 20              | -     | -                              | -                    | -                              | -                         |
|                                  | SD   | 1               | -     | -                              | -                    | -                              | -                         |

<sup>1</sup>Mean change is calculated separately from the mean body weights using individual body weights (See Appendix VI).

(-) = No data available due to mortality.

\*Statistically different from the control group at  $p < 0.05$  (Dunnett's t-test).\*\*Statistically different from the control group at  $p < 0.01$  (Dunnett's t-test).

- 22 -

TABLE 2

Page 2

Mean Body Weight (g) from a Northern Bobwhite Acute Dietary Toxicity Study with PFOS

| Experimental Group<br>(ppm a.i.) |                    | Post-Exposure Period |        |                                 |        |                                  |                                     |
|----------------------------------|--------------------|----------------------|--------|---------------------------------|--------|----------------------------------|-------------------------------------|
|                                  |                    | Day 8                | Day 15 | Change <sup>1</sup><br>Day 8-15 | Day 22 | Change <sup>1</sup><br>Day 15-22 | Total<br>Change <sup>1</sup> (8-22) |
| Control                          |                    |                      |        |                                 |        |                                  |                                     |
| 0                                | Mean               | 37                   | 59     | 23                              | 82     | 22                               | 45                                  |
|                                  | SD                 | 6                    | 10     | 5                               | 13     | 3                                | 8                                   |
| Treatment                        |                    |                      |        |                                 |        |                                  |                                     |
| 18.3                             | Mean               | 40                   | 68     | 24                              | 87     | 23                               | 47                                  |
|                                  | SD                 | 6                    | 8      | 3                               | 7      | 3                                | 3                                   |
| 36.6                             | Mean               | 38                   | 65     | 26                              | 89     | 24                               | 50                                  |
|                                  | SD                 | 3                    | 5      | 2                               | 7      | 2                                | 4                                   |
| 73.2                             | Mean               | 35                   | 60     | 24                              | 79     | 20                               | 44                                  |
|                                  | SD                 | 3                    | 4      | 2                               | 4      | 2                                | 2                                   |
| 146                              | Mean               | 34*                  | 58     | 24                              | 79     | 21                               | 45                                  |
|                                  | SD                 | 3                    | 3      | 1                               | 2      | 2                                | 1                                   |
| 293                              | Mean               | 21                   | 35     | 14                              | 55     | 20                               | 34                                  |
|                                  | <sup>2</sup><br>SD | -                    | -      | -                               | -      | -                                | -                                   |
| 586                              | Mean               | -                    | -      | -                               | -      | -                                | -                                   |
|                                  | SD                 | -                    | -      | -                               | -      | -                                | -                                   |
| 1171                             | Mean               | -                    | -      | -                               | -      | -                                | -                                   |
|                                  | SD                 | -                    | -      | -                               | -      | -                                | -                                   |

<sup>1</sup>Mean change is calculated separately from the mean body weights using individual body weights (See Appendix VI).<sup>2</sup>n=1, could not be evaluated statistically with Dunnett's t-test.

(-) = No data available due to mortality.

\*Statistically different from the control group at  $p < 0.05$  (Dunnett's t-test).

- 23 -

TABLE 3  
Page 1Mean Feed Consumption (g/bird/day) from a Northern Bobwhite Acute Dietary  
Toxicity Study with PFOS

| Experimental Group (ppm a.i.)             |      | Exposure Period | Post-Exposure Period |
|-------------------------------------------|------|-----------------|----------------------|
|                                           |      | Day 0-5         | Day 6-8              |
| <b>Control</b>                            |      |                 |                      |
| 0                                         | Mean | 9               | 10                   |
|                                           | SD   | 2               | 2                    |
| <b>Treatment</b>                          |      |                 |                      |
| 18.3                                      |      | 9               | 11                   |
| 36.6                                      |      | 8               | 12                   |
| 73.2                                      |      | 10              | 13                   |
| 146                                       |      | 9               | 10                   |
| 293                                       |      | 5               | 9                    |
| 586                                       |      | 6               | 19                   |
| 1171                                      |      | 4               | -                    |
| (-) = No data available due to mortality. |      |                 |                      |

- 24 -

TABLE 3  
Page 2Mean Feed Consumption (g/bird/day) from a Northern Bobwhite Acute Dietary  
Toxicity Study with PFOS

| Experimental<br>Group<br>(ppm a.i.)       | Post-Exposure Period |           |
|-------------------------------------------|----------------------|-----------|
|                                           | Day 8-15             | Day 15-22 |
| <b>Control</b>                            |                      |           |
| 0                                         | Mean 9               | 13        |
|                                           | SD 2                 | 1         |
| <b>Treatment</b>                          |                      |           |
| 18.3                                      | 10                   | 12        |
| 36.6                                      | 14                   | 15        |
| 73.2                                      | 13                   | 15        |
| 146                                       | 11                   | 14        |
| 293                                       | 8                    | 9         |
| 586                                       | -                    | -         |
| 1171                                      | -                    | -         |
| (-) = No data available due to mortality. |                      |           |



- 25 -

TABLE 4  
Group Gross Pathological Observations  
from a Northern Bobwhite Acute Dietary Toxicity Study with PFOS

Birds that died during the course of the study

| Finding                                     | Control<br>N=1 | Male, Female, and Undetermined<br>PPM A.I. |            |             |              |
|---------------------------------------------|----------------|--------------------------------------------|------------|-------------|--------------|
|                                             |                | 146<br>N=2                                 | 293<br>N=8 | 586<br>N=10 | 1171<br>N=10 |
| Abdominal cavity, some autolysis            | 0              | 0                                          | 2          | 2           | 4            |
| Abdominal cavity, autolysis throughout      | 0              | 0                                          | 0          | 1           | 1            |
| Crop, empty                                 | 0              | 0                                          | 2          | 5           | 2            |
| Emaciated                                   | 0              | 0                                          | 2          | 5           | 8            |
| Fractured leg                               | 1              | 1                                          | 0          | 0           | 0            |
| G.I. tract empty                            | 0              | 0                                          | 1          | 1           | 0            |
| Gizzard contents bile stained               | 0              | 0                                          | 2          | 5           | 1            |
| Heart, anterior portion mottled white color | 0              | 0                                          | 1          | 0           | 0            |
| Heart, pale                                 | 0              | 0                                          | 0          | 2           | 1            |
| Intestinal contents tar-like                | 0              | 0                                          | 0          | 2           | 0            |
| Keel, prominent                             | 0              | 0                                          | 1          | 3           | 10           |
| Kidneys, pale                               | 0              | 0                                          | 0          | 2           | 0            |
| Liver, pale and mottled                     | 0              | 1                                          | 0          | 0           | 0            |
| Loss of muscle mass                         | 0              | 0                                          | 4          | 7           | 9            |
| Muscular-skeletal, pale                     | 0              | 1                                          | 0          | 0           | 0            |
| Small in stature                            | 0              | 0                                          | 3          | 0           | 0            |
| Spleen, black                               | 0              | 0                                          | 0          | 1           | 0            |
| Spleen, dark                                | 0              | 0                                          | 0          | 0           | 2            |
| Spleen, grey                                | 0              | 0                                          | 0          | 1           | 0            |
| Spleen, grey-brown                          | 0              | 0                                          | 0          | 0           | 1            |
| Spleen, pale                                | 0              | 0                                          | 1          | 0           | 1            |
| Spleen, small                               | 0              | 0                                          | 0          | 0           | 1            |
| Spleen, small and pale                      | 0              | 0                                          | 0          | 3           | 0            |
| Thin                                        | 0              | 0                                          | 0          | 4           | 2            |
| Not Remarkable                              | 0              | 0                                          | 1          | 0           | 0            |

- 26 -

Table 5  
Cumulative Mortality (Estimated Cumulative Dose, mg/kg<sup>1</sup>) from a Northern Bobwhite  
Acute Dietary Toxicity Study with PFOS

| Experimental Group<br>(ppm a.i.) | No. Dead Per No. Exposed (Cumulative Dose, mg/kg) |       |           |           |           | No. Dead Per No. Exposed |            |       |       |                    |
|----------------------------------|---------------------------------------------------|-------|-----------|-----------|-----------|--------------------------|------------|-------|-------|--------------------|
|                                  | Exposure Period                                   |       |           |           |           | Post-Exposure Period     |            |       |       |                    |
|                                  | Pen                                               | Day 0 | Day 1     | Day 2     | Day 3     | Day 4                    | Day 5      | Day 6 | Day 7 | Day 8 <sup>2</sup> |
| Control                          |                                                   |       |           |           |           |                          |            |       |       |                    |
| 0                                | 1                                                 | 0/5   | 0/5       | 0/5       | 0/5       | 0/5                      | 0/5        | 0/5   | 0/5   | 0/5                |
| 0                                | 2                                                 | 0/5   | 0/5       | 0/5       | 0/5       | 0/5                      | 0/5        | 0/5   | 0/5   | 0/5                |
| 0                                | 3                                                 | 0/5   | 0/5       | 0/5       | 0/5       | 0/5                      | 0/5        | 0/5   | 0/5   | 0/5                |
| 0                                | 4                                                 | 0/5   | 0/5       | 0/5       | 0/5       | 0/5                      | 0/5        | 0/5   | 0/5   | 0/5                |
| 0                                | 5                                                 | 0/5   | 0/5       | 0/5       | 0/5       | 0/5                      | 0/5        | 1/5   | 1/5   | 1/5                |
| 0                                | 6                                                 | 0/5   | 0/5       | 0/5       | 0/5       | 0/5                      | 0/5        | 0/5   | 0/5   | 0/5                |
| Treatment                        |                                                   |       |           |           |           |                          |            |       |       |                    |
| 18.3                             | 1                                                 | 0/5   | 0/5 (7)   | 0/5 (14)  | 0/5 (21)  | 0/5 (28)                 | 0/5 (35)   | 0/5   | 0/5   | 0/5                |
| 18.3                             | 2                                                 | 0/5   | 0/5 (6)   | 0/5 (12)  | 0/5 (18)  | 0/5 (24)                 | 0/5 (30)   | 0/5   | 0/5   | 0/5                |
| 36.6                             | 1                                                 | 0/5   | 0/5 (13)  | 0/5 (26)  | 0/5 (39)  | 0/5 (52)                 | 0/5 (65)   | 0/5   | 0/5   | 0/5                |
| 36.6                             | 2                                                 | 0/5   | 0/5 (11)  | 0/5 (22)  | 0/5 (33)  | 0/5 (44)                 | 0/5 (55)   | 0/5   | 0/5   | 0/5                |
| 73.2                             | 1                                                 | 0/5   | 0/5 (33)  | 0/5 (66)  | 0/5 (99)  | 0/5 (132)                | 0/5 (165)  | 0/5   | 0/5   | 0/5                |
| 73.2                             | 2                                                 | 0/5   | 0/5 (32)  | 0/5 (64)  | 0/5 (96)  | 0/5 (128)                | 0/5 (160)  | 0/5   | 0/5   | 0/5                |
| 146                              | 1                                                 | 0/5   | 0/5 (46)  | 0/5 (92)  | 0/5 (138) | 0/5 (184)                | 0/5 (230)  | 0/5   | 1/5   | 1/5                |
| 146                              | 2                                                 | 0/5   | 0/5 (49)  | 0/5 (98)  | 1/5 (147) | 1/5 (196)                | 1/5 (245)  | 1/5   | 1/5   | 1/5                |
| 293                              | 1                                                 | 0/5   | 0/5 (66)  | 0/5 (132) | 0/5 (198) | 0/5 (264)                | 1/5 (330)  | 1/5   | 4/5   | 4/5                |
| 293                              | 2                                                 | 0/5   | 0/5 (101) | 0/5 (202) | 0/5 (303) | 0/5 (404)                | 1/5 (505)  | 3/5   | 4/5   | 4/5                |
| 586                              | 1                                                 | 0/5   | 0/5 (213) | 0/5 (429) | 1/5 (639) | 2/5 (852)                | 2/5 (1065) | 4/5   | 5/5   | 5/5                |
| 586                              | 2                                                 | 0/5   | 0/5 (178) | 0/5 (356) | 0/5 (534) | 0/5 (712)                | 3/5 (890)  | 4/5   | 5/5   | 5/5                |
| 1171                             | 1                                                 | 0/5   | 0/5 (256) | 0/5 (512) | 3/5 (768) | 5/5                      | 5/5        | 5/5   | 5/5   | 5/5                |
| 1171                             | 2                                                 | 0/5   | 0/5 (256) | 0/5 (512) | 0/5 (768) | 5/5                      | 5/5        | 5/5   | 5/5   | 5/5                |

The LC50 value was calculated to be approximately 220 ppm a.i. with a 95% confidence interval of 164 to 289 ppm a.i.

\*. Bird was euthanized due to a broken leg.

<sup>1</sup>. Estimated cumulative dose is based upon the average body weight and feed consumption over the 5-day exposure period, and serves as a rough approximation of the actual amount of test substance consumed.

<sup>2</sup>. No mortalities occurred in any of the treatment or control groups after Day 8.

## APPENDIX I

## Certificate Of Analysis

FC-95, Lot 217

March 9, 2000

Richard M. Payfer

This sample was analyzed using LC/MS,  $^1\text{H}$ -NMR,  $^{19}\text{F}$ -NMR, and elemental analyses techniques. The results of these tests show the sample to contain the following weight percent composition:

|                                                           |         |
|-----------------------------------------------------------|---------|
| $\text{C}_2\text{F}_5\text{SO}_2\text{K}^+$               | 0.04 %  |
| $\text{C}_3\text{F}_7\text{SO}_2\text{K}^+$               | 0.83 %  |
| $\text{C}_4\text{F}_9\text{SO}_2\text{K}^+$               | 1.38 %  |
| $\text{C}_5\text{F}_{11}\text{SO}_2\text{K}^+$            | 1.30 %  |
| $\text{C}_6\text{F}_{13}\text{SO}_2\text{K}^+$            | 3.71 %  |
| $\text{C}_7\text{F}_{15}\text{SO}_2\text{K}^+$            | 1.19 %  |
| $\text{C}_8\text{F}_{17}\text{SO}_2\text{K}^+$            | 90.49 % |
| $\text{C}_9\text{F}_{19}\text{SO}_2\text{K}^+$            | 0.49 %  |
| $\text{C}_{10}\text{F}_{21}\text{SO}_2\text{K}^+$         | 0.13 %  |
| $\text{C}_{11}\text{F}_{23}\text{SO}_2\text{K}^+$         | 0.04 %  |
| $\text{CF}_3\text{-CO}_2\text{K}^+$                       | 0.05 %  |
| $\text{SF}_6\text{C}_4\text{F}_{16}\text{SO}_2\text{K}^+$ | 0.35 %  |

Additionally, the isomer distribution of the sample was determined using  $^{19}\text{F}$ -NMR techniques and found to contain the following mole percent composition:

|                                                                                                                                                                                 |       |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|
| $\text{CF}_3(\text{CF}_2)_x\text{-SO}_2\text{K}^+$<br>(Normal chain, where x is mainly 7)                                                                                       | 70.5% |
| $\text{CF}_3(\text{CF}_2)_x\text{-CF}(\text{CF}_3)\text{-(CF}_2)_y\text{-SO}_2\text{K}^+$<br>(Internal monomethyl branch,<br>where x+y is mainly 5, and x $\neq$ 0, y $\neq$ 0) | 17.1% |
| $(\text{CF}_3)_2\text{CF}(\text{CF}_2)_x\text{-SO}_2\text{K}^+$<br>(Isopropyl branch, where x is mainly 5)                                                                      | 10.3% |
| $\text{C}_4\text{F}_{2x+1}\text{-CF}(\text{CF}_3)\text{-SO}_2\text{K}^+$<br>(Alpha branch, where x is mainly 6)                                                                 | 1.6%  |
| $(\text{CF}_3)_3\text{C}(\text{CF}_2)_x\text{-SO}_2\text{K}^+$<br>(t-butyl branch, where x is mainly 4)                                                                         | 0.2%  |
| $\text{CF}_3\text{-(CF}_2)_x\text{-C}(\text{CF}_3)_2\text{-(CF}_2)_y\text{-SO}_2\text{K}^+$<br>(Internal gem-dimethyl branch, where x+y<br>is mainly 4, and x $\neq$ 0)         | 0.2%  |

## APPENDIX II

DIET FORMULATION  
WILDLIFE INTERNATIONAL LTD. GAME BIRD RATION<sup>1</sup>

| INGREDIENTS                            | PERCENT (%) |
|----------------------------------------|-------------|
| Fine Corn Meal                         | 44.83       |
| Soy Bean Meal, 48% Protein             | 30.65       |
| Wheat Midds                            | 6.50        |
| Protein Base                           | 6.00        |
| Agway Special, 60% Protein             | 4.00        |
| Alfalfa Meal, 20% Protein              | 3.00        |
| Dried Whey                             | 2.50        |
| Ground Limestone                       | 0.90        |
| Eastman CalPhos                        | 0.60        |
| Methionine Premix + Liquid             | 0.35        |
| Vitamin and Mineral Premix (see below) | 0.32        |
| GL Ferm (Fermatco) <sup>2</sup>        | 0.25        |
| Salt Iodized                           | 0.10        |
| Total                                  | 100.00      |

## VITAMIN AND MINERAL PREMIX

## AMOUNT ADDED PER TON

|                                                     |                  |
|-----------------------------------------------------|------------------|
| Vitamin D <sub>3</sub>                              | 2,000,000 I.C.U. |
| Vitamin A                                           | 7,000,000 I.U.   |
| Riboflavin                                          | 6 grams          |
| Niacin                                              | 40 grams         |
| Pantothenic Acid                                    | 10 grams         |
| Vitamin B <sub>12</sub>                             | 8 mgs            |
| Folic Acid                                          | 600 mgs          |
| Biotin                                              | 64 mgs           |
| Pyridoxine                                          | 1.2 grams        |
| Thiamine                                            | 1.2 grams        |
| Vitamin E                                           | 20,000 I.U.      |
| Vitamin K (Menadione Dimethylpyrimidinol Bisulfite) | 5.8 grams        |
| Manganese                                           | 102 grams        |
| Zinc                                                | 47 grams         |
| Copper                                              | 6.8 grams        |
| Iodine                                              | 1.5 grams        |
| Iron                                                | 51 grams         |
| Selenium                                            | 182 mgs          |

<sup>1</sup>The guaranteed analysis is a minimum of 27% protein, a minimum of 2.5% crude fat and a maximum of 5% crude fiber.

<sup>2</sup>Fermentation By-Products (Source of Unidentified Growth Factors).

APPENDIX III  
ANALYTICAL METHODS AND RESULTS

## APPENDIX III

Table 1

## Typical LC/MS Operational Parameters

|                                      |                                                                                                                                                                                                                          |
|--------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| INSTRUMENT:                          | Hewlett-Packard Model 1100 High Performance Liquid Chromatograph with a Perkin-Elmer API 100LC Mass Spectrometer equipped with a Perkin-Elmer TurboIonSpray ion source. Operated in selective ion monitoring mode (SIM). |
| ANALYTICAL COLUMN:                   | Keystone Betasil C <sub>18</sub> column (100 mm x 2 mm I.D., 3 µm particle size)                                                                                                                                         |
| OVEN TEMPERATURE:                    | 30°C                                                                                                                                                                                                                     |
| STOP TIME:                           | 10.0 minutes                                                                                                                                                                                                             |
| FLOW RATE:                           | 0.220 mL/minute                                                                                                                                                                                                          |
| MOBILE PHASE:                        | 72.0% Methanol : 28.0% NANOpure® Water containing 0.1% Formic Acid                                                                                                                                                       |
| INJECTION VOLUME:                    | 25.0 µL                                                                                                                                                                                                                  |
| PFOS RETENTION TIME:                 | Approximately 7.0 minutes                                                                                                                                                                                                |
| INTERNAL STANDARD<br>RETENTION TIME: | Approximately 4.8 minutes                                                                                                                                                                                                |
| PFOS MONITORED MASS:                 | 498.6 amu                                                                                                                                                                                                                |
| INTERNAL STANDARD<br>MONITORED MASS: | 426.7 amu                                                                                                                                                                                                                |

## APPENDIX III

Table 2

Matrix Blanks Analyzed Concurrently During Sample Analysis

| Sample               |              | Measured Concentration of<br>PFOS <sup>1</sup><br>(ppm a.i.) |
|----------------------|--------------|--------------------------------------------------------------|
| Number<br>(454-103-) | Type         |                                                              |
| MAB-1                | Matrix Blank | < LOQ                                                        |
| MAB-2                | Matrix Blank | < LOQ                                                        |

<sup>1</sup>The limit of quantitation (LOQ) was 1.15 ppm a.i. based upon the product of the lowest calibration standard analyzed (0.00229 mg a.i./L) and the dilution factor of the matrix blank samples (500).

## APPENDIX III

Table 3

Matrix Fortifications Analyzed Concurrently During Sample Analysis

| Sample Number<br>(454-103-) | Concentrations of PFOS<br>(ppm a.i.) |          | Percent<br>Recovered |
|-----------------------------|--------------------------------------|----------|----------------------|
|                             | Fortified                            | Measured |                      |
| MAS-1A                      | 4.57                                 | 4.54     | 99.2                 |
| MAS-4A                      | 4.57                                 | 4.79     | 105                  |
| MAS-2                       | 183                                  | 176      | 96.1                 |
| MAS-5                       | 183                                  | 162      | 88.3                 |
| MAS-3                       | 1830                                 | 1576     | 86.1                 |
| MAS-6                       | 1830                                 | 1716     | 93.7                 |
| Mean =                      |                                      |          | 94.7                 |
| Standard Deviation =        |                                      |          | 6.99                 |
| CV =                        |                                      |          | 7.38                 |
| N =                         |                                      |          | 6                    |

Note: Results and corrections for new test substance purity were generated using MacQuan version 1.5 software and manual calculations. Values have been rounded for reporting purposes.



- 33 -

## APPENDIX III

Table 4

## Homogeneity of PFOS in Avian Diet

| Nominal Concentration (ppm a.i.) | Sample Number (S-454-103-) | Location Sampled in Mixing Vessel | PFOS Measured Concentration (ppm a.i.) | Mean Measured Concentration ( $\bar{x}$ )<br>Standard Deviation (SD)<br>Coefficient of Variation (CV) <sup>1</sup> | Mean Percent of Nominal |
|----------------------------------|----------------------------|-----------------------------------|----------------------------------------|--------------------------------------------------------------------------------------------------------------------|-------------------------|
| 18.3                             | 2                          | Top Left                          | 18.5                                   | $\bar{x}$ = 19.5 ppm a.i.<br>SD = 2.13 ppm a.i.<br>CV = 10.9%                                                      | 107                     |
|                                  | 3                          | Top Right                         | 23.4                                   |                                                                                                                    |                         |
|                                  | 4                          | Middle Left                       | 18.3                                   |                                                                                                                    |                         |
|                                  | 5                          | Middle Right                      | 17.3                                   |                                                                                                                    |                         |
|                                  | 6                          | Bottom Left                       | 19.4                                   |                                                                                                                    |                         |
|                                  | 7                          | Bottom Right                      | 19.9                                   |                                                                                                                    |                         |
|                                  |                            |                                   |                                        |                                                                                                                    |                         |
| 1171                             | 18                         | Top Left                          | 1239                                   | $\bar{x}$ = 1196 ppm a.i.<br>SD = 70.2 ppm a.i.<br>CV = 5.87%                                                      | 102                     |
|                                  | 19                         | Top Right                         | 1221                                   |                                                                                                                    |                         |
|                                  | 20                         | Middle Left                       | 1118                                   |                                                                                                                    |                         |
|                                  | 21                         | Middle Right                      | 1301                                   |                                                                                                                    |                         |
|                                  | 22                         | Bottom Left                       | 1163                                   |                                                                                                                    |                         |
|                                  | 23                         | Bottom Right                      | 1133                                   |                                                                                                                    |                         |

Note: Results and corrections for new test substance purity were generated using MacQuan version 1.5 software and manual calculations. Values have been rounded for reporting purposes.

<sup>1</sup>Coefficient of variation was calculated using full precision of mean and standard deviation results.

- 34 -

## APPENDIX III

Table 5

## Verification of PFOS Concentrations in Avian Diet

| Nominal Concentration (ppm a.i.) | Sample Number (S-454-103-) | Sampling Interval (Day) | PFOS Measured Concentration <sup>1</sup> (ppm a.i.) | Percent of Nominal | Mean Measured Concentration (ppm a.i.) | Mean Percent of Nominal |
|----------------------------------|----------------------------|-------------------------|-----------------------------------------------------|--------------------|----------------------------------------|-------------------------|
| 0.0                              | 1                          | 0                       | < LOQ                                               | —                  | —                                      | —                       |
| 18.3                             | —                          | —                       | —                                                   | —                  | 19.5 <sup>2</sup>                      | 107 <sup>2</sup>        |
| 36.6                             | 8                          | 0                       | 45.7                                                | 125                | 40.2                                   | 110                     |
|                                  | 9                          | 0                       | 34.6                                                | 94.5               |                                        |                         |
| 73.2                             | 10                         | 0                       | 77.8                                                | 106                | 74.5                                   | 102                     |
|                                  | 11                         | 0                       | 71.2                                                | 97.3               |                                        |                         |
| 146                              | 12                         | 0                       | 176                                                 | 120                | 174                                    | 119                     |
|                                  | 13                         | 0                       | 172                                                 | 117                |                                        |                         |
| 293                              | 14                         | 0                       | 274                                                 | 93.8               | 291                                    | 99.3                    |
|                                  | 15                         | 0                       | 307                                                 | 105                |                                        |                         |
| 586                              | 16                         | 0                       | 550                                                 | 93.9               | 537                                    | 91.6                    |
|                                  | 17                         | 0                       | 523                                                 | 89.4               |                                        |                         |
| 1171                             | —                          | —                       | —                                                   | —                  | 1196 <sup>2</sup>                      | 102 <sup>2</sup>        |

Note: Results and corrections for new test substance purity were generated using MacQuan version 1.5 software and manual calculations. Values have been rounded for reporting purposes.

<sup>1</sup>The limit of quantitation (LOQ) was 1.15 ppm a.i. based upon the product of the lowest calibration standard analyzed (0.00229 mg a.i./L) and the dilution factor of the matrix blank samples (500).

<sup>2</sup>Result obtained from Table 4.

- 35 -

## APPENDIX III

Table 6


Ambient Stability of PFOS in Avian Diet During the Northern Bobwhite LC50 Study

| Nominal<br>Concentration<br>(ppm a.i.) | Day 0 <sup>1</sup>               |                                              | Day 5                         |                                  |                                                      |                                              |
|----------------------------------------|----------------------------------|----------------------------------------------|-------------------------------|----------------------------------|------------------------------------------------------|----------------------------------------------|
|                                        | Sample<br>Number<br>(S-454-103-) | Mean Measured<br>Concentration<br>(ppm a.i.) | Mean Percent<br>of<br>Nominal | Sample<br>Number<br>(S-454-103-) | Measured<br>Concentration <sup>2</sup><br>(ppm a.i.) | Mean Measured<br>Concentration<br>(ppm a.i.) |
| 0                                      | 1                                | --                                           | --                            | 24                               | < LOQ                                                | --                                           |
| 18.3                                   | 2-7                              | 19.5                                         | 107                           | 25<br>26                         | 19.2<br>19.9                                         | 19.6                                         |
| 36.6                                   | 8, 9                             | 40.2                                         | 110                           | 27<br>28                         | 44.4<br>53.8                                         | 49.1                                         |
| 73.2                                   | 10, 11                           | 74.5                                         | 102                           | 29<br>30                         | 76.4<br>77.9                                         | 77.2                                         |
| 146                                    | 12, 13                           | 174                                          | 119                           | 31<br>32                         | 177<br>174                                           | 176                                          |
| 293                                    | 14, 15                           | 291                                          | 99.3                          | 33<br>34                         | 318<br>315                                           | 317                                          |
| 586                                    | 16, 17                           | 537                                          | 91.6                          | 35<br>36                         | 560<br>665                                           | 613                                          |
| 1171                                   | 18-23                            | 1196                                         | 102                           | 37<br>38                         | 1260<br>1187                                         | 1224                                         |
|                                        |                                  |                                              |                               |                                  |                                                      | 102                                          |

<sup>1</sup>Day 0 results obtained from Table 4 and Table 5.<sup>2</sup>The limit of quantitation (LOQ) was 1.15 ppm a.i. based upon the product of the lowest calibration standard analyzed (0.00229 mg a.i./L) and the dilution factor of the matrix blank samples (500).

- 36 -

## APPENDIX III

**METHOD OUTLINE FOR THE ANALYSIS OF PFOS  
IN AVIAN DIET**

**Figure 1.** Analytical method flowchart for the analysis of PFOS in avian diet.

- 37 -

## APPENDIX III




Figure 2. A typical calibration curve for PFOS. Slope = 2.77397; Intercept = 0.01894;  $r = 0.9981$ . Curve is weighted ( $1/x$ ).

- 38 -

## APPENDIX III




Figure 3. A representative ion chromatogram of a low-level (0.00229 mg a.i./L) PFOS standard.

- 39 -

## APPENDIX III




Figure 4. A representative ion chromatogram of a high-level (0.0457 mg a.i./L) PFOS standard.

- 40 -

## APPENDIX III




Figure 5. A representative chromatogram of a matrix blank sample (454-103-MAB-1).  
The arrow indicates the retention time of PFOS.



- 41 -

## APPENDIX III




Figure 6. A representative chromatogram of a matrix fortification sample (454-103-MAS-1A).

- 42 -

## APPENDIX III




Figure 7. A representative chromatogram of a test sample (454-103-2).

- 43 -

APPENDIX IV  
DIET PREPARATION

Weight and volume of constituents used to prepare test diets:

| Nominal<br>Concentrations<br>(ppm a.i.) | Test<br>Substance<br>(g) | Basal<br>Ration<br>(g) |
|-----------------------------------------|--------------------------|------------------------|
| 0                                       | -                        | 9000.0                 |
| 18.3                                    | 0.1818                   | 8999.8                 |
| 36.6                                    | 0.3638                   | 8999.6                 |
| 73.2                                    | 0.7282                   | 8999.3                 |
| 146                                     | 1.4659                   | 8998.5                 |
| 293                                     | 2.9123                   | 8997.1                 |
| 586                                     | 5.8239                   | 8994.2                 |
| 1171                                    | 11.6483                  | 8988.4                 |

Diets were prepared as follows:

- 5000.0 g of basal ration was weighed into a tared Hobart mixing bowl.
- The test substance was weighed in a tared weigh boat.
- Approximately 100 g of basal ration was taken from the mixing bowl and placed in a Waring blender.
- The test substance was added to the blender and the weigh boat was rinsed with additional ration, with the rinse also being placed in the blender.
- The blender contents were blended for approximately 60 seconds and transferred to the mixing bowl. The blender was rinsed with additional ration, with the rinse also being placed in the mixing bowl.
- The bowl was placed on a Hobart mixer and the contents were mixed for approximately six minutes. The remaining ration as added to the bowl and the contents were mixed for six more minutes.
- The diet was transferred to a labelled paper feed bag.

- 44 -

APPENDIX V  
Cumulative Mortality by Pen from a Northern Bobwhite Acute Dietary Toxicity Study with PFOS

| Experimental Group<br>(ppm a.i.) | Pen | No. Dead Per No. Exposed |       |       |       |       | No. Dead Per No. Exposed |       |       |                    |
|----------------------------------|-----|--------------------------|-------|-------|-------|-------|--------------------------|-------|-------|--------------------|
|                                  |     | Exposure Period          |       |       |       |       | Post-Exposure Period     |       |       |                    |
|                                  |     | Day 0                    | Day 1 | Day 2 | Day 3 | Day 4 | Day 5                    | Day 6 | Day 7 | Day 8 <sup>1</sup> |
| Control                          |     |                          |       |       |       |       |                          |       |       |                    |
| 0                                | 1   | 0/5                      | 0/5   | 0/5   | 0/5   | 0/5   | 0/5                      | 0/5   | 0/5   | 0/5                |
| 0                                | 2   | 0/5                      | 0/5   | 0/5   | 0/5   | 0/5   | 0/5                      | 0/5   | 0/5   | 0/5                |
| 0                                | 3   | 0/5                      | 0/5   | 0/5   | 0/5   | 0/5   | 0/5                      | 0/5   | 0/5   | 0/5                |
| 0                                | 4   | 0/5                      | 0/5   | 0/5   | 0/5   | 0/5   | 0/5                      | 0/5   | 0/5   | 0/5                |
| 0                                | 5   | 0/5                      | 0/5   | 0/5   | 0/5   | 0/5   | 0/5                      | 1/5*  | 1/5   | 1/5                |
| 0                                | 6   | 0/5                      | 0/5   | 0/5   | 0/5   | 0/5   | 0/5                      | 0/5   | 0/5   | 0/5                |
| Treatment                        |     |                          |       |       |       |       |                          |       |       |                    |
| 18.3                             | 1   | 0/5                      | 0/5   | 0/5   | 0/5   | 0/5   | 0/5                      | 0/5   | 0/5   | 0/5                |
| 18.3                             | 2   | 0/5                      | 0/5   | 0/5   | 0/5   | 0/5   | 0/5                      | 0/5   | 0/5   | 0/5                |
| 36.6                             | 1   | 0/5                      | 0/5   | 0/5   | 0/5   | 0/5   | 0/5                      | 0/5   | 0/5   | 0/5                |
| 36.6                             | 2   | 0/5                      | 0/5   | 0/5   | 0/5   | 0/5   | 0/5                      | 0/5   | 0/5   | 0/5                |
| 73.2                             | 1   | 0/5                      | 0/5   | 0/5   | 0/5   | 0/5   | 0/5                      | 0/5   | 0/5   | 0/5                |
| 73.2                             | 2   | 0/5                      | 0/5   | 0/5   | 0/5   | 0/5   | 0/5                      | 0/5   | 0/5   | 0/5                |
| 146                              | 1   | 0/5                      | 0/5   | 0/5   | 0/5   | 0/5   | 0/5                      | 0/5   | 0/5   | 1/5                |
| 146                              | 2   | 0/5                      | 0/5   | 0/5   | 1/5*  | 1/5   | 1/5                      | 1/5   | 1/5   | 1/5                |
| 293                              | 1   | 0/5                      | 0/5   | 0/5   | 0/5   | 0/5   | 1/5                      | 1/5   | 4/5   | 4/5                |
| 293                              | 2   | 0/5                      | 0/5   | 0/5   | 0/5   | 0/5   | 0/5                      | 3/5   | 4/5   | 4/5                |
| 586                              | 1   | 0/5                      | 0/5   | 0/5   | 1/5   | 2/5   | 2/5                      | 4/5   | 5/5   | 5/5                |
| 586                              | 2   | 0/5                      | 0/5   | 0/5   | 0/5   | 0/5   | 3/5                      | 4/5   | 5/5   | 5/5                |
| 1171                             | 1   | 0/5                      | 0/5   | 0/5   | 3/5   | 5/5   | 5/5                      | 5/5   | 5/5   | 5/5                |
| 1171                             | 2   | 0/5                      | 0/5   | 0/5   | 0/5   | 5/5   | 5/5                      | 5/5   | 5/5   | 5/5                |

The LC50 value was calculated to be 220 ppm a.i. with a 95% confidence interval of 164 ppm a.i. to 289 ppm a.i.

\* - Bird was euthanized due to a broken leg.

1 - No mortalities occurred in any of the control or treatment groups after Day 8.

- 45 -

## APPENDIX VI

Individual Body Weights (g) from a Northern Bobwhite  
 Acute Dietary Toxicity Study with PFOS  
 Page 1

| Experimental<br>Group<br>(ppm a.i.) | Bird | Day 0 | Day 5 | Change<br>Day 0-5 | Day 8 | Change<br>Day 5-8 | Total<br>Change |
|-------------------------------------|------|-------|-------|-------------------|-------|-------------------|-----------------|
| 0                                   | 1    | 19    | 24    | 5                 | 27    | 3                 | 8               |
|                                     | 2    | 20    | 27    | 7                 | 34    | 7                 | 14              |
|                                     | 3    | 21    | 32    | 11                | 42    | 10                | 21              |
|                                     | 4    | 22    | 32    | 10                | 40    | 8                 | 18              |
|                                     | 5    | 23    | 39    | 16                | 50    | 11                | 27              |
|                                     | Mean | 21    | 31    | 10                | 39    | 8                 | 18              |
|                                     | SD   | 2     | 6     | 4                 | 9     | 3                 | 7               |
| 0                                   | 1    | 19    | 26    | 7                 | 35    | 9                 | 16              |
|                                     | 2    | 20    | 25    | 5                 | 35    | 10                | 15              |
|                                     | 3    | 20    | 29    | 9                 | 37    | 8                 | 17              |
|                                     | 4    | 22    | 33    | 11                | 43    | 10                | 21              |
|                                     | 5    | 22    | 31    | 9                 | 39    | 8                 | 17              |
|                                     | Mean | 21    | 29    | 8                 | 38    | 9                 | 17              |
|                                     | SD   | 1     | 3     | 2                 | 3     | 1                 | 2               |
| 0                                   | 1    | 19    | 23    | 4                 | 30    | 7                 | 11              |
|                                     | 2    | 20    | 26    | 6                 | 29    | 3                 | 9               |
|                                     | 3    | 20    | 29    | 9                 | 32    | 3                 | 12              |
|                                     | 4    | 22    | 30    | 8                 | 37    | 7                 | 15              |
|                                     | 5    | 22    | 35    | 13                | 40    | 5                 | 18              |
|                                     | Mean | 21    | 29    | 8                 | 34    | 5                 | 13              |
|                                     | SD   | 1     | 5     | 3                 | 5     | 2                 | 4               |
| 0                                   | 1    | 19    | 28    | 9                 | 37    | 9                 | 18              |
|                                     | 2    | 19    | 27    | 8                 | 35    | 8                 | 16              |
|                                     | 3    | 19    | 30    | 11                | 36    | 6                 | 17              |
|                                     | 4    | 21    | 32    | 11                | 42    | 10                | 21              |
|                                     | 5    | 22    | 33    | 11                | 43    | 10                | 21              |
|                                     | Mean | 20    | 30    | 10                | 39    | 9                 | 19              |
|                                     | SD   | 1     | 3     | 1                 | 4     | 2                 | 2               |

- 46 -

## APPENDIX VI

Individual Body Weights (g) from a Northern Bobwhite  
 Acute Dietary Toxicity Study with PFOS  
 Page 2

| Experimental<br>Group<br>(ppm a.i.) | Bird | Day 0 | Day 5 | Change<br>Day 0-5 | Day 8 | Change<br>Day 5-8 | Total<br>Change |
|-------------------------------------|------|-------|-------|-------------------|-------|-------------------|-----------------|
| 0                                   | 1    | 19    | 28    | 9                 | 35    | 7                 | 16              |
|                                     | 2    | 20    | 27    | 7                 | 36    | 9                 | 16              |
|                                     | 3    | 21    | 32    | 11                | 44    | 12                | 23              |
|                                     | 4    | 20    | 30    | 10                | -     | -                 | -               |
|                                     | 5    | 22    | 34    | 12                | 44    | 10                | 22              |
|                                     | Mean | 20    | 30    | 10                | 40    | 10                | 19              |
|                                     | SD   | 1     | 3     | 2                 | 5     | 2                 | 4               |
| 0                                   | 1    | 18    | 30    | 12                | 39    | 9                 | 21              |
|                                     | 2    | 21    | 35    | 14                | 46    | 11                | 25              |
|                                     | 3    | 19    | 29    | 10                | 37    | 8                 | 18              |
|                                     | 4    | 21    | 32    | 11                | 41    | 9                 | 20              |
|                                     | 5    | 22    | 32    | 10                | 42    | 10                | 20              |
|                                     | Mean | 20    | 32    | 11                | 41    | 9                 | 21              |
|                                     | SD   | 2     | 2     | 2                 | 3     | 1                 | 3               |
| Group                               | Mean | 20    | 30    | 10                | 38    | 8                 | 18              |
| Total                               | SD   | 1     | 4     | 3                 | 5     | 2                 | 4               |

- 47 -

## APPENDIX VI

Individual Body Weights (g) from a Northern Bobwhite  
 Acute Dietary Toxicity Study with PFOS  
 Page 3

| Experimental<br>Group<br>(ppm a.i.) | Bird | Day 0 | Day 5 | Change<br>Day 0-5 | Day 8 | Change<br>Day 5-8 | Total<br>Change |
|-------------------------------------|------|-------|-------|-------------------|-------|-------------------|-----------------|
| 18.3                                | 1    | 19    | 26    | 7                 | 34    | 8                 | 15              |
|                                     | 2    | 19    | 33    | 14                | 39    | 6                 | 20              |
|                                     | 3    | 21    | 27    | 6                 | 35    | 8                 | 14              |
|                                     | 4    | 22    | 34    | 12                | 46    | 12                | 24              |
|                                     | 5    | 22    | 36    | 14                | 46    | 10                | 24              |
|                                     | Mean | 21    | 31    | 11                | 40    | 9                 | 19              |
|                                     | SD   | 2     | 4     | 4                 | 6     | 2                 | 5               |
| 18.3                                | 1    | 19    | 28    | 9                 | 38    | 10                | 19              |
|                                     | 2    | 19    | 28    | 9                 | 37    | 9                 | 18              |
|                                     | 3    | 20    | 29    | 9                 | 38    | 9                 | 18              |
|                                     | 4    | 22    | 34    | 12                | 44    | 10                | 22              |
|                                     | 5    | 22    | 35    | 13                | 44    | 9                 | 22              |
|                                     | Mean | 20    | 31    | 10                | 40    | 9                 | 20              |
|                                     | SD   | 2     | 3     | 2                 | 3     | 1                 | 2               |
| Group                               | Mean | 21    | 31    | 11                | 40    | 9                 | 20              |
| Total                               | SD   | 1     | 4     | 3                 | 5     | 2                 | 3               |

- 48 -

APPENDIX VI  
Individual Body Weights (g) from a Northern Bobwhite  
Acute Dietary Toxicity Study with PFOS  
Page 4

| Experimental<br>Group<br>(ppm a.i.) | Bird | Day 0 | Day 5 | Change<br>Day 0-5 | Day 8 | Change<br>Day 5-8 | Total<br>Change |
|-------------------------------------|------|-------|-------|-------------------|-------|-------------------|-----------------|
| 36.6                                | 1    | 18    | 26    | 8                 | 35    | 9                 | 17              |
|                                     | 2    | 19    | 30    | 11                | 39    | 9                 | 20              |
|                                     | 3    | 19    | 29    | 10                | 37    | 8                 | 18              |
|                                     | 4    | 21    | 30    | 9                 | 38    | 8                 | 17              |
|                                     | 5    | 21    | 34    | 13                | 43    | 9                 | 22              |
|                                     | Mean | 20    | 30    | 10                | 38    | 9                 | 19              |
|                                     | SD   | 1     | 3     | 2                 | 3     | 1                 | 2               |
| 36.6                                | 1    | 19    | 29    | 10                | 37    | 8                 | 18              |
|                                     | 2    | 19    | 29    | 10                | 37    | 8                 | 18              |
|                                     | 3    | 21    | 32    | 11                | 40    | 8                 | 19              |
|                                     | 4    | 22    | 35    | 13                | 44    | 9                 | 22              |
|                                     | 5    | 23    | 36    | 13                | 44    | 8                 | 21              |
|                                     | Mean | 21    | 32    | 11                | 40    | 8                 | 20              |
|                                     | SD   | 2     | 3     | 2                 | 4     | 0                 | 2               |
| Group<br>Total                      | Mean | 20    | 31    | 11                | 39    | 8                 | 19              |
|                                     | SD   | 2     | 3     | 2                 | 3     | 1                 | 2               |



- 49 -

## APPENDIX VI

Individual Body Weights (g) from a Northern Bobwhite  
 Acute Dietary Toxicity Study with PFOS  
 Page 5

| Experimental<br>Group<br>(ppm a.i.) | Bird | Day 0 | Day 5 | Change<br>Day 0-5 | Day 8 | Change<br>Day 5-8 | Total<br>Change |
|-------------------------------------|------|-------|-------|-------------------|-------|-------------------|-----------------|
| 73.2                                | 1    | 18    | 26    | 8                 | 32    | 6                 | 14              |
|                                     | 2    | 19    | 26    | 7                 | 33    | 7                 | 14              |
|                                     | 3    | 20    | 29    | 9                 | 36    | 7                 | 16              |
|                                     | 4    | 21    | 32    | 11                | 39    | 7                 | 18              |
|                                     | 5    | 21    | 30    | 9                 | 36    | 6                 | 15              |
|                                     | Mean | 20    | 29    | 9                 | 35    | 7                 | 15              |
|                                     | SD   | 1     | 3     | 1                 | 3     | 1                 | 2               |
| 73.2                                | 1    | 18    | 27    | 9                 | 34    | 7                 | 16              |
|                                     | 2    | 20    | 31    | 11                | 38    | 7                 | 18              |
|                                     | 3    | 21    | 30    | 9                 | 37    | 7                 | 16              |
|                                     | 4    | 21    | 32    | 11                | 39    | 7                 | 18              |
|                                     | 5    | 22    | 32    | 10                | 41    | 9                 | 19              |
|                                     | Mean | 20    | 30    | 10                | 38    | 7                 | 17              |
|                                     | SD   | 2     | 2     | 1                 | 3     | 1                 | 1               |
| Group                               | Mean | 20    | 30    | 9                 | 37    | 7                 | 16              |
| Total                               | SD   | 1     | 2     | 1                 | 3     | 1                 | 2               |

- 50 -

## APPENDIX VI

Individual Body Weights (g) from a Northern Bobwhite  
Acute Dietary Toxicity Study with PFOS

Page 6

| Experimental<br>Group<br>(ppm a.i.) | Bird | Day 0 | Day 5 | Change<br>Day 0-5 | Day 8 | Change<br>Day 5-8 | Total<br>Change |
|-------------------------------------|------|-------|-------|-------------------|-------|-------------------|-----------------|
| 146                                 | 1    | 19    | 28    | 9                 | 35    | 7                 | 16              |
|                                     | 2    | 19    | 28    | 9                 | 32    | 4                 | 13              |
|                                     | 3    | 20    | 24    | 4                 | -     | -                 | -               |
|                                     | 4    | 22    | 33    | 11                | 38    | 5                 | 16              |
|                                     | 5    | 23    | 28    | 5                 | 32    | 4                 | 9               |
|                                     | Mean | 21    | 28    | 8                 | 34    | 5                 | 14              |
|                                     | SD   | 2     | 3     | 3                 | 3     | 1                 | 3               |
| 146                                 | 1    | 18    | 24    | 6                 | 31    | 7                 | 13              |
|                                     | 2    | 19    | 27    | 8                 | 35    | 8                 | 16              |
|                                     | 3    | 19    | 27    | 8                 | 33    | 6                 | 14              |
|                                     | 4    | 21    | -     | -                 | -     | -                 | -               |
|                                     | 5    | 23    | 24    | 1                 | 28    | 4                 | 5               |
|                                     | Mean | 20    | 26    | 6                 | 32    | 6                 | 12              |
|                                     | SD   | 2     | 2     | 3                 | 3     | 2                 | 5               |
| Group                               | Mean | 20    | 27*   | 7*                | 33*   | 6**               | 13**            |
| Total                               | SD   | 2     | 3     | 3                 | 3     | 2                 | 4               |

(-) = No data available due to mortality.

\*Statistically different from the control group at  $p < 0.05$  (Dunnett's t-test).\*\*Statistically different from the control group at  $p < 0.01$  (Dunnett's t-test).

- 51 -

## APPENDIX VI

Individual Body Weights (g) from a Northern Bobwhite  
Acute Dietary Toxicity Study with PFOS  
Page 7

| Experimental<br>Group<br>(ppm a.i.) | Bird | Day 0 | Day 5 | Change<br>Day 0-5 | Day 8 | Change<br>Day 5-8 | Total<br>Change |
|-------------------------------------|------|-------|-------|-------------------|-------|-------------------|-----------------|
| 293                                 | 1    | 18    | 20    | 2                 | 21    | 1                 | 3               |
|                                     | 2    | 19    | 17    | -2                | -     | -                 | -               |
|                                     | 3    | 20    | 18    | -2                | -     | -                 | -               |
|                                     | 4    | 22    | -     | -                 | -     | -                 | -               |
|                                     | 5    | 22    | 20    | -2                | -     | -                 | -               |
|                                     | Mean | 20    |       | -1                | 21    | 1                 | 3               |
|                                     | SD   | 2     |       | 2                 | -     | -                 | -               |
| 293                                 | 1    | 19    | 19    | 0                 | 15    | -4                | -4              |
|                                     | 2    | 19    | -     | -                 | -     | -                 | -               |
|                                     | 3    | 20    | 20    | 0                 | -     | -                 | -               |
|                                     | 4    | 20    | 17    | -3                | -     | -                 | -               |
|                                     | 5    | 21    | 16    | -5                | -     | -                 | -               |
|                                     | Mean | 20    | 18    | -2                | 15    | -4                | -4              |
|                                     | SD   | 1     | 2     | 2                 | -     | -                 | -               |
| Group                               | Mean | 20    | 18**  | -2**              | 18**  | -2**              | -1**            |
| Total                               | SD   | 1     | 2     | 2                 | 4     | 4                 | 5               |

(-) = No data available due to mortality.  
 \*Statistically different from the control group at  $p < 0.05$  (Dunnett's t-test).  
 \*\*Statistically different from the control group at  $p < 0.01$  (Dunnett's t-test).

- 52 -

## APPENDIX VI

Individual Body Weights (g) from a Northern Bobwhite  
Acute Dietary Toxicity Study with PFOS  
Page 8

| Experimental<br>Group<br>(ppm a.i.) | Bird | Day 0 | Day 5 | Change<br>Day 0-5 | Day 8 | Change<br>Day 5-8 | Total<br>Change |
|-------------------------------------|------|-------|-------|-------------------|-------|-------------------|-----------------|
| 586                                 | 1    | 18    | -     | -                 | -     | -                 | -               |
|                                     | 2    | 20    | 13    | -7                | -     | -                 | -               |
|                                     | 3    | 21    | 18    | -3                | -     | -                 | -               |
|                                     | 4    | 21    | -     | -                 | -     | -                 | -               |
|                                     | 5    | 22    | 17    | -5                | -     | -                 | -               |
|                                     | Mean | 20    | 16    | -5                | -     | -                 | -               |
|                                     | SD   | 2     | 3     | 2                 | -     | -                 | -               |
| 586                                 | 1    | 19    | 16    | -3                | -     | -                 | -               |
|                                     | 2    | 20    | -     | -                 | -     | -                 | -               |
|                                     | 3    | 20    | 16    | -4                | -     | -                 | -               |
|                                     | 4    | 19    | -     | -                 | -     | -                 | -               |
|                                     | 5    | 21    | -     | -                 | -     | -                 | -               |
|                                     | Mean | 20    | 16    | -4                | -     | -                 | -               |
|                                     | SD   | 1     | 0     | 1                 | -     | -                 | -               |
| Group                               | Mean | 20    | 16**  | -4**              | -     | -                 | -               |
| Total                               | SD   | 1     | 2     | 2                 | -     | -                 | -               |

(-) = No data available due to mortality.  
\*\*Statistically different from the control group at  $p < 0.01$  (Dunnett's t-test).

- 53 -

## APPENDIX VI

Individual Body Weights (g) from a Northern Bobwhite  
 Acute Dietary Toxicity Study with PFOS  
 Page 9

| Experimental<br>Group<br>(ppm a.i.) | Bird | Day 0 | Day 5 | Change<br>Day 0-5 | Day 8 | Change<br>Day 5-8 | Total<br>Change |
|-------------------------------------|------|-------|-------|-------------------|-------|-------------------|-----------------|
| 1171                                | 1    | 18    | -     | -                 | -     | -                 | -               |
|                                     | 2    | 20    | -     | -                 | -     | -                 | -               |
|                                     | 3    | 20    | -     | -                 | -     | -                 | -               |
|                                     | 4    | 21    | -     | -                 | -     | -                 | -               |
|                                     | 5    | 21    | -     | -                 | -     | -                 | -               |
|                                     | Mean | 20    | -     | -                 | -     | -                 | -               |
|                                     | SD   | 1     | -     | -                 | -     | -                 | -               |
| 1171                                | 1    | 18    | -     | -                 | -     | -                 | -               |
|                                     | 2    | 20    | -     | -                 | -     | -                 | -               |
|                                     | 3    | 20    | -     | -                 | -     | -                 | -               |
|                                     | 4    | 22    | -     | -                 | -     | -                 | -               |
|                                     | 5    | 21    | -     | -                 | -     | -                 | -               |
|                                     | Mean | 20    | -     | -                 | -     | -                 | -               |
|                                     | SD   | 1     | -     | -                 | -     | -                 | -               |
| Group                               | Mean | 20    | -     | -                 | -     | -                 | -               |
| Total                               | SD   | 1     | -     | -                 | -     | -                 | -               |

(-) = No data available due to mortality.

- 54 -

## APPENDIX VI

Individual Body Weights (g) from a Northern Bobwhite  
 Acute Dietary Toxicity Study with PFOS  
 Page 10

| Experimental Group (ppm a.i.) | Bird | Day 8 | Day 15 | Change Day 8-15 | Day 22 | Change Day 15-22 | Total Change |
|-------------------------------|------|-------|--------|-----------------|--------|------------------|--------------|
| 0                             | 1    | 27    | 42     | 15              | 61     | 19               | 34           |
|                               | 2    | 34    | 56     | 22              | 77     | 21               | 43           |
|                               | 3    | 42    | 69     | 27              | 92     | 23               | 50           |
|                               | 4    | 40    | 63     | 23              | 84     | 21               | 44           |
|                               | 5    | 50    | 78     | 28              | 104    | 26               | 54           |
|                               | Mean | 39    | 62     | 23              | 84     | 22               | 45           |
|                               | SD   | 9     | 14     | 5               | 16     | 3                | 8            |
| 0                             | 1    | 35    | 55     | 20              | 74     | 19               | 39           |
|                               | 2    | 35    | 53     | 18              | 74     | 21               | 39           |
|                               | 3    | 37    | 64     | 27              | 85     | 21               | 48           |
|                               | 4    | 43    | 74     | 31              | 103    | 29               | 60           |
|                               | 5    | 39    | 59     | 20              | 80     | 21               | 41           |
|                               | Mean | 38    | 61     | 23              | 83     | 22               | 45           |
|                               | SD   | 3     | 8      | 6               | 12     | 4                | 9            |
| 0                             | 1    | 30    | 54     | 24              | 79     | 25               | 49           |
|                               | 2    | 29    | 45     | 16              | 63     | 18               | 34           |
|                               | 3    | 32    | 54     | 22              | 77     | 23               | 45           |
|                               | 4    | 37    | 55     | 18              | 75     | 20               | 38           |
|                               | 5    | 40    | 68     | 28              | 95     | 27               | 55           |
|                               | Mean | 34    | 55     | 22              | 78     | 23               | 44           |
|                               | SD   | 5     | 8      | 5               | 11     | 4                | 8            |
| Group                         | Mean | 37    | 59     | 23              | 82     | 22               | 45           |
| Total                         | SD   | 6     | 10     | 5               | 13     | 3                | 8            |

- 55 -

## APPENDIX VI

Individual Body Weights (g) from a Northern Bobwhite  
 Acute Dietary Toxicity Study with PFOS  
 Page 11

| Experimental<br>Group<br>(ppm a.i.) | Bird | Day 8 | Day 15 | Change<br>Day 8-15 | Day 22 | Change<br>Day 15-22 | Total<br>Change |
|-------------------------------------|------|-------|--------|--------------------|--------|---------------------|-----------------|
| 18.3                                | 1    | 34    | 53     | 19                 | 78     | 25                  | 44              |
|                                     | 2    | 39    | 65     | 26                 | 85     | 20                  | 46              |
|                                     | 3    | 35    | 60     | 25                 | 86     | 26                  | 51              |
|                                     | 4    | 46    | 73     | 27                 | 95     | 22                  | 49              |
|                                     | 5    | 46    | 71     | 25                 | 92     | 21                  | 46              |
|                                     | Mean | 40    | 64     | 24                 | 87     | 23                  | 47              |
|                                     | SD   | 6     | 8      | 3                  | 7      | 3                   | 3               |

| Experimental<br>Group<br>(ppm a.i.) | Bird | Day 8 | Day 15 | Change<br>Day 8-15 | Day 22 | Change<br>Day 15-22 | Total<br>Change |
|-------------------------------------|------|-------|--------|--------------------|--------|---------------------|-----------------|
| 36.6                                | 1    | 35    | 60     | 25                 | 81     | 21                  | 46              |
|                                     | 2    | 39    | 65     | 26                 | 91     | 26                  | 52              |
|                                     | 3    | 37    | 65     | 28                 | 91     | 26                  | 54              |
|                                     | 4    | 38    | 61     | 23                 | 83     | 22                  | 45              |
|                                     | 5    | 43    | 72     | 29                 | 97     | 25                  | 54              |
|                                     | Mean | 38    | 65     | 26                 | 89     | 24                  | 50              |
|                                     | SD   | 3     | 5      | 2                  | 7      | 2                   | 4               |

- 56 -

## APPENDIX VI

Individual Body Weights (g) from a Northern Bobwhite  
Acute Dietary Toxicity Study with PFOS  
Page 12

| Experimental Group (ppm a.i.) | Bird | Day 8 | Day 15 | Change Day 8-15 | Day 22 | Change Day 15-22 | Total Change |
|-------------------------------|------|-------|--------|-----------------|--------|------------------|--------------|
| 73.2                          | 1    | 32    | 54     | 22              | 76     | 22               | 44           |
|                               | 2    | 33    | 58     | 25              | 77     | 19               | 44           |
|                               | 3    | 36    | 62     | 26              | 79     | 17               | 43           |
|                               | 4    | 39    | 65     | 26              | 87     | 22               | 48           |
|                               | 5    | 36    | 59     | 23              | 78     | 19               | 42           |
|                               | Mean | 35    | 60     | 24              | 79     | 20               | 44           |
|                               | SD   | 3     | 4      | 2               | 4      | 2                | 2            |

| Experimental Group (ppm a.i.) | Bird | Day 8 | Day 15 | Change Day 8-15 | Day 22 | Change Day 15-22 | Total Change |
|-------------------------------|------|-------|--------|-----------------|--------|------------------|--------------|
| 146                           | 1    | 35    | 58     | 23              | 81     | 23               | 46           |
|                               | 2    | 32    | 55     | 23              | 77     | 22               | 45           |
|                               | 3    | -     | -      | -               | -      | -                | -            |
|                               | 4    | 38    | 63     | 25              | 81     | 18               | 43           |
|                               | 5    | 32    | 57     | 25              | 78     | 21               | 46           |
|                               | Mean | 34*   | 58     | 24              | 79     | 21               | 45           |
|                               | SD   | 3     | 3      | 1               | 2      | 2                | 1            |

(-) = No data available due to mortality.

\*Statistically different from the control group at  $p < 0.05$  (Dunnett's t-test).



- 57 -

## APPENDIX VI

Individual Body Weights (g) from a Northern Bobwhite  
Acute Dietary Toxicity Study with PFOS  
Page 13

| Experimental<br>Group<br>(ppm a.i.) | Bird | Day 0 | Day 5 | Change<br>Day 0-5 | Day 8 | Change<br>Day 5-8 | Total<br>Change |
|-------------------------------------|------|-------|-------|-------------------|-------|-------------------|-----------------|
| 293                                 | 1    | 21    | 35    | 14                | 55    | 20                | 34              |
|                                     | 2    | -     | -     | -                 | -     | -                 | -               |
|                                     | 3    | -     | -     | -                 | -     | -                 | -               |
|                                     | 4    | -     | -     | -                 | -     | -                 | -               |
|                                     | 5    | -     | -     | -                 | -     | -                 | -               |
|                                     | Mean | 21    | 35    | 14                | 55    | 20                | 34              |
|                                     | SD   | -     | -     | -                 | -     | -                 | -               |

(-) = No data available due to mortality.

- 58 -

## APPENDIX VII

Feed Consumption (g/bird/day) by Pen from a Northern Bobwhite Acute  
Dietary Toxicity Study with PFOS  
Page 1

| Experimental Group<br>(ppm a.i.) | Pen  | Exposure Period | Post-Exposure Period |
|----------------------------------|------|-----------------|----------------------|
|                                  |      | Day 0-5         | Day 6-8              |
| Control                          | 1    | 9               | 12                   |
|                                  | 2    | 7               | 12                   |
|                                  | 3    | 8               | 7                    |
|                                  | 4    | 12              | 12                   |
|                                  | 5    | 9               | 9                    |
|                                  | 6    | 10              | 9                    |
|                                  | Mean | 9               | 10                   |
|                                  | SD   | 2               | 2                    |

- 59 -

## APPENDIX VII

Feed Consumption (g/bird/day) by Pen from a Northern Bobwhite Acute  
Dietary Toxicity Study with PFOS

Page 2

| Experimental Group<br>(ppm a.i.)            | Pen  | Exposure Period | Post-Exposure Period |
|---------------------------------------------|------|-----------------|----------------------|
|                                             |      | Day 0-5         | Day 6-8              |
| 18.3                                        | 1    | 9               | 10                   |
|                                             | 2    | 8               | 11                   |
|                                             | Mean | 9               | 11                   |
| 36.6                                        | 1    | 8               | 14                   |
|                                             | 2    | 7               | 9                    |
|                                             | Mean | 8               | 12                   |
| 73.2                                        | 1    | 10              | 12                   |
|                                             | 2    | 10              | 14                   |
|                                             | Mean | 10              | 13                   |
| 146                                         | 1    | 7               | 8                    |
|                                             | 2    | 10              | 12                   |
|                                             | Mean | 9               | 10                   |
| 293                                         | 1    | 4               | 8                    |
|                                             | 2    | 6               | 10                   |
|                                             | Mean | 5               | 9                    |
| 586                                         | 1    | 6               | 16                   |
|                                             | 2    | 5               | 22                   |
|                                             | Mean | 6               | 19                   |
| 1171                                        | 1    | 4               | -                    |
|                                             | 2    | 4               | -                    |
|                                             | Mean | 4               | -                    |
| - No data available due to adult mortality. |      |                 |                      |

- 60 -

## APPENDIX VII

Feed Consumption (g/bird/day) by Pen from a Northern Bobwhite Acute  
Dietary Toxicity Study with PFOS  
Page 3

| Experimental Group<br>(ppm a.i.) | Pen  | Exposure Period | Post-Exposure Period |
|----------------------------------|------|-----------------|----------------------|
|                                  |      | Day 8-15        | Day 15-22            |
| Control                          | 1    | 10              | 14                   |
|                                  | 2    | 10              | 12                   |
|                                  | 3    | 7               | 12                   |
|                                  | Mean | 9               | 13                   |
|                                  | SD   | 2               | 1                    |

| Experimental Group<br>(ppm a.i.) | Pen | Exposure Period | Post-Exposure Period |
|----------------------------------|-----|-----------------|----------------------|
|                                  |     | Day 8-15        | Day 15-22            |
| 18.3                             | 1   | 10              | 12                   |
| 36.6                             | 1   | 14              | 15                   |
| 73.2                             | 1   | 13              | 15                   |
| 146                              | 1   | 11              | 14                   |
| 293                              | 1   | 8               | 9                    |

- 61 -

#### APPENDIX VIII

#### CHANGES TO PROTOCOL

1. The protocol was amended to indicate that bile will be collected from all study birds. The protocol was clarified to indicate the collection of liver from birds that died during the course of the study
2. Blood samples were collected on Day 8 and Day 22 in non-heparinized 5 ml borosilicate glass test tubes. The protocol indicated that heparinized vacutainers would be used.
3. The protocol was amended to change the test concentrations from 0, 20, 40, 80, 160, 640 and 1280 ppm a.i., to 0, 18.3, 36.6, 73.2, 146, 293, 586 and 1171 ppm a.i. Test concentrations were changed to reflect the test substance purity given in the new certificate of analysis.
4. The temperatures from several brooder units on Day 4, 7, 8, and all brooder units on Day 22 were not recorded
5. The afternoon observations were inadvertently not recorded for 2 birds in the 640 ppm a.i. treatment group on April 26, 1999.

- 62 -

## APPENDIX IX

## PERSONNEL INVOLVED IN THE STUDY

The following key Wildlife International Ltd. personnel were involved in the conduct or management of this study:

- (1) Mark Jaber, Wildlife Toxicologist
- (2) Joann B. Beavers, Director, Avian Toxicology
- (3) Sean P. Gallagher, Senior Biologist
- (4) Courtney Casey, M.S., Senior Biologist
- (5) Willard B. Nixon, Ph.D., Manager, Analytical Chemistry
- (6) Timothy Z. Kendall, Supervisor, Analytical Chemistry
- (7) Raymond L. Van Hoven, Ph.D., Scientist
- (8) Ellen Mank, Chemist