

# The New England Journal of Medicine

Established in 1812 as The NEW ENGLAND JOURNAL OF MEDICINE AND SURGERY

VOLUME 271

OCTOBER 22, 1964

NUMBER 17



## Original Articles

- Pelvic Angiography in the Diagnosis of Malignant Trophoblastic Disease . . . . . 859  
*J. P. de V. Hendrickse, W. P. Cockshott, K. T. E. Evans and C. J. Barton*

- Carbohydrate Metabolism in Pregnancy. V. The Interrelations of Glucose, Insulin and Free Fatty Acids in Late Pregnancy and post Partum . . . . . 866  
*Sheldon J. Bleicher, John B. O'Sullivan and Norbert Freinkel*

- Benzene and Leukemia . . . . . 872  
*Enrico C. Vigliani and Giulio Saita*

- Reversal of Digitalis Intoxication by Beta-Adrenergic Blockade with Pronethalol . . . . . 877  
*R. R. Taylor, C. I. Johnston and Anthony D. Jose*

## Special Article

- The Training of the Physician: Continuing Education — Postgraduate Medical Institute . . 883  
*Laurence B. Ellis*

## Medical Progress

- Abdominal Surgery (To be concluded) . . . . 885  
*Claude E. Welch*

## Medical Intelligence

- Current Concepts: Treatment of Ulcerative Colitis . . . . . 891  
*Louis Zetzel*

- Ventricular Fibrillation during Interim Pacing of the Heart by a Bipolar Endocardial Electrode: Report of a Case . . . . . 895  
*S. Moulopoulos, K. Kehayoglou, S. Stamatelopoulos and D. Sideris*

## Case Records of the Massachusetts General Hospital

- Hemolytic Anemia and Valvular Heart Disease 898  
*William Dameshek and Sanford I. Roth*

## Editorials

- "Sex Information and Education Council" . . 906  
"Varium et Mutabile Semper Femina" . . . 907  
Metropolitan Gets out the Vote . . . . . 907  
A Hospital is a Hospital . . . . . 907

- Massachusetts Medical Society . . . . . 908

- Massachusetts Department of Public Health . . . . . 908

## Correspondence

- Evaluation of Lymphangiography . . . . . 908  
Macroglossia in Acromegaly . . . . . 909  
Equal Opportunity . . . . . 909  
Suppression of Lipid Deposition . . . . . 909  
Training for General Practice . . . . . 910  
Osteolytic vs. Osteoblastic . . . . . 910  
Hypertension in Families of Toxic Patients . 911  
Trishydroxy Methyl Amino Methane (That Is) . 911  
Granby Coppers . . . . . 912  
Autopsies — Another View . . . . . 912  
Logomachy . . . . . 912  
Training for General Practice . . . . . 912  
Scleral Icterus . . . . . 913

- Books Received . . . . . 913

- Nonclinical Notes . . . . . 914

- Notices . . . . . xxx

Owned and Published by The Massachusetts Medical Society

Second-Class postage paid at Boston, Mass. and at additional mailing office.  
Published weekly at 8 Fenway, Boston, 02215. Domestic, \$8.00 per year.



unaltered disposition of glucose despite concurrent elevations of plasma insulin are interpreted as consistent with peripheral resistance to insulin during late pregnancy. Contrainsulin factor or factors of a lipolytic nature have been implicated. The metabolic interactions are designated as a form of "accelerated starvation," and the parasitization of maternal glucose and gluconeogenic precursors by the conceptus is emphasized.

It is postulated that the contributions from contrainsulin factors would be additive to the effects of intraplacental insulin degradation in the gestational challenges to maternal insulin economy.

We are indebted to Miss Carolann D'Amico for technical assistance and to Miss Claire Mahan for assistance in the statistical evaluation of the data.

### REFERENCES

- Freinkel, N., and Goodner, C. J. Insulin metabolism and pregnancy. *Arch. Int. Med.* 109:235-244, 1962.
- Kyle, G. C. Diabetes and pregnancy. *Ann. Int. Med.* 59:1-82, 1963.
- Freinkel, N., and Goodner, C. J. Carbohydrate metabolism in pregnancy. I. Metabolism of insulin by human placental tissue. *J. Clin. Investigation* 39:116-131, 1960.
- Goodner, C. J., and Freinkel, N. Carbohydrate metabolism in pregnancy: degradation of insulin by extracts of maternal and fetal structures in pregnant rat. *Endocrinology* 65:957-967, 1959.
- Idem. Carbohydrate metabolism in pregnancy. IV. Studies on permeability of rat placenta to I-131-insulin. *Diabetes* 10:383-392, 1961.
- Idem. Carbohydrate metabolism in pregnancy: turnover of I-131-insulin in pregnant rat. *Endocrinology* 67:862-872, 1960.
- Freinkel, N. Effect of pregnancy on insulin homeostasis. *Diabetes* 13:260-267, 1964.
- Johnson, D. G., and Bonsnes, R. W. Intravenous glucose tolerance test in pregnancy. *J. Clin. Investigation* 27:745-748, 1948.
- Silverstone, F. A., Solomons, E., and Rubricius, J. Rapid intravenous glucose tolerance test in pregnancy. *J. Clin. Investigation* 40:2180-2189, 1961.
- Nelson, N. Photometric adaptation of Somogyi method for determination of glucose. *J. Biol. Chem.* 153:375-380, 1944.
- Dole, V. P. Relation between non-esterified fatty acids in plasma and metabolism of glucose. *J. Clin. Investigation* 35:150-154, 1956.
- Yalow, R. S., and Berson, S. A. Immunoassay of endogenous plasma insulin in man. *J. Clin. Investigation* 39:1157-1175, 1960.
- Freinkel, N., et al. Alcohol hypoglycemia. I. Carbohydrate metabolism of patients with clinical alcohol hypoglycemia and experimental reproduction of syndrome with pure ethanol. *J. Clin. Investigation* 42:1112-1133, 1963.
- Berson, S. A., Yalow, R. S., Bauman, A., Rothschild, M. A., and Newerly, K. Insulin-<sup>125</sup>I metabolism in human subjects: demonstration of insulin binding globulin in circulation of insulin treated subjects. *J. Clin. Investigation* 35:170-190, 1956.

## BENZENE AND LEUKEMIA\*

ENRICO C. VIGLIANI, M.D.,† AND GIULIO SAITTA, M.D.‡

MILAN, ITALY

YPOREGENERATIVE and aplastic types of hemopathy brought about by exposure to benzene were described in the nineteenth century, and the question of leukemic myelosis from the same cause has been debated since the first description of

- Snedecor, G. W. *Statistical Methods: Applied to experiments in agriculture and biology*. Fifth edition. 534 pp. Iowa: Iowa State Univ. Press, 1956.
- Grumbach, M. M., and Kaplan, S. L. Amino acid and alpha-keto acid-induced hyperinsulinism in leucine-sensitive type of infantile and childhood hypoglycemia. *J. Pediatr.* 57:346-362, 1960.
- Mabry, C. C., DiGeorge, A. M., and Auerbach, V. H. Leucine-induced hypoglycemia. II. Studies concerning other amino acid and leucine metabolites. *J. Pediatr.* 57:539-547, 1960.
- Spellacy, W. N., Goetz, F. C., and Greenberg, B. Z. Placental transfer of insulin. *Clin. Research* 11:300, 1963.
- Hagen, A. Blood sugar findings during pregnancy in normal and possible pre-diabetics. *Diabetes* 10:438-444, 1961.
- Burt, R. L. Plasma nonesterified fatty acids in normal pregnancy and puerperium. *Obst. & Gynec.* 15:460-464, 1960.
- Idem. Peripheral utilization of glucose in pregnancy and puerperium. *Obst. & Gynec.* 4:58-66, 1954.
- Leake, N. H., and Burt, R. L. Insulin-like activity in serum during pregnancy. *Diabetes* 11:419-421, 1962.
- Spellacy, W. N., and Goetz, F. C. Plasma insulin in normal late pregnancy. *New Eng. J. Med.* 268:988-991, 1963.
- Freinkel, N. Carbohydrate metabolism in pregnancy. VI. Possible contributions of conceptus to changing pattern of maternal metabolism. In *The Nature and Treatment of Diabetes Mellitus*. New York: Excerpta Medica Foundation, 1964 (in press).
- Bleicher, S. J., Moldow, C. F., Scherrer, J., and Goldner, M. O. Lipid-mobilizing substance in serum of pregnant women, of probable placental origin. *Metabolism* 13:583-586, 1964.
- Fritz, I. B. Factors influencing rates of long-chain fatty acid oxidation and synthesis in mammalian systems. *Physiol. Rev.* 41:52-129, 1961.
- Hagerman, D. D., and Villee, C. A. Transport functions of placenta. *Physiol. Rev.* 40:313-330, 1960.
- Van Duyne, C. M., and Havel, R. J. Plasma unesterified fatty acid concentration in fetal and neonatal life. *Proc. Soc. Exper. Biol. & Med.* 102:599-602, 1959.
- Roth, J., Glick, S. M., Yalow, R. S., and Berson, S. A. Hypoglycemia: potent stimulus to secretion of growth hormone. *Science* 140:987, 1963.
- Unger, R. H., Eisentraut, A. M., and Madison, L. L. Effects of total starvation upon levels of circulating glucagon and insulin in man. *J. Clin. Investigation* 42:1031-1039, 1963.
- Josimovich, J. B., and MacLaren, J. A. Presence in human placenta and term serum of highly lactogenic substance immunologically related to pituitary growth hormone. *Endocrinology* 71:209-224, 1962.
- Kaplan, S. L., and Grumbach, M. M. Studies of human and simian placental hormone with growth hormone-like and prolactin-like activities. *J. Clin. Endocrinol.* 24:80-100, 1964.
- Greenwood, F. C., Hunter, W. M., and Klopfer, A. Assay of human growth hormone in pregnancy, at parturition and in lactation: detection of growth-hormone-like substance from placenta. *Brit. Med. J.* 1:22-24, 1964.
- Randle, P. J., Garland, P. B., Hales, C. N., and Newsholme, E. A. Glucose fatty-acid cycle: its role in insulin sensitivity and metabolic disturbances of diabetes mellitus. *Lancet* 1:785-789, 1963.
- Hales, C. N., and Randle, P. J. Effects of low-carbohydrate diet and diabetes mellitus on plasma concentrations of glucose, non-esterified fatty acid, and insulin during oral glucose tolerance test. *Lancet* 1:790-794, 1963.
- Schalch, D. S., and Kipnis, D. M. Impairment of carbohydrate tolerance by elevated plasma free fatty acids. *J. Clin. Investigation* 43:1283, 1964.
- Madison, L. L., Mebane, D., Unger, R. H., and Lochner, A. Hypoglycemic action of ketones. II. Evidence for stimulatory feedback of ketones on pancreatic beta cells. *J. Clin. Investigation* 41:408-415, 1964.
- Wulff, J., and Madison, L. Role of liver and peripheral tissues in reduced glucose tolerance of prolonged starvation. *Clin. Research* 12:283, 1964.

\*From the Clinica del Lavoro "Luigi Devoto" of the University of Milan.

†Professor of industrial medicine, University of Milan.

‡Member, Institute of Occupational Health, Clinica del Lavoro.

a case of leukemia referred to the action of benzene.<sup>1</sup>

A paper by Cronkite<sup>2</sup> (1961) summarizes the natural history of "benzene leukemia." From this and other American papers on benzene poisoning it seems that the European and especially the Italian literature on benzene leukemias is poorly known. Therefore, we thought that it would be of some interest to give a short account of the cases of leukemia attrib-

uted to benzene in Italy, those seen at the Clinica del Lavoro of Milan.

In 1938 Penati and others, in a review of the world literature, pointed out that, in so far as benzene was concerned, the diagnosis of hyperplastic bone marrow was difficult.

Although all types of leukemia have been attributed to benzene, many of the cases reported in 1938 lack reliable confirmation. Since 1941, 13 cases of leukemia have been reported in Italy, 12 at the Clinica del Lavoro of Milan and 1 at the Institute of Occupational Health of Padua.

Of these 13 cases, 9 were of the myeloblastic, according to the morphological picture. The remainder were of the myelocytic (2 cases), erythremia (2 cases). In all cases in which the clinical and severe exposure to benzene was hemocytometrically confirmed by a period of aplasia or leukopenia.

In the province of Mantova, in the case of benzene poisoning arising from the use of benzene in the manufacture of nitrobenzene, which was also at workers' homes, the Institute of Occupational Health of the University of Padua, in the last three years 41 cases of leukemia, in connection with hemopoietic disorders — more precisely myeloblastic and myelocytic — came from factories where benzene was used. It also occurred and when the percentage of benzene in the air was high. In the factories, workers with leukemia were found well above the ma-

### CASES SEEN AT THE CLINICA DEL LAVORO

Between 1942 and 1956, 13 cases of leukemia and leukopenia were seen at the Clinica del Lavoro, 12 of which were of the myeloblastic type, 2 of the myelocytic type, and 1 of the erythremic type.

We present the clinical picture of 6 cases of leukemia.

Case 1.\* M.G., a 49-year-old man, employed in a factory in 1933 as a worker applying resins dissolved in benzene. The working conditions were bad, and the air was very polluted. After 1 year the worker developed pallor, with mucosae and skin becoming dry. In 1940, he had repeated attacks of fever and was admitted to our clinic. He had previously reported by Vigliani.

uted to benzene in Italy, with special emphasis on those seen at the Clinica del Lavoro of the University of Milan.

In 1938 Penati and Vigliani<sup>3</sup> reviewed the available world literature on benzene myelopathy and pointed out that, in some cases, the pathology hardly bore out the diagnosis of aplastic myelosis — that is, hyperplastic bone marrow and leukemoid reaction.

Although all types of leukemia have been attributed to benzene, many cases reported between 1928 and 1938 lack reliable data on exposure to benzene.

Since 1941, 13 cases of "benzene leukemia" have been reported in Italy, besides the cases of the Clinica del Lavoro of Milan and the Institute of Occupational Health of Pavia, which will be considered later.

Of these 13 cases, 9 were hemocytoblastic (or myeloblastic, according to Naegeli's terminology) in type. The remainder were chronic myeloid leukemia (1 case), erythremia (2 cases) and erythroleukemia (1 case). In all cases in which a history of prolonged and severe exposure to benzene could be obtained, the leukemia was hemocytoblastic in type, often preceded by a period of apparently aplastic anemia with leukopenia.

In the province of Pavia many cases of chronic benzene poisoning arose from the use of glues containing benzene in the manufacturing of shoes. Shoes are manufactured in many small factories, and very often also at workers' homes. The Institute of Occupational Health of the University of Pavia saw in the last three years 41 cases of chronic benzene intoxication with hemopathy. Five of them were leukemias — more precisely, 3 hemocytoblastic, 1 paramyeloblastic and 1 myeloblastic. All these cases came from factories where cases of aplastic anemia also occurred and where the glues contained a high percentage of benzene. According to the determinations of benzene vapor carried out in some factories, workers with glue were exposed to concentrations well above the maximum allowable value.

#### CASES SEEN AT THE CLINICA DEL LAVORO

Between 1942 and 1963 we observed 47 patients with benzene hemopathy. Two showed anemia, 7 anemia and leukopenia, 5 anemia and thrombocytopenia, 26 pancytopenia, 6 leukemia, and 1 leukemoid reaction.

We present the clinical and laboratory details of the 6 cases of leukemia.

CASE 1.\* M.G., a 49-year-old man, entered a leather-goods factory in 1933 as a spreader and calender operator, applying resins dissolved in benzene and acetone. Working conditions were bad, and many workers had fallen ill with leukemia. After 1 year the patient complained of asthenia and pallor, with mucosanguineous diarrhea, and from 1938 and 1940, he had repeated epistaxis. His condition worsened, and he was admitted to our department in January, 1942.

\*Previously reported by Vigliani and Saita.<sup>4</sup>

Physical examination showed pallor, an enlarged liver and a spleen palpable 2 fingerbreadths below the left costal margin.

Examination of the blood revealed a red-cell count of 1,410,000, with a hemoglobin of 34 per cent (Sahli), and a white-cell count of 4800, with 19 per cent hemocytoblast forms (including many microhemocytoblast forms), and occasional neutrophilic myeloblast forms, promyelocytes and myelocytes. The platelet count was 56,000. Sternal-marrow puncture yielded 71 per cent hemocytoblast forms, of which 36 per cent were microhemocytoblast forms.

No improvement occurred in spite of treatment. Hemorrhages, fever and leukocytosis (white-cell count of 34,300, with 50 per cent hemocytoblast forms) supervened, and death followed after 1½ months.

CASE 2.† In 1940 R.F., a 38-year-old man, became an assistant operator in a rotogravure firm, where the inks contained 40 per cent benzene. The benzene concentration in the department where he worked varied between 0.60 and 2.10 mg. per liter. Asthenia and pallor developed in 1944. He had pain in the bones, with fever, in 1945, when he entered the Clinica with pallor, slight splenomegaly and some ecchymoses and petechiae.

The red-cell count was 1,450,000, with a hemoglobin of 38 per cent (Sahli), and the white-cell count 3700, with 9 per cent hemocytoblast forms and some immature myelocytes; the platelet count was 50,000. The red-cell count fell to 800,000, and the white-cell count to 2200, with 33 per cent hemocytoblast forms, mostly micro forms. The spleen enlarged to the transverse umbilical line, and there were diffuse hemorrhages. The patient died in May, 1945. Post-mortem findings were myeloid (hemocytoblastic) metaplasia of the liver and spleen; the sternal, costal and vertebral bone marrow was atrophic. Other cases of benzene anemia and a case of leukemia (Case 4) occurred in the same factory.

CASE 3. T.S., a 29-year-old man, for 8 years was a spray varnisher with nitrocellulose varnishes dissolved in solvents containing 60 per cent benzene. In 1956 he began to complain of asthenia and loss of appetite. In July, 1958, high fever and diffuse pain in the bones developed. Our diagnosis was hemocytoblastic leukemia with leukopenia. Remission followed treatment with cortisone. In November, 1958, his condition appeared fairly good: the spleen was scarcely palpable; the red-cell count was 4,800,000, with a hemoglobin of 90 per cent (Sahli), and the white-cell count 47,800, with 5 per cent hemocytoblast forms and numerous neutrophilic myelocytes and metamyelocytes; the platelet count was 150,000. Sternal-marrow puncture revealed a high concentration of hemohistioblast and hemocytoblast forms.

On treatment with 6-mercaptopurine and prednisone the white-cell count fell to 3600, with a few myelocytes and metamyelocytes. Cessation of antimetabolites was followed in a short time by a rise in the count to 200,000, with 64 per cent hemocytoblast forms, which could be controlled by renewed treatment with 6-mercaptopurine. The patient was discharged in a satisfactory condition with a good hematopoietic response, a normal white-cell count and very few hemocytoblast forms. He was then lost to follow-up study.

CASE 4. At 24 years of age, in 1938, M.A. began to work in the same rotogravure department as Case 2. In 1945, when his mate, R.F., died from benzene leukemia and other workers also had signs of benzene poisoning, this man was not examined. During 1949 he showed a slight tendency to leukopenia (count of 4000) and relative neutropenia (count of 50 per cent). After 1949 benzene was replaced by toluene, xylene, hexane and organic acetates, and no further cases of benzene poisoning have since been found. In the course of routine examinations M.A. was seen every 3 months and was reported normal until February, 1961, when he began to complain of asthenia, sporadic bouts of fever and pallor. In March the red-cell count was 1,600,000, with a hemoglobin of 29 per cent (Sahli), and the white-cell count 30,000, with 81 per cent hemocytoblast forms, mainly of the micro type. Sternal-marrow puncture gave evi-

†Previously reported by Saita.<sup>5</sup>

dence of morphologically similar cells of hemocytoblastic type, with rare occurrence of granuloblastic and erythroblastic types. The platelet count was 83,000; the coagulation time (Howell method) was prolonged, and prothrombin utilization was markedly reduced.

Transfusion, prednisone and 6-mercaptopurine led to marked improvement; by May, 1961, the red-cell count was 4,200,000, with a hemoglobin of 80 per cent (Sahli), and the white-cell count 10,000, with rare hemocytoblast forms (3 per cent), neutrophilic myelocytes and metamyelocytes.

Examination in February, 1962 (over a year after discharge from the hospital), after a relapse, showed a red-cell count of 2,000,000, with a hemoglobin of 30 per cent (Sahli); and a white-cell count of 83,000, with 58 per cent hemocytoblast forms and very few platelets. The patient had hemorrhagic and feverish bouts. Treatment with antibiotic transfusions, prednisone and 50 mg. of 6-mercaptopurine per day led to another transitory remission (white-cell count of 4000, with 22 per cent hemocytoblast forms), but fever reappeared, with bronchopneumonic foci at the lung bases and a rise in the white-cell count to 63,000, with 57 per cent hemocytoblast forms, many of which were very atypical. At no time was either splenomegaly or lymphadenopathy discovered. Death occurred in April, 1962, more than 1 year after the clinical appearance of the disease. No autopsy was performed.

CASE 5. M.A., a 50-year-old man, from 1943 to the spring of 1962 had a job of sticking rubber ribbons to the edge of billiards with glue containing pure benzene, spread on the ribbons with a pencil. He did this work for 2 or 3 hours a day. After the spring of 1962 the benzene in the glue was replaced with a nonaromatic solvent. In November, 1961, he experienced fever, asthenia and pallor; in June examination of the blood in a town hospital showed a red-cell count of 2,650,000, with a hemoglobin of 50 per cent (Sahli), and a white-cell count of 2000, with 20 per cent neutrophils and 80 per cent mononuclear leukocytes; sternal-marrow puncture disclosed many atypical hemocytoblast forms and very few normal cells of the red and white series. He improved considerably with blood transfusions and corticosteroids and was discharged in good condition. A checkup at the Clinica del Lavoro in January, 1963, showed a red-cell count of 3,900,000, with a hemoglobin of 83 per cent (Sahli), and a white-cell count of 3900, with 42 per cent neutrophils, 4 per cent eosinophils, 48 per cent lymphocytes and 6 per cent monocytes; the platelet count was 60,000. Sternal-marrow puncture showed a normal bone marrow, with only a slight increase of basophilic erythroblast forms. During that winter the white-cell count fell progressively to 2500, and the neutrophils to 14 per cent, but no immature forms were seen in the circulating blood. At the beginning of March bouts of fever appeared; and the count fell to 600, with 15 per cent hemocytoblast forms, 1 per cent myeloblast forms, 5 per cent neutrophils, 78 per cent lymphocytes and 1 per cent monocytes. Sternal-marrow puncture disclosed very few cells, mostly of the hemocytoblast type. Corticosteroid treatment was started; the white-cell count remained between 1600 and 2400, and the hemocytoblast forms in the blood smear between 5 and 32 per cent, with no other immature form. Platelets diminished to the positivity of vascular fragility tests; the red-cell count was 2,800,000, and the hemoglobin 60 per cent (Sahli). At the end of April the patient went home and was lost sight of. At no time was enlargement of the spleen or lymph nodes noticed.

CASE 6. L.A., a 53-year-old woman, was an artificial-flower maker for 3 years, using a glue dissolved in a solvent containing 25 per cent benzene. In June, 1963, she had weakness, nausea and pallor. On admission to our department, examination of the blood showed a red-cell count of 1,800,000, with a hemoglobin of 40 per cent (Sahli); and a white-cell count of 2800, with 30 per cent neutrophils, 2 per cent eosinophils, 65 per cent lymphocytes and 3 per cent monocytes, with no pathologic cells; there were 16 erythroblast forms for every 100 white cells. The platelet count was 110,000. Sternal-marrow puncture gave a normal pattern except for a high number of mitoses. This hypoplastic anemia was treated with repeated blood transfusions.

In October there were attacks of fever, with a maximum temperature of 102.2°F. (39°C.). On October 20 the white-cell count was 51,000, with 4 per cent hemocytoblast and 8 per cent myeloblast forms, 14 per cent myelocytes, 4 per cent metamyelocytes, 29 per cent neutrophils, 2 per cent eosinophils, 1 per cent basophils, 13 per cent lymphocytes and 25 per cent monocytes. The hemocytoblast and myeloblast forms showed atypical features, with large cytoplasm and monocytoïd nucleus. A total of 3 normoblast forms for every 100 white cells were seen. Sternal-marrow puncture performed on October 24 showed that almost all cells in the marrow were hemocytoblast and myeloblast forms.

The patient's condition deteriorated rapidly in spite of therapy with 6-mercaptopurine and prednisone, and she died on October 28 with large pharyngeal and buccal ulcers. The spleen was never palpable. Autopsy demonstrated complete leukemic metaplasia of the spleen, with leukemic infiltrations of the liver, lymph nodes, kidneys, colon, lungs and brain. The leukemic cells looked like undifferentiated stem cells.

## DISCUSSION

The attribution of the cases seen at the Clinica del Lavoro to the exposure to benzene cannot be doubted. Cases 2 and 4 occurred in the same workshop, and both patients were exposed to concentrations of benzene ten times greater than the maximum allowable values generally accepted. In Cases 1, 2 and 4, which occurred in factories where other workers were affected, aplastic or hyporegenerative hemopathy developed. Cases 3 and 6 handled a diluent containing a dangerous percentage of benzene. Case 5 handled a glue dissolved in pure benzene.

The following significant points are worth emphasis:

Hematologically, the 6 cases described were similar. They were all hemocytoblastic leukemias with, sometimes, absence of elevated white-cell counts, moderate or absent splenomegaly, thrombocytopenia and invasion of the bone marrow by intensive hemocytoblastic proliferation. Thus, the picture was one of undifferentiated stem-cell leukemia, with myeloid characteristics as shown, for example, by a positive peroxidase reaction.

Leukemias of this kind, even a few years ago, ran an acute or a subacute course, with, at most, a few weeks' survival. The terminal septic and hemorrhagic phenomena today can be prevented by medicinal means so that fairly long remissions have been obtained in some cases.

It is known that microhemocytoblastic leukemias are especially sensitive to the folic acid antagonists, to antimetabolites and to cortisone, a fact that may explain the longer and more frequent remissions of infantile leukemia in which microhemocytoblast forms are more prevalent than in adult cases. On the other hand, hemocytoblastic leukemia with extremely large or atypical cells is less responsive to these therapeutic measures. Cases 3 and 4 showed many microhemocytoblast cells, with few atypical or large-sized cells. In Case 4 the latter appeared in large numbers only in the terminal stages. Case 6 had a rapid course, probably owing to the highly atypical

aspect of the and paramyelo monocytoid c

Five of the at the outset apy. In Case throughout, a leukopenia m finding leads mia may not hemocytoblast terminal phas

In Case 6 hyporegenerat also occurred Occupational described in 1963. It is p kemia develo anemia.

Thus, Case or and sporad two years. C bone marrow 4 had mild le years before th

The particu the onset of a preneoplastic, better underst carried out re cases of huma of which had cells obtained the mitoses bl of colchicine. somes of 342

cent of the cell The large me two groups of dence of delet few cells. So chromosomal c Some D, or ci merary chrom tagenetic effec and may help mia during th anemia. Actual especially the an abnormal l and perhaps re study of the cells in chroni clinically and t tered chromosc

We have ne or lymphatic benzene: all ca

aspect of the hemocytoblast (parahemocytoblast and paramyeloblast) forms, with an abundance of monocytoid cells in the circulating blood.

Five of the patients had low white-cell counts at the outset that were independent of any therapy. In Cases 2 and 5 leukopenia was a feature throughout, and we consider it probable that the leukopenia may have escaped us in Case 4. This finding leads us to think that leukopenic leukemia may not be an infrequent feature of benzene hemocytoblastic leukemias, leukocytosis being the terminal phase in many cases.

In Case 6 benzene leukemia was preceded by a hyporegenerative or aplastic phase. Such an event also occurred in some cases seen at the Institut of Occupational Health in Pavia and in the case described in the United States by De Gowin<sup>6</sup> in 1963. It is possible that many more cases of leukemia developed on a background of aplastic anemia.

Thus, Case 1 complained for eight years of palmar and sporadic epistaxis. Case 3 was asthenic for two years. Case 2 showed diffuse atrophy of the bone marrow (sternal, costal and vertebral). Case 4 had mild leukopenia and neutropenia for twelve years before the late appearance of the leukemia.

The particular aspect of these cases, in which the onset of a frank leukemia was preceded by a preneoplastic, hyporegenerative phase, may be better understood in the light of an investigation carried out recently by Pollini and Colombi<sup>7</sup> in 6 cases of human aplastic anemia due to benzene, 1 of which had a fatal outcome. They cultivated the cells obtained by sternal puncture, and observed the mitoses blocked in the metaphase by means of colchicine. Then they mapped the chromosomes of 342 dividing cells and saw that 70 per cent of the cells showed abnormalities in karyotype. The large metacentric chromosomes of the first two groups of the Denver classification gave evidence of deletions; polyploidism was limited to a few cells. Some other cells showed structural chromosomal changes, as the presence of a chromosome D, or chromosomal fragments, or supernumerary chromosomes. This finding suggests a mutagenetic effect of benzene on the blood cells, and may help to explain the appearance of leukemia during the course of a benzene hypoplastic anemia. Actually, it is known that leukemic cells, especially the most undifferentiated, often possess an abnormal karyotype. It would be interesting, and perhaps rewarding, to undertake a systematic study of the chromosomes of the bone-marrow cells in chronic benzene poisoning and to follow clinically and hematologically the patients with altered chromosome pattern.

We have never seen a case of chronic myeloid or lymphatic leukemia in workers poisoned with benzene: all cases seen by us and at the Institute

of Pavia, in which harmful and prolonged exposure to benzene vapors was ascertained, were hemocytoblastic in type. Whether chronic types of leukemia can be induced by benzene is still an open question, but we must emphasize our view that in some of these cases reported in the literature the occupational history was not convincing. The number of persons occasionally exposed to benzene or exposed to very low benzene concentrations was so high that some cases of chronic leukemia could have occurred among them as among any other working population.

We are fully aware of the fact that no final statement about the existence of a true "benzene leukemia" may be made, without a statistical analysis of the incidence of leukemia among workers exposed to benzene as compared with that among a control group of the same age, sex and living habits. Unfortunately, this analysis is particularly difficult in our cases because of the large number of shoe and other factories handling benzene, the artisan work carried out at home by many factory-employed workers and the frequent and unsuspected changes in benzene content of the glues and solvents used by the workers.

In the biennial period 1959-1961 the incidence of all forms of leukemia among the general population of Milan was 0.01 per cent; 46 per cent of these were acute leukemias, 28 per cent chronic myeloid leukemias and 26 per cent chronic lymphatic leukemias. Thus, each year 1 case of acute leukemia occurred among 20,000 people.

There is reason to believe that in the Milan and Pavia provinces no more than 5000 workers were exposed to contact with benzene and probably less than 3000 were exposed to dangerous concentrations of benzene vapors.

According to the statistics of the National Institute for Insurance against Accidents and Occupational Diseases from January, 1960, to December, 1963, cases of blood dyscrasias due to chronic benzene poisoning occurred in the provinces of Milan and Pavia as shown in Table 1.

The incidence of acute leukemias in 1962-63 showed a sharp rise, coinciding with the increase in the number of cases of benzene intoxication, and was about twenty times higher than expected,

TABLE 1. *Blood Dyscrasias Due to Chronic Benzene Poisoning in the Provinces of Milan and Pavia, 1960-1963.*

| YEAR   | NO. OF CASES | NO. OF DEATHS | ANEMIA       |               | LEUKEMIA     |               |
|--------|--------------|---------------|--------------|---------------|--------------|---------------|
|        |              |               | NO. OF CASES | NO. OF DEATHS | NO. OF CASES | NO. OF DEATHS |
| 1960   | 4            | 1             | 4            | 1             | 0            | 0             |
| 1961   | 7            | 2             | 6            | 1             | 1            | 1             |
| 1962   | 36           | 11            | 31           | 7             | 5            | 4             |
| 1963   | 21           | 12            | 15           | 6             | 6            | 6             |
| Totals | 68           | 26            | 56           | 15            | 12           | 11            |

even assuming that 5000 people were exposed to benzene. Moreover, if one considers the cases of benzene poisoning that actually occurred, 11 cases of leukemia among 68 of blood dyscrasias cannot be overlooked. The incidence of leukemia is even more striking if one considers the fatal cases: out of 26 deaths, 11 were due to leukemia, and 15 to aplastic anemia.

Our figures correspond fairly well to those collected in France by Cavignaux<sup>8</sup>; according to him, during 1960-61, 42 cases of benzene myelopathy occurred, 25 in men and 17 in women; 6 of these 42 cases were leukemias. From 1947 to 1961, 45 fatal cases of chronic benzene poisoning were recorded in France; 15 patients had acute leukemia, 22 aplastic anemia, and 2 lymphatic and 6 myeloid leukemia. These data point out the high incidence of leukemias among the cases of benzene poisoning in France and the prominent position of acute stem-cell leukemias among the cases of leukemia attributed to benzene. It may be interesting to compare the action of benzene and of ionizing radiation in producing leukemia: they seem to stand in contrasting position in that the former more readily causes the acute hemocytoblastic type, and the latter the chronic myeloid type.

Our Case 4, with a latent period of twelve years after cessation of exposure to benzene, does not permit us to attribute the disease to persistence of benzene in the bone marrow. A similar case, with a latent period of fifteen years, was recently presented by De Gowin.<sup>6</sup> On the basis of the initiation-promotion theory of the induction of neoplasms, we might regard benzene as an initiator of the leukemia process, but we have no suggestion of a possible promoter.

Hyporegenerative and pancytopenic types of hemopathy are more frequently associated with benzene than the hyperplastic-anaplastic types. The cases of hypoplastic hemopathy are almost always reversible. In our series of 31 cases of involutional myelopathy, only 2 were fatal, and 1 of these from complicating typhoid. Leukosis, on the other hand, is always fatal. We suspect that, formerly, many fatal cases diagnosed as acute or subacute hemorrhagic aleukia were really cases of acute leukopenic or aleukemic leukemias.

#### SUMMARY AND CONCLUSIONS

In the last twenty years 47 cases of benzene hemopathy have been seen at the Clinica del Lavoro of Milan; 40 were hypoplastic in type, and 2 of them were fatal. Six were hemocytoblastic leukemia. No case of chronic typical myeloid or lymphatic leukemia was seen.

Two of the 6 cases of leukemia, seen in 1942-45, were rapidly fatal. Three of the remainder, seen recently, were treated with corticosteroids and with

6-mercaptopurine and showed remissions. These favorable responses to modern therapy may have been due to the presence of microhemocytoblast forms and to the scarcity of atypical cells: in the patient who died rapidly in spite of the therapy, the hemocytoblast cells were grossly atypical.

Attention is called to the initial leukopenia in benzene leukosis and the frequency of hyporegenerative or aplastic stages before the development of a frank leukemia.

In 1 of the 6 cases, the leukemia appeared twelve years after cessation of the exposure to benzene.

A summing up of the cases of benzene myelopathy seen at the Institute of Occupational Health of Pavia and in France shows approximately the same incidence of leukemias (12 to 15 per cent) as seen at our institute and the high prevalence of stem-cell (hemocytoblastic and myeloblastic) types.

Leukemia, mainly hemocytoblastic or myeloblastic in type, may be due to the action of benzene. Great caution must be exercised before admitting the benzene etiology of chronic myeloid or lymphatic types of leukemia.

We are indebted to Dr Maurice W. Goldblatt, who translated the Italian manuscript into English and helped us with suggestions and criticisms, and to Professor Salvatore Maugeri, director of the Institute of Occupational Health of the University of Pavia, who gave us information about the cases of benzene myelopathy seen at his institute.

#### REFERENCES

1. Delore, P. and Borgomano, C. Leucémie aigue au cours de l'intoxication benzénique: sur l'origine toxique de certaines leucémies aigues et leurs relations avec les anémies graves. *3. de méd. de Lyon* 9:227-233, 1926.
2. Cronkite, E. P. Evidence for radiation and chemicals as leukemogenic agents. *Arch. Environ. Health* 3:297-303, 1961.
3. Penati, F., and Vigliani, E. C. Sul problema delle mielopatie aplastiche, pseudoplastiche e leucemiche da benzolo. *Rassegna med. indust.* 9:345-361, 1958.
4. Vigliani, E. C., and Saita, G. Leucemia emocitoblastica da benzolo. *Med. d. lavoro* 34:182-191, 1943.
5. Saita, G. Mielosi aplastica e successiva mielosi leucemica. Leucopenia, provocata da benzolo. *Med. d. lavoro* 36:143-158, 1945.
6. De Gowin, R. L. Benzene exposure and aplastic anemia and leukemia 15 years later. *J.A.M.A.* 185:748-751, 1963.
7. Pollini, G., and Colombi, R. Il danno cromosomico midollare nell'anemia aplastica benzolica. *Med. d. lavoro* 55:241-255, 1964.
8. Cavignaux, L. LCS intoxications confirmées. *Cnh. mrd. Interprof.* 2:28-31, 1962.

#### ADDITIONAL REFERENCES

- Curletto, R., and Ciconali, M. Sull'emopatia benzolica. *Med. d. lavoro* 53:505-546, 1962.
- De Dominicus, G., and Buccini, G. Contributo clinico allo studio delle emopatie da idrocarburi benzenici e problemi di prevenzione. *Minerva med.* 54:3524-3529, 1963.
- Di Guglielmo, G. Relazione introduttiva alle leucemie acute. *Atti XII Congresso Nazionale Società Ital. Emat.* Bologna 1954. Rome: Ermes, 1955.
- Di Guglielmo, G. Les hémopathies aigues dans le benzénisme chronique. Fifth Congress of Société internationale d'hématologie. *Song* 2:244-246, 1955.
- Di Guglielmo, R. Particolari aspetti clinico-morfologici di alcuni casi di mielopatie da benzolismo. *Rassegna Giuc Med.* 15:71-78, 1959.
- Galavotti, B., and Troisi, F. XI. Erythro-leukemic myelosis in benzene poisoning. *Brit. J. Indust. Med.* 7:79-81, 1950.
- Gallinelli, R., and Traldi, A. L'emopatia benzenica: tre casi di benzolismo cronico di cui due mortali (leucemia acuta, panmielotisti acuta). *Med. d. lavoro* 54:169-182, 1963.
- Paterni, T., Teodori, S., and Dotta, F. Emopatia benzolica: contributo clinico (eritromielopatia aneritremica) e concetti generali. *Politico (sez. prat.)* 61:905-913, 1954.
- Rosso, N. T. Mielopatia leucemica da benzolo. *Minerva med.* 50:600, 1941.
- Vannucchi, V. Mielosi acuta da benzolismo cronico con aspetto iniziale di aplasia e aspetto successivo di leucemia o di eritro-leucemia. *Riv. crit. clin. med.* 57:51-78, 1957.

RE

THE  
pat  
rhythm  
the pro  
talis is  
laved E  
either i  
pletion  
The  
system  
groups,  
vasocon  
vey carc  
nipheral  
blockad  
sympati  
pre sing  
protere  
animals  
digitalis  
for clini  
and mc  
has been  
venously  
to digit.  
and. by  
our expo  
patients

The 8  
mias att.  
by slow  
mg. or t  
being gi  
trocardio  
injection  
In all  
stopped  
chloride,  
by mouth  
course o  
subseque.

From th  
Hos. tal.  
Supporte  
of Australia.  
Smith K  
Research  
Director,  
Hospital.  
Kindly su  
chemically,