

POULTRY SCIENCE

November, 1965, Vol. 44, No. 6

Contents

	Page
1. Protein and Lysine Requirements of Developing Turkeys as Influenced by Pelletting <i>L. S. Jensen, G. O. Ranit, R. K. Wagstaff and J. McGinnis</i>	1435
2. Selection for Chick Comb Weight with Androgen and Gonadotrophin Stimulation <i>K. E. Nestor and R. G. Jaap</i>	1441
3. Cardiovascular Changes in Restrained Chickens <i>G. C. Whitlow, P. D. Sturkie and G. Stein, Jr.</i>	1452
4. Studies of the Chick Edema Disease. 3. <i>D. F. Flick, R. G. O'Dell and V. A. Childs</i>	1460
5. Avian Tracheal Rings <i>W. J. Mathey</i>	1465
6. The Effect of a Lysine Deficiency on Growth Characteristics, Age at Sexual Maturity, and Reproductive Performance of Meat-Type Pullets <i>E. P. Singesen, J. Nagel, S. G. Patrick, and L. D. Matterson</i>	1467
7. Effect of Varying Dietary Calcium:Phosphorus Ratios on Tibia and Femur Composition of the Chick <i>B. C. Dilworth and E. J. Day</i>	1474
8. Effect of Autoxidation on the Nutritive Value of Acidulated Soybean Soapstock in Chicks <i>B. Lipstein, P. Budowski and S. Bornstein</i>	1480
9. The Relation Between α -Tocopherol Content of the Breeder Diet and That of the Newly Hatched Chick <i>I. Bartov, P. Budowski and S. Bornstein</i>	1489
10. A Hydrostatic Tester for Egg Shell Strength <i>S. J. Sluka, E. L. Besch and A. H. Smith</i>	1494
11. Bromide and Fluoride Toxicities in the Chick <i>A. R. Doberenz, A. A. Kurnick, B. J. Hulet and B. L. Reid</i>	1500
12. Supplementation of Low Quality Fish and Meat Meals with L-lysine Hydrochloride Plus DL-Methionine <i>A. Anwar</i>	1505
13. The Influence of Calcium on Phosphorus Utilization by Chicks <i>T. S. Nelson, W. A. Hargus, N. Storer and A. C. Walker</i>	1508
14. The Relationship of Egg Shell Weight to Cracked Eggs <i>E. H. McNally</i>	1513
15. Studies on the Acidulation of Soft Phosphate <i>P. W. Waldroup, C. B. Ammerman and R. H. Hurms</i>	1519
16. Effect of Some Antibiotics and Furazolidone on Performance of Laying Chickens <i>B. W. Heywang</i>	1523
17. Heritability of Semen Quantity, Concentration and Motility in White Rock Roosters, and Their Genetic Correlation with Rate of Gain <i>M. Soller, N. Snapir and H. Schindler</i>	1527
18. Social Grouping, Stress and Resistance to Coliform Infection in Cockerels <i>H. S. Siegel and W. B. Gross</i>	1530
19. The Detection of Acute Avian Leucosis as a Contagious Disease <i>W. R. Dunlop, S. D. Kottaridis, J. R. Gallagher, S. C. Smith and R. G. Strout</i>	1537
20. Effect of Calcium on the Incorporation of Proline into Hydroxyproline of Collagen Fractions from Normal and Lathyrtic Chick Embryos <i>E. C. Naber, K. Scott and R. M. Johnson</i>	1540
21. The Fluoride Content of Commercial Broiler Backs and Necks <i>M. F. Pool, W. J. Tango and A. A. Klose</i>	1545
22. Chemical Analyses of the Blood of Chicks Infected or Endointoxicated with <i>Escherichia coli</i> <i>J. R. Cole, Jr., and F. M. Boyd</i>	1551
23. The Relationship of Egg Shape to Shell Strength <i>J. F. Richards and M. H. Swanson</i>	1555
24. Monolayer Growth of Chicken Neuroglial Cells in Tissue Culture <i>E. V. Adams</i>	1558
25. Surgical Procedure for Exteriorization of the Ureteral Openings of the Hen <i>L. Ainsworth</i>	1561

The official publication of the Poultry Science Association. A bi-monthly journal published for the purpose of advancing the scientific study of poultry. Manuscripts for publication should be sent to the Editor, University of Guelph, Guelph, Ont., Canada. All manuscripts should be typewritten and double spaced. An original and a copy, together with an abstract, are required. References should be arranged alphabetically. Tables and legends for figures must be typed on separate sheets. A charge of five dollars per printed page or fraction thereof is made to partially cover composition costs.

TERMS OF SUBSCRIPTION

Fifteen dollars (15.00) annually in advance, single copies \$2.85. The journal is sent to all members of the Poultry Science Association. All subscriptions should be sent to the Secretary-Treasurer, The Poultry Science Association, Texas A and M University, College Station, Texas. Advertising rates \$100 per page. Copyright, 1966 by the Poultry Science Association. Postage paid at Menasha, Wisconsin. Acceptance for mailing at special rate of postage provided for in Act of October 3, 1917, authorized April 4, 1932. Printed in U.S.A.



Studies of the Chick Edema Disease

3. SIMILARITY OF SYMPTOMS PRODUCED BY FEEDING CHLORINATED BIPHENYL*

D. F. FLICK, ROGER G. O'DELL AND VIRGINIA A. CHILDS
*Nutrition Research Branch, Division of Nutrition, Food and Drug Administration,
Washington, D.C. 20204*

(Received for publication February 26, 1965)

INTRODUCTION

IN EARLY attempts to chemically identify the chick edema disease factor, a number of investigators (Brew *et al.*, 1959; Wootton and Alexander, 1959; Friedman *et al.*, 1959; Harman *et al.*, 1960; and Yartzoff *et al.*, 1961) reported that the factor was heavily chlorinated. More recent reports (Friedman, 1962; Wootton *et al.*, 1962; Firestone *et al.*, 1963; and Wootton and Courchene, 1964) have confirmed and extended the early studies and have shown that the factor is quite toxic for the chick. In the latter report, Wootton and Courchene (1964) "estimated that 5 μ g. of one of the toxic factors is enough to kill one chick."

McCune *et al.* (1962) reported that a chlorinated hydrocarbon used as a plasticizer for an epoxy-resin cage coating material elicited symptoms similar to those found when chicks were fed the chick edema factor or fats containing the factor. Using Vantress-White Rock chicks, these workers reported that hydropericardium, edema (subcutaneous and abdominal), hemorrhages of internal organs and enteritis resulted when the active material, chlorinated biphenyl, was fed at a level of 0.01% or higher.

It was, therefore, considered of interest to study in more detail the responses of

chicks after they had been fed chlorinated biphenyl at various dietary levels.

EXPERIMENTAL

Day-old S. C. White Leghorn cockerels were used for these studies. On day of receipt from the hatchery, the chicks were randomized in the shipping carton, individually wing banded and weighed. They were divided into groups of twelve and placed in battery brooders with raised wire mesh bottoms. Room temperature (23.9°C.) and relative humidity (50%) were kept constant by automatic control. Incandescent lights were controlled automatically for ten hours of darkness and fourteen hours of light. Chicks were immediately placed on diets containing 0, 200 or 400 p.p.m. chlorinated biphenyl¹ (CBP). They were fed and watered *ad libitum* and were weighed twice each week.

A practical type of ration was used for these studies. The composition of the ration is reported elsewhere (Flick *et al.*, 1965). The CBP was added to cottonseed oil as carrier and thoroughly mixed, and the carrier was then mixed into the dry components of the ration.

At weekly intervals of a three-week feeding regimen six or seven chicks were randomly removed from each group, weighed, and anesthetized with ether, and

* Portions of this report were presented at the 48th Annual Meeting of the Federation of American Societies for Experimental Biology, Chicago, Illinois, April 12-17, 1964.

¹ The chlorinated biphenyl, sold as Aroclor 1242, was obtained from Monsanto Chemical Company, St. Louis, Missouri.

TABLE 1.—Mortality and pathological observations of chicks fed chlorinated biphenyl (CBP) for three weeks

CBP (p.p.m.)	No. chicks on diet	Deaths	No. chicks autopsied	Edema incidence			Tissue and organ pathology ²
				HP ¹	Abdomi- nal	Subcu- taneous	
0	21	0	6	0	0	0	None
200	24	0	10	1	0	0	Pancreas (8), kidney (5), spleen (3), skin and feathers (2) and adrenal (1).
400	24	3 ³	7	5	4	3	Pancreas (7) ⁴ , adrenal (6), kidney (5), spleen (1), heart (1) ⁵ , skin and feathers (1).

¹ Only HP volumes of >0.21 ml. were included.

² Organ changes observed included: hemorrhages, enlargement, paleness, smallness (spleen), and dermatitis and/or defeathering. Numbers in parentheses are incidence of observations/organ or tissue change.

³ Two deaths occurred during the second week and one during the third week. No autopsies on these chicks.

⁴ All animals in this group had a pale cream-colored pancreas.

⁵ Pronounced heart enlargement of a bird having hemoglobin level of 4.05 g. % and hematocrit of 19.1

CBP had a pale cream-colored pancreas. The kidneys were extensively hemorrhagic and enlarged, and the adrenals (6/7) in this group were also enlarged and hemorrhagic.

The results of HP volume determination are shown in Figure 2. At the first and second weeks, the mean HP volumes of the three groups were about the same, except that the chicks fed 400 p.p.m. CBP had

slightly elevated HP volumes at two weeks. At three weeks, the mean HP volume of the group fed 400 p.p.m. CBP was approximately 10-fold higher than that of the group fed 200 p.p.m., and the latter, in turn, was somewhat higher than that of the controls.

The results of the blood studies are shown in Table 2. Blood hemoglobin levels decreased weekly among control chicks. A comparison of levels at each week revealed that the birds fed CBP had lower hemoglobin levels than controls. The birds fed 200 p.p.m. CBP showed a slight but steady decrease in hemoglobinemia throughout the 3 week study; those fed 400 p.p.m. CBP showed a marked hypohemoglobinemia at the second week but recovered somewhat at the third week. This recovery may have been due to the development of a compensatory mechanism to overcome the inhibitory effect of CBP on hematopoiesis manifested in the earlier days of the experiment.

Hematocrit levels (Table 2) followed a pattern similar to hemoglobin levels. Maximal depression of hematocrit levels was found at the second week among the chicks fed 400 p.p.m. CBP. As with hemoglobin levels, the increased hematocrit at the third




FIG. 2. Effect of chlorinated biphenyl (CBP) on hydropericardial response (ml. $\pm$ S. E. of the mean).

compound (isomer) responsible for the chick edema disease, but to elucidate the observed relationships between toxicity and potency of CEF and the toxic nature of chlorinated aromatic compounds.

SUMMARY

Chlorinated biphenyl (CBP), fed to S. C. White Leghorn cockerels at 200 or 400 p.p.m. in a practical type of ration for three weeks, produced signs which partially simulated those of the chick edema disease factor, a chlorinated aromatic hydrocarbon.

CBP feeding was accompanied by growth depression in direct proportion to the dietary level. Edema formation in the pericardial sac and in the abdominal and subcutaneous areas was pronounced only at the 400 p.p.m. level of CBP. Transient changes in blood glucose, hemoglobin and hematocrit levels accompanied CBP feeding, particularly at the 400 p.p.m. level.

Pathological changes observed were as follows in decreasing order of incidence: cream-colored pancreas, enlarged hemorrhagic kidney, enlarged adrenal, small spleen, dermatitis and defeathering.

Relationships between the toxic nature and relative potency of CBP and the chick edema factor are discussed.

REFERENCES

- Brew, W. B., J. B. Dore, J. H. Benedict, G. C. Potter and E. Sipos, 1959. Characterization of a type of unidentified compound producing edema in chicks. *J. Assoc. Offic. Agr. Chemists*, 42: 120-128.
- Douglass, C. D., and D. F. Flick, 1961. Collaborative bioassay for chick edema factor. *J. Assoc. Offic. Agr. Chemists*, 44: 449-456.
- Firestone, D., W. Ibrahim and W. Horwitz, 1963. Chick edema factor III. Application of microcoulometric gas chromatography to detection of chick edema factor in fats or fatty acids. *J. Assoc. Offic. Agr. Chemists*, 46: 384-396.
- Firestone, D., 1964, Division of Food Chemistry, Food and Drug Administration, Washington, D. C. 20204, personal communication.
- Flick, D. F., C. D. Douglass and L. Gallo, 1963. Studies of the chick edema disease. 1. Body water distribution and effect of diet. *Poultry Sci.* 42: 855-862.
- Flick, D. F., D. Firestone and J. P. Marliac, 1965. Studies of the chick edema disease 2. Purification and biological effects of the chick edema factor. *Poultry Sci.* 44: 1214-1222.
- Flick, D. F., V. A. Childs and R. G. O'Dell, unpublished observations.
- Friedman, L., D. Firestone, W. Horwitz, D. Banes, M. Anstead and G. Shue, 1959. Studies of the chick edema disease factor. *J. Assoc. Offic. Agr. Chemists*, 42: 129-140.
- Friedman, L., 1962. Progress in the chick edema problem. *Feedstuffs*, 34 (11): 18.
- Harman, R. E., G. E. Davis, W. H. Ott, N. G. Brink and F. A. Kuehl, Jr., 1960. The isolation and characterization of the chick edema factor. *J. Am. Chem. Soc.* 82: 2078-2079.
- McCune, E. L., J. E. Savage and B. L. O'Dell, 1962. Hydropericardium and ascites in chicks fed a chlorinated hydrocarbon. *Poultry Sci.* 41: 295-299.
- Natelson, S., 1961. *Microtechniques of Clinical Chemistry*, 2nd Ed. Charles C. Thomas, Springfield, Ill., 74-79.
- Negherbon, W. O., 1959. *Handbook of Toxicology. III. Insecticides, a Compendium*, Aero Medical Laboratory, Wright Air Development Center, Air Research and Development Command, United States Air Force, Wright-Patterson Air Force Base, Ohio.
- Pudelkiewicz, W. J., R. V. Boucher, E. W. Callenbach and R. C. Miller, 1958. Some physiological responses of Broad Breasted Bronze poults to chlorinated naphthalene. *Poultry Sci.* 37: 185-187.
- Pudelkiewicz, W. J., R. V. Boucher, E. W. Callenbach and R. C. Miller, 1959. Some physiological responses of New Hampshire chickens to a mixture of penta- and hexachloronaphthalenes. *Poultry Sci.* 38: 424-430.
- Wootton, J. C., and J. C. Alexander, 1959. Some chemical characteristics of the chicken edema disease factor. *J. Assoc. Offic. Agr. Chemists*, 42: 141-148.
- Wootton, J. C., N. R. Artman and J. C. Alexander, 1962. Isolation of three hydropericardium-producing factors from a toxic fat. *J. Assoc. Offic. Agr. Chemists*, 45: 739-746.
- Wootton, J. C., and W. L. Courchene, 1964. A contribution to the knowledge of the structure