

MESOTHELIOMAS IN RATS

FOLLOWING THE INTRA-PLEURAL INOCULATION OF ASBESTOS

J. C. WAGNER, G. BERRY and V. TIMBRELL

Pneumoconiosis Research Unit of the Medical Research Council, Penarth, Glamorgan

Epidemiological studies have suggested that mesotheliomas of the pleura and peritoneum are related to exposure to asbestos dust. Pleural mesotheliomas may be produced in rats following intra-pleural inoculation of asbestos dust and experiments in animals may help in the determination of the factors influencing the occurrence of such tumours. In this paper the results of three experiments in which asbestos was inoculated intra-pleurally into rats will be given.

EXPERIMENTS

EXPERIMENT A: COMPARISON OF TYPES OF ASBESTOS

The treatments were amosite, chrysotile, crocidolite, oil-extracted crocidolite and a saline control. There were 96 S.P.F. rats per treatment and the experiment was duplicated with a similar number of Standard rats. The dose was 20 mg. of dust per rat and the experiment was started at the end of 1962.

EXPERIMENT B: VARYING DOSE EXPERIMENT

This experiment involved only chrysotile and crocidolite each at 5 doses, 0.5, 1, 2, 4 and 8 mg. per rat with 12 rats per dose per dust. Injection was during March 1965.

EXPERIMENT C: COMPARISON OF CANADIAN CHRYSOTILES

Samples of chrysotile from seven Canadian mines were used, the same chrysotile as in experiments A and B and a saline control. The dose was 20 mg. per rat, and there were 16 rats per Canadian chrysotile, 32 for the original sample of chrysotile and 48 controls. Injection was during December 1966.

All the experiments were survival experiments, i.e. the animals were only killed if they appeared to be distressed. Experiments A and B are complete but 20% of the rats in Experiment C were still living at the end of January 1969 so that the results presented of this experiment are not final.

A necropsy examination was carried out on each animal. Of the rats inoculated with asbestos an appreciable proportion developed mesotheliomas in all the experiments but we have never found this type of tumour with the control treatments.

Thus a mesothelioma can be said to be a direct result of inoculation with asbestos.

METHOD OF ANALYSIS

The treatment comparisons to be given are based on the incidence of mesotheliomas. The proportion of animals developing mesotheliomas may be used for comparison within an experiment but it has to be realized that this proportion is the result of two factors, the first being the natural mortality experienced by a group of animals whether injected with asbestos or not, and the second the increased mortality due to the risk of developing a mesothelioma. It is the latter component we are interested in and we would like to separate it from the former. With large groups of animals such as in experiment A this can be achieved by life table methods and the result of the calculations depicted by a survival curve showing the proportion of animals alive at a given time when the animals were at risk of developing a mesothelioma but the risk of natural death has been eliminated. If we knew the mathematical form of this curve we could estimate its parameters and effectively produce a smoother version of the survival curve. This would be particularly useful for experiments B and C where small groups of animals were used and it would be expected that the calculated survival curves would show large irregularities.

A model relating the induction rate of tumours with time was given by Nordling (1953) and developed by Armitage & Doll (1954) and Pike (1966) and has been used both for analysing epidemiological data and animal experiments. In our situation, if m is the age-specific death rate of animals dying with a mesothelioma at time t after inoculation the model is

$$m = ck(t-w)^{k-1} \text{ for } t > w$$

where c , k and w are constants. Two theoretical approaches lead to this form. First it may be considered as the third asymptotic extreme value distribution arising if a large

OF ASBESTOS

Penarth, Glamorgan

can be said to be a direct
with asbestos.

OF ANALYSIS

Comparisons to be given are
of mesotheliomas. The
rats developing meso-
d for comparison within
has to be realized that
the result of two factors,
the natural mortality
group of animals whether
or not, and the second
ity due to the risk of
elioma. It is the latter
interested in and we
ate it from the former.
of animals such as in
an be achieved by life
he result of the calcula-
survival curve showing
imals alive at a given time
re at risk of developing
the risk of natural death
d. If we knew the
of this curve we could
s and effectively produce
the survival curve. This
y useful for experiments
l groups of animals were
expected that the calcu-
es would show large

the induction rate of
was given by Nordling
d by Armitage & Doll
(1966) and has been used
pidemiological data and
In our situation, if m is
h rate of animals dying
at time t after inocula-

w^{k-1} for $t > w$

w are constants. Two
lead to this form. First
as the third asymptotic
bution arising if a large

number of cells are considered at risk of
malignancy and a cancer cell is formed when
the first such cell succumbs. Secondly the
model would hold if a cancer cell was the end
result of k successive cellular changes. With
both approaches the constant c is related to
the dose and w is the induction or latent
period. In our analysis we will also describe
the age-specific natural death rate as
 $\exp(a + bt)$ which gives a satisfactory fit to
our control groups and has been used in
epidemiological studies.

RESULTS

Experiment A. The percentages of rats
developing mesotheliomas are given in Table
1. We note that the SPF and Standard rats

TABLE 1: PERCENTAGE OF RATS DEVELOPING MESO-
THELIOMA—EXPERIMENT A

	SPF	Standard
Amosite	40	31
Chrysotile	64	69
Crocidolite	59	68
Oil-extracted Crocidolite	59	64

gave similar results, that amosite gave fewest
mesotheliomas and that the oil-extracted
crocidolite gave similar values to the natural
crocidolite. Calculation of the survival curve
eliminating natural death, i.e. considering only




Fig. 1. Experiment A: Survival of S.P.F.
rats with mesotheliomas after eliminating effect
of mortality due to other causes.

mesotheliomas as causing death, gives Fig. 1
for SPF rats. For convenience of presentation
the oil-extracted crocidolite has been excluded
since its results were so similar to the natural

crocidolite. For each dust the pattern was
similar, consisting of an initial period during
which no mesotheliomas were found followed
by a rapid onset of cases. However, the
length of the initial period, i.e. the induction
period, was dependent on the dust. For
Standard rats similar results were obtained
but the difference between chrysotile and
crocidolite was smaller.

The model given earlier has been fitted to
this experiment and generally gave good
fits. The constant k had values near to 3 and
may be taken as 3 without any loss of
precision (this value has been assumed to
hold for experiments B and C also). As an




Fig. 2. Experiment A: Comparison of observed
and expected distribution of deaths for croci-
dolite in Standard rats. The expected deaths
are calculated from the model given in the text
with $a = -10.44$, $b = 0.00586$, $c = 1.54 \times$
 10^{-6} , $k = 3$, $w = 332$.

illustration of the fit of the model, Fig. 2
shows the cumulative number of deaths with
and without mesotheliomas for crocidolite in
Standard rats; there is close agreement
between the observed and expected distribu-
tions.

Experiment B. Averaged over doses, 36%
and 19% of rats developed mesotheliomas for
chrysotile and crocidolite respectively. For
each dust the constant w was assumed
independent of dose. This is required on
theoretical grounds but with such a small
experiment the data cannot be used to check
this assumption very rigorously. There is a
relationship between the estimate of the
constant c and dose. With only 12 rats per
dose the scatter is large but it has been shown
that the relationship may be taken as linear,
i.e. at any age the risk of developing a
mesothelioma is proportional to amount of
dust injected. The induction period was 100
days longer for crocidolite than chrysotile.

Fig. 3 shows the observed and expected numbers of mesotheliomas. If we extrapolate the results of this experiment to a dose of 20 mg, we find that for both dusts this would lead to a higher rate of mesotheliomas than actually occurred in experiment A, the difference being significant for chrysotile but not for crocidolite.




Fig. 3. Experiment B: Observed and expected numbers of mesotheliomas (out of 12 rats) on the basis that the risk of developing a mesothelioma at any age is proportional to amount injected.

Experiment C. At this stage 69% of the rats injected with the original chrysotile have developed mesotheliomas and between 19 and 56% for the 7 samples from different Canadian mills.

Again w was assumed constant for the different forms of chrysotile and in fact the best value was almost identical with that of experiment B. Comparing the treatments in terms of the parameter c , this parameter was significantly larger for the original chrysotile than the Canadian samples and the value fitted in with what would be extrapolated from experiment B.

In Table 2 are shown the values of c , and the brucite, cobalt and chromium contents of the dusts. There is correlation between cobalt and c , but mine C, which has the lowest value

of c but an average cobalt content, does not fit into the same trend as the other mines. In fact the correlation is significant only if mine C is excluded. The correlation between c and chromium does not suffer from this disadvantage since mine C has a low chromium content. The relationship between c and

TABLE 2: VALUES OF c AND BRUCITE CONTENT EXPERIMENT C

Mine	$c \times 10^6$	Brucite Content
B	9.1	20%
D	8.7	Trace
F	8.5	Trace
E	6.9	Trace to 5%
A	5.4	Trace to 5%
H	4.3	Trace
C	2.0	Absent

chromium is shown in Table 3. The correlation is high (0.91) and significant ($p < 0.01$). Nickel, scandium and iron have also been looked at, Nickel gives a similar pattern to cobalt except that mine C is even more divergent from the relationship shown by the other 6 mines. The relationships of c with scandium and iron showed nothing of interest. Mine C is in western Canada, the other 6 in a relatively small area of eastern Canada.

TABLE 3: INTRA-PLEURAL INOCULATION OF CANADIAN CHRYSOTILES
VALUES OF c (CARCINOGENICITY) AND CHEMICAL PROPERTIES

Mine	$c \times 10^6$	Brucite Content	Cobalt (ppm)	Chromium (ppm)
Original Sample	17.7	—	—	—
B	9.1	20%	110	780
D	8.7	Trace	78	930
F	8.5	Trace	78	730
E	6.9	Trace	57	440
A	5.4	Trace to 5%	63	520
H	4.3	Trace	43	480
C	2.0	Absent	60	120

DISCUSSION

All these experiments were started before the U.I.C.C. Reference Samples became available but the seven Canadian dusts used in experiment C came from 7 of the 8 mines from which material was supplied to form the Canadian chrysotile sample. Also, the sample of chrysotile used in all three experiments was a super-fine grade from one of these

