

300 OceanGate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



March 6, 1992

Susan Belcher
5960 Canyon Ferry Road
Helena, MT 59601

Dear Ms. Belcher:

While you were on board the S/S Arco Prudhoe Bay from January 22 through January 28, 1992, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Prudhoe Bay showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Prudhoe Bay 18 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from less than 0.001 to 0.005 f/cc.

The complete report is on board the S/S Arco Prudhoe Bay and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec



March 6, 1992

Randy Ingraham
20950 SW Rock Creek Road
Sheridan, OR 97378

Dear Mr. Ingraham:


While you were on board the S/S Arco Prudhoe Bay from January 22 through January 28, 1992, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Prudhoe Bay showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Prudhoe Bay 18 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from less than 0.001 to 0.005 f/cc.

The complete report is on board the S/S Arco Prudhoe Bay and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



March 6, 1992

Edward Marz
1156 Salina Avenue
Port Charlotte, FL 33948

Dear Mr. Marz:


While you were on board the S/S Arco Prudhoe Bay from January 22 through January 28, 1992, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Prudhoe Bay showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Prudhoe Bay 18 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from less than 0.001 to 0.005 f/cc.

The complete report is on board the S/S Arco Prudhoe Bay and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



March 6, 1992

Jill Willey
24 E. Milan Street
Bath, NE 04530

Dear Ms. Willey:


While you were on board the S/S Arco Prudhoe Bay from January 22 through January 28, 1992, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Prudhoe Bay showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Prudhoe Bay 18 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from less than 0.001 to 0.005 f/cc.

The complete report is on board the S/S Arco Prudhoe Bay and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



March 6, 1992

Solomon Marini
6 Winster Fax
Williamsburg, VA 23185

Dear Mr. Marini:

While you were on board the S/S Arco Prudhoe Bay from January 22 through January 28, 1992, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Prudhoe Bay showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Prudhoe Bay 18 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from less than 0.001 to 0.005 f/cc.

The complete report is on board the S/S Arco Prudhoe Bay and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



March 6, 1992

Gerlach Gerardus
1761 Hammer Avenue #42
Norco, CA 91760

Dear Mr. Gerardus:

While you were on board the S/S Arco Prudhoe Bay from January 22 through January 28, 1992, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Prudhoe Bay showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Prudhoe Bay 18 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from less than 0.001 to 0.005 f/cc.

The complete report is on board the S/S Arco Prudhoe Bay and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



March 6, 1992

David Hanstein
P. O. Box 2352
Leucadia, CA 92024

Dear Mr. Hanstein:

While you were on board the S/S Arco Prudhoe Bay from January 22 through January 28, 1992, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Prudhoe Bay showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Prudhoe Bay 18 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from less than 0.001 to 0.005 f/cc.

The complete report is on board the S/S Arco Prudhoe Bay and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



March 6, 1992

Jean Jernigan
P. O. Box 66
Selma, OR 97538

Dear Ms. Jernigan:

While you were on board the S/S Arco Prudhoe Bay from January 22 through January 28, 1992, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Prudhoe Bay showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Prudhoe Bay 18 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from less than 0.001 to 0.005 f/cc.

The complete report is on board the S/S Arco Prudhoe Bay and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



March 6, 1992

David Littlefield
RR1, Box 244L
Gouldsboro, NE 04607

Dear Mr. Littlefield:


While you were on board the S/S Arco Prudhoe Bay from January 22 through January 28, 1992, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Prudhoe Bay showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Prudhoe Bay 18 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from less than 0.001 to 0.005 f/cc.

The complete report is on board the S/S Arco Prudhoe Bay and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

ARCO Marine, Inc.
300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



March 6, 1992

Stephen Szeibert
14650 Figueras Road
La Mirada, CA 90638

Dear Mr. Szeibert:


While you were on board the S/S Arco Prudhoe Bay from January 22 through January 28, 1992, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Prudhoe Bay showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Prudhoe Bay 18 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from less than 0.001 to 0.005 f/cc.

The complete report is on board the S/S Arco Prudhoe Bay and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



March 6, 1992

Christopher Weber
412 "12th" Street
Galveston, TX 77550

Dear Mr. Weber:

While you were on board the S/S Arco Prudhoe Bay from January 22 through January 28, 1992, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Prudhoe Bay showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Prudhoe Bay 18 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from less than 0.001 to 0.005 f/cc.

The complete report is on board the S/S Arco Prudhoe Bay and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

ARCO Marine, Inc.
300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



March 6, 1992

Todd Barr
126 Great Lake Drive
Annapolis, MD 21403

Dear Mr. Barr:

While you were on board the S/S Arco Prudhoe Bay from January 22 through January 28, 1992, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Prudhoe Bay showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Prudhoe Bay 18 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from less than 0.001 to 0.005 f/cc.

The complete report is on board the S/S Arco Prudhoe Bay and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



March 6, 1992

Donald Clarkson
410 Pasadena Way
Vancouver, WA 98661

Dear Mr. Clarkson:


While you were on board the S/S Arco Prudhoe Bay from January 22 through January 28, 1992, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Prudhoe Bay showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Prudhoe Bay 18 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from less than 0.001 to 0.005 f/cc.

The complete report is on board the S/S Arco Prudhoe Bay and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



March 6, 1992

Mark Donnell
P. O. Box 2771
Big Bear Lake, CA 92314

Dear Mr. Donnell:

While you were on board the S/S Arco Prudhoe Bay from January 22 through January 28, 1992, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Prudhoe Bay showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Prudhoe Bay 18 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from less than 0.001 to 0.005 f/cc.

The complete report is on board the S/S Arco Prudhoe Bay and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



March 6, 1992

Kevin Garnier
753 West 36th Street
San Pedro, CA 90731

Dear Mr. Garnier:

While you were on board the S/S Arco Prudhoe Bay from January 22 through January 28, 1992, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Prudhoe Bay showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Prudhoe Bay 18 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from less than 0.001 to 0.005 f/cc.

The complete report is on board the S/S Arco Prudhoe Bay and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



March 6, 1992

Robert Pierce
Box 249
Hyannis, MA 02601

Dear Mr. Pierce:


While you were on board the S/S Arco Prudhoe Bay from January 22 through January 28, 1992, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Prudhoe Bay showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Prudhoe Bay 18 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from less than 0.001 to 0.005 f/cc.

The complete report is on board the S/S Arco Prudhoe Bay and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



March 6, 1992

John Piotrowski
2377 Rose Lane
Camarillo, CA 93012

Dear Mr. Piotrowski:

While you were on board the S/S Arco Prudhoe Bay from January 22 through January 28, 1992, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Prudhoe Bay showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Prudhoe Bay 18 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from less than 0.001 to 0.005 f/cc.

The complete report is on board the S/S Arco Prudhoe Bay and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



March 6, 1992

Christopher Rowe
P. O. Box 352537
Palm Coast, FL 32135

Dear Mr. Rowe:

While you were on board the S/S Arco Prudhoe Bay from January 22 through January 28, 1992, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Prudhoe Bay showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Prudhoe Bay 18 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from less than 0.001 to 0.005 f/cc.

The complete report is on board the S/S Arco Prudhoe Bay and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



March 6, 1992

Robert Wells
1534 Glen Cove Lane
Bellingham, WA 98226

Dear Mr. Wells:

While you were on board the S/S Arco Prudhoe Bay from January 22 through January 28, 1992, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Prudhoe Bay showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Prudhoe Bay 18 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from less than 0.001 to 0.005 f/cc.

The complete report is on board the S/S Arco Prudhoe Bay and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

ARCO Marine, Inc.
300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



March 6, 1992

Paul Shortino
Arco Marine, Inc.
c/o Arco Spirit

Dear Mr. Shortino:

While you were on board the S/S Arco Prudhoe Bay from January 22 through January 28, 1992, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Prudhoe Bay showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Prudhoe Bay 18 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from less than 0.001 to 0.005 f/cc.

The complete report is on board the S/S Arco Prudhoe Bay and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



March 6, 1992

William Kay
Arco Marine, Inc.
c/o Arco Prudhoe Bay

Dear Mr. Kay:

While you were on board the S/S Arco Prudhoe Bay from January 22 through January 28, 1992, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Prudhoe Bay showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Prudhoe Bay 18 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from less than 0.001 to 0.005 f/cc.

The complete report is on board the S/S Arco Prudhoe Bay and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



March 6, 1992

Christopher Ortiz
Arco Marine, Inc.
c/o Arco Prudhoe Bay

Dear Mr. Ortiz:

While you were on board the S/S Arco Prudhoe Bay from January 22 through January 28, 1992, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Prudhoe Bay showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Prudhoe Bay 18 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from less than 0.001 to 0.005 f/cc.

The complete report is on board the S/S Arco Prudhoe Bay and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

ARCO Marine, Inc.
300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



March 6, 1992

Robert Bean
Arco Marine, Inc.
c/o Arco California

Dear Mr. Bean:

While you were on board the S/S Arco Prudhoe Bay from January 22 through January 28, 1992, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Prudhoe Bay showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Prudhoe Bay 18 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from less than 0.001 to 0.005 f/cc.

The complete report is on board the S/S Arco Prudhoe Bay and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



March 6, 1992

Donald Woodward
Arco Marine, Inc.
c/o Arco Prudhoe Bay

Dear Mr. Woodward:

While you were on board the S/S Arco Prudhoe Bay from January 22 through January 28, 1992, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Prudhoe Bay showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Prudhoe Bay 18 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from less than 0.001 to 0.005 f/cc.

The complete report is on board the S/S Arco Prudhoe Bay and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



March 6, 1992

Dan Keyes
Arco Marine, Inc.
c/o Arco Prudhoe Bay

Dear Mr. Keyes:

While you were on board the S/S Arco Prudhoe Bay from January 22 through January 28, 1992, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Prudhoe Bay showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Prudhoe Bay 18 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from less than 0.001 to 0.005 f/cc.

The complete report is on board the S/S Arco Prudhoe Bay and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



March 6, 1992

Paul Preciado
Arco Marine, Inc.
c/o Arco Prudhoe Bay

Dear Mr. Preciado:

While you were on board the S/S Arco Prudhoe Bay from January 22 through January 28, 1992, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Prudhoe Bay showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Prudhoe Bay 18 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from less than 0.001 to 0.005 f/cc.

The complete report is on board the S/S Arco Prudhoe Bay and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



March 6, 1992

Jack Hendrix
Arco Marine, Inc.
c/o Arco Prudhoe Bay

Dear Mr. Hendrix:

While you were on board the S/S Arco Prudhoe Bay from January 22 through January 28, 1992, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Prudhoe Bay showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Prudhoe Bay 18 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from less than 0.001 to 0.005 f/cc.

The complete report is on board the S/S Arco Prudhoe Bay and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

ARCO Marine, Inc.
300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



March 6, 1992

Scott Williams
16142 Twinkle Circle
Huntington Beach, CA 92649

Dear Mr. Williams:

While you were on board the S/S Arco Prudhoe Bay from January 22 through January 28, 1992, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Prudhoe Bay showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Prudhoe Bay 18 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from less than 0.001 to 0.005 f/cc.

The complete report is on board the S/S Arco Prudhoe Bay and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec