FILE NAME: Newspaper & Magazine Articles (NMA)

DATE: 1968 Oct 12

DOC#: NMA012

DOCUMENT DESCRIPTION: The New Yorker - The Magic Mineral

A REPORTER AT LARGE

THE MAGIC MINERAL

BACK in the nineteen-twenties, when the word "ASBESTOS" frequently appeared in large block letters on the fireproof drop curtains that closed off the proscenium openings of theatre stages, it was customary in certain parts of the country for youngsters attending Saturday-matinée performances of "Charley's Aunt" or "The Vagabond King" or "Mrs. Wiggs of the Cabbage Patch" to greet the raising of the asbestos curtain with the chant "All school boys eat stewed tomatoes on Saturday!" This was not, of course, an exercise in mnemonics but an expression of youthful exuberance, and, to a lesser extent, perhaps, a reflection of the tendency in an era celebrated for parochialism and flippancy to dismiss strange and exotic substances with jokes. Among older people, it was certainly common knowledge that asbestos was noncombustible, for when "Joe and Asbestos"—the syndicated comic strip about horse racing—first appeared in the Baltimore *Sun*, in 1925, the stableboy named Asbestos explained to the readership that he had acquired his odd moniker because "no matter how hot my tips are, they'll never burn a hole in your pocketbook." Such knowledge was hardly universal, however, for it was also true that a popular put-on of the flapper age consisted of solemnly assuring a susceptible theatre companion that the funny-looking word on the curtain meant "Welcome" in Latin. The fact of the matter is that although asbestos was linked one way or another in the public mind with fireproof theatre curtains, many people thought its name was a trademark, few people knew that it was being utilized in a wide variety of industrial products, only a handful of medical researchers were just then beginning to suspect that it was affecting the health of the workers who handled it, and virtually no one had any idea that it would one day prove to be one of the most important industrial causes of cancer in this country.

ASBESTOS, which comes from the Greek for inextinguishable or unquenchable, is a broad term embracing a number of fibrous silicate minerals that are found in practically every country in the world. According to geologists, chrysotile asbestos—a variety of the mineral that accounts for ninety-five per cent of the world's production—had its origin millions of years ago, when subterranean upheavals provided fractures in serpentine rock which were penetrated by water heated in the earth's interior. Subsequently, these waters dissolved some of the serpentine, and as the solutions cooled, fibrous silicate crystals began to grow within the fractures. The two other major varieties of asbestos—crocidolite and amosite—originated through somewhat different dynamic and thermal processes in metamorphic rock, and their fibres, though similar to chrysotile, possess different structural and chemical properties. Although known to man as early as the Stone Age, when it was used in pottery, asbestos was considered a novelty until the late eighteen-seventies, when it began to be mined on a commercial basis. Since then, it has achieved enormous industrial importance because of its unique and astonishing physical properties. It appears to be highly combustible, yet it can withstand the fiercest heat. It seems as perishable as grass, but by virtue of being almost immune to the forces of corrosion and decay under almost every condition of temperature and moisture (and of being resistant as well to the action of most acids, alkalies, and other chemicals) it is just about indestructible. It looks extremely fragile, yet its fibres have a tensile strength equal to that of piano wire. Apparently as light and feathery as thistle- or eiderdown, it is actually as heavy and dense as the rock from which it is extracted. In one sense, then, it is a fibre of stone. In another sense, however, it is a mineralogical vegetable whose fibres are so soft and flexible that they can be carded, spun, and woven as easily as fibres of cotton or flax.

Asbestos is the only mineral that can be woven into cloth, and its fibrous structure is, if anything, even more amazing than its remarkable ability to withstand heat. In fact, if it were not for the electron microscope, the extent to which asbestos is fibrous would be difficult to believe, for there are approximately a million individual fibrils lying side by side in a linear inch of chrysotile asbestos, whereas only thirty-eight hundred glass fibrils, such as those found in various insulation materials, or six hundred and thirty human hairs can be aligned along the same distance. Moreover, in addition to their extreme fineness, high tensile strength, unusual flexibility, spinnability, and resistance to heat and the elements, asbestos fibres possess great powers to adsorb and to



10/12/68

filter. Small wonder, then, that when asbestos, which had been known to the ancients as "the magic mineral," was, in effect, rediscovered less than a hundred years ago in the age of industrial expansion, it was put to work.

Since that time, the demand for asbestos has risen almost as fast as the stuff could be mined from the earth and milled from its host rock. In 1879, when the world's first commercial asbestos mine was opened at Thetford, in the Province of Quebec, only three hundred tons of the mineral was produced. By 1910, annual world production had jumped to thirty thousand tons; by the middle thirties, it had increased to five hundred thousand tons; and today the total output is more than four million tons a year. Up until a few years ago, asbestos was often called "the mineral of a thousand uses" by those who sold it, but this label has become obsolete, especially in the United States, where it is estimated that asbestos may have as many as three thousand different industrial applications. The United States is the largest consumer of asbestos in the world, importing nearly a million tons a year, mostly chrysotile from Canada, and since fully ninety-five per cent of this amount is used in combination with other raw materials—such as Portland cement, asphalt, and vinyl and other plastics—asbestos is present as an invisible and anonymous ingredient in a myriad of industrial products, many of which are never identified as containing asbestos even though it may constitute up to fifty per cent of their content.

The asbestos-cement industry, which is the principal consumer of the mineral, makes shingles for roofing and siding, sheets for exterior and interior walls, insulation board, clapboard, casings for electric motors, pipes to carry water, sewage, and gas, and numerous other products. Asbestos textiles are used not only in fireproof theatre curtains but also in lagging and other insulating wrapping, in conveyor belting and safety clothing, in pot holders, ironing-board covers, draperies, rugs, and motion-picture screens, as filters in gas masks and in the processing of fruit juices, acids, beer, and medicines, and as mailbags, prison-cell padding, airplane fittings, stove and lamp wicks, sparkplugs, and fire hose. The electrical equipment industry has depended for years on asbestos yarn and tape for insulating conductors ranging from fine gauge wire to heavy cable, and makes asbestos friction materials—such as brake linings and clutch fa-

ngs—are found in cars, trucks, tractors, power shovels, hoists, and other industrial machines, as are asbestos-both packings, seals, and gaskets. Asbestos papers and felts are used for roofing, for piano padding, for lining stoves, heaters, filing cabinets, military helmets, and the mufflers and hoods of automobiles, and they also are found in armored cars, carpets, and cartridges, and as boiler jackets and radiator covers. Asbestos millboard, which is a heavier version of asbestos paper, is used for acoustical ceilings, as plasterboard and fireproof wallboard, and in electric switch boxes, safes, table pads, stove mats, ovens, and dry kilns. Vinyl and asphalt floor tiles contain anywhere from thirty to fifty per cent, by weight, of short asbestos fibres, and the floor-tile industry has become the second-largest user of the mineral. Since the development of phenol-formaldehyde resins in 1909, asbestos has been used in large quantities as a reinforcement and filler by the plastics industry, and today there are countless plastic products that contain asbestos, ranging from frying-pan handles to rocket nose cones. Asbestos is also an ingredient of many paints and sealants, of compounds used for roof coating and road building, of putties, caulking, and other crack fillers, of clays used for pottery and sculpture, of artificial snow, of plaster and stucco, of fireproof insulation sprayed on structural steel, and of undercoating sprayed on automobile bodies.

As a result of these multiple and miscellaneous applications, asbestos has become practically ubiquitous in modern society. There is not an automobile, airplane, train, ship, missile, or engine of any sort that does not contain asbestos in some form or other, and it has found its way into literally every building, factory, home, and farm across the land. And, because its minuscule fibres are eminently respirable, asbestos has also found its way into the lungs of man, where, by remaining as indestructible as it does in nature, it can wreak terrible havoc.

THE adverse biological effects of asbestos were observed as early as the first century by the Greek geographer Strabo and by the Roman naturalist Pliny the Elder, both of whom mentioned in passing a sickness of the lungs in slaves whose occupation was to weave asbestos into cloth. Strabo and Pliny were calling attention for the first time in history to a disease that would one day be known as asbestosis—a form of pneumoconiosis (the

general term for all dust diseases of t
lungs) caused by the inhalation of fin
fibres and particles of asbestos—but th
ancients were much too awed by th
magic of the mineral to be concerne
with the possibility that it might con
stitute a health hazard. Indeed, the
attitude toward asbestos sometimes bo
dered on veneration. Strabo and Pl
tarch noted that the "perpetual" wick
used in the sacred and inextinguish
ble lamps of the vestal virgins we
made of asbestos, and Pliny describe
asbestos cloth, which had been used
for centuries in cremations, as the rare
and costly "funeral dress of kings." It
was generally assumed in ancient times
that asbestos was of vegetable origin—
Pliny called it "*linum vivum*," and
the second-century Greek topographer
Pausanias referred to it as "the only
linen that is not consumed by fire"—
and this misconception endured for
centuries. (Only a few years ago, a
major shareholder of an English as-
bestos company who was visiting the
source of its supplies in South Africa
asked the resident manager where the
plantations were.) The Romans were
especially impressed by the fact that
when cloth made of asbestos was ex-
posed to flame it always came out
whiter than before—hence the Latin
word for asbestos, *amiantus*, meaning
unpolluted or undefiled—and they are
said to have cleaned asbestos dinner
napkins by tossing them into the fire.

The use of asbestos in Europe ap-
pears to have diminished greatly during
the Dark Ages, although it is said that
the Emperor Charlemagne convinced
some warrior guests from a rival king-
dom that he possessed magic powers by
throwing an asbestos tablecloth into
the fire and then withdrawing it, un-
scathed, from the flames. In the latter
part of the thirteenth century, when the
indefatigable Venetian traveller Mar-
co Polo was traversing a part of Si-
beria then known as the Great Em-
pire of Tartary, he was shown some
cloth that would not burn, and was
told by his Tartar hosts that it was
made from salamander's wool. Too
wily to be taken in by a story like that,
Marco examined the material carefully
and, after making some inquiries,
learned that there was a mineral in the
mountains of the district that con-
tained threads like those of wool. In
1720, asbestos deposits were discovered
in the Ural Mountains, and forty years
later a factory was built to manu-
facture asbestos products. Because the
known uses of the mineral were few
and the demand was limited, this first
commercial attempt to revive the long-


dormant asbestos industry was a failure, but the second attempt, which took place slightly more than a century later, succeeded beyond the wildest dreams of its backers. As a result, the fact that asbestos could produce lung disease, which had been forgotten since Strabo and Pliny first recorded it around the time of Christ, soon manifested itself again.

Modern knowledge of asbestosis dates from the year 1900, when a post-mortem examination was performed by Dr. H. Montague Murray, a physician in London's Charing Cross Hospital, on the body of a thirty-three-year-old man who had worked for fourteen years in an asbestos-textile factory. The patient, found to have been suffering from severe pulmonary fibrosis, which is scarring of the lungs, had been the last survivor of a group of ten men who were working in the carding room of the factory in 1886, and since Dr. Murray found spicules of asbestos in the lung tissues at autopsy, he was able to establish a presumptive connection between the man's occupation and the disease that killed him. A few years later, an inspector in the Department of Labor at Caen, whose name was Auribault, reported the results of what was undoubtedly the first study of mortality among asbestos workers. According to Auribault, whose account appeared in the *Bulletin de l'Inspection du Travail et de l'Hygiène Industrielle*, an asbestos-weaving mill had been established in 1890 at Condé-sur-Noireau, in Calvados, and during its first five years of operation, when there was no artificial ventilation and the employees were heavily exposed to dust from the looms, fifty of them died. Moreover, the director of the factory, who had previously owned a cotton mill at Gonneville, in Manche, had recruited seventeen workers from his former staff, of whom sixteen were wiped out by what Auribault, who was not a physician, mistakenly described as chalicosis—a pneumoconiosis common among stonecutters.

Unfortunately, neither Murray nor Auribault realized the importance of their findings. Testifying before a Departmental Committee on Compensation for Industrial Diseases, in 1906, Murray said, "One hears, generally speaking, that considerable trouble is now taken to prevent the inhalation of dust, and so the disease is not so likely to occur as heretofore," and the concluding recommendation of Auribault's report was simply that people under eighteen years of age should not be allowed to work in dusty conditions. Be-

cause of ignorance about the cause of the disease, and because it manifested itself slowly and insidiously over a period of from ten to twenty years or more, asbestosis was probably more often than not misdiagnosed in the early part of the century as pulmonary tuberculosis, silicosis, or fibrosing pneumonia. In addition, coyness on the part of industry, timidity on the part of labor, and obtuseness on the part of the medical profession often combined to make early investigations into the health of asbestos workers meaningless. As a result, even though the asbestos industry was growing very rapidly in all the industrialized countries of the world, almost nothing was done to control dust levels in asbestos factories, and almost no medical information on asbestosis was gathered during the quarter of a century that followed the disclosures of Murray and Auribault—although asbestos workers were probably dying like flies of it.

It was 1924 before the first clear case of death due to asbestosis appeared in medical literature. In that year, Dr. W. E. Cooke, an English physician, who gave the disease its name, performed a post-mortem examination on a thirty-three-year-old woman patient who had started working at the age of thirteen in an asbestos-textile factory in which there was no provision for the removal of dust. By 1917, after thirteen years of exposure, she had been coughing and in bad health, and from then on her attendance at work had been intermittent because of shortness of breath and lassitude. The autopsy showed extensive pulmonary fibrosis and dense strands of abnormal fibrous tissue connecting the lungs and the pleural membranes surrounding them. In addition, Cooke noted the presence of solid yellowish-brown particles, which he called "curious bodies," in the areas of fibrosis. "We have never seen anything parallel to this in pneumoconiosis due to other dusts, nor have we been able to find such occurrence in literature," he declared, and he added, "We cannot think there is any reasonable doubt that the particles in the lung are the heavy, brittle, iron-containing fragments of asbestos fibre."



Cooke's discovery, which was published in the *British Medical Journal* of July 26, 1924, and republished in 1927, provided the point of departure for a scientific journey that started on an apparently straightforward path of occupational hygiene and that has since branched out into epidemiology, oncology, physical chemistry, physiology, and experimental pathology. As a result of studies made between 1927 and 1931, a whole literature relating to asbestosis appeared in Great Britain, which included clinical and X-ray findings, descriptions of the histology of the disease, and the hypothesis (soon widely accepted) that the strange-looking yellowish-brown bodies noticed by Cooke were asbestos fibres that had been altered by the reaction of lung tissue, and coated with a colloidal substance rich in iron. In fact, Cooke's "curious bodies" proved to be so abundant in the lungs of autopsied asbestos workers, many of which contained literally billions of them, that they quickly came to be called "asbestos bodies."

The most important investigation of this period was conducted by Dr. E. R. A. Merewether, a medical inspector for the Factory Department of the British Home Office. Between 1928 and 1929, he examined three hundred and sixty-three asbestos-textile workers out of an estimated total of twenty-two hundred asbestos workers employed in Great Britain at the time, and found that ninety-five, or slightly more than twenty-five per cent, showed evidence of suffering from pulmonary fibrosis. He also demonstrated that the incidence of fibrosis increased in direct proportion to the number of years of exposure, reaching eighty-one per cent in the group of workers who had been employed in the industry for twenty years or more. As a result of Merewether's report, Parliament passed legislation in 1931 that made asbestosis a compensable disease, required improved methods of exhaust ventilation and dust suppression in asbestos-textile factories, and instituted periodical medical examinations for workers engaged in particularly dusty processes in the asbestos-textile industry.

After this promising burst of activity, however, progress in the field of asbestosis was depressingly slow, for, as things turned out, the doctors had only begun to scratch the surface of a problem of bewildering complexity. Many investigators assumed that the new regulations would prove effective in controlling the disease, thus failing to appreciate the fact that, once inhaled,

asbestos, which is virtually indestructible, continues to react within the lungs for a lifetime, and to heed the warning of Merewether, who, when asked if two years of exposure was sufficient to cause asbestosis in a young girl, replied, "Yes, if she lives long enough." Moreover, techniques of dust sampling were primitive and inaccurate, and dust levels varied so greatly—not only from factory to factory but from area to area within a given factory—that it would have been almost impossible to establish criteria upon which to recommend a safe level of exposure. But as time went by, the generally lowered dust levels that did result from the regulations of 1931 appeared to be slowing down the attack rate of the disease, at least in Great Britain, and the life expectancy of the average asbestos-textile worker was further increased by the advent of antibiotics, which drastically reduced the death rate from infectious pulmonary diseases that often had to be feared as a complication of asbestosis. The totally unforeseeable consequence of all this was that asbestos workers, instead of dying of asbestosis, began living long enough to develop asbestos-induced cancer.

The first published suggestion of an association between asbestosis and malignant disease was made in 1935 by Dr. Kenneth M. Lynch, professor of pathology at the Medical College of South Carolina, who described the case of a fifty-seven-year-old asbestos weaver who died of asbestosis and in whose lungs cancer was also found. Because the regulations of 1931 required British coroners to conduct autopsies on men who had been suspended from work with asbestos on account of pulmonary disability, or when their length of service rendered it likely that their occupation was a contributory cause of death, pathologists in Britain were able to investigate the suggested link between asbestosis and lung cancer much more easily than their counterparts in the United States, where few states had workmen's compensation laws covering dust diseases, and where there were no specific regulations requiring post-mortem examinations of asbestos workers. The British investigations were not made the subject of official comment until 1949, however, when Dr. Merewether, who was then Chief Inspector of Factories, published a review of data that had been accumulated from 1924 to 1946, which revealed that lung cancer was present in thirty-one out of two hundred and thirty-five deaths that were caused by asbestosis or in which asbestosis was

found at autopsy. This was an incidence of slightly more than thirteen per cent, and Merewether found it highly significant, for lung cancer, according to its incidence in the general population, should only have been present in approximately one per cent of the cases. At the time, no formal claim of an association between the two diseases was made, but during the next few years an increasing amount of evidence began to establish one. Then, in 1955, Dr. Richard S. Doll, director of the Statistical Research Unit of the British Medical Research Council, completed a study of a hundred and thirteen consecutive autopsies of men with more than twenty years of exposure in a large asbestos-textile factory, which showed that the extent to which they had risked lung cancer over the whole period was eleven times as great as that of men in the general population of England and Wales. From then on, many doctors were convinced that a causal relationship between asbestos exposure and lung cancer had been demonstrated, although almost no part of the vast asbestos industry either acknowledged the relationship or found it advisable to aid the independent researchers who were investigating it.

AT this point, many epidemiologists had concluded that the hazards to human health presented by asbestos were limited to the relatively small group of heavily exposed workers engaged in making asbestos textiles. The trouble was that very few studies had been made of the tens of thousands of workers engaged in other branches of the industry, which was then expanding at an enormous rate all over the world, so that no one really had any way of knowing whether there was any danger from moderate or light occupational exposure—or, for that matter, from indirect, non-occupational exposure. Such studies were under way in a number of countries, however, and when the results began to appear, in the late fifties, it became evident that the hazard posed to man by asbestos had once again been underestimated.

The most dramatic development was the discovery of an association between exposure to asbestos and mesothelioma, an almost always malignant tumor of the pleura, which is the delicate membrane that encases the lungs, or of the peritoneum, a similar membrane that lines the abdominal cavity. Previously, mesothelioma had been so rare that it was considered a pathological curiosity. It was found in only one out of every ten thousand autopsies, it was

listed in most medical books, and it was not even coded separately in the World Health Organization's International Classification of Diseases. In 1956 and 1957, however, sixteen cases of this rare tumor—tantamount to an epidemic—were diagnosed at the Pneumoconiosis Research Unit in Johannesburg, South Africa, by a pathologist named J. Christopher Wagner. Two circumstances led Dr. Wagner and his associates to suggest that asbestos might play some part in the mysterious outbreak: asbestos bodies had been found at autopsy in the lungs of the very first case, and ten of the cases had come from a hospital to which patients suspected of suffering from tuberculosis were referred from a large crocidolite-asbestos mining area.

At the beginning, this hypothesis seemed rather tenuous, for none of the sixteen patients who suffered from mesothelioma had ever worked in asbestos mines or mills. Within a short period, however, cases of mesothelioma began to crop up elsewhere in South Africa, and when Wagner and his associates discovered that the father of one of their new mesothelioma patients had been the manager of an asbestos mine, and that the patient had played around the mine and on the adjacent dumps as a child, they undertook to reinvestigate the sixteen original cases and to obtain detailed life histories in all future cases. By the end of 1961, they had diagnosed a total of eighty-seven mesotheliomas of the pleura and two mesotheliomas of the peritoneum. Only twelve of the cases proved to have had industrial exposure to asbestos, but all the rest came from the arid, windswept region of the Cape asbestos fields, where they could easily have sustained exposure simply by living in the vicinity of asbestos mills and dumps. A particularly striking case was that of a woman who was born in a town on the asbestos fields, left the region at the age of five, and died of a pleural mesothelioma at the age of fifty-seven. Detailed questioning about her childhood revealed that she had attended a kindergarten near an asbestos dump and that she and her playmates (two of whom also died of mesothelioma) used to enjoy sliding down the dump on their way home from school. Another example was provided by the youngest patient of the series—a man who died at the age of twenty-one. He had been exposed to asbestos from the age of six weeks until he was weaned—a period during which his mother took him along with her when she went to work at a dobbing site, which is a place where

asbestos is separated from its host rock. Even among those patients who had worked in asbestos mines, the actual exposure appeared to have been slight, for in most cases the lungs showed no evidence of asbestosis but only a few asbestos bodies and fibres in the air spaces. From this, Wagner drew the chilling conclusion that mesothelioma might result from an exposure to asbestos dust that was considerably less than that needed to produce fibrosis—the condition that had always been considered essential for the development of asbestos lung cancer. (In some areas of South Africa, occupational exposure to asbestos dust was so heavy, however, that fibrosis of the lungs was almost certainly killing people before cancer could develop. According to Dr. G. W. H. Schepers, a native of South Africa who is now director of the Bureau of Laboratories in the Department of Health in Washington, D.C., conditions of industrial hygiene in the asbestos mines and factories were often "deplorable." In 1949, when Schepers was sent into the northeastern Transvaal to make the first official radiological and clinical survey of the asbestos industry for the South African government, he found barefoot Bantu children spending their days inside large jute shipping bags, where they were forced by overseers armed with whips to trample down fluffy amosite asbestos as it came cascading over their heads into the bags. Schepers found that a number of these children had developed asbestosis with cor pulmonale—right-sided heart failure, which often accompanies the disease—before the age of twelve, and he has said that he would be very surprised if any of them had lived long enough to develop either lung cancer or mesothelioma.)

In 1960, the year that Wagner published the first of his disturbing reports about mesothelioma, there were two additional developments that caused grave concern among scientists investigating the asbestos problem. One of them occurred in Finland, where Dr. Raimo Kiviluoto, a radiologist, had been surveying the general population for tuberculosis; he detected deposits of calcium on the pleura, which had previously been seen almost exclusively in asbestos workers, in four hundred and ninety-nine out of sixty-three hundred and twelve people (mostly farmers and their wives) who lived in a county where asbestos had been mined and milled for many years, and no pleural calcifications in seventy-one hundred and one people who lived in a neighboring county where there was no as-

bestos industry. The other development took place back in South Africa, where Dr. J. G. Thomson, professor of pathology at the Medical School of the University of Cape Town, was examining lung-fluid smears from five hundred consecutive autopsies to determine the extent to which the ordinary urban dweller was exposed to the inhalation of asbestos; he found asbestos bodies in twenty-five per cent of the total. Thomson's discovery astonished the medical world, for up until that time no one suspected that asbestos bodies might be present in the lungs of people who did not work in asbestos mines or factories. He pointed out, however, that there had been a tremendous increase in the use of asbestos throughout the world over the previous twenty years—three million tons were marketed in 1960—and that asbestos was being incorporated into an increasing number and variety of industrial products with which the ordinary urban dweller came into intimate daily contact. He also suggested that the air of city streets was a likely source of asbestos exposure, and that the average motor vehicle, which wears out three or four sets of asbestos brake linings and one or two asbestos clutch facings in its lifetime, might well be a major cause of the liberation of asbestos dust. Thomson reminded his colleagues that, compared to a half-life of twenty-eight years for strontium 90, the half-life of asbestos fibre was an infinity of years, and he concluded that, because of the continuous disintegration of asbestos products such as brake linings, floor tiles, and roofing materials, as well as the demolition of buildings containing numerous other asbestos products, asbestos was constantly finding its way into the atmosphere as an everlasting contaminant. He even went so far as to predict that in future decades, as the accumulation of asbestos in urban areas led inexorably to higher concentrations of asbestos fibres in the lungs of urban dwellers, asbestos-induced neoplasms would rival cigarette-induced lung cancer as a cause of death.



As might be expected, the disclosures made by Wagner, Kiviluoto, and Thomson provided a powerful stimulus for further investigation of the problem of asbestos exposure. During the next few years, a hunt for mesotheliomas and asbestos bodies was undertaken by pathologists all over the world. Thomson's observations were confirmed by studies made in Miami, in London, and in Belfast, while in Pittsburgh asbestos bodies were found in the lungs of forty-six of a hundred consecutive autopsies at Presbyterian-University Hospital. And Wagner's findings were corroborated by investigations that associated mesothelioma with exposure to asbestos in half a dozen cities in Europe and the United States. Perhaps the most striking confirmation of Wagner's data came from London, where Dr. Muriel L. Newhouse and Dr. Hilda Thompson, of the Department of Occupational Health at the London School of Hygiene and Tropical Medicine, investigated the life histories of seventy-six cases of mesothelioma that had been confirmed by autopsy or biopsy in London Hospital. To no one's surprise, thirty-one of the seventy-six patients had worked with asbestos, but eleven of those who had died had simply lived within half a mile of an asbestos factory, and nine others—seven women and two men—were relatives of asbestos workers. Most of these women had washed their husbands' work clothes regularly. The investigators learned that one of the husbands, a docker, had come home "white with asbestos" every evening for three or four years, and that his wife had brushed him down. Both of the men in this group, when boys of eight or nine years, had had sisters who worked in asbestos-textile factories. One of the sisters had been employed as a spinner from 1925 to 1936, and had died of asbestosis in 1947, at which time it was determined at an inquest that "she used to return home from work with dust on her clothes." Her brother, who had apparently sustained no other exposure to asbestos in his lifetime, died in 1956 of a pleural mesothelioma.

Further corroboration of Wagner's findings came from southeastern Pennsylvania, where Dr. Jan Lieben, then head of the Division of Occupational Health at the State Department of Health, in Harrisburg, investigated the backgrounds of forty-two people who had died of mesothelioma between 1958 and 1963. Of the forty-two patients, ten proved to have had definite occupational exposure to asbestos, three were relatives of asbestos workers, eight either lived or were employed in the vicinity of an asbestos factory, and ten

others were discovered to have had at least some minor exposure to the mineral. The ten people with minor exposure included a self-employed maker of Swiss cheese who used a five-hundred-gallon boiler covered with asbestos insulation that flaked off on contact, a salesman who had mixed and applied asbestos-cement insulation to boilers in his home, and a fourteen-year-old boy who at one time had helped his father saw up and install plasterboard during an extensive remodelling of their house.

From an epidemiological point of view, the investigations that tended to support the findings of Wagner and Thomson were weak, because they lacked a defined population as a base against which the data could be measured. On the other hand, the presence of asbestos bodies in the lungs of such a large percentage of consecutive autopsies in so many widely distributed cities of the world was sufficient to indicate that asbestos fibres were probably being inhaled by a majority of urban dwellers everywhere, and the association between asbestos and mesothelioma was striking enough to suggest that a history of exposure to the mineral was likely to be found whenever a positive diagnosis of the tumor was made. As a result, epidemiologists were forced to revise their idea that asbestos was only an industrial hazard, and to give serious consideration to Thomson's prediction that it might prove to be dangerous for untold numbers of unsuspecting people in the general community.

IN October of 1964, in order to review the data collected in the past and to discuss the problems awaiting solution in the future, the New York Academy of Sciences sponsored an international Conference on the Biological Effects of Asbestos, at the Waldorf-Astoria. The conference enabled more than four hundred scientists from all over the world to discuss the immensely complicated epidemiological and clinical problems that had appeared in the forty years since Dr. Cooke described the first clear case of asbestosis, and its planning was largely the work of one of its co-chairmen, Dr. Irving J. Selikoff, who is head of the Division of Environmental Medicine at the City University's Mount Sinai School of Medicine, director of its Environmental Sciences Laboratory, and a pioneer in the field of modern asbestos epidemiology.

Like many of his colleagues at the conference, Dr. Selikoff became inter-

ested in asbestos in a roundabout
A native of New York City, he
terned at Beth Israel Hospital in Ne
ark; did his pathology at Mount Sin
where he has been a member of t
staff since 1947; and became a che
physician at the Sea View Hospital o
Staten Island, specializing in tubercu
losis. In 1951, he participated in the
basic research on isoniazid—the anti
biotic drug that, by effectively killing
tubercle bacilli, has provided a cure for
tuberculosis—for which he receive
the American Public Health Associ
tion's Albert Lasker Award in Med
cine. In 1953, Dr. Selikoff founded
medical clinic in Paterson, New Jerse
where, by chance, seventeen of h
early patients were men referred fo
chest-disease consultations who worke
in a local factory that had been mak
ing asbestos-insulation materials sin
1940. When the factory closed down
in 1954, the seventeen men went int
other work, and at that point Dr. Seli
koff decided to continue his observa
tion of these patients with X-ray ex
aminations and pulmonary-function
tests in order to determine the histor
and the natural course of asbestosis i
men who would not be subjected t
further exposure.

At the time, Dr. Selikoff was chiefly
interested in asbestosis because, as he
has explained, "I was still unconvinced
of the relationship between asbestos and
lung cancer. There wasn't much data,
and what there was scarcely seemed to
indicate that it would prove to be
much of a problem." As things turned
out, he changed his mind. In 1954,
all seventeen men from the Paterson
factory were able-bodied and work
ing. By 1961, however, six of them
were dead, and today only six of them
are alive. Of the eleven who died,
four were victims of lung cancer, three
of other cancers, one of mesothelioma,
and two of asbestosis. Of the six who
are still alive, two are disabled becaus
of respiratory insufficiency caused by
fibrosis of the lungs, and one has re
covered from surgery for cancers of
the lung and larynx. Dr. Selikoff
shook his head in describing the fate
of these patients, but as early as 19
their ranks had been thinned out s
drastically that, as he said, "I w
forced to the awful realization of wh
could happen to men who were oc
pationally exposed to asbestos." Af
the company that owned the fact
said it was unable to make its empl
ment records available, he wrote
several other large asbestos manuf
turers in the United States, inquiring
about the health experiences in th

plants, and was unable to obtain information from any of them. Meanwhile, he and Dr. Jacob Churg, chief pathologist at Barnert Memorial Hospital, in Paterson, who had been finding asbestosis and lung cancer in a large number of workers from the Paterson factory, took their data to Dr. Roscoe P. Kandle, Commissioner of the New Jersey State Department of Health, and Dr. Kandle applied to the United States Public Health Service for funds to undertake a study of the Paterson factory and to make a statewide survey to determine how many people were being occupationally exposed to asbestos. However, the request was denied for lack of resources.

Since Dr. Selikoff already knew that men who had worked in the factory that made insulation materials were dying of asbestosis and lung cancer at an alarming rate, he considered that men who were installing such materials might also risk disease, and early in 1962 he made contact with officials of New York Local 12 and Newark Local 32 of the International Association of Heat and Frost Insulators and Asbestos Workers. The asbestos insulators had been trying for years to interest doctors and various government agencies in their medical problems, so they were only too glad to coöperate, and urged Dr. Selikoff to make a study in depth of the effects of asbestos exposure among their members.

During the next six months, Dr. Selikoff and his administrative assistant, Mrs. Janet S. Kaffenburgh, pored through the unions' records and compiled a list of the names and addresses of the six hundred and thirty-two men who had been members of either of the metropolitan locals on December 31, 1942, and of the eight hundred and ninety men who had joined between then and December 31, 1962. From the unions' employment records, they obtained detailed work histories of the total membership of fifteen hundred and twenty-two men, including data on their withdrawal from work for other employment, war service, retirement, or illness. This enabled them to calculate the onset and duration of exposure to asbestos for each worker. Records of the unions' health and welfare funds provided them with the dates and places of death of the two hundred and sixty-two workers who had died between 1942 and 1962, and copies of the death certificates of all but one of them were obtained. In addition, autopsy protocols, histological specimens, and hospital records were reviewed by Dr. Selikoff and Dr. Churg in those

aths (approximately one-half of th
tal) which occurred in hospitals.

In the next phase of the study, Dr
likoff and Dr. Churg were joine
Dr. E. Cuyler Hammond, Vice
President for Epidemiology and Sta
stics of the American Cancer Society
ho had participated in an analysis o
ne medical effects of the atomic ex
osions that devastated Hiroshima an
agasaki, in 1945, and whose large
le epidemiological studies of more
an a million men and women pro-
ded a major basis for the conclusion
awn in the 1964 Surgeon General's
port on the effects of cigarette smok-
g. Since previous studies had sug-
sted that lung cancer associated with
bestosis seldom develops until twenty
ars after initial exposure to asbestos
ust, Dr. Selikoff and Dr. Hammond
cided to limit their first report to the
hundred and thirty-two men who
ere on the union rolls as of Decem-
r 31, 1942. Taking the men's ages
to consideration, Dr. Selikoff and Dr.
ammond then set about comparing
e number and causes of death among
em with those of the general male
pulation in the United States. The
sults were depressing. According to
e standard mortality tables, two hun-
red and three deaths could have been
pected among the six hundred and
irty-two workers. Instead, there
ere two hundred and fifty-five, not
unting seven men who had died be-
re incurring twenty years of expo-
re—an excess of twenty-five per cent.
he reason for the excess was not hard
find. The fact that twelve of the
aths were attributed to asbestosis was
t particularly surprising, but only six
seven deaths from cancer of the lung,
ura, or trachea were to be expected,
hereas there were forty-five. And
here nine or ten gastro-intestinal can-
rs were to be expected there were
tually twenty-nine. Since the lung-
ncer death rate was known to
more than ten times as high
mong cigarette smokers as among
n-smokers, Dr. Selikoff and Dr.
ammond realized that they would
ve to take the smoking habits of the
bestos-insulation workers into account
their findings were to have any va-
ity. It was, of course, impossible for
em to ascertain this with accuracy in
e cases of the two hundred and fifty-
e men who had died, so, for purposes
calculation, they assumed that all six
ndred and thirty-two men had
oked a pack or more of cigarettes
ch day, and they demonstrated that
n if this had been the case it would
ve produced a lung-cancer death rate

only three and a half tu[illegible] general male popula[illegible] smoking, therefore, co[illegible] the fact that in this gro[illegible] insulation worke[illegible] from lung canc[illegible] expected rate.

Because of [illegible] and its thoroughness, this study [illegible] ducted by Drs. Selikoff, [illegible], and Hammond had a great impact on the medical community when it was published, in the spring of 196[illegible], in the *Journal of the American Medical Association*, and when it was discussed before the Conference on the Biological Effects of Asbestos that October. It was the first study ever made that had taken a large enough group of asbestos workers from a point far enough back in time and followed them long enough to determine unequivocally what their health experience had been. Unlike almost all the previous investigations, which simply indicated that there was an association between asbestos and various kinds of cancer, it was based upon the incidence of disease within a defined population, and thus answered a fundamental epidemiological question of how many cancers had developed out of how many persons exposed. In doing so, it furnished the first incontrovertible evidence that industrial exposure to asbestos was hazardous; it established sound methodology for future studies; and it marked a turning point in the views held by doctors and health officials around the world.

IN the autumn of 1962, Dr. Selikoff—using donations from private sources and a grant awarded by the Health Research Council of New York City, headed by Dr. George S. Mirick—had set up informal clinics at union halls in New York and Newark and started giving X-ray examinations, pulmonary-function tests, and blood tests to the members of the two asbestos workers' locals. During the next year, he examined eleven hundred and seventeen workers and he also questioned them closely about their smoking habits, the kinds of materials they used, and the conditions under which they worked. "The insulation workers, an intelligent and highly skilled group of craftsmen, patiently educated me concerning their trade," he once said. "I learned that they had had comparatively light and intermittent exposure as asbestos workers go, because they often worked out-of-doors on construction projects, and because most of the materials they used had an asbestos content of less than

seen per cent. I also found out how their problems had been neglected. Most of them had never seen a labor inspector in their lives, and only a handful could remember seeing anyone make a dust-level count in the area where they were at work." Dr. Selikoff also learned that although some of the workers knew they ran a risk of developing asbestosis (which they usually referred to as silicosis) almost none of them had any idea that they also risked cancer because of their occupation. As things turned out, he found radiological evidence of asbestosis in fully half of the eleven hundred and seventeen men. The extent of the fibrosis varied directly with the duration of exposure. Of three hundred and forty-six men whose exposure had begun less than ten years before, only ten per cent showed any abnormality, and of three hundred and seventy-nine men who had been exposed for between ten and nineteen years more than half still had normal X-rays; in neither group was there any discernible evidence of cancer. On the other hand, Dr. Selikoff found that among three hundred and ninety-two men with more than twenty years of exposure three hundred and thirty-nine had developed asbestosis which had by that time become moderate or extensive in more than fifty per cent of the cases; he also found that eleven of the men had developed cancers of the lung or pleura.

Aided by yearly grants from the Health Research Council, Dr. Selikoff has maintained a close watch on the asbestos workers' health, paying particular attention to those men with more than twenty years of exposure by examining them once or twice a year. As a result, he has been able to detect symptoms of illness in many of the men much earlier than they might have been detected and to observe with great accuracy the mortality experience of the three hundred and seventy whom he found in 1963 to be survivors of the original six hundred and thirty-two members of the union in 1942. It continues to be disastrous. Between January of 1963 and March of 1968, fifty deaths would normally have been expected to occur among these men. Instead, there were a hundred and thirteen. Fifteen of them were caused by asbestosis, and, as with the earlier study, the rest of the excess turned out to be caused by cancer of various kinds. Two or three bronchogenic cancers would have been normal, but twenty-eight occurred, and where the actuarial tables predicted only two gastrointestinal cancers there were eight. A

pectedly sharp rise in the number
esotheliomas was observed. In th
arlier study, there had been only fo
two hundred and fifty-five death
tween 1963 and 1968, howeve
hirteen of these rare tumors we
ound in a hundred and thirteen death
Dr. Selikoff suspects that the late
eriod for mesothelioma may be long
an that for either asbestosis or lun
ancer, but so far he regards the heav
ncrease of mesothelioma among th
nsulation workers as something of
mystery.

Dr. Selikoff and Dr. Hammon
ound special significance in the fa
hat all twenty-eight bronchogen
arcinomas occurred among asbest
workers who smoked cigarettes regu
larly. None were found among eighty
even men who never smoked cig
rettes. Dr. Selikoff and Dr. Hammond
have reservations about these data,
which suggest that asbestos worker
who smoke cigarettes are ninety time
more likely to die of lung cancer than
men who neither work with asbest
nor smoke, because they are base
upon the experience of a relatively
mall number of people. They do con
clude from it, however, that an im
portant co-carcinogenic effect exists be
tween asbestos exposure and cigarette
moking, and in a recent article pub
hed in the *Journal of the American
Medical Association* they recommend
"that asbestos exposure be minimized,
hat asbestos workers who do not smok
hould never start, and that those now
moking should stop immediately." It
as been suggested in some quarter
hat asbestos fibres, which are known
o be highly adsorbent as well as vir
ally indestructible, may retain the
ubstance known as benzo [a] pyren
rom cigarette smoke for long periods
f time in lung tissue, thus tending to
roduce lung cancer by a synergistic
ffect. No one knows for sure, however,
nd Dr. Selikoff will not hazard a gues
bout the matter until more informa
ion is gathered. He expresses certainty
bout the effects of asbestos exposure
nly when he has epidemiological proof.
He knows, for example, that one out
f every five asbestos-insulation work
rs who die in the New York-New
ersey metropolitan area dies of lung
ancer, and that almost half of the
hree hundred and sixty-eight deaths he
as recorded among these men in the
ast twenty-five years have been caused
y cancer of one type or another. Last
March, citing these figures before a
House subcommittee on labor, he
arned that "if they are projected to
he country as a whole, and if no im-

Total Recall


This Fujica Single-8 Instant Load camera has a photographic "memory". It records what you see just as you see it. Clear, bright, true-to-life.

Unique features not found in any other instant load movie system make this possible. The camera has a built-in pressure plate. You have nothing to do with it . . . but it does something important for you . . . automatically locks the film flat behind the lens to give you the sharpest movies you've ever seen outside a theatre screen.

Even if you've never taken a movie before, you get professional results easily. Drop in cartridge . . . aim . . . shoot! A full 50 continuous feet if you wish with no film flipover. Zoom in or zoom out! Fully automatic electric eye sets correct exposure. Reflex lens gets you exactly what you see. Less than $125. Other Fujica Single-8s less than $80 . . . at camera stores or write for the colorful facts.

FUJI PHOTO FILM U.S.A., Inc.
U.S. Distributor: Ehrenreich Photo-Optical Industries, Inc., Garden City, N.Y. 11530
In Canada: R. & H. Products, Ltd.

FUJICA
INSTANT LOAD
Single-8
Reflex Zoom P-300




provement in the health hazards these men face is achieved, we may look forward to perhaps six thousand deaths from lung cancer and to perhaps fifteen thousand deaths from cancer in general among men in the single asbestos trade alone." Considering that the total membership of the International Association of Heat and Frost Insulators and Asbestos Workers is only eighteen thousand men, and that there are about the same number of non-union asbestos workers, his prediction has staggering implications.

I RECENTLY spent several days with Dr. Selikoff in the Environmental Sciences Laboratory at Mount Sinai, where he has assembled a team of thirty men and women to work on various aspects of the asbestos problem. The rooms include laboratories for histology, pathology, electron microscopy, chemistry, and mineralogy, and among the imposing array of equipment they contain is an electron microprobe that is capable of identifying particles of infinitesimal size. Dr. Selikoff's office is large and modern. Its walls are lined with shelves and cabinets containing hundreds of bound volumes of research papers and reports on asbestos, and at one end are a large conference table and a blackboard. The place gives an immediate impression of rather cold efficiency, but this is tempered to a great degree by the affability of the director. As we talked, Dr. Selikoff, a large, soft-spoken man in his early fifties, was frequently interrupted by telephone calls and by questions from staff members, but he generally resumed—often with an enthusiastic "Where were we?"—and rattled off statistics about asbestos disease the way some baseball fanatics recite the batting averages of big-league ballplayers. On the day of my first visit, however, he received a call from a doctor in New Jersey who informed him that a lung cancer had just been diagnosed in another of the insulation workers. Dr. Selikoff told the physician that the patient's last X-rays showed moderate asbestosis, that he had had more than twenty years of exposure, and that he smoked a pack of cigarettes a day. When he hung up the phone, he was visibly upset, and fell silent. A few moments later, he told me that he had taken more calls like that than he cared to remember. "I don't relish the role of Cassandra, but I feel we've got to forge ahead on this problem as quickly as possible," he said. "There are about thirty-six thousand insulation workers—union and non-union—in the

J. PRESS INC.

—Squire Style—
Matching Jackets and Waistcoats



Power Loom Irish Tweed $85

Handwoven Irish Tweed $90

Irish 'Supersax' Hunting Checks $95

Tailored in Ireland Solely For J. PRESS in long center vent & steep side vented models with matching waistcoat.

Sample Swatches & New Brochure Upon Request



Turlough Finn
DUBLIN NEW YORK
SUTLER AND MANCIPLE TO THE SPORTING GENTRY
IRISH TWEED HATS & CAPS
$8.00 to $12.00
FOR RACE WATCHING, HUNTING FISHING, STROLLING, GOLF, SKATING AND ALL FASHIONABLE SPORT EVENTS
SEND FOR ORDER FORM AND PRICE LIST GRATIS
Turlough Finn
520 FIFTH AVENUE, N.Y.C., N.Y.

Pipe of Peace
P.O. BOX 418 / NASSAU / BAHAMAS
PIPES FROM AROUND THE WORLD
DUNHILL COMOY SASIENI PETERSON
BARLING KRISWILL STANWELL LONGCHAMP



Please send me your FREE Illustrated Catalogue
Name: .....
Address: .....
City: .....
State: ..... Zip: .....
Airmail to Nassau 15¢
PIPE of PEACE



Weatherproof Syl-mer treated leather boot by Joyce.

You don't have to worry about how this Sherlock Holmes corduroy coat will look after sleuthing in the rain. The SYL-MER™ brand silicones give it durable water repellency, crush and stain resistance. And a luxurious hand. And this protection lasts. So all you have to worry about is the next case, Watson.

RAINCHEETAHS by NAMAN—a great fashion look in Cone Mills Regina 100% cotton corduroy. Brown, bone. Sizes 8-14. About $65. At Lord & Taylor, New York and branches.

SYL-MER
SILICONE FINISH

Dow Corning Corporation, Midland, Michigan

United States. All in all, including asbestos-textile workers and people engaged in manufacturing other asbestos products, there are probably more than a hundred thousand workers who are regularly and directly exposed. The hazard would be serious enough if this were the only group at risk, but it's obvious that the problem extends further, simply because floating asbestos fibres do not respect job classifications. Dr. William H. Stewart, who is the Surgeon General of the United States, told the House subcommittee that the asbestos-insulation workers are probably sharing their exposure to some extent with more than three and a half million other construction workers—steam fitters, electricians, welders, carpenters, ironworkers, plumbers, masons, tile setters, and the like—who are employed on the same projects, often in the same room or general area, and who therefore must be inhaling the same dusts."


Because of their concern with the possible effects of such indirect occupational exposure, Dr. Selikoff and his assistants have spent months compiling the names, addresses, and work histories of twelve hundred tile setters, three thousand roofing workers, and three thousand tunnel, foundation, and caisson workers in the New York metropolitan area. They plan to examine these men for possible health problems as they did the insulation workers, and to determine whether they have significant indirect occupational exposure, and, if they do, whether it is associated with risk of asbestos disease. "We will study defined populations that have been selected in terms of what appear to be diminishing levels of exposure to a known risk," Dr. Selikoff told me. "It's as if a rock had been thrown into a pond and we're watching how far the ripples extend. Since the pond in our case is the general community, you can see how difficult the epidemiological detective work will be. If we discover, for example, that the tile setters run a significant risk of developing asbestos disease—and there are some preliminary indications that they might—then we must look into the question of fibre fallout from the construction sites where they're employed. We're not going to wait for the results of the tile-setter study, however, because the problem is too urgent. We're initiating a whole cluster of studies designed to provide information on whether environmental asbestos pollution carries a risk for the general population. At the same time, by analyzing the ripples, we can determine which rocks thrown into the pond



The beautiful lizards

darting along with tweeds, furs, leathers—ours alone, Newton Elkin's pumps in brown, black or grey with a twist of gilt metal. 38.00

Designer shoe collections at all Lord & Taylor stores



NEW YORK'S BELOVED

Algonquin

Superb pre-theatre dining . . . and just a stroll to your theatre. After-theatre drinks and Supper Buffet with "the talk of the town"

HOTEL Algonquin
59 WEST 44th ST. NEW YORK • MU 7-4400

Indoor parking (no charge) for dinner guests and weekend visitors.

JULES PODELL PRESENTS

JACK JONES

(DOUG TALBERT, Musical Director)

MARTY BARRIS

COPACABANA


FULL COURSE DINNERS
SERVED 6:30 P.M.-9:30 P.M.
2 SHOWS NIGHTLY 8 & 12
3 SHOWS SATS 8, 11:30 & 2

10 E. 60 • PL 8-0900

AIR CONDITIONED

make the biggest splash. By thus identifying the sources of contamination, appropriate control measures will become feasible. Our plan of attack is to discover where asbestos-associated diseases may occur, and to control the sources in such a way as to eliminate them as a hazard. It must be a team effort, of course, including epidemiologists, basic scientists, engineers, public-health officials, and industry and labor."

One of Dr. Selikoff's studies will examine the possible health problems arising from interrupted exposure by tracing thirteen hundred and sixty-seven men who worked at the asbestos-insulation plant in Paterson—closed down in 1954—where Dr. Selikoff's first seventeen patients were employed. Another will look into the matter of neighborhood exposure by tracing and investigating the health experience of seventy-five hundred people who lived within a half-mile radius of the plant in 1942. "The question of neighborhood exposure was emphasized as a result of the mesotheliomas discovered by Christopher Wagner in South Africa, and later by Molly Newhouse and Hilda Thompson in London and by Jan Lieben in Pennsylvania," Dr. Selikoff told me. "Since current evidence suggests that the tumor rarely occurs without a history of exposure to asbestos, epidemiologists consider it a 'marker' disease, and we'll be looking for it when we trace the medical history of the people in Paterson. Incidentally, there's one preliminary indication that leads us to suspect we may find it. In 1955, the proprietor of a junk yard next to the Paterson insulation plant died of a pleural tumor." A third study in Dr. Selikoff's cluster will aim at the protection of relatives of asbestos workers. As in the case of neighborhood exposure, previous investigators of this group had no population base against which to measure the occurrence of the disease they discovered. Dr. Selikoff, however, has a definitely established group to work with. It is composed of the wives and children of all the insulation workers, and their health problems will be carefully scrutinized, especially for evidence of mesothelioma. In Dr. Selikoff's opinion, this study will be of tremendous importance, because its subjects have been far removed from direct exposure. "If the families of insulation workers turn out to be in trouble, the implications will be very serious indeed," he said. "On the other hand, if we find no trouble among them—and this is my hope—we'll be on safer ground. Our observations will also enable us




Adaptation of a traditional votive candlestand in the spirit of Ch'ien Lung bronzes. 41 inches. About $100.*

The Stiffel Lamp

For brochure of selected Stiffel lamps, send 25¢ to The Stiffel Company, Chicago 60610.

*Slightly higher west and south.



violets

A perennial reminder of spring . . . violet plant abundant with oxidized green flowers and leaves. Hand-crafted in metal and mounted on a wooden base.

12" high, 12" spread. $50.

Add $3.00 for shipping outside regular delivery area.

america house

THE FINEST IN AMERICAN CRAFTS

44 W. 53rd St., N.Y. 10019 • PL 7-9494

Sometime this week—take a quiet walk along our canal in Bucks County.

Mid-week is a wonderful time to stay with us at the 1740 House, an early American inn overlooking the Delaware River. When you get here, the 1740 House will be exactly as you had hoped it would be . . . quiet, charming and memorable.

24 air-conditioned rooms, each with bath and terrace. Swimming, fishing, boating and tennis.


Please write for our brochure and driving directions. Then make reservations for a refreshing, mid-week respite from crowds and cares.

1740 HOUSE

LUMBERVILLE, PA. 18933 BUCKS COUNTY

Tel: 215-297-5661

ENJOYING STEAK AND LOBSTER IS THE GREAT INDOOR SPORT AT...



AL SCHACHT'S

10 E. 52ND ST., N.Y.C. PL 9-8570

7 DAYS A WEEK TILL 1 A.M.



Great legs—

would you know they're wearing Belle Sharmeer's Whisper Sheer support stockings? Seamless nylon-and-spandex, to own in suntan, nude, pale beige, misty taupe, dark brown or off-black —all in proportioned leg sizes, 4.95 the pair
All Lord & Taylor stores

to recommend whether special facilities should be provided so that insulation workers can change their work clothes before going home to their families."

With the aid of a grant from the United States Public Health Service, Dr. Selikoff and his associates are currently conducting an investigation that covers an even wider range, by studying samples of lung tissue from three thousand consecutive autopsies that have been performed at three different hospitals in New York City and tracing the work histories of each of the dead people. So far, they have examined two thousand lung samples, and have found asbestos bodies in nearly half of them. According to Dr. Victor Baden, a histologist at the Environmental Sciences Laboratory, from one to five or more asbestos bodies are found in fifty per cent of the tissue slides he examines. Since each slide is composed of seven sections that together represent less than one half-millionth of an entire lung, and since it is widely assumed that asbestos bodies are evenly distributed throughout a lung, the finding of just one of them can be projected to mean that half a million may be present.

Dr. Selikoff and his fellow-epidemiologists do not know whether the relatively small amount of asbestos now being inhaled by the general public carries a risk of inducing mesothelioma, or whether the presence of what appear to be asbestos bodies in the lungs of many people in the general population represents a serious public-health problem. Nor do they know whether relatively light inhalation of asbestos is associated with a special lung-cancer risk in cigarette smokers. "At present, we simply have no way of telling what the dose-disease relationships are," Dr. Selikoff said. "Is it better to have breathed in only fifty asbestos fibres than fifty thousand or fifty million? Does each fibre have the potential of causing a cell with which it comes in contact to turn cancerous? The answers to these immensely complicated questions will have to come from epidemiology over a long period of time. An eightfold increase in asbestos production over the last thirty years would appear to make it likely that asbestos cancers will increase, but only the future will tell. Remember that current cases of mesothelioma must be associated with inhalation that took place twenty, thirty, or more years ago, when world consumption of asbestos was only about five hundred thousand tons a year, and that the neoplasms associated with today's production of over four million tons

De Busschere
Belgian Shoes
MR. HENRI BENDEL
Owner-President
HAND MADE · HAND SEWN



DAISY
Black or blue calf
with lizard trim
$65

Classic, timeless styling.
Brochure on request.

BELGIAN SHOES
[illegible] 56 St., New York 10022
[illegible] Plaza 5-7372
[illegible] J. Marcus & Co.

VIRTUTE NON VERBIS
Internationally Famous Restaurant with a tradition of Elegance
HOTEL Shelburne
BOARDWALK, ATLANTIC CITY
NEW YORK PHONE (Local Call) RE 2-5450
or ATlantic City 344-8131



LADIES GIFT HANDKERCHIEF
Personalized with her full name (or 3 initials), this handkerchief is made of fine Irish linen with the added glamour of inch wide non-shrinking lace from France. Price $4.75 each. Box of three of the same name $13.50 ppd. Write for free Gift Brochure.

year will not be evident until the nineteen-nineties. Moreover, we have found that all varieties of asbestos fibres appear to be implicated in mesothelioma, and not just crocidolite, as some of us had hoped following Wagner's observations in South Africa. The problem would have been simple if only one variety of asbestos had been related to this tumor. Control procedures could have been narrowly focussed. As it is, however, a control program will have to be concerned with asbestos in general."

I asked Dr. Selikoff if he thought that asbestos air pollution was responsible for the occurrence of asbestos bodies in the lungs of people in the general population, and he replied that although there was relatively little direct confirmation, he felt it was at least partly responsible. Pointing out again that direct and indirect occupational exposure were much more common than had been suspected, especially in the construction and shipbuilding industries, he went on to say, "The possible sources of environmental contamination are numerous. Mining and milling operations, manufacturing processes, and the delivery and storage of finished products constitute just one part of it. The constant use of asbestos materials in the construction industry is another, and the destruction and deterioration of asbestos products in the community at large constitutes a third. There's no question in my mind that unsuspecting people are being exposed to small quantities of fibres from such sources as torn ironing-board covers and pot holders, and who knows how much asbestos is inhaled by hobbyists engaged in do-it-yourself projects around the house—like sawing up wallboard, mixing crack-filler compounds, insulating pipes with asbestos materials, and repairing furnaces and boilers with asbestos cements? I agree with those who suggest that small numbers of fibres may be liberated from many products in widespread use, such as asbestos-cement pipe and asbestos filters, which are employed in the food and pharmaceutical industries, but we have no information that this carries a disease potential. What troubles me is that relatively few people are aware of the fact that it can be dangerous to inhale asbestos. Not long ago, we discovered that children in a number of kindergartens in New York, California, and other states were being allowed to mix raw asbestos with water to make a cheap form of play dough. We have also learned that the powders in some children's cosmetics kits and in certain



Holidays are happy days in Germany

Germany has probably the most modern cities in Europe. But many old fashioned things like the well in the picture still persist. The Germans cling to the relics and reminders of the past—their visitors are charmed by them. The old and the new stand side by side... contrasting, yet blending. But more important is the spirit of the cities and of the towns, villages, hamlets and resorts. This spirit, too, is a wonderful combination of old traditions and strong, vital contemporary ideas and thinking. It's in the air, as are music and song and the gay laughter of your friendly hosts, the Germans.

Come to Germany where holidays are happy days.

Send coupon for illustrated brochures. See Your Travel Agent

GERMAN NATIONAL TOURIST OFFICE B

500 Fifth Avenue, New York 10036 323 Geary Street, San Francisco 94102
11 S. LaSalle Street, Chicago 60603 1176 Sherbrooke Street W., Montreal, Can.

Please send me free and postpaid illustrated travel brochures.

Name_____

Street Address_____

City & State_____ Zip Code _____



Anything can happen when you wear Fame



FAME de CORDAY

An imported French perfume. To 40.00

©1967 PARFUMS CORDAY, INC. AVAILABLE IN CANADA.

powder-water-mix jewelry kits contain asbestos. In addition, within the past couple of years two different thoracic surgeons have published articles in leading medical journals which advocate the sprinkling of asbestos on surgical incisions in order to promote the adhesion and healing of tissue. This is very bad advice. Why, Dr. William E. Smith, who conducts animal studies at Fairleigh Dickinson University, has readily induced mesotheliomas in hamsters by injections of asbestos fibres into their pleural cavities!"

Dr. Selikoff went on to say that he understands full well that asbestos is an extremely valuable material, and that it has become essential in modern industrial society. "We epidemiologists recognize and study its dangers so that we may devise ways of minimizing or avoiding them," he said. "Improvements in the practice of industrial hygiene are needed to eliminate unnecessary exposure among asbestos workers and men in other building trades, and awareness of the potential risks of environmental contamination is needed until our studies indicate just how far out into the general population the problem extends. All I can say is that the public should be made aware of the problem, and that no one, particularly children and young people, should be unduly exposed. After all, today's children are almost certainly inhaling more asbestos than their parents did, and we now know that if a person inhales significant amounts of asbestos dust, he carries a burden that will provide a latent potential for the development of cancer for the rest of his life. Here's an example I'll never forget. A couple of years ago, just after Cuyler Hammond, Jack Churg, and I published an article about asbestos and mesothelioma in the *New England Journal of Medicine*, I received a phone call from a doctor in Massachusetts who said that when he read the piece he immediately told his forty-three-year-old wife about it, because her sister had died of mesothelioma the year before. 'Imagine my horror,' this physician went on, 'when my wife proceeded to tell me that in 1943, while she and her sister were students in college, they had responded to a patriotic appeal and spent six weeks in a shipyard insulating pipes in destroyer escorts.'" Dr. Selikoff paused. "Two sisters with six weeks of exposure in 1943," he murmured. "Twenty-three years later, one of them dies of mesothelioma. And today, the other one—the doctor's wife—is being examined every few months to keep track of



Demi-lune handles

of gilded metal dress this handsome small bag by Block: black or brown leather or plastic patent, or in black alligator-grain leather. 30.00

At all Lord & Taylor stores

One of the 7 Wonderful Inns of the world!

Luxurious accommodations for 600
Year 'round resort activities
Write for Color Brochure & Weekend Plans

Cherry Hill Inn

Route 38 at Haddonfield Rd.
15 min. from downtown Phila.
N.J. Turnpike Exit 4 • Cherry Hill, N.J.
Reservations: (609) NO 2-7200

Café Carlyle



Currently Appearing

BOBBY SHORT

Complete Dinner $6.50 served to 9 p.m.
Entertainment from 9 p.m. to 2 a.m. Cover $3 p.p.
Minimum $3 p.p.

HOTEL CARLYLE
MADISON AVE. AT 76TH ST.
NEW YORK • TEL. RH 4-1000

some suspicious-looking shadows on her X-rays."

AS I was arriving at the Environmental Sciences Laboratory one day, I happened to notice a newspaper clipping tacked up on a bulletin board near the entrance. The story, which was from the *Times* of September 2, 1967, started out as follows: "The scene looked wintry on Madison Avenue, between 57th and 58th Streets, yesterday. Flurries of what seemed to be snowflakes fell on the area from between the 25th and 30th floors of the General Motors Building, which is nearing completion on the site of the former Savoy Plaza Hotel." The article went on to say that residents of the neighborhood had complained to the contractor, and that one of the project engineers had explained that the white flakes were fallout from fireproofing material that was being sprayed on the structural steel girders, beams, and decks of the fifty-story skyscraper. According to the *Times*, the engineer would not identify the insulating material, which was being applied with spray guns, "because I don't want to read newspaper stories about how we are messing up city streets." He was also quoted as saying, "This is a job condition all over the city." The last paragraph of the article read, "The city's Buildings Department said that it had not received any complaints about the Madison Avenue 'snowfall.' The department said the use of tarpaulins was required for spraying operations in construction work to prevent inconvenience to neighborhoods."

Beneath the article was this typed notation: "Examined in our laboratory, the material was found to contain amorphous glass fibres (presumably 'rock wool'), and chrysotile asbestos. It is known that other sprayed insulation materials contain amosite asbestos plus rock wool, while still others contain asbestos fibres alone."

When I asked Dr. Selikoff about the bulletin-board items, he said that the mineralogist in the laboratory, Dr. Arthur M. Langer, had collected a sample of the material and identified its contents by means of electron diffraction and electron microscopy, and with the aid of the electron microprobe. Then he went to a shelf and took down a small glass jar that contained a grayish wad of fibrous-looking stuff, about the size of a half dollar. "This is the sample Dr. Langer brought back and analyzed," he told me. "It comes from a type of insulation material that is about twenty-five per cent asbestos,



Hunge
Daily

CARMENCIT
FILIPINA. A
children. Fat
blind. Unable
suspect. Wash
only occasion
$12.82 per m
pig-sty is woo
by flattened
Hole in roof se
ture. Few kit
aginable priv
gry, ragged. Pa
cannot feed, c
dren adequat
cita means h

Thousands of ch
anxiously await "a
Choose a boy or g
Viet Nam, Hong
Colombia, Ecuad
grant helps provid
your Foster Child
In addition, PLAN
cal care when ca
clothing and ho
emphasis on educ
come self-supporting citizens. Since 1937, more than 110,000 c
PLAN's program. You receive a case history and photograph. Ea
a letter (original and translation). These letters will tell you
the entire family. Soon, through the regular letters and PLAN
child develop a warm, loving relationship.

CHECK YOUR CHARITY! We eagerly offer our financial sta
see that your contribution *truly benefits* the child for whom it

PLAN is a non-political, non-profit, non-sectarian, independe
under No. VFA019 with the Advisory Committee on Voluntary
International Development.

Foster Parents Pl


352 PARK AVENUE SOUTH, NEW YORK, N.Y.

PARTIAL LIST OF SPONSORS AND FOSTER PARENTS

Steve Allen
Julie Andrews
Sen. Paul H. Douglas
Helen Hayes
Conrad N. Hilton
Sen. Jacob K. Javits
Art Linkletter
Amb. & Mrs.
Henry Cabot Lodge
Garry Moore
Sen. William Proxmire
Dr. Howard A. Rusk
Mr. & Mrs.
Robert W. Sarnoff
Gov. & Mrs.
William W. Scranton
Sen. John G. Tower

FOSTER PARENTS PLAN, I
352 PARK AVENUE SOUTH, NEW Y
In Canada: P.O. Box 65, Sta. B, Mon

A. I wish to become a Foster Parent of
If possible, sex....., age.....
I will pay $16 a month for one year
Payment will be monthly ( ), quarte
annually ( ).
I enclose herewith my first payment
B. I cannot "adopt" a child but I wo
$.....
Name.....
Address.....
City.....
Date.....



and in this tiny wad alone there are literally billions of fibres. Of course, the 'snowfall' those people saw and complained about was not what may have been hazardous. What may have been hazardous was what they couldn't see—an untold number of sub-microscopic and highly respirable particles. The 'snowfall' was a perfect example of how asbestos gets into the ambient air, and that engineer was right when he said it's a job condition all over the city. You'll see the same material floating down from many other building projects in town. It's sprayed on by men in the Plasterers and Cement Masons Union. Incidentally, as far as I know, no one has ever made a study of *their* health."

It was a blustery day, filled with intimations of the season's changing, and when I left the hospital I took a Fifth Avenue bus downtown. I sat by a window, and was looking idly out at people strolling along the sidewalks when the bus came to a stop in the upper Forties beside a dump truck parked in front of a small office building that was being gutted. Several workmen were carrying canvas sacks of refuse across the sidewalk and heaving them up to a tall, well-built Negro who stood in the rear of the truck, emptying the sacks at his feet. As he did, clouds of dust rose into the air around him. Some of it showered the window beside me. It appeared to be ordinary cement dust, but then I saw that the pile of debris on which the man was standing contained broken pieces of acoustical panelling and wallboard, both of which often contain asbestos, and lengths of pipe coated with crumbling white insulation material that looked very much like asbestos.

When I got off the bus, I walked west along Forty-fourth Street. Halfway down the block, I saw two workmen slide a large rectangular container from a pickup truck and carry it into a building. The container was full of white powder, some of which was being blown away by the wind, and the men's clothes were covered with white dust. When I looked into the truck, I saw half a dozen large paper sacks labelled "Fireproof Plaster." One of the workmen returned for his tools, and I asked him if the plaster contained asbestos. "Sure," he replied. "That's what makes it fireproof."

I walked on to the corner of Sixth Avenue, where a forty-four-story office building had been under construction for several months. During that period, I had passed the building nearly every day, but now, for the first time, I



This charming fountain in lead will make a lovely addition to any terrace or informal garden setting. Complete with recirculating pump available in oxidized lead patina or Pompeian green finish. 37" high, 20" diameter shell. $375.00 F.O.B. New York City.

Send 25¢ in stamps or coins for 40 page Catalogue illustrating many other complete fountains and garden ornaments.

The FLORENTINE CRAFTSMEN Inc.
650-654 FIRST AVE. (37th St.) NEW YORK, N.Y. 10016
Phone LExington 2-3926



EMILY SHAW'S INN
POUND RIDGE, N. Y.
Finest type of American Food and Cocktails
Merritt Parkway, exit 35, north on 137, or Route 22 to Bedford Village.
John F. Shaw, Pres.
CLOSED MONDAYS • Pound Ridge 4-5779
VISIT OUR WALK IN WINE CELLAR

ABBE LANE
sets the world on desire
from October 2-October 22
Performances at 9:15 & 12:15
For reservations call
PLaza 9-3000
PERSIAN ROOM
AT THE PLAZA

looked up and scrutinized it carefully. About halfway up its side, I saw some tarpaulins billowing in the wind. Above the tarpaulins, the steel girders and beams were reddish brown; between the tarpaulins and the ground, however, they had been sprayed with asbestos insulation, which clung to them like gray frosting. Then, all at once, I saw wads of the same gray stuff falling out of the area that had been curtained off by the tarpaulins. Some of the pieces were carried away on the wind; others that were heavier came tumbling down around me. One fell on the sidewalk at my feet. I picked it up, crumbled it between my fingers, and saw dozens of tiny fibres float away. It was the same color and texture as the material that Dr. Selikoff had shown me half an hour before. The sidewalk was covered with it, and even as I stood there more of it came showering down. A woman hurrying to make the light with a friend brushed some off the sleeve of her coat, and said, "What the devil is this stuff?" I watched her cross the avenue, and saw asbestos insulation blowing over the pavement like thistledown across a field. The wind was southerly, and most of the asbestos was travelling north, in the direction of Mount Sinai Hospital, where Dr. Selikoff and his associates are still following the trail Dr. Cooke set out on in 1924. But I was not thinking about Dr. Selikoff or the difficulties of epidemiology just then. I was thinking about the children who play in Central Park every day, and about all the other children who live in this city and breathe its air. —PAUL BRODEUR

•

CRIES WE DOUBT EVER GOT CRIED OUT, RAPTURE OR NO RAPTURE

[*From the St. Moritz Courier*]

When GIOVANNI SEGANTINI first came up to the Engadine from Lake Como, he cried out, full of rapture:


"I bow before this earth which is hallowed by beauty; I kiss the grasses and the flowers under this blue vault of heaven, and while the birds sing and meet together for flight and the bees suck honey from the open blooms, I drink at these pure springs, where beauty is ever renewed, where love is renewed that gives life to all things."

•

Western observers believed Brezhnev's remarks would aid anti-American propaganda abroad while leaving the Kremlin free to deal with almost any future American President, except possibly Richard Nivox.—*Washington Post.*



The compact carry-all. Ready to work with a r
checkbook holder, 5" frame coin pocket, and
pocket. A 9-section vinyl card holder takes ca
else. A swarm of honey kid colors, about $12.
ST. THOMAS At Himelhoch Bros. & Co., Detroit; Frost Bros.,
B. Altman & Co., New York City and all branch
St. Thomas Inc., St. Thomas Place, Gloversville, N.Y. and all other fine stores.



Two confident candidates from our exciting collection of original pin designs.
Donkey, about $17.50. At fine stores. Ciner Jewelers, 20 West 37th St., New York