

(December) 1977

Modern Concepts in Brain Tumor Therapy:
Laboratory and Clinical Investigations

DHEW Publication No. (NIH) 77-1236
U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE
Public Health Service
National Institutes of Health
NATIONAL CANCER INSTITUTE, BETHESDA, MARYLAND 20014

OLI 6784

Chemical- and Virus-Induced Brain Tumors¹

James A. Swenberg^{2,3}

ABSTRACT—Experimental animal models resembling most human brain tumor types can be induced by exposure to oncogenic viruses or chemical carcinogens: Astrocytomas and glioblastoma multiforme can be produced experimentally by intracerebral injection of oncornaviruses, whereas medulloblastomas, choroid plexus papillomas, and ependymomas can be induced by the papovaviruses. Adenoviruses have been utilized to cause medullopitheliomas, neuroblastomas, and retinoblastomas. All three groups of viruses can result in sarcoma production. Gliomas represent the primary tumor type induced in the brain by chemical carcinogens. These autochthonous tumor systems are reviewed, with emphasis on methods, tumor type, latency period, advantages, and disadvantages. In addition, recent investigations of molecular events involved in neoplastic transformation by chemical carcinogens are summarized.—*Natl Cancer Inst Monogr* 48: 3-10, 1977.

A variety of experimental brain tumor models resembling most human tumor types have been developed during the past decade. Brain tumors can be induced with intracerebral injection of viruses or chemicals, or by transplacental, parenteral, oral, or even topical exposure to chemical carcinogens. Depending on the model, such tumors arise weeks to years after exposure. The

availability of suitable animal models has stimulated increased research on brain tumor kinetics, immunology, chemotherapy, and histogenesis. Recently, these models have been used in the study of molecular events responsible for neoplastic transformation. This review is intended to summarize the experimental brain tumor systems available today and to highlight some of the exciting research being done with these systems.

VIRUS-INDUCED BRAIN TUMORS

Work on experimental brain tumor induction with viruses was reviewed comprehensively in

¹ Presented at the Symposium on Modern Concepts in Brain Tumor Therapy: Laboratory and Clinical Investigations, held in Atlanta, Ga., February 26-28, 1976, and sponsored by the Clinical Investigation Branch and the Cancer Clinical Investigations Review Committee, Division of Cancer Treatment, National Cancer Institute, Bethesda, Md. 20014.

² Pathology and Toxicology Research Unit, The Upjohn Company, Kalamazoo, Mich. 49001.

³ Dr. Darell Bigner provided many helpful suggestions during the preparation of this manuscript.

TABLE 1.—Specificity of experimental brain tumor induction with oncogenic viruses

Experimental brain tumor type	Inducing viruses	
	Group	Type ^a
Anaplastic astrocytoma or glioblastoma multiforme	RNA-Oncornaviruses	ASV, MuSV, SSV
Medulloblastoma	DNA-Papova	Human papova JC
Neuroblastomas or retinoblastomas	DNA-Adenovirus	Human adenovirus type 12
Ependymomas or choroid plexus papillomas	DNA-Papova	SSV40, human papova PML-1, JC, and BK
Sarcomas or meningeal tumors	DNA-Adenovirus	SA7 and CELO
	RNA-Oncornavirus	ASV and MuSV
	DNA-Papova	Bovine papilloma, murine polyoma, human papova JC
	DNA-Adenovirus	SA7

^a ASV = avian sarcoma virus; MuSV = murine sarcoma virus; SSV = simian sarcoma virus; BK = human papovavirus BK; SA7 = simian adenovirus 7.

several publications (1-6). Intracranial tumors have been induced by representatives of most oncogenic tumor virus groups, including the adenoviruses, papovaviruses, and oncornaviruses. To date, techniques for induction and morphologic descriptions were emphasized in most studies on these models. Future investigations will probably de-emphasize morphologic classification and instead turn toward utilizing such models in basic and applied research. Morphologic studies have led to several generalizations that are summarized in table 1. For example, only oncornaviruses induce astrocytomas and glioblastomas, whereas the human papovavirus JC is the only virus known to cause the development of medulloblastomas. Choroid plexus papillomas are induced by simian virus 40 (SV40), the SV-PML papovaviruses isolated from patients with progressive multifocal leukoencephalopathy (PML) and the avian adenovirus (CELO), whereas medullopitheliomas and retinoblastomas are only elicited with human adenoviruses. In contradistinction, sarcomas were induced by members of most tumor virus groups.

RNA Tumor Viruses

Avian Sarcoma Virus

Mammalian neuro-oncogenicity of ASV was first described by Rabotti and Raine (7). Subsequently, gliomas and intracranial sarcomas were induced in cats, dogs, guinea pigs, mice, rabbits, rats, and subhuman primates. The relative proportion of glial and mesenchymal elements in these brain tumors varies considerably among species. Homogeneous glial tumors have clearly been produced in the rat and dog with ASV. Bigner et al. (8) induced gliomas in 27 of 30 dogs, 19 of which had gliomas only, with no evidence of sarcoma formation. They later demonstrated that the site of ASV inoculation influenced tumor type; i.e., injection of ASV deep in periventricular zones resulted exclusively in astrocytomas, whereas only sarcomas were induced if the virus was injected superficially over the cerebellar vermis (9). Well-differentiated astrocytomas also developed in F344 (10) and Sprague-Dawley (11) rats after neonatal intracerebral injections of ASV. In addition, Copeland et al. (12) demonstrated that rats remained susceptible to the neuro-oncogenic effects of ASV at 100 days of age. Although the incidence of brain tumors in rats inoculated with ASV as adults was only 50%, nearly all were astrocytic.

In contrast to the rat and dog, a large percentage of the hamster, guinea pig, mouse, cat, and subhuman primate tumors induced with intracerebral ASV have a mesenchymal nature. Whereas neoplastic astrocytic elements can be observed, prominent reticulin fibers are also present. Tumors composed of large balloon-like cells (13) and "giant cell glioblastomas" (14, 15) usually contain such reticulin. This cell type probably represents a histiocytic element in a mesenchymal or mixed glial-mesenchymal tumor.

Several other features of ASV-induced brain tumors deserve comment: 1) There is generally no replication of infectious or noninfectious virus in mammalian tumor systems. However, virus structural antigens and virus-related transplantation antigens are usually produced in transformed mammalian cells. One can "rescue" ASV from such cells by growing the "nonpermissive" transformed cells with normal avian cells and allowing cell fusion to occur. 2) The dose of virus required to elicit intracranial tumors is lower than that for extracranial tumor induction. This is probably due to a greater immunologic response against the latter. 3) Improvements in the techniques for growing and concentrating ASV have allowed

large pools of standardized virus to be produced (16). Using this virus, researchers have obtained reproducible survival curves for rats bearing a 100% incidence of astrocytomas. This system provides the first primary glioma model that is well suited for chemotherapy trials (17). Preliminary results of these trials have been reported (5, 16-18).

Other Oncomaviruses

MuSV has been reported to cause brain tumors in mice and rats following intracerebral inoculation (5). Gliomas, meningiomas, and hemangioendotheliomas represent the most common tumor types induced with MuSV. In contrast to ASV-induced mammalian tumors, replicating virions were easily detectable in these mouse tumors by electron microscopy (19). Recently, a 100% incidence of extremely vascular brain tumors containing neoplastic astrocytes and extensive endothelial proliferation was induced in Wistar-Furth rats with the Kirsten MuSV (20). No C-type virions were detected by electron microscopy.

Intracerebral inoculation of neonatal marmosets with SSV induced solitary tumors in 6 of 10 animals (21). The tumors were composed of pleomorphic neuroglial cells, areas of necrosis, hemorrhage, palisading, and prominent endothelial proliferation. These SSV-induced marmoset tumors closely resemble the human glioblastoma multiforme. Infectious virus was isolated from brain and cerebral spinal fluid of tumor-bearing animals.

DNA Tumor Viruses

Papovaviruses

Several members of the papovavirus group cause tumors when inoculated intracerebrally into animals (5). Bovine papilloma virus induced meningiomas, fibromas, and fibrosarcomas in hamsters and calves following latency periods ranging from 20 days to nearly a year. No virus particles were detected in the tumors by electron microscopy. Fibrosarcomas were also detected in the leptomeninges after polyoma virus was injected intracerebrally into newborn rats, hamsters, and rabbits. Intracerebral inoculations of SV40 into newborn hamsters induced fibrosarcomas in the leptomeninges and choroid plexus papillomas in the ventricles.

Of greatest interest, however, are the tumors that developed in hamsters after intracerebral

injections of JC virus, a human papovavirus isolated from a patient with PML (22-24). Brain tumors developed in 50 of 63 hamsters inoculated with JC virus, with medulloblastomas representing the most common tumor type. These appear to arise from the internal granule cell layer of the cerebellum and frequently are multifocal. Several other tumor types have been induced with JC virus, including primitive gliomas, papillary ependymomas, meningiomas, and pineocytomas. Virus was recovered from 5 of 7 tumors tested. The JC virus is morphologically indistinguishable from SV40 and SV40-PML strain 1, but differs from them in its cell culture host range and antigenic properties (5, 25).

Adenoviruses

The experimental induction of neurogenic tumors in rats, hamsters, and mice after the animals received intracerebral, intraocular, or ip inoculations of human adenovirus 12 provides an excellent model for future research on neuroblastomas, medulloepitheliomas, and retinoblastomas (26-31). Medulloepitheliomas were induced in 88% of the rats given intracerebral injections of human adenovirus 12. These appeared to develop from the subependymal plate and were characterized by the formation of rosettes, high levels of cholinesterase following in vitro cultivation, and the presence of cilia containing 9+0 tubules (a hallmark of normal sensory neuronal cells). Similar 9+0 tubules were demonstrated ultrastructurally in the cilia of retinoblastomas induced in rats and hamsters following neonatal intraocular injections of human adenovirus 12.

Other members of the adenovirus group that elicited intracranial tumors (5) include the simian and avian adenoviruses (SA7, SV20, and CELO). SA7 induced poorly differentiated tumors of the choroid plexus, trigeminal nerve, and dura mater. Injection of SV20 into hamsters caused extremely undifferentiated tumors, which had the same morphology whether induced sc or intracerebrally. Intracerebral tumors produced with CELO had a morphology distinctly different from neoplasms induced with either the human or the simian adenoviruses; they were composed of papillary arrangements of cuboidal and columnar cells that filled the ventricles. Although originally thought to be ependymal tumors, the CELO tumors probably represent choroid plexus papillomas.

MODERN CONCEPTS IN BRAIN TUMOR THERAPY

CHEMICALLY INDUCED TUMORS OF THE CENTRAL NERVOUS SYSTEM (CNS)

The susceptibility of neuroectodermal tissue to the oncogenic effects of chemical carcinogens was first demonstrated in 1939 (32). Since then, brain tumors have been induced by local implantation of several polycyclic hydrocarbons (PCH). More recently, additional classes of carcinogens were shown to be oncogenic for the nervous system following systemic exposure. These ranged from agents such as 2-acetylaminofluorene, which caused a few tumors of the nervous system along with many tumors elsewhere, to the nitrosoureas that caused neuroectodermal tumors in 100% of the animals. Several reviews (33-36) and a monograph (5) have been published on the chemical induction of brain tumors. In addition to reports by some scientists on the induction and characterization of various brain tumor models, other investigators used animal models for neuro-oncogenesis to study the molecular basis for neoplastic transformation. Whether similar mechanisms are involved in human brain tumors is unknown. It has been estimated that 80-90% of all human cancer is caused by chemicals; however, no chemical carcinogen has been clearly linked to human brain tumors.

In the section to follow, various brain tumor models have been arranged by the class of carcinogen used to induce them.

Polycyclic Hydrocarbons

Over 200 years ago, Sir Percival Pott first theorized the potential carcinogenic hazard of PCH. Therefore, it was fitting that the first experimental tumors of the nervous system were also induced with PCH (32). Since that time, brain tumors have been produced with several PCH, including 3-methylcholanthrene, benzo[a]pyrene, dibenzanthracene, dimethylbenz[a]anthracene (DMBA), and trimethyl benzanthracene. Such tumors have been induced in frogs, toads, mice, rats, hamsters, and dogs. However, birds, guinea pigs, rabbits, cats, and monkeys were resistant to the neuro-oncogenic effect of PCH. In general, brain tumors occur only when the PCH are in direct contact with the CNS. To accomplish this, the carcinogens are directly implanted in the brain. The incidence and type of tumor produced by PCH are greatly influenced by the positioning of the pellet (37). Few gliomas but many sarcomas and meningiomas result from superficial placement of the pellet near the dura. Ependymomas were primarily induced when the

pellet was placed in the ventricles, whereas oligodendrogliomas were associated with implantation in the frontal white matter. Astrocytomas arose near pellets placed in the subcortical regions of the parietal lobe, and medulloblastomas formed after pellet implantation into the cerebellum.

Although several investigators have successfully induced brain tumors in rats with PCH, mice appear to be the most susceptible animal species (34). Tumor incidence varies from less than 10 to nearly 100%, with an average of 40-60% of the animals developing tumors. Most animals die between 200 and 300 days; however, latency periods have varied from as short as 68 to as long as 750 days. The primary advantage of this model is that the site of tumor formation can be localized. Disadvantages of the system include the variable incidence and latency period of the experimental tumors. The model is poorly suited for quantitative dose-response studies and investigations of early biochemical processes, since the time of exposure can vary from minutes to months because of the extremely long half-life of the pellet. The importance of unique doughnut-shaped virions detected in macrophages surrounding pellets of carcinogens prior to tumor formation requires further elucidation (38). Whether these virions represent activated viruses that play an important role in tumor induction remains unknown. However, they have never been detected in experimental gliomas and have induced fibrosarcomas rather than gliomas when they were injected intracerebrally into mice.

When pregnant rats were exposed to DMBA iv on day 21 (39) or orally on days 14, 17, and 20 of gestation (40), the most frequently detected neoplasms induced in the offspring were tumors of the nervous system. Transplacental exposure to additional PCH will be necessary to determine whether this is a general phenomenon or a specific effect of DMBA. In any event, it should now be possible to design pulse exposure experiments and to demonstrate dose-response relationships for DMBA.

Nitroso Compounds

A landmark in experimental neuro-oncology occurred in 1964 when Druckrey and co-workers (41) selectively induced tumors of the rat nervous system after iv injections of methylnitrosourea (MNU). This provided the first reproducible brain tumor model that did not traumatize the brain. Since then, several additional carcinogens have been discovered that primarily induce neuroectodermal tumors following systemic administration.

These animal models represent some of the most powerful tools for modern research in chemical neuro-oncogenesis.

Methylnitrosourea

This acyl-alkyl-nitrosamide is among the most potent of all the chemical carcinogens known today. When administered iv at repeated low doses, the nervous system clearly represents the target organ. The highest incidence of neurogenic tumors is obtained with repeated iv injections of MNU at a dose schedule of 5 mg/kg/week for 32-36 weeks (42). This dosage has consistently produced a 90-100% incidence of grossly detectable brain tumors in Sprague-Dawley rats. Well-differentiated neurinomas are also induced; however, the incidence of these is much lower. This low incidence of tumors of the peripheral nervous system (PNS) can be reversed if F344 rats are used instead of Sprague-Dawley (43). The rat and the rabbit appear most susceptible to brain tumor induction with MNU (5). Dogs have been more variable; some brain and peripheral nervous system tumors have been induced in various breeds. Mice were quite resistant to neuro-oncogenesis with MNU. Denlinger et al. (44) induced 2 gliomas in 19 C3H mice exposed iv to 25 mg/kg every 4 weeks (total dose, 175 mg/kg). In spite of observation periods exceeding 6 years, attempts to induce brain tumors in rhesus monkeys have been unsuccessful (5). When cats were exposed to iv or oral doses of 25 or 20 mg/kg, respectively, all animals died within 10 days from severe bone marrow toxicity and septicemia (35). Additional resistant species include guinea pigs, sheep, and swine.

Anaplastic gliomas, mixed gliomas, and oligodendrogliomas represent the most common brain tumors induced with MNU. These are frequently located near periventricular regions, in subcortical white matter, and in the hippocampus. If the dose of MNU is raised from 5 mg/kg/week to 10 or 20 mg/kg twice a week, a shift from glial to mesenchymal cell types becomes apparent, with most tumors being sarcomas or gliosarcomas (45). MNU is also a potent carcinogen when administered locally, as shown clearly in experiments utilizing the sc, ip, or oral routes. Tumors develop at the site of administration and to a lesser extent in the nervous system. The lower incidence of tumors detected in the nervous system can best be explained by the decreased levels of circulating MNU. In addition to tumors at the site of injection and in the nervous system, a high incidence

of thymic lymphomas can be induced after oral or sc administration of 20 mg/kg twice a week for 9 weeks. These types of experiments have clearly shown that MNU is not "neurospecific." Rather, MNU is a potent carcinogen for many tissues. Potential mechanisms involved in determining which target site is affected by this and related carcinogens will be discussed later.

Ethylnitrosourea (ENU)

The exquisite sensitivity of the fetal rat's nervous system to the oncogenic effects of ENU was first demonstrated in the experiments of Ivan-kovic and Druckrey (46). A 100% incidence of neurogenic tumors developed in offspring exposed to a single dose of 20 mg/kg or more during the last trimester of pregnancy. Clear dose-response relationships exist with single doses of 1, 5, 20, or 50 mg/kg inducing a 12, 79, 100, and 100% incidence of neurogenic tumors, respectively (47). While the tumor incidence increased, the latency period for tumor induction decreased from a mean of 655 days for 1 mg/kg to 211 days for 50 mg/kg. Age of the individual greatly influences the susceptibility to the neuro-oncogenic effects of ENU. The rat fetus is resistant to the carcinogenic effects of ENU before the 12th day of gestation; however, it is susceptible to teratogenic effects before this stage of development. Susceptibility to neuro-oncogenesis increases from the 12th day of gestation to birth, after which it begins decreasing (33). By 30 days of age, the rat has a susceptibility comparable to the adult; i.e., the nervous system no longer represents the target organ and much higher doses of ENU are required to induce a similar incidence of neoplasia.

ENU has usually been administered iv to pregnant rats, although it can be given orally, ip, or sc. Neurogenic tumors were even reported in 38% of the offspring whose mothers had 50 mg/kg of ENU applied topically to the skin (48). It has also been possible to induce tumors by administration of precursors of ENU in the food and water of pregnant rats. Under the mildly acidic conditions of the rat's stomach, ethylurea and sodium nitrite undergo nitrosation and form ENU *in vivo* (49).

Rats are the species most sensitive to the neuro-oncogenic effects of ENU. Hamsters, mice, and opossums have also developed neurogenic tumors. In hamsters, tumors have been confined to the PNS, whereas in mice a low incidence of CNS and PNS tumors develops. Mixed gliomas and oligodendrogliomas represent the most common

brain tumors induced with ENU, whereas anaplastic neurinomas are the type occurring most frequently in the PNS. Anaplastic gliomas and sarcomas are rare. Animals bearing anaplastic neurinomas die significantly earlier than those bearing gliomas. When the sequential development of these tumors was investigated, evidence of neoplastic proliferation was detected in trigeminal nerves as early as 3 weeks after exposure to ENU (50). Comparable changes were not evident in the brain until 4 months after ENU exposure. Mechanisms involved in this differential triggering of neoplastic proliferation between glial and Schwann cells remain unknown.

Other Nitroso Compounds

Additional nitroso compounds possessing neuro-oncogenic properties have been described since the original discovery of MNU (5, 33), including dimethylnitrosourea, trimethylnitrosourea, propylnitrosourea, butylnitrosourea, dinitrosopiperazine, nitrosopiperidine, nitrosomorpholine, ethylnitrosobiuret, methylnitrosobiuret, ethylnitrosourethane, and methylnitrosourethane. In general, these offer no major advantages over MNU and ENU and will not be discussed in detail.

Other Carcinogens

Triazenes

Several of the dialkyl-aryl-triazenes have been shown to be carcinogenic for the nervous system as well as for other tissues (51). Neuro-oncogenesis has been successful both in adult rats and transplacentally. As noted for MNU and ENU, the methyl compounds were most effective in adults, whereas the corresponding ethyl compounds were highly oncogenic to the fetal rat's nervous system.

1,2-Diethylhydrazine, Azoxethane, and Azoxyethane

Transplacental neuro-oncogenesis has been induced in the rat with these three carcinogens (33). In contrast to the triazenes, adult rats developed multiple intestinal tumors instead of neurogenic neoplasms when the methyl counterpart was administered.

INVESTIGATIONS ON THE MECHANISMS OF CHEMICAL NEURO-ONCOGENESIS

Scientists conducting studies at several laboratories have suggested a possible mechanism for

the neurospecificity of some chemical carcinogens. That chemical carcinogens are electrophilic reagents has been known for several years (52). Those carcinogens that are not electrophilic must be metabolized to form electrophiles, which interact with macromolecules such as DNA, RNA, and protein. Neoplastic transformation represents a heritable change at the cellular level, with the most likely site for this change being the DNA of the originally transformed cell (53).

Investigations of several carcinogens and tissues demonstrated alkylation of DNA bases. The greatest amount of this alkylation occurred at the N-7 position on guanine, with lesser amounts being detected at the N-3 position of adenine and the O⁶-position of guanine. When the extent of N-7 alkylguanine formation was determined in brains and livers of adult and 10-day-old rats given injections of ENU, no correlation was found between DNA alkylation and carcinogenicity (54). Subsequent studies determined the degree of purine alkylation at various times after ENU exposure. Goth and Rajewsky (55, 56) demonstrated surprising differences between the repair rates of O⁶-ethylguanine and those of N-7 ethylguanine or N-3-ethyladenine. O⁶-Ethylguanine persisted for long periods in the target organ (brain) but was repaired much more rapidly in the liver. N-7-Ethylguanine was removed rapidly in both tissues. If one determines the brain:liver ratio for each of the major sites of purine alkylation, a high degree of correlation is evident between alkylation at the O⁶-position of guanine and carcinogenicity.

Further support for this hypothesis has been reported by Kleihues and co-workers (57-59), who demonstrated persistence of O⁶-methylguanine in brains of rats treated with MNU and methylmethanesulfonate (MMS). Both methylating agents caused brain tumors in rats; however, MMS has a much weaker neuro-oncogenic effect. When equimolar doses of MNU and MMS were administered to rats, only 0.05 as much O⁶-methylguanine was present after exposure to MMS. Thus the extent of initial O⁶-alkylguanine formation and its persistence over time correlates with carcinogenicity. A possible mechanism for this correlation was suggested by Loveless (60). Alkylation at the O⁶-position of guanine forces the base into the enol form, which causes anomalous base pairing during DNA replication. Instead of pairing with cytosine, O⁶-alkylguanine pairs with thymine. Fixation of anomalous base pairing requires that at least one cell replication take place before repair of the damaged DNA. The

lack of this replication in end-stage neurons may explain why no neuronal tumors have been induced with MNU, even though DNA alkylation occurs to a greater extent in neurons than in glia (61). It also helps to explain the predilection sites for ENU- and MNU-induced gliomas, since these areas contain the glia with the greatest propensity for cell replication. Enzymic repair systems for O⁶-alkylguanine have recently been demonstrated (62, 63). If these systems are deficient in rat brain, the exquisite susceptibility of the fetal rat brain to carcinogens may thus be explained. Over half the 35 or more compounds that caused transplacental carcinogenesis induced tumors of the rat nervous system.

In summary, pathogenetic mechanisms involved in experimental neuro-oncogenesis seem to be as well or better known than those of other neoplasms. Delayed repair of O⁶-alkylguanine appears to be a major factor in sensitizing the rat nervous system to carcinogenesis. Whether this mechanism is responsible for human brain tumors remains to be investigated. Sensitivity to neuro-oncogenesis may be a general phenomenon for many carcinogens if effective tissue concentrations reach the nervous system. At present, the somatic mutation theory of cancer induction best explains chemical neuro-oncogenesis.

REFERENCES

- (1) JOHN DS: Oncogenic viruses: Expectations and applications in neuropathology. *Prog Exp Tumor Res* 17:74-92, 1972
- (2) IKUTA F, KUMANISHI T: Experimental virus-induced brain tumors. *Prog Neuropathol* 253-334, 1973
- (3) JOHNSON RT, NARAYAN O: Experimental neurological diseases of animals caused by viruses. In *Models of Human Neurological Diseases* (Klawans HL, ed). Amsterdam: Excerpta Medica, 1974, pp 43-82
- (4) BIGNER DD, PEGRAM C: A review of virus-induced experimental brain tumors and of the putative associations of viruses with human brain tumors. *Adv Neurol* 13:57-83, 1976
- (5) JANISCH W, SCHREIBER D: Experimental Tumors of the Central Nervous System. [Bigner D, Swenberg J, eds (first English edition)]. Kalamazoo: The Upjohn Co., 1977
- (6) RABOTTI GF: Experimental intracranial tumors of viral etiology. In *The Experimental Biology of Brain Tumors* (Kirsch WM, Grossi-Paoletti E, Paoletti P, eds). Springfield, Ill.: Charles C Thomas, 1972, pp 148-180
- (7) RABOTTI GF, RAINE WA: Brain tumors induced in hamsters inoculated intracerebrally at birth with Rous sarcoma virus. *Nature* 204:898-899, 1964
- (8) BIGNER DD, ODOM GL, MAHALEY MS JR, et al: Brain tumors induced in dogs by the Schmidt-Ruppin strain of Rous sarcoma virus: Neuropathological and immunological observations. *J Neuropathol* 28:648-680, 1969

- (9) BIGNER DD, KVEDAR JP, SHAFFER TC, et al: Factors influencing the cell type of brain tumors induced in dogs by Schmidt-Ruppin-Rous sarcoma virus. *J Neuropathol Exp Neurol* 31:583-595, 1972
- (10) WILFONG RF, BIGNER DD, SELF DJ, et al: Brain tumor types induced by the Schmidt-Ruppin strain of Rous sarcoma virus in inbred Fischer rats. *Acta Neuropathol (Berlin)* 25:196-206, 1973
- (11) SWENBERG JA, BIGNER DD, HALL TL: Experimental chemotherapy of viral-induced brain tumors. *Proc Am Assoc Cancer Res* 15:43, 1974
- (12) COPELAND DD, VOGEL FS, BIGNER DD: The induction of intracranial neoplasms by the inoculation of avian sarcoma virus in perinatal and adult rats. *J Neuropathol Exp Neurol* 34:340-358, 1975
- (13) BURGER PC, BIGNER DD, SELF DJ: Morphologic observations of brain tumors in PD 4 hamsters induced by four strains of avian sarcoma virus. *Acta Neuropathol (Berlin)* 26:1-21, 1973
- (14) KUMANISHI T: Brain tumors induced with Rous sarcoma virus, Schmidt-Ruppin strain. I. Induction of brain tumors in adult mice with Rous chicken sarcoma cells. *Jpn J Exp Med* 37:461-474, 1967
- (15) KUMANISHI T, IKUTA F, NISHIDA K, et al: Brain tumors induced in adult monkeys by Schmidt-Ruppin strain of Rous sarcoma virus. *Gann* 64:641-643, 1973
- (16) BIGNER DD, SELF DJ, FREY J, et al: Refinement of the avian oncornavirus-induced primary rat brain tumor model and its potential for therapeutic screening. *Recent Results Cancer Res* 51:20-34, 1975
- (17) CRAFTS D, WILSON CB: Animal models of brain tumors. *Natl Cancer Inst Monogr* 46:11-17, 1977
- (18) SWENBERG JA, BIGNER DD, HALL TL: Experimental chemotherapy of viral-induced brain tumors. *Proc Am Assoc Cancer Res* 15:43, 1974
- (19) DUFFY PE: Virus-induced cerebral sarcoma. *J Neuropathol Exp Neurol* 29:370-391, 1970
- (20) YUNG WK, BLANK NK, VICK NA: Glioblastoma. Induction of a reproducible autochthonous tumor in rats with murine sarcoma virus. *Neurology* 26:76-83, 1976
- (21) JOHNSON L, WOLFE LG, WHISLER W, et al: Induction of gliomas in marmosets with simian sarcoma virus, type 1 (SSV-1). *Proc Am Assoc Cancer Res* 16:119, 1975
- (22) PADGETT BL, WALKER DL, ZU RHEIN GM, et al: Cultivation of papova-like virus from human brain with progressive multifocal leukoencephalopathy. *Lancet* 1:1257-1260, 1971
- (23) WALKER DL, PADGETT BL, ZU RHEIN GM, et al: Human papovavirus (JC) induction of brain tumors in hamsters. *Science* 181:674-676, 1973
- (24) ZU RHEIN GM, VARAKIS J: Morphology of brain tumors induced in Syrian hamsters after inoculation with JC virus, a new human papovavirus. In *Proceedings VII International Congress on Neuropathology* (Környey S, Tariska S, Gosztönyi G, eds), vol 1. Amsterdam: Excerpta Medica; Budapest: Akadémiai Kiado, 1975, pp 479-481
- (25) BECKER LE, NARAYAN O, JOHNSON RT: Demonstration of papovavirus tumor antigen in brain tumors. *J Neuropathol Exp Neurol* 34:95, 1975
- (26) OGAWA K, HAMAYA K, FUJII Y, et al: Tumor induction by adenovirus type 12 and its target cells in the central nervous system. *Gann* 60:383-392, 1969
- (27) MUKAI N, KOBAYASHI S: Human adenovirus-induced medulloepitheliomatous neoplasms in Sprague-Dawley rats. *Am J Pathol* 73:671-690, 1973
- (28) ———: Primary brain and spinal cord tumors induced by human adenovirus Type 12 in hamsters. *J Neuropathol Exp Neurol* 32:523-541, 1973
- (29) MUKAI N, KOBAYASHI S, OGURI M: Ultrastructural studies of human adenovirus-produced retinoblastoma-like neoplasms in Sprague-Dawley rats. *Invest Ophthalmol* 13:593-601, 1974
- (30) MUKAI N, MURAO T: Retinal tumor induction by ocular inoculation of human adenovirus in 3-day-old rats. *J Neuropathol Exp Neurol* 34:28-37, 1975
- (31) KOBAYASHI S, MUKAI N: Retinoblastoma-like tumors induced by human adenovirus type 12 in rats. *Cancer Res* 34:1646-1651, 1974
- (32) SELIGMAN AM, SHEAR MJ: Studies in carcinogenesis. VIII. Experimental production of brain tumors in mice with methylcholanthrene. *Am J Cancer* 37:364-395, 1939
- (33) DRUCKREY H, IVANKOVIC S, PREUSSMANN R, et al: Selective induction of malignant tumors of the nervous system by resorptive carcinogens. In *The Experimental Biology of Brain Tumors* (Kirsch WM, Grossi-Paoletti E, Paoletti P, eds). Springfield, Ill.: Charles C Thomas, 1972, pp 85-147
- (34) KIRSCH WM, SCHULZ D: The chemical induction of brain tumors with carcinogenic hydrocarbons and miscellaneous agents. In *The Experimental Biology of Brain Tumors* (Kirsch WM, Grossi-Paoletti E, Paoletti P, eds). Springfield, Ill.: Charles C Thomas, 1972, pp 181-193
- (35) SWENBERG JA: Chemical induction of brain tumors. In *Neoplasia in the Central Nervous System* (Thompson RA, Green JR, eds). New York: Raven Press, 1976, pp 85-99
- (36) KLEIHUES P, IANTOS PL, MAGEE PN: Chemical carcinogenesis in the nervous system. *Int Rev Exp Pathol* 15:153-232, 1976
- (37) ZIMMERMAN HM: The nature of experimental gliomas. *Clin Neurosurg* 7:247-260, 1959
- (38) IKUTA F, ZIMMERMAN HM: Virus particles in reactive cells induced by intracerebral implantation of dibenzanthracene. *J Neuropathol Exp Neurol* 24:225-243, 1965
- (39) ALEXANDROV VA, NAPALOV NP: Carcinogenic effect of DMBA in rats exposed transplacentally. *Vopr Onkol* 20:50-54, 1974 (Russ)
- (40) JOSHI SR, SHENEFELT RE, RICE JN: Transplacental carcinogenicity of DMBA in the rat. *Proc Am Assoc Cancer Res* 15:27, 1974
- (41) DRUCKREY H, IVANKOVIC S, PREUSSMANN R: Selective induction of brain tumors in rats with MNU. *Naturwissenschaften* 51:144, 1964 (Ger)
- (42) SWENBERG JA, KOESTNER A, WECHSLER W: The induction of tumors of the nervous system with intravenous methylnitrosourea. *Lab Invest* 26:74-85, 1972
- (43) DENLINGER RH, KOESTNER A, SWENBERG JA: An experimental model for selective production of neoplasms of the PNS. *Acta Neuropathol (Berlin)* 23:219-228, 1973
- (44) DENLINGER RH, KOESTNER A, WECHSLER W: Induction of neurogenic tumors in C3HeB/FeJ mice by nitrosourea derivatives: Observations by light microscopy, tissue culture, and electron microscopy. *Int J Cancer* 13:559-571, 1974
- (45) SWENBERG JA, KOESTNER A, WECHSLER W, et al: Differential oncogenic effects of methylnitrosourea. *J Natl Cancer Inst* 54:89-96, 1975

- (46) IVANKOVIC S, DRUCKREY H: Transplacentare Erzeugung maligner Tumoren des Nervensystems. I. Äthylnitrosoharnstoff (ANH) an BD XI-Ratten. *Z. Krebsforsch* 71:320-360, 1968
- (47) SWENBERG JA, KOESTNER A, WECHSLER W, et al: Quantitative aspects of transplacental tumor induction with ethylnitrosourea in rats. *Cancer Res* 32:2656-2660, 1972
- (48) GRAW J, ZELLER WJ, IVANKOVIC S: Entstehung von neurogenen Malignomen bei den Nachkommen von Sprague-Dawley Ratten nach perkutaner Applikation von Äthylnitrosoharnstoff (ANH) während der Tragezeit. *Z. Krebsforsch* 81:169-172, 1974
- (49) KOESTNER A, WECHSLER W, DENLINGER RH: Induction of nervous system and lymphoid neoplasms by feeding threshold levels of methylnitrosourea and ethylnitrosourea precursors to adult rats. *Food Cosmet Toxicol* 13:605-609, 1975
- (50) SWENBERG JA, CLENDENON N, DENLINGER R, et al: Sequential development of ethylnitrosourea-induced neurinomas: Morphology, biochemistry, and transplantability. *J Natl Cancer Inst* 55:147-152, 1975
- (51) PREUSSMANN R, IVANKOVIC S, LANDSCHUTZ C, et al: Carcinogene Wirkung von 13 Aryldialkyltriazenen an BD-Ratten. *Z. Krebsforsch* 81:285-310, 1974
- (52) MILLER JA: Carcinogenesis by chemicals: An overview. (The G. H. A. Clowes Memorial Lecture). *Cancer Res* 30:559-576, 1970
- (53) IRVING CC: Interaction of chemical carcinogens with DNA. In *Methods in Cancer Research VII* (Busch H, ed). New York: Academic Press, 1973, pp 189-244
- (54) GOTH R, RAJEWSKY MF: Ethylation of nucleic acids by ethylnitrosourea- l - ^{14}C in the fetal and adult rat. *Cancer Res* 32:1501-1505, 1972
- (55) ———: Molecular and cellular mechanisms associated with pulse-carcinogenesis in the rat nervous system by ENI. Ethylation of nucleic acids and elimination rates of ethylated bases from the DNA of different tissues. *Z. Krebsforsch* 82:37-64, 1974
- (56) ———: Persistence of O⁶-ethylguanine in rat-brain DNA. Correlation with nervous system-specific carcinogenesis by ethylnitrosourea. *Proc Natl Acad Sci USA* 71:639-643, 1974
- (57) KLEIHUES P, MARGISON GP: Carcinogenicity of N-methyl-N-nitrosourea: Possible role of excision repair of O⁶-methylguanine from DNA. *J Natl Cancer Inst* 53:1839-1841, 1974
- (58) KLEIHUES P, MAGEE PN: Alkylation of rat brain nucleic acids by N-methyl-N-nitrosourea and methylmethanesulphonate. *J Neurochem* 20:595-606, 1973
- (59) KLEIHUES P, PATZSCHKE K, MARGISON GP, et al: Reaction of methylmethanesulphonate with nucleic acids of fetal and newborn rats in vivo. *Z. Krebsforsch* 81:273-283, 1974
- (60) LOVELESS A: Possible relevance of O⁶ alkylation of deoxyguanosine to the mutagenicity and carcinogenicity of nitrosamines and nitrosamides. *Nature* 223:206-207, 1969
- (61) KLEIHUES P, MAGEE PN, AUSTOKER J, et al: Reaction of N-methyl-N-nitrosourea with DNA of neuronal and glial cells in vivo. *FEBS Lett* 32:105-108, 1973
- (62) KIRTIKAR DM, GOLDTHWAIT DA: The enzymic release of O⁶-methylguanine and 3-methyladenine from DNA reacted with the carcinogen N-methyl-N-nitrosourea. *Proc Natl Acad Sci USA* 71:2022-2026, 1974
- (63) LINDAHL T: New class of enzymes acting on damaged DNA. *Nature* 259:64-66, 1976