

4.

epidemiological techniques to their content and most of the authors do not make claim to having done so. What has happened is that succeeding authors have drawn conclusions and generalized beyond the scope of the works which they quote. Nowhere, for example, have we found references to a population of asbestos workers, although several authors who have quoted the observed incidence of lung cancer in autopsies of persons who also had asbestosis imply that this incidence applies to asbestos workers, generally. We have likewise been unable to find any study which actually calculated the incidence of lung cancer among a population of persons who had asbestosis, and not just those who came to autopsy. With the exception of a paper by Doll⁽²⁸⁾, none of those reviewed gave any data on exposure and dust concentrations, and even Doll's paper merely mentions "scheduled" areas, by which is meant, "those areas where processes are carried on which were scheduled under the Asbestos Industry Regulations of 1931 as being dusty."

There is, furthermore, a complete lack of definition of terms as used in the published literature. For example, the term "asbestosis", as used, may refer to changes observable only by microscopic examination of the lung tissue, or it may mean a radiologically detectable condition.

Most of the published reports obviously included women among their cases, but some of them do not give the number or proportion of women involved in the study.

There is also a lack of uniformity as to what type of exposure most studies have dealt with. Of 99 cases enumerated by Hueper⁽⁵⁸⁾ in 1955, only