

INDUSTRIAL HYGIENE FEATURES

Data on mass screening for tuberculosis and missed cases are presented in Abst. No. 952.

Information on the smoking patterns of Canadian university students is given in Abst. No. 956.

A report on the pathologic effects of smoking tobacco on the trachea and bronchial mucosa is the subject of Abst. No. 958.

See Abst. No. 963 for a discussion of rubber glove dermatitis.

Consult Abst. No. 965 for the management of extensive cutaneous burns.

Two years' experience in a Canadian poison control center is the theme of Abst. No. 966.

Information on an automatic hydrogen fluoride recorder for industrial hygiene and stack monitoring is given in Abst. No. 970.

Abst. No. 973 is concerned with a survey of results of investigations on some organic mercury compounds used as fungicides.

The toxicity of bromochloromethane is considered in Abst. No. 977.

Abst. No. 979 is concerned with a histogenetic study of chronic interstitial pneumonitis.

Hypertension due to inhalation of amorphous silica is the topic of Abst. No. 980.

Factors to be considered regarding discharge electrodes and electrostatic precipitators are discussed in Abst. No. 985.

Is humidity important in determining temperature comfort range? Abst. No. 987.

The topic of Abst. No. 988 is the effect of cold and vascular diseases on cold injury.

An account of the design and use of the A. I. S. sampler is presented in Abst. No. 989.

For odor determination in air pollution control, see Abst. No. 1008.

FOUNDATION FACTS

NEW MEMBERS OF I-H-F. The Foundation welcomes to membership:

Autometric Corporation
Van Straaten Chemical Company

SILVER ANNIVERSARY MEETING THIS MONTH. The immediate response to the invitations for our 25th Annual Meeting which will be held here at Mellon Institute, Pittsburgh, on October 26-27, has been most gratifying. We trust that every member company will be represented at these conferences and are confident that those in attendance will find the meeting both profitable and interesting.

TWO NEW REPRINTS. "Chronic Interstitial Pneumonitis. A Histogenetic Study", by Paul Gross, Arch. Pathol. 69, 706-715, (June 1960); and "The Pulmonary Response to Tuberculin", by Paul Gross, M. L. Westrick, and J. M. McNerney, Arch. Pathol. 70, 233-243 (August 1960). These reprints are available from the Foundation upon request.

- 981 On a Case of Rapidly Evolving Silicosis. B. Milić and Vida Miletić.
Med. lavoro 50, 679-682 (Nov. 1959). Italian.

A case is reported of rapidly evolving silicosis in a 25-year-old worker who was employed for 13 months in the digging of a tunnel for the Mavrovo hydroelectric power station, in the People's Republic of Macedonia. The tunneling was performed by the dry process with no protective measures and it produced a great amount of dust. The rapidity of the onset and the further course of the disease can be explained by the massive inhalation of silica and by a predisposition to the disease. Such predisposition may be due either to a deficient pulmonary auto-depuration or to a particular state of immunological hyperreactivity of the subject as was shown in this case by the behavior of serum proteins. It is concluded that detailed periodical controls with clinical and laboratory investigations of the workers exposed to silicosis hazards may be very useful.

-- Cond. from Bull. Hyg.

- 982 Asbestosis and Lung Carcinoma. 1. I. A. Böhme.
Arch. Gewerbepathol. Gewerbehyg. 17, 384-395 (1959). German.

Previous reports in the literature of a lung carcinoma associated with an asbestosis condition, are reviewed and the results collected in a table in the text. Of a total of 894 tissue specimens examined from cases of known asbestosis, 143 indicated the presence also of lung carcinoma. Using the original groupings, as originally reported, the lung cancer incidence ranged from 15 to 20%. One of these groups is the result of a survey of world literature by Hueper in 1955 where 738 examinations indicated the presence of 143 lung carcinomas or an incidence of 15.5%. It is noted that contained within this latter large grouping are several small ones where the individual reports would indicate an incidence of 31 to 40%. Asbestosis develops slowly following a long period of exposure, and the lung carcinoma appears to make its appearance in the latter part of this period. The author presents detailed case histories of 6 patients from a group of 62, and later 100 asbestosis patients on whom the author and his associates made studies over the period of 1936-1958. In 4 of the 6 patients a lung carcinoma was shown to exist. The interrelationship of the 2 pathological conditions is discussed briefly. The necessity for good pathological differentiations of the 2 conditions is stressed.

-- APCA Absts.

- 983 A Routine Sampler for Detecting Variations in the Emission of Dust and Grit. B. Lees and M. C. Morley. J. Inst. Fuel 33, 90-95 (Feb. 1960).

The Report of the Committee on Air Pollution, 1954, recommended that simpler standard methods of sampling flue gases to determine grit emission should be devised. The sampler described has been developed for routine sampling on a comparative basis. It is extremely simple, yields consistent results and requires only a few minutes per day for routine control. Its application is described for assessing on a routine basis the efficiency of collection of grits by mechanical arrestors and the performance efficiency of electrostatic precipitators on boilers fired with pulverized coal.

-- Public Health Eng. Absts.

- 984 A Study of the Problem of Particle-Size Measurement. H. E. Rose.
Chem. Ingr. Tech. 31, 183-191 (Mar. 1959). German.

The problems of the measurement of the size of particles are considered somewhat in relation to the form and other aspects of the particle, but more especially in relation to particle distribution. The determination of the average-size particles of a dust or powder sample is discussed according to the following methods: gas permeability, light intensity, adsorption in a monomolecular layer, reflection, and bulk density measurements. The methods are considered in detail and several pictures of the various apparatus, or diagrammatic representations of them, are given in the text. Measurement of the particle-size/particle-distribution function is also discussed under headings of screen or sieve analysis, measurement of the velocity of falling bodies, and of particle size in suspension.

-- APCA Absts.