

# TECHNICAL REPORT SUMMARY

File TDB

FC-95

TO: TECHNICAL COMMUNICATIONS CENTER - 201-2C-12

(Important - If report is printed on both sides of paper, send two copies to TCC.) Guidelines on reverse side.

|                                                                                       |                                                |                                     |
|---------------------------------------------------------------------------------------|------------------------------------------------|-------------------------------------|
| Division<br><b>Environmental Laboratory (EE &amp; PC)</b>                             |                                                | Dept. Number<br><b>0535</b>         |
| Project<br><b>New Methods Development</b>                                             |                                                | Project Number<br><b>9970030000</b> |
| Report Title<br><b>Effect of Potassium Perfluorooctanesulfonate on Survival, etc.</b> |                                                | Report Number<br><b>#011</b>        |
| To<br><b>R. L. Bohon</b>                                                              | Period Covered or Date<br><b>2/13/84</b>       |                                     |
| Author(s)<br><b>A. N. Welter</b>                                                      | Employee Number(s)<br><b>09362</b>             |                                     |
| Notebook Reference                                                                    | No. of Pages Including Coversheet<br><b>12</b> |                                     |

SECURITY ☒ Open Report & Summary (Company Confidential) ☐ Closed Report - Open Summary (Special Authorization) 3M CHEMICAL REGISTRY ☐ Check box if new chemicals are reported. Use Chemical Registry Form 6092 to report all new substances.

|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |                                                                                                           |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------|
| KEYWORDS:<br>Lab Code<br><br>Other Keywords<br><br><b>EE &amp; PC</b><br><b>ENV. LAB.</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   | CURRENT OBJECTIVE:<br><br><b>Technology Transfer: To develop a daphnid chronic life cycle capability.</b> |
| REPORT ABSTRACT: This abstract information is distributed by the Technical Communications Center to alert 3M'ers to Company R&D. It is Company confidential material.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |                                                                                                           |
| Abstract - Acute static and chronic renewal <u>Daphnia magna</u> tests were conducted to evaluate this organisms response to varying concentrations of potassium perfluoro-octanesulfonate. The acute 48-h EC <sub>50</sub> was 27 mg/L, with 95% C.L. of 25 to 28 mg/L. In 14, 21, 28-day chronic exposure studies the NOEC was 7 mg/L. At the highest concentration tested (18 mg/L) reproductive indices; viz: cumulative live young per adult, average number of young per brood, and average number of broods per adult, were inhibited to a significant degree $p < 0.05$ . This concentration of potassium perfluorooctanesulfonate was several orders of magnitude greater than levels anticipated under normal use conditions. The survival, growth, and reproduction of this species will not be impacted negatively by chronic 14, 21, or 28-day exposure to potassium perfluorooctanesulfonate. |                                                                                                           |
| This study should be repeated since the two highest concentrations tested, 7 and 18 mg/L represent the NOEC and significant effect levels respectively. There is a need to develop information on concentrations lying between these two extremes.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |                                                                                                           |
| Key Words: Toxicity, acute, chronic, <u>Daphnia magna</u>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |                                                                                                           |

cc: M. T. Elnabarawy  
E. A. Reiner

Information Liaison  
Initials: *Am*

3M CONFIDENTIAL

**Exhibit  
2601**

 State of Minnesota v. 3M Co.,  
Court File No. 27-CV-10-28862

## INTRODUCTION

Ubel et al<sup>(1)</sup> reported that perfluorinated chemicals had been found in the serum of manufacturing plant workers. One of these materials has been the subject of animal toxicity studies<sup>(2)</sup>. Data on a similar compound, potassium perfluorooctanesulfonate, a off-white solid, indicates that this material is water soluble and preliminary information suggests that it will neither readily biodegrade nor bioconcentrate. Preliminary aquatic testing of potassium perfluorooctanesulfonate indicated that toxic signs could be elicited during acute exposure conditions.

Of the aquatic species available for use as test organisms, the crustacean, Daphnia magna is being utilized widely in acute and chronic effects testing. Advantages associated with the use of daphnids are their size, comparatively short life cycle, ecological importance, ease of laboratory culturing, sensitivity, etc.<sup>(3,4,5)</sup> Guidelines for conducting acute 48-h tests have been published<sup>(6)</sup>, and chronic exposure protocols are being developed or have been adopted<sup>(7)</sup>. A broad database consisting of both acute and chronic studies utilizing D. magna presently exists. Daphnids were used, therefore, to evaluate the effect of acute (48-h) and 14, 21, or 28-day chronic exposures to varying concentrations of potassium perfluorooctanesulfonate.

## METHODS AND MATERIALS

### Water Diluent

The dilution water used in all tests was an unchlorinated, carbon filtered well water, aerated to saturation before use. Mean values for some water quality parameters were: pH 7.6 (+0.1), total hardness as mg/L CaCO<sub>3</sub> of 256 (247-265), alkalinity as mg/L CaCO<sub>3</sub> of 224 (219-229) and conductivity as umhos/cm at 25°C of 349 (325-373).

### Test Species

The laboratory stock culture of D. magna was maintained under intermittent illumination, 16h light and 8h dark, at a temperature of 20±2°C. Stock and organisms used during chronic exposure tests were fed blended fish food of known nutritional quality and yeast at the rate of 30 mg solids/L of test solution.

### Acute Tests

During these tests the dilution water temperature averaged 22±1°C, while a photoperiod of 16-h light and 8-h darkness was maintained throughout the test. For the 48-h tests, 20 daphnids 12 ± 12-h old were placed in duplicate<sup>(6)</sup> test beakers, 250 ml capacity, each containing 200 ml of test solution. All acute tests were run without food additions. Five concentrations of potassium perfluorooctanesulfonate, and an untreated control group were run in duplicate. The endpoint used in these tests was daphnid immobilization as characterized by a lack of coordinated movement. The calculated EC<sub>50</sub> value was used to define the range of concentrations to be used in the life-cycle renewal test.

### Chronic Tests

The chronic daphnid studies using first instars 12 + 12-h old were conducted in a temperature controlled growth chamber at  $23 \pm 1^{\circ}\text{C}$  with photoperiod automatically fixed to a 16-h light, and 8-h dark cycle. This static renewal chronic test was conducted using ten 250 ml beakers containing 200 ml test solution at each of five concentrations of the test material plus an untreated control set of 20 beakers. Each test concentration was replicated three times with five Daphnia magna first instars per beaker for survival data and seven times with but one daphnid per beaker for reproductive measurements. Stock solutions were prepared in deionized water. Fresh solutions were made up thrice weekly (MWF) with food also being replenished at this time. Disposable glass pipettes having an I.D. of 5 mm were used to transfer the adult daphnids to fresh solution. The young daphnids were then counted and discarded.

During the test the following data were tabulated: day of initial brood release, number of young per adult per day, number of broods per adult, mortality of young or adult daphnids. All data are based on nominal concentrations of potassium perfluorooctanesulfonate.

### Statistics

The 48-h, acute and 14, 21, and 28-day chronic  $\text{EC}_{50}$  values and their associated 95% confidence limits were determined by either probit analysis<sup>(8)</sup> or the moving angle average method<sup>(9)</sup>. Reproductive and survival data were analyzed by analysis of variance. Throughout the text "significant" will refer to statistical significance, having a p value of .05.

## RESULTS

### Acute Test

The calculated 48-h  $\text{EC}_{50}$  value was 27 mg/L of potassium perfluorooctanesulfonate with a 95% C.L. ranging from 25 to 28 mg/l. Based on this value the range of test concentrations selected for the daphnid chronic test were 0.26, 1.0, 2.6, 7.0, and 18 mg/l of potassium perfluorooctanesulfonate. Temperature and pH varied minimally while the dissolved oxygen level did not fall below 70% saturation.

### Chronic Test - 14, 21, and 28-Day Duration

Indices of reproductive performance and survival of Daphnia magna exposed to the above listed concentrations of the test material are tabulated in Tables 1-4.

Untreated control reproductive and survival indices as shown in Table 1 are in agreement with reported literature values<sup>(10)</sup>. Literature values cited for the cumulative number of young per adult ranged from 30-122 at temperatures of  $20-25^{\circ}\text{C}$  while at temperatures of  $18-20^{\circ}\text{C}$  Daphnia magna survival varied from 80-92%.

The effect on reproductive indices elicited by chronic exposure to potassium perfluorooctanesulfonate are illustrated (Table 1). Over the 28-day exposure period and at chemical concentrations varying from 0.26 to 7.0 mg/L the mean values of the above cited parameters varied minimally when compared to control data. Thus, the absence of a concentration dependent response up to 7.0 mg/L is apparent.

Interference with the normal reproductive process did occur at a toxicant concentration between 7.0 and 18.0 mg/L. At this latter concentration, the cumulative number of live young per adult were reduced by 80%, average number of live young per brood was inhibited by 55%, and the number of broods per adult was reduced by 62%.

It has been recommended<sup>(7)</sup> that three brood studies replace 21 or 28 day chronic exposure tests (Table 2). At potassium perfluorooctanesulfonate concentrations up to and including 7.0 mg/L the time to three brood release was either 14 or 15 days after initiation of the study. At 18 mg/L it required approximately 17 days for release of the three broods, a statistically significant difference when compared to control values.

Calculated values for 50% reproductive impairment at 14, 21, and 28 days of exposure to potassium perfluorooctanesulfonate are shown (Table 3). Adult mortality  $EC_{50}$  values could not be determined as they exceeded the highest test concentration used in this study. The 28-day  $EC_{50}$  values calculated for all reproductive parameters were similar, ranging from 11.4 to 16.1 mg/l. Overlap of the 95% C.L. is indicative of the absence of statistically significant differences in these values.

In this study statistically significant changes in all reproductive indices did occur at a concentration of 18 mg/L potassium perfluorooctanesulfonate (Table 4). Exposure to concentrations of 0.26, 1.0, 2.6, and 7.0 mg/L of the test substance did not affect these indices or mortality to a significant degree. Thus, a concentration of 7.0 mg/L of potassium perfluorooctanesulfonate is considered to be the chronic NOEC or no observable effect concentration.

Table 5 illustrates the derived acute/chronic ratios for those reproductive indices evaluated throughout the potassium perfluorooctanesulfonate exposure period. The 28-day acute/chronic ratios for all parameters studied were identical and were equivalent to the 14 and 21-day acute/chronic ratios derived from cumulative number of young per adult data. This may indicate a concentration dependent process which is not affected by increasing the exposure duration.

Table 6 shows the chronic renewal toxicity of potassium perfluorooctanesulfonate expressed as the NOEC, MATC, and  $EC_{50}$  which values were derived from reproductive indices. As in the preceding table, the derived values are chemical dependent and time independent. Thus, the NOEC and MATC are identical for all parameters measured at all exposure periods. The  $EC_{50}$  values differ slightly however there is no statistical difference in these values based on variable exposure durations.

### DISCUSSION

Reproductive measurements have been used as indicators of toxicant effects following exposure to sublethal concentrations of PCB, p,p' DDT, or Atrazin®. Furthermore, Parkhurst et al<sup>(3)</sup> investigated the reproducibility of the Daphnia magna chronic test and concluded that, of the toxicity criteria studied, the number of young produced per brood or per adult were sensitive indicators of a chronic toxicant effect.

Our results are in agreement with those reported by Maki and Johnson<sup>(13)</sup> and Parkhurst et al<sup>(3)</sup> that changes in indices of reproduction are sensitive parameters of a toxicant effect. Changes occurring at concentration levels of 18 mg/L of potassium perfluorooctanesulfonate included reductions in: 1) the average number of broods per parent, 7 to 3; 2) number of live young per parent, 190 to 37; 3) average number of young per brood 27 to 13. An increase in the number (to 54) of dead young per parent was also noted. When combining the number of live and dead young a 52% reduction in total number of young released was observed. The most noteworthy result of daphnid exposure to high levels of potassium perfluorooctanesulfonate has been the attenuation of all reproductive indices. On the other hand, at this concentration maturation was not delayed nor was longevity of the parent affected.

Water quality guidelines have specified the use of two protective concentrations, namely a 24-h average equivalent to the no observable effect concentration (NOEC) and either a 24-h maximum or 48-h EC<sub>50</sub> level. Neither of these protective concentrations should be exceeded over the indicated time period. Utilizing this concept, protective concentrations of 7.0 and 27 mg/L would be established for potassium perfluorooctanesulfonate. Over the 28-day exposure period no observable effects occurred using either mortality or reproductive indices as the response criteria. A NOEC of 7 mg/L and the toxicant level which elicited an effect, 18 mg/L, give us a gross approximation of the GM-MATC, equal to 11 mg/L. Conversely, during a 28-day chronic exposure to 18 mg/L of potassium perfluorooctanesulfonate, reproductive impairment was noted. This phenomenon was observed as early as the second week and remained at the same EC<sub>50</sub> value for the duration of the study.

Based on these findings and considering the concentrations of test chemical necessary to elicit a chronic effect, it is concluded that potassium perfluorooctanesulfonate will not impact negatively on the survival, growth and reproduction of Daphnia magna under present environmental release conditions.

### REFERENCES

1. Ubel, F. A., Sorenson, S. D., and Roach, D. E. 1980. Health Status of plant workers exposed to fluorochemicals-a preliminary report. J. Am. Ind. Hyg. Assoc. 41:584-589.
2. Griffith, F. D. and J. E. Long, 1980. Animal Toxicity Studies with Ammonium Perfluorooctanesulfonate. Am. Ind. Hyg. Assoc. 41:576-583.

3. Parkhurst, B. R., J. L. Forte and G. P. Wright, 1981. Reproducibility of a Life-Cycle Toxicity Test with *Daphnia magna*. Bull. Environ. Contam. Toxicol 26:1-8.
4. Presing, M. 1981. On the Effects of Dikonirt (sodium salt of 2,4-Dichlorophenoxy-acetic acid) on the Mortality and Reproduction of *Daphnia magna* Hydrobiologia 83:511-516.
5. Adema, D. M. M., Kuiper, J., Hantsveit, A. O., and Canton, J. H. 1982. Consecutive System of Tests for Assessment of the Effects of Chemical Agents in the Aquatic Environment. National Technical Information Service, Springfield, VA.
6. American Society for Testing Materials, 1981, Proposed Standard practice for conducting renewal life-cycle toxicity tests with the *Daphnia magna*. American Society for Testing and Materials, Philadelphia, PA.
7. OECD. 1981. *Daphnia* sp., 14-day reproduction test (including an acute immobilisation test). Organization of Economic Cooperation and Development. Paris, France.
8. Finney, D. J. 1971. Probit Analysis. University Press, Cambridge.
9. Stephan C. E. 1977. Methods for calculating an LC<sub>50</sub>. In F. L. Mayer and J.L. Hamelink, eds. Aquatic toxicology and hazard evaluation. American Society for Testing and Materials, Philadelphia, pp. 65-84.
10. U.S. EPA 1982. Environmental Effects Test Guidelines, EPA 560/6-82-002 ES-1, PP. 1-42.
11. Nebeker, A. V., C. K. McAuliffe, R. Mshar, and G. Stevens, 1983. Toxicity of silver to steelhead and rainbow trout, fathead minnows and *Daphnia magna*. Environ, Toxicol. Chem. 2:95-104.
12. Maki, A. W., and H. E. Johnson, 1975. Effects of PCB (Aroclor R54) and p,p'DDT on production and survival of *Daphnia magna* Strauss. Bull Environ Contam. Toxicol 13:412-416.
13. Schober, V. and W. Lampert: 1977. Effects of Sublethal Concentrations of the herbicide Atrazin® on growth and reproduction of *Daphnia Pulex*. Bull. Environ Contam. Toxicol: 269-277.

TABLE 1

Chronic Exposure of Daphnia magna to  
Potassium Perfluorooctanesulfonate:  
Young Reproduction<sup>a</sup>

| Treatment<br>mg/L | n  | Cumulative Live<br>Young/Adult<br>Number | Average Number<br>Young/Brood | Avg. Number<br>Broods/Adult |
|-------------------|----|------------------------------------------|-------------------------------|-----------------------------|
| Control           | 11 | 190<br><u>+36</u>                        | 27<br><u>+4</u>               | 7<br><u>+1</u>              |
| 0.26              | 11 | 169<br><u>+33</u>                        | 26<br><u>+4</u>               | 7<br><u>+1</u>              |
| 1.0               | 7  | 183<br><u>+40</u>                        | 25<br><u>+4</u>               | 7<br><u>+1</u>              |
| 2.6               | 7  | 186<br><u>+29</u>                        | 26<br><u>+4</u>               | 7<br><u>+1</u>              |
| 7.0               | 6  | 199<br><u>+19</u>                        | 27<br><u>+3</u>               | 7<br><u>+1</u>              |
| 18.0              | 7  | 37 <sup>b</sup><br><u>+28</u>            | 12 <sup>b</sup><br><u>+8</u>  | 3 <sup>b</sup><br><u>+1</u> |

<sup>a</sup> 28-day exposure data. Mean  $\pm$  S.D.

<sup>b</sup>  $p \leq 0.05$ .

TABLE 2

Chronic Exposure of Daphnia magna to Potassium  
Perfluorooctanesulfonate: Comparison of 1st and  
3rd Brood Release Days

| Treatment<br>mg/L |                | Initial Brood<br>Release Day | Third Brood<br>Release Day   |
|-------------------|----------------|------------------------------|------------------------------|
| Control           | n <sup>a</sup> | 9<br><u>+1</u>               | 15<br><u>+1</u>              |
| 0.26              | 17             | 10<br><u>+2</u>              | 15<br><u>+2</u>              |
| 1.0               | 10             | 9<br><u>+1</u>               | 15<br><u>+2</u>              |
| 2.6               | 10             | 9<br><u>+0.3</u>             | 15<br><u>+2</u>              |
| 7.0               | 9              | 9<br><u>+0.3</u>             | 14<br><u>+1</u>              |
| 18.0              | 10             | 10<br><u>+2</u>              | 17 <sup>b</sup><br><u>+2</u> |

<sup>a</sup> Beakers in test.

<sup>b</sup>  $p \leq 0.05$ .

Table 3

Acute and Chronic Toxicity of Potassium Perfluorooctanesulfonate  
to Daphnia magna: Acute and Chronic EC<sub>50</sub>

| <u>Exposure<br/>Period</u> | <u>Immobilization</u> | <u>Reproduction EC<sub>50</sub> mg/L (95% C.L.)</u> |                                 |                                  |
|----------------------------|-----------------------|-----------------------------------------------------|---------------------------------|----------------------------------|
|                            |                       | <u>Cumulative<br/>Young/Adult</u>                   | <u>Live Young<br/>Per Brood</u> | <u>Avg. No.<br/>Broods/Adult</u> |
| 48-hr                      | 26.6<br>(25-28)       |                                                     |                                 |                                  |
| 14 Day                     |                       | 14.7<br>(12-18)                                     | a                               | a                                |
| 21 day                     |                       | 12.4<br>(11-14)                                     |                                 |                                  |
| 28 day                     |                       | 11.4<br>(10-13)                                     | 16.1<br>(13-20)                 | 14.7<br>(13-18)                  |

<sup>a</sup> Not calculated.

TABLE 4

Comparison of the Chronic Effects of 7 and 18 mg/L  
Potassium Perfluorooctanesulfonate to Daphnia magna

|                                       | <u>14-Day</u> | <u>21-Day</u> | <u>28-Day</u> |
|---------------------------------------|---------------|---------------|---------------|
| A. 7.0 mg/L:                          |               |               |               |
| Number Live Young /Adult              | 59            | 129           | 199           |
| Number Dead/Adult                     | 3             | 3             | 4             |
| Total Number Young                    | 369           | 793           | 1220          |
| Adult Mortality %                     | 0             | 4.5           | 4.5           |
| B. 18.0 mg/L:                         |               |               |               |
| Number Live Young /Adult <sup>a</sup> | 24            | 33            | 37            |
| Number Dead/Adult <sup>a</sup>        | 12            | 51            | 54            |
| Total Number Young <sup>a</sup>       | 249           | 591           | 641           |
| Adult Mortality %                     | 4.5           | 9.1           | 18.2          |

<sup>a</sup> Comparison of 7.0 and 18.0 mg/L  $p \leq 0.05$ .

TABLE 5

Acute/Chronic EC<sub>50</sub> Ratios Following Exposure of  
Daphnia magna to Potassium Perfluorooctanesulfonate<sup>a</sup>

| <u>Day of<br/>Exposure</u> | <u>Cumulative<br/>Young/Adult</u> | <u>Cumulative<br/>Broods/Adult</u> | <u>Number of<br/>Young/Adult</u> |
|----------------------------|-----------------------------------|------------------------------------|----------------------------------|
| 14                         | 1.5                               |                                    |                                  |
| 21                         | 1.5                               |                                    |                                  |
| 28                         | 1.5                               | 1.5                                | 1.5                              |

<sup>a</sup> Acute/Chronic EC<sub>50</sub> Ratio = 
$$\frac{48\text{h Concentration Eliciting an Acute Effect in 50\% of Organisms}}{(\text{Exposure Duration}) \text{ Concentration Eliciting a Chronic Effect in 50\% of Organisms}}$$

TABLE 6

Chronic Renewal Toxicity of Potassium  
Perfluorooctanesulfonate to *Daphnia magna*<sup>a</sup>

| Days                          | 14              | Cumulative<br>Young/Adult<br>21 | 28              | Cumulative<br>Broods/Adult<br>28 | Number of<br>Young/Adult<br>28 |
|-------------------------------|-----------------|---------------------------------|-----------------|----------------------------------|--------------------------------|
| NOEC <sup>b</sup>             | 7               | 7                               | 7               | 7                                | 7                              |
| MATC <sup>c,d</sup>           | 11.2            | 11.2                            | 11.2            | 11.2                             | 11.2                           |
| EC <sub>50</sub> <sup>e</sup> | 14.7<br>(12-18) | 12.4<br>(11-14)                 | 11.4<br>(10-13) | 14.7<br>(13-18)                  | 16.1<br>(13-20)                |

<sup>a</sup> Data in mg/L.

<sup>b</sup> No observable effect concentration.

<sup>c</sup> Maximum acceptable toxicant concentration.

<sup>d</sup> MATC = Geometric mean between highest test concentration having no significant ( $p \geq 0.05$ ) effect and the lowest test concentration having a significant ( $p \leq 0.05$ ) effect.

<sup>e</sup> Dose at which 50% of organisms are affected by the toxicant. Expressed as Mean (95% C.L.).