



E. I. DU PONT DE NEMOURS & COMPANY
WILMINGTON, DELAWARE 19896

FABRICS AND FINISHES DEPARTMENT
RESEARCH DIVISION

cc: J. A. Zapp - Haskell Lab.

C.C.S.-7

file

4-13

April 3, 1973

R. J. MUSIAK
HASKELL LAB.

MESOTHELIOMA IN RELATION TO ASBESTOS

The attached tables are from a report by Dr. W. E. Smith of the Health Research Institute at Fairleigh Dickinson. We do not have the full report but presume that it is available if you wish to order it.

E. E. SWAIN, JR.

EES:ps

RECEIVED

1345

APR 11 1973

HASKELL LABORATORY

BETTER THINGS FOR BETTER LIVING

DUP 0903156

120970

DU 024538

Vanderbilt Technical
Data

Vanderbilt Company Inc. • 230 Park Avenue • New York, N.Y. 10017 • (212) 686-6666

A-132

TALC AND THE ASBESTOS DUST STANDARDS

DUP 0903157

The recommendations for use of our materials are based upon tests carried out to date. However, we do not guarantee the results of our tests.

120971

DU 024539

Exhibit 1

"ASBESTOS" MINERALS AS LISTED IN FEDERAL REGISTER

<u>Asbestos or Fibrous Variety</u>	<u>Chemical Composition</u>	<u>Non-Asbestos or Non-Fibrous Variety</u>
Chrysotile	$Mg_3(Si_2O_5)(OH)_4$	Antigorite, Lizardite
— Crocidolite	$Na_2Fe_3Fe_2(Si_3O_{22})(OH)_2$	Riebeckite
Amosite	$(Mg, Fe)_7(Si_8O_{22})(OH)_2$	Cummingtonite
()	$(Mg, Fe)_7(Si_8O_{22})(OH, F)_2$	Anthophyllite
()	$Ca_2Mg_5(Si_8O_{22})(OH, F)_2$	Tremolite
()	$Ca_2(Mg, Fe)_5(Si_8O_{22})(OH, F)_2$	Actinolite

DUP 0903158

120972

DU 024540

A-134

Exhibit 8

The source of the data on the following pages, 3 and 4, is:

TERMINAL PROGRESS REPORT: GRANT NO. 3R01 EC-00226-0451

TITLE: MESOTHELIOMA IN RELATION TO ASBESTOS

PERIOD: 03/01/66 - 08/31/70

Principal Investigator: William E. Smith, M.D.
Health Research Institute
Fairleigh Dickinson University
Madison, N.J. 07940

The aim of this project was to test 23 preparations of asbestos or other minerals for carcinogenicity by means of intrapleural injections into hamsters. Each sample was tested in a group of 50 hamsters.

DUP 0903159

120973

DU 024541

F-3-

Exhibit 8

Table 1. FINDINGS IN HAMSTERS AFTER INTRAPLEURAL INJECTION
 50 hamsters per group
 (Cumulative data to 650 days except where noted)

SAMPLE	Mean Fiber Length microns	Dose mg.	Mesothel- ioma	Sarcoma Injection Site	Lymphoid Tumors	Remarks
Group No.						
1. Soft Chrysotile	32.54	25	8	2	1	
2. Harsh Chrysotile	38.85	25	9	3	1	
3. Soft Chrys. (heated)	32.54	25	10	0	0	
4. Soft Chrysotile	.76	25	0	0	2	700 days
5. Harsh Chrysotile	1 (approx.)	25	0	1	3	700 days
6. Soft Chrys. (heated)	1 (approx.)	25	0	1	0	700 days
7. Harsh Chrysotile	38.85	10	4	1	1	
8. Harsh Chrysotile	38.85	1	0	1	0	700 days
9. Amosite	17	10	3	2	0	
10. Amosite	17	1	0	0	0	
11. Serpentine Rock Dust (Antigorite)		25	1	0	0	700 days
12. Microquartz Fibers	55.7	10	4	1	3	
13. Nematite	53.1	25	0	0	2	

DUP 0903160

120974

DU 024542

- 136

Exhibit 8

Table 1 (continued) FINDINGS IN HAMSTERS AFTER INTRAPLEURAL INJECTION
 50 hamsters per group
 (Cumulative data to 650 days except where noted)

SAMPLE Group No.	Mean Fiber Length microns	Dose mg.	Mesothel- ioma	Sarcoma Injection Site	Lymphoid Tumors	Remarks
14. Tremolite Talc		25	0	0	0	700 days
15. Diatomaceous Earth		25	3	2	2	
15A. Diatomaceous Earth (Calcined)		25	4	2	1	
16. Fiberglass with Phenol Formal Binder	24.9	10	0	0	5	
17. Nickel Powder	8	10	4	4	0	
18. Crocidolite	UICC	10	10	5	1	
19. Crocidolite	UICC	1	2	0	4	700 days
20. Anthophyllite	Long (UICC)	10	3	0	2	
21. Anthophyllite	Short (3.7)	10	1	0	0	
22. Glass Fibers	16.9	10	0	0	5	
23. Untreated		none	0	0	0	700 days
24. Untreated		none	0	0	4	660 days (survivors are being followed to 700 days)

DUP 0903161

120975

DU 024543

R.T. Vanderbilt Company, Inc. 230 Park Avenue New York, N.Y. 10017 212-686-6862

TALC AND THE ASBESTOS DUST STANDARDS

1015

The recommendations for use of our material are based upon tests conducted to date, and are subject to change without notice.

DUP 0903162

DU 024544

Exhibit 1

"ASBESTOS" MINERALS AS LISTED IN FEDERAL REGISTER

<u>Asbestos or Fibrous Variety</u>	<u>Chemical Composition</u>	<u>Non-Asbestos or Non-Fibrous Variety</u>
Chrysotile	$Mg_3(Si_2O_5)(OH)_4$	Antigorite, Lizardite
— Crocidolite	$Na_2Fe_3Fe_2(Si_3O_{22})(OH)_2$	Riebeckite
Amosite	$(Mg, Fe)_7(Si_8O_{22})(OH)_2$	Cummingtonite
()	$(Mg, Fe)_7(Si_8O_{22})(OH, F)_2$	Anthophyllite
()	$Ca_2Mg_5(Si_8O_{22})(OH, F)_2$	Tremolite
()	$Ca_2(Mg, Fe)_5(Si_8O_{22})(OH, F)_2$	Actinolite

DUP 0903163

DU 024545

Exhibit B

The source of the data on the following pages, 3 and 4, is:

TERMINAL PROGRESS REPORT: GRANT NO. 3R01 EC-00226-0451

TITLE: MESOTHELIOMA IN RELATION TO ASBESTOS

PERIOD: 03/01/66 - 08/31/70

Principal Investigator: William E. Smith, M.D.
Health Research Institute
Fairleigh Dickinson University
Madison, N.J. 07940

The aim of this project was to test 23 preparations of asbestos or other minerals for carcinogenicity by means of intrapleural injections into hamsters. Each sample was tested in a group of 50 hamsters.

DUP 0903164

DU 024546

Exhibit 8

Table 1. FINDINGS IN HAMSTERS AFTER INTRAPLEURAL INJECTION
 50 hamsters per group
 (Cumulative data to 650 days except where noted)

SAMPLE Group No.	Mean Fiber Length microns	Dose mg.	Mesothel- ioma	Sarcoma Injection Site	Lymphoid Tumors	Remarks
1. Soft Chrysotile	32.54	25	8	2	1	
2. Harsh Chrysotile	38.85	25	9	3	1	
3. Soft Chrys. (heated)	32.54	25	10	0	0	
4. Soft Chrysotile	.76	25	0	0	2	700 days
5. Harsh Chrysotile	1 (approx.)	25	0	1	3	700 days
6. Soft Chrys. (heated)	1 (approx.)	25	0	1	0	700 days
7. Harsh Chrysotile	38.85	10	4	1	1	
8. Harsh Chrysotile	38.85	1	0	1	0	700 days
9. Amosite	17	10	3	2	0	
10. Amosite	17	1	0	0	0	
11. Serpentine Rock Dust (Antigorite)		25	1	0	0	700 days
12. Microquartz Fibers	56.7	10	4	1	3	
13. Nematite	53.1	25	0	0	2	

DUP 0903165

DU 024547

Exhibit 8

Table 1 (continued) FINDINGS IN HAMSTERS AFTER INTRAPLEURAL INJECTION
50 hamsters per group
(Cumulative data to 650 days except where noted)

SAMPLE	Mean Fiber Length microns	Dose mg.	Mesothel- ioma	Sarcoma Injection Site	Lymphoid Tumors	Remarks
Group No.						
14. Tremolite Talc		25	0	0	0	700 days
15. Diatomaceous Earth		25	3	2	2	
15A. Diatomaceous Earth (Calcined)		25	4	2	1	
16. Fiberglass with Phenol Formal Binder	24.9	10	0	0	5	
17. Nickel Powder	8	10	4	4	0	
18. Crocidolite	UICC	10	10	5	1	
19. Crocidolite	UICC	1	2	0	4	700 days
20. Anthophyllite	Long (UICC)	10	3	0	2	
21. Anthophyllite	Short (3.7)	10	1	0	0	
22. Glass Fibers	16.9	10	0	0	5	
23. Untreated		none	0	0	0	700 days
24. Untreated		none	0	0	4	660 days (survivors are being followed to 700 days)

DUP 0903166

DU 024548

1. Verderer Company Inc. - 230 Park Avenue - New York NY 10017 - (212) 688-6881.

A. 52

TALC AND THE ASBESTOS DUST STANDARDS

DUP 0903167

1997-1998-1999-2000-2001-2002-2003-2004-2005-2006-2007-2008-2009-2010-2011-2012-2013-2014-2015-2016-2017-2018-2019-2020-2021-2022-2023-2024-2025-2026-2027-2028-2029-2030-2031-2032-2033-2034-2035-2036-2037-2038-2039-2040-2041-2042-2043-2044-2045-2046-2047-2048-2049-2050-2051-2052-2053-2054-2055-2056-2057-2058-2059-2060-2061-2062-2063-2064-2065-2066-2067-2068-2069-2070-2071-2072-2073-2074-2075-2076-2077-2078-2079-2080-2081-2082-2083-2084-2085-2086-2087-2088-2089-2090-2091-2092-2093-2094-2095-2096-2097-2098-2099-2100-2101-2102-2103-2104-2105-2106-2107-2108-2109-2110-2111-2112-2113-2114-2115-2116-2117-2118-2119-2120-2121-2122-2123-2124-2125-2126-2127-2128-2129-2130-2131-2132-2133-2134-2135-2136-2137-2138-2139-2140-2141-2142-2143-2144-2145-2146-2147-2148-2149-2150-2151-2152-2153-2154-2155-2156-2157-2158-2159-2160-2161-2162-2163-2164-2165-2166-2167-2168-2169-2170-2171-2172-2173-2174-2175-2176-2177-2178-2179-2180-2181-2182-2183-2184-2185-2186-2187-2188-2189-2190-2191-2192-2193-2194-2195-2196-2197-2198-2199-2200-2201-2202-2203-2204-2205-2206-2207-2208-2209-2210-2211-2212-2213-2214-2215-2216-2217-2218-2219-2220-2221-2222-2223-2224-2225-2226-2227-2228-2229-2230-2231-2232-2233-2234-2235-2236-2237-2238-2239-2240-2241-2242-2243-2244-2245-2246-2247-2248-2249-2250-2251-2252-2253-2254-2255-2256-2257-2258-2259-2260-2261-2262-2263-2264-2265-2266-2267-2268-2269-2270-2271-2272-2273-2274-2275-2276-2277-2278-2279-2280-2281-2282-2283-2284-2285-2286-2287-2288-2289-2290-2291-2292-2293-2294-2295-2296-2297-2298-2299-2300-2301-2302-2303-2304-2305-2306-2307-2308-2309-2310-2311-2312-2313-2314-2315-2316-2317-2318-2319-2320-2321-2322-2323-2324-2325-2326-2327-2328-2329-2330-2331-2332-2333-2334-2335-2336-2337-2338-2339-2340-2341-2342-2343-2344-2345-2346-2347-2348-2349-2350-2351-2352-2353-2354-2355-2356-2357-2358-2359-2360-2361-2362-2363-2364-2365-2366-2367-2368-2369-2370-2371-2372-2373-2374-2375-2376-2377-2378-2379-2380-2381-2382-2383-2384-2385-2386-2387-2388-2389-2390-2391-2392-2393-2394-2395-2396-2397-2398-2399-2400-2401-2402-2403-2404-2405-2406-2407-2408-2409-2410-2411-2412-2413-2414-2415-2416-2417-2418-2419-2420-2421-2422-2423-2424-2425-2426-2427-2428-2429-2430-2431-2432-2433-2434-2435-2436-2437-2438-2439-2440-2441-2442-2443-2444-2445-2446-2447-2448-2449-2450-2451-2452-2453-2454-2455-2456-2457-2458-2459-2460-2461-2462-2463-2464-2465-2466-2467-2468-2469-2470-2471-2472-2473-2474-2475-2476-2477-2478-2479-2480-2481-2482-2483-2484-2485-2486-2487-2488-2489-2490-2491-2492-2493-2494-2495-2496-2497-2498-2499-2500-2501-2502-2503-2504-2505-2506-2507-2508-2509-2510-2511-2512-2513-2514-2515-2516-2517-2518-2519-2520-2521-2522-2523-2524-2525-2526-2527-2528-2529-2530-2531-2532-2533-2534-2535-2536-2537-2538-2539-2540-2541-2542-2543-2544-2545-2546-2547-2548-2549-2550-2551-2552-2553-2554-2555-2556-2557-2558-2559-2560-2561-2562-2563-2564-2565-2566-2567-2568-2569-2570-2571-2572-2573-2574-2575-2576-2577-2578-2579-2580-2581-2582-2583-2584-2585-2586-2587-2588-2589-2590-2591-2592-2593-2594-2595-2596-2597-2598-2599-2600-2601-2602-2603-2604-2605-2606-2607-2608-2609-2610-2611-2612-2613-2614-2615-2616-2617-2618-2619-2620-2621-2622-2623-2624-2625-2626-2627-2628-2629-2630-2631-2632-2633-2634-2635-2636-2637-2638-2639-2640-2641-2642-2643-2644-2645-2646-2647-2648-2649-2650-2651-2652-2653-2654-2655-2656-2657-2658-2659-2660-2661-2662-2663-2664-2665-2666-2667-2668-2669-2670-2671-2672-2673-2674-2675-2676-2677-2678-2679-2680-2681-2682-2683-2684-2685-2686-2687-2688-2689-2690-2691-2692-2693-2694-2695-2696-2697-2698-2699-2700-2701-2702-2703-2704-2705-2706-2707-2708-2709-2710-2711-2712-2713-2714-2715-2716-2717-2718-2719-2720-2721-2722-2723-2724-2725-2726-2727-2728-2729-2730-2731-2732-2733-2734-2735-2736-2737-2738-2739-2740-2741-2742-2743-2744-2745-2746-2747-2748-2749-2750-2751-2752-2753-2754-2755-2756-2757-2758-2759-2760-2761-2762-2763-2764-2765-2766-2767-2768-2769-2770-2771-2772-2773-2774-2775-2776-2777-2778-2779-2780-2781-2782-2783-2784-2785-2786-2787-2788-2789-2790-2791-2792-2793-2794-2795-2796-2797-2798-2799-2800-2801-2802-2803-2804-2805-2806-2807-2808-2809-2810-2811-2812-2813-2814-2815

120971

DU 024549

Exhibit 1

"ASBESTOS" MINERALS AS LISTED IN FEDERAL REGISTER

<u>Asbestos or Fibrous Variety</u>	<u>Chemical Composition</u>	<u>Non-Asbestos or Non-Fibrous Variety</u>
Chrysotile	$Mg_3(Si_2O_5)(OH)_4$	Antigorite, Lizardite
— Crocidolite	$Na_2Fe_3Fe_2(Si_3O_{22})(OH)_2$	Riebeckite
Amosite	$(Mg, Fe)_7(Si_8O_{22})(OH)_2$	Cummingtonite
()	$(Mg, Fe)_7(Si_8O_{22})(OH, F)_2$	Anthophyllite
()	$Ca_2Mg_5(Si_8O_{22})(OH, F)_2$	Tremolite
()	$Ca_2(Mg, Fe)_5(Si_8O_{22})(OH, F)_2$	Actinolite

DUP 0903168

120972

DU 024550

ن ج - م

Exhibit 8

The source of the data on the following pages, 3 and 4, is:

TERMINAL PROGRESS REPORT: GRANT NO. 3R01 EC-00226-0451

TITLE: MESOTHELIOMA IN RELATION TO ASBESTOS

PERIOD: 03/01/66 - 08/31/70

Principal Investigator: William E. Smith, M.D.
Health Research Institute
Fairleigh Dickinson University
Madison, N.J. 07940

The aim of this project was to test 23 preparations of asbestos or other minerals for carcinogenicity by means of intrapleural injections into hamsters. Each sample was tested in a group of 50 hamsters.

DUP 0903169

120973

DU 024551

Exhibit 8

Table 1. FINDINGS IN HAMSTERS AFTER INTRAPLEURAL INJECTION
 50 hamsters per group
 (Cumulative data to 650 days except where noted)

SAMPLE Group No.	Mean Fiber Length microns	Dose mg.	Mesothel- ioma	Sarcoma Injection Site	Lymphoid Tumors	Remarks
1. Soft Chrysotile	32.54	25	8	2	1	
2. Harsh Chrysotile	38.85	25	9	3	1	
3. Soft Chrys. (heated)	32.54	25	10	0	0	
4. Soft Chrysotile	1.76	25	0	0	2	700 days
5. Harsh Chrysotile	1 (approx.)	25	0	1	3	700 days
6. Soft Chrys. (heated)	1 (approx.)	25	0	1	0	700 days
7. Harsh Chrysotile	38.85	10	4	1	1	
8. Harsh Chrysotile	38.85	1	0	1	0	700 days
9. Amosite	17	10	3	2	0	
10. Amosite	17	1	0	0	0	
11. Serpentine Rock Dust (Antigorite)		25	1	0	0	700 days
12. Microquartz Fibers	56.7	10	4	1	3	
13. Nemo-lite	53.1	25	0	0	2	

DUP 0903170

120974

DU 024552

Exhibit 8

Table 1 (continued) FINDINGS IN HAMSTERS AFTER INTRAPLEURAL INJECTION
 50 hamsters per group
 (Cumulative data to 650 days except where noted)

SAMPLE Group No.	Mean Fiber Length microns	Dose mg.	Mesothelioma	Sarcoma Injection Site	Lymphoid Tumors	Remarks
14. Tremolite Talc		25	0	0	0	700 days
15. Diatomaceous Earth		25	3	2	2	
15A. Diatomaceous Earth (Calcined)		25	4	2	1	
16. Fiberglass with Phenol Formal Binder	24.9	10	0	0	5	
17. Nickel Powder	8	10	4	4	0	
18. Crocidolite	UICC	10	10	5	1	
19. Crocidolite	UICC	1	2	0	4	700 days
20. Anthophyllite	Long (UICC)	10	3	0	2	
21. Anthophyllite	Short (3.7)	10	1	0	0	
22. Glass Fibers	16.9	10	0	0	5	
23. Untreated		none	0	0	0	700 days
24. Untreated		none	0	0	4	660 days (survivors are being followed to 700 days)

128975

DUP 0903171

DU 024553

X
DRAFT

EXPOSURE TO ASBESTOS DUST

In the interest of protecting the health of its employees, the Du Pont Company and your Plant Management have established an acceptable level for asbestos exposure. The recommendation for the acceptable level is based upon the information that no health problems related to asbestos will result in anyone who is exposed to this concentration for their working lifetime. The reason this level has been made mandatory is naturally to protect the health of people who are occupationally exposed to asbestos. I am sure some of you have seen Dr. Hubiak from Haskell Lab checking air samples around the shop areas while you were doing your normal jobs. The purpose for these air checks was to determine the concentrations of asbestos fibers in the air and to determine what changes in jobs or equipment were necessary to get and keep the air levels of asbestos at a minimum and an acceptable level. These changes probably have had some effect upon the way in which you perform your job, but they were done to protect your health as well as the health of your fellow employees.

Now, we have talked about what your Plant Management is doing to be sure that you have a place to work that is free from recognized hazards from asbestos. Efforts to protect your health will be ineffective unless you try to minimize your own exposure to asbestos. There are

DUP 0903172.

120976

DU 024554

DRAM

EXPOSURE TO ASBESTOS DUST

(Continued)

several things you can do to protect yourself from unacceptable exposure to asbestos dust.

- 1.) Follow established standard practices for handling asbestos.
- 2.) Stop smoking if you do; don't start if you don't. In my opinion, it has been conclusively proven that cigarette smoking is a cause of irritation to the lungs and bronchial tubes and can cause cancer of the lungs and chronic bronchitis and emphysema without any other factors being present.

Significant asbestos exposure also has its primary effect on the lungs, as we will discuss further, so the use of cigarettes and their effects on the lungs could compound any problem that developed.

- 3.) You should wash your hands, arms, and face, before eating to prevent the inhalation of dust that might be on you.
- 4.) Do not eat in shop area where asbestos dust accumulates.
- 5.) Clean off loose dust from clothing before leaving the plant.

- 2 -

DUP 0903173

120977

DU 024555

DRAFT

EXPOSURE TO ASBESTOS DUST

(Continued)

6.) Use and wear all personal protective equipment provided you.

Company standards and protective equipment provided by the Company and designed to protect you from asbestosis and other complications are only as effective as your willingness to do everything possible to protect yourself.

Asbestos has been known to man since the stone ages, having been used to make pottery. The term asbestos comes from the Greek word that means inextinguishable or unquenchable and is a term that embraces a number of fibrous silicate minerals found throughout the world. There are 3 major types of asbestos; the type being determined by the fiber length and physical properties. Chrysotile asbestos accounts for 95% of the asbestos production and is the type most used in the United States. Crocidolite and Amosite are the other two major types. Asbestos is the only mineral that can be woven into cloth. Commercial mining of asbestos began in the late 1870's and it is estimated that there are about 3000 different industrial applications for asbestos in the United States today. This ranges from insulation material to brake shoes, in cement, floor tile, and many other uses.

Now let's talk a few minutes about asbestosis. The recognition of asbestosis as a disease goes back to the first century when the Greek Geographer, Strabo, and the Roman

- 3 -

DUP 0903174

120978

DU 024556

DRAFT

EXPOSURE TO ASBESTOS DUST

(Continued)

Naturalist, Pliny the Elder, each mentioned a sickness of the lungs in slaves whose occupation was to weave asbestos into cloth. The cloth at that time was used in cremation and was described as the "funeral dress of kings". Very little constructive medical information was obtained or pursued about asbestosis following this. The modern knowledge of asbestosis dates to about 1900 when Dr. E. Montague Murray of London described the fibrosis of asbestosis in the lungs of a man who had worked in the carding room of a factory for 14 years. A few years later, an inspector in the Department of Labor of England, described the illness in a group of employees in an asbestos weaving mill. This was the first report of the epidemiology of asbestosis. Unfortunately, neither he nor Dr. Murray realized the full significance of their findings. The first written report of the illness in the modern medical literature was in 1924, when Dr. W. E. Cooke, an English physician, described a case in a woman who developed symptoms after working for 13 years in an asbestos textile mill. He was also first to describe the difference between asbestosis and other pneumoconioses.

Between 1928 and 1929, Dr. E.R.A. Merewether examined 363 asbestos-textile workers and found that slightly more than 25% showed evidence of pulmonary fibrosis with symptoms.

- 4 -

DUP 0903175

120979

DU 024557

DRAFT

EXPOSURE TO ASBESTOS DUST

(Continued)

He also showed that the incidence of fibrosis increased in direct proportion to the number of years of exposure, reaching 81% in the group of workers employed in the industry for 20 years or more. In 1931, as a result of this study, Great Britain passed regulations to control the dust and exposures better than had been done previously. This was the first time that attempts were made to control the exposure of humans to asbestos dust.

On a survey in 1949 in South Africa, a larger mining area for asbestos, an American physician found children developing asbestosis before the age of 12. In checking on their activity, he found that they were employed in a mill and their job was to stand in large jute bags and pack the asbestos fiber down with their feet as the material was being put into the bag.

In 1960, 2 physicians working independently, one in Finland, and one in South Africa, discovered asbestos bodies in the lungs of people who lived near asbestos factories but did not work in them. This was the first indication that asbestos could be inhaled without direct occupational exposure. Many studies have been done since this to substantiate the prevalence of asbestos fibers in the atmosphere.

- 3 -

DUP 0903175

120980

DU 024558

DRAFT

EXPOSURE TO ASBESTOS DUST

(Continued)

There is still much work being done on asbestos exposure and asbestosis and much work still needs to be done. Medical authorities and industry are becoming more concerned over exposure to asbestos; and better protection and precautions are being developed.

The exact mechanism of asbestosis is unknown, but we believe that asbestos fibers are inhaled and become trapped in the small air sacs of the lungs. Because of the movement of the bronchial tubes with breathing, the fibers work into the lining of the bronchial tubes. They create a chronic irritation here and the body tries to get rid of the irritation and deposits various materials around the fiber to make it less irritating and to try to absorb it. The body mechanisms are not able to remove the fibers so the conglomerate of asbestos fibers and defense mechanisms make up an asbestos body. The chronic irritation to the lung continues and once the asbestos body is deposited, it is always present. Because of the chronic irritation, scarring and scar tissue develop in the lung and reduce the lung's ability to move air. This makes the lung no longer effective as a means of exchanging oxygen; and then chronic lung disease called asbestosis develops. This usually first shows up as shortness of breath. A dry cough then develops and then production of blood-streaked sputum. An occasional inhaled fiber will not cause significant scarring in the lungs to interfere with their function, but constant inhalation of large numbers

DUP 0903177

120981

DU 024559

DRAFT

EXPOSURE TO ASBESTOS DUST

(Continued)

of fibers would; and that is the basis for the permissible air concentration being established. Asbestosis does not predispose someone to tuberculosis as does silicosis and anthracosis.

Calculations suggest that asbestos workers who smoke have about 90 times the risk of developing lung complications from asbestos exposure as do men who neither smoke nor work with asbestos. This is the reason for recommending stopping smoking if you do, don't start smoking if you don't.

I think that with the program the Company and your Plant Management have started, and with your cooperation and help, there should not be any health hazards related to asbestos through your working lifetime.

Bruce W. Karsh, M.D.

April 4, 1973

BWK/bjd

- 7 -

DUP 0903178

120982

DU 024560