

standard to 500,000 Fibers per cubic meter of air. In 1976, NIOSH recommended lowering the standard to 100,000 fibers.

NIOSH and OSHA established an Asbestos Work Group in the fall of 1979. Its charge was to review all health effects data since the 1976 NIOSH recommended standard was transmitted to OSHA and to determine whether any changes to that recommendation were warranted. Was there health damage occurring at the level of the current standard? The Work Group consisted of four experts from NIOSH and three from OSHA.

The Work Group has completed its report and NIOSH has formally recommended that OSHA adopt the Work Group's findings. A copy of the report is in your press kit.

The Group reconfirmed that there is no safe exposure level for asbestos. Although data suggest that lower exposures result in lower risks of developing cancer, there is no known level below which asbestos-related diseases do not occur.

The Group reconfirmed that all commercial and several non-commercial forms of asbestos cause disease. There has at times been debate over which types of asbestos fibers cause disease. The Group found strong evidence that all forms are dangerous, including chrysotile; crocidolite; and fibrous cummingtonite-grunerite including amosite, fibrous tremolite, fibrous actinolite, and fibrous anthophyllite. The Work Group found no basis for regulating one type of fiber and not another.

With these two premises in mind, let me enumerate some of the specific findings of the Work Group:

- o Optical microscopy continues to be the most reliable and economically feasible method for determining airborne levels of

-more-