

- on hepatic drug-metabolizing enzymes and porphyria related to contamination with chlorinated dibenzo-p-dioxins and dibenzofurans. *Biochem Pharm* 1977; 26:1549-1557.
66. Calvert GM, Sweeney MH, Fingerhut MA, Hornung RW, and Halperin WE. Evaluation of porphyria cutanea tarda in U.S. workers exposed to 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Am J Industrial Med* 1994; 25:559-571.
 65. Doss MO and Colombi AM. Chronic hepatic porphyria induced by chemicals: the example of dioxin. In *Occupational and Environmental Chemical Hazards*, Fao V, Emmett EA, Maroni M, and Colombi A. Ellis Horwood 1987.
 67. Van Bergelin APJM, Fase KM, Van der Kolk J, Poiger H, Brouwer A, Seinen W, and Van den Berg M. Synergistic effect of 2,2',4,4',5,5'-Hexachlorobiphenyl and 2,3,6,7-tetrachlorobibenzodioxin on hepatic prothyrin levels in female Sprague-Dawley rats. In *Organohalogen Compounds* 25: 105-110, 1995.
 68. Noweir MH and Pfitzer EA. Evaluation of Coproporphyrin in urine from workers exposed to lead. *Amer Indust Hyg Assoc J* 1970; x:492-500
 69. Bombick DW, Matsumura G, Madhukar BV. Low density lipoprotein receptor activities in the hepatic plasma membrane of the guinea pig and rat. *Biochem Biophys Res Comm* 1984; 118:548-554.
 70. Brewster DW and Matsumura F. 2,3,7,8-TCDD reduces lipoprotein lipase activity in the adipose tissue of the guinea pig. *Biochem Biophys Res Comm* 1984; 122:810-817.
 71. Haring Sweeney M, Fingerhut MA, Arezzo JC, Hornung RW, and Connally B. Peripheral Neuropathy After Occupational Exposusre to 2,3,7,8-TCDD. *Am J Industrial Med* 1993; 23:845-858.
 72. Klawans HL. Dystonia and Tremor Following Exposure to 2,3,7,8-Tetrachlorodibenzo-p-dioxin. *Movement Disorders* 1987; 2:255-261.
 73. Fingerhut MA, Haring-Sweeney M, Patterson DG, Piacitelli LA, Morris JA, Marolow DA, Hornung RW, Cameron LW, Connally LB, Needam LL, and Halperin WE. Levels of 2,3,7,8-TCDD in the