

**ATTACHMENT TO LETTER TO C. AUER DATED MAY 4, 2000: PUBLISHED
STUDIES ON PERFLUOROOCETANESULFONATES**

Environmental Studies

Key, B.L.; Howell, R.D.; Criddle, C.S. "Fluorinated Organics in the Biosphere," *Environ. Sci. Technol.*, 1997, 31: 2445-2454

Key, B.L.; Howell, R.D.; Criddle, C.S. "Defluorination of Organofluorine Sulfur Compounds by *Pseudomonas* Sp. Strain D2," *Environ. Sci. Technol.*, 1998, 32:2283-2287

Toxicology Studies

Belisle J, Hagan DF (1978). *Anal Biochem.* 87, 545 (Note Error: In this report the blank was erroneously reported as 0.02 mg; it should be 0.02µg.)

Belisle, J (1981). *Science* 212, pp. 1509-1510.

Guy, WS (1972). Ph.D. Thesis, University of Rochester, Rochester, NY.

Guy, WS, Taves, DR, Brey, Jr., WS (1976). Organic fluorocompounds in human plasma: prevalence and characterization. (In) *Biochemistry Involving Carbon-Fluorine Bond*, pages 117-134.

Haughom, B. and Spydevold, O. 1992. The mechanism underlying the hypolipemic effect of perfluorooctanoic acid (PFOA), perfluorooctane sulphonic acid (PFOSA) and clofibrilic acid. *Biochim. Biophys. Acta.* 1128, 65-72.

Ikeda, T., Fukuda, K., Mori, I., Enomoto, M., Komai, T. and Suga, T. 1987. Induction of cytochrome P-450 and peroxisome proliferation in rat liver by perfluorinated octanesulfonic acid. In: *Peroxisomes in Biology and Medicine*, H.D. Fahimi and H. Sies, Eds. Springer Verlag, New York, 304-308.

Johnson, J.D., Gibson, S.J. and Ober, R.E. 1984. Cholestamine-enhanced fecal elimination of carbon-14 in rats after administration of ammonium [¹⁴C]perfluorooctanoate or potassium [¹⁴C]perfluorooctanesulfonate. *Fund. Appl. Toxicol.* 4, 972-976.

Nabbefeld D., Butenhoff J., Bass N. and Seacat A. 1998. Displacement of a fluorescently labeled fatty acid analogue from fatty acid carrier proteins by wyeth-14,643, ammonium perfluorooctanoate, potassium perfluorooctane sulfonate and other known peroxisome proliferators. (SOT Abstract. Accepted, *Toxicologist* 1998).

Olsen, GW, Burris, JM, Mandel, JH and Zobel, LR (1999). Serum perfluorooctane sulfonate and hepatic and lipid clinical chemistry tests in fluorochemical production employees. *JOEM* 41:799-805.

Pothapragada V, (1975). Determination of total fluorine in serum and other biological materials by oxygen bomb and reverse extraction techniques. *Analytical Biochem* 68:512-521.

Pothapragada V, Singer R, Armstrong WD (1971). Determination of ionic (plus ionizable) fluoride in biological fluids. Procedure based on adsorption of fluoride ion on calcium phosphate. *Anal Biochem* 42:350-359.

Singer L and Armstrong WD (1959). Determination of fluoride in blood serum. *Analytical Chem* 31:105-109..

Singer L and Ophaug RH (1979). Concentrations of ionic, total, and bound fluoride in plasma. *Clin Chem* 25:523-525.

Sohlenius, A-K., Eriksson, A.M., Högström, C., Kimland, M. and DePierre, J.W. 1993. Perfluorooctane sulfonic acid is a potent inducer of peroxisomal fatty acid B-oxidation and other activities known to be affected by peroxisome proliferators in mouse liver. *Pharmacol. Toxicol.* 72, 90-93.

Taves D (1968a). Evidence that there are two forms of fluoride in human serum. *Nature* 217:1050-1051.

Taves D (1968b). Electrophoretic mobility of serum fluoride. *Nature* 220:582-583.

Taves D, Guy W, Brey W (1976) . Organic fluorocarbons in human plasma: Prevalence and characterization. In: Filler R, eds. *Biochemistry Involving Carbon-Fluorine Bonds*. Washington, DC:American Chemical Society, pages 117-134.

Ubel FA, Sorenson SD, Roach DE (1980). Health status of plant workers exposed to fluorochemicals, a preliminary report. *Am Ind Hyg Assoc J.* 41:584-589.

Yamamoto G, Yoshitake K, Sato T, Kimura T and Ando T (1989). Distribution and forms of fluorine in whole blood of human male. *Analytical Biochem* 182:371-376.