

FILE NAME: Reynolds Metals (RM)

DATE: 1966 Dec 21

DOC#: RM102

DOCUMENT DESCRIPTION: Memo - A Reminder on Marinite

1966 most likely
cites week old report

(filed at end of RMZ)

Marinite

INTERDEPARTMENTAL CORRESPONDENCE

FORM 2 58

FROM Ed Largent	AT Richmond	TO	AT
COPIES TO			
SUBJECT A REMINDER ON MARINITE			DATE December 21, 1966

Dust from Marinite should be treated with great caution (see Industrial Hygiene Bulletin No. 7, Asbestos, September, 1963). The indictment of asbestos as a cancer producing chemical continues to grow. Evidence of this can be seen in the Asbestos (December 15, 1966) shown below.

CANCER-ASBESTOS LINK REPORTED BY PUBLIC HEALTH SERVICE:

Exposure to high levels of asbestos dust 20 to 30 years ago is probably responsible for a 21 percent increase in the death rate among certain workers in three industries using asbestos, according to a study reported by the Public Health Service to a Conference on Asbestos Research in Cincinnati. The excess in deaths above the normal rate was due to cancer of the lung and gastrointestinal tract and to asbestosis. The death rates were much lower in other industries using asbestos where dust levels were relatively even, even though there was long-term exposure.

"While dust levels have been sharply reduced in the asbestos textile industry, where the excess mortality was found, the finding adds to the growing evidence that asbestos exposure is associated with cancer of the lung," said Dr. Murray C. Brown, chief of the USPHS Division of Occupational Health.

The report was based on research by Dr. Philip Enterline of McGill University and Dr. Mildred Kendrick of USPHS. Their data was derived from Social Security Administration records for 21,755 men who had worked in the three industries and from death certificates.

The conferees agreed on the necessity for a much more concentrated attack on asbestos as an environmental contaminant. High on the list of research priorities were studies of population groups with both heavy and light exposures to asbestos fibers. It was pointed out that it is still not known whether asbestos mineral fibers are vehicles or carriers of carcinogenic chemicals or are active themselves.

The group stressed the need for expanded laboratory animal studies on the effects of different types of asbestos dusts, alone and in combination with various metals and hydrocarbons.

000011

A REMINDER ON MARINITE

V. G. KNEESKERN	EXO
D. W. JOHNSON	LONGVIEW
R. E. COLE	LISTERHILL RED.
H. C. WILLMS	PATTERSON
W. E. CAMPBELL	TROUTDALE
D. E. THRASHER	JONES MILLS
D. P. KELLER	SAN PATRICIO
G. F. BRADLEY	MASSENA
R. H. HENKE	EXO
R. B. NEWMAN	MCCOOK
D. H. HIPP	LISTERHILL ALLOYS
L. H. GRAVES, JR.	EXO
E. G. DREWRY	PHOENIX EXT.
C. C. HUDGINS	GRAND RAPIDS
A. M. MURPHY	BELLWOOD EXT.

(Plant Managers)

Dr. J. M. MacMillan	EXO
---------------------	-----



000010