YOUNGER LABORATORIES

Biochemists . . . Pharmacologists . . . Analysts

5620 TELEGRAPH ROAD
ST. LOUIS 29, MISSOURI

PHONE: TILDEN 6-2340

Certificate of Analysis

C+2AM

June 25th, 1962

SUBJECT -

Toxicological Investigation Of: Aroclor 1262
Monsanto Sample Number 40
Monsanto Project Number Y-62-38

STUDY CONDUCTED FOR -

Monsanto Chemical Company, St. Louis, Missouri

EXPERIMENTAL PROCEDURE -

A) Oral LD_{50} (Rats)

The diluted compound was fed by stomach tube to Sprague-Dawley strain albino rats.

After the approximate Minimum Lethal Dose was determined, groups of male and female rats were fed in increasing doses at increments of 0.1 fractional log intervals at four levels designed to blanket the toxicity range thereby supplying data for calculation of the LD_{50} which was done according to a modification of the method of E. J. de Beer.

Observations were made for toxic symptoms and the viscera of the animals that succumbed were examined macroscopically.

The data, together with the dilution at which the compound was fed, are shown in Table I.

B) Skin Absorption MLD (Rabbits)

The diluted compound was applied in increasing doses at increments of 0.2 fractional log intervals to the closely clipped, intact skin of New Zealand white female rabbits.

The treated areas were covered with plastic strips and the animals placed in wooden stocks for periods up to twenty-four hours, after which time they were assign to individual cages.

Observations were made for toxic symptoms and the viscera of the animals that succumbed were examined macroscopically.

The data, together with the dilution at which the compound was applied, are shown in Table II.

GBRN002553

This confidential report is not to be used in any form o[illegible] [illegible]ermission

EXHIBIT
K-31

COPY

To: Monsanto Chemical Company
St. Louis, Missouri
Younger Laboratories Certificate of Analysis - Page 2 (6/25/62) - Y-62-38

SUMMARY -

Aroclor 1262

A) Oral LD_{50} (Rats)

The Oral LD_{50} for male and female rats was placed at 11,300 milligrams per kilogram with lower and upper limits of 9940 to 12,680 milligrams per kilogram.

The compound was classed as practically non-toxic by oral ingestion in male and female rats.

B) Skin Absorption MLD (Rabbits)

The Minimum Lethal Dose by skin absorption in female rabbits was found to be greater than 1260 milligrams per kilogram and less than 3160 milligrams per kilogram.
There was one death and one survival at 2000 milligrams per kilogram.

The compound was classed as mildly toxic by skin absorption in female rabbits.

YOUNGER LABORATORIES

Fred M. Younger

BY: FRED M. YOUNGER

The material in this report is to be used in development of the product and may be given to responsible sales contacts, but it is not to be used by them in advertising copy. The source of this material is not to be divulged until it appears in formal publications. No exceptions to the established rule may be made without the approval of the Medical Department in St. Louis. Customers' inquiries regarding matters of toxicity are to be referred as before to the Medical Department in St. Louis for reply.

— Monsanto Chemical Company

GBRN002554

LEXOLDMON002174

TABLE I

THE ORAL LD_{50} OF 'Aroclor 1262' FOR RATS

Sample Fed As A 50.0% Solution In Corn Oil

Animal No. - Sex	Weight Gm.	Dose M./Kg.	Fate
1- Female	230	7,940	Survived
2- Female	215	7,940	Survived
3- Female	220	7,940	Survived
4- Male	235	7,940	Survived
5- Male	250	7,940	Survived
6- Male	230	10,000	Died
7- Female	215	10,000	Survived
8- Female	225	10,000	Died
9- Female	220	10,000	Survived
10- Male	230	10,000	Survived
11- Male	245	12,600	Died
12- Male	225	12,600	Died
13- Female	210	12,600	Survived
14- Female	220	12,600	Died
15- Female	225	12,600	Survived
16- Male	235	15,800	Died
17- Male	245	15,800	Died
18- Male	240	15,800	Died
19- Female	215	15,800	Died
20- Male	230	15,800	Died

DISCUSSION -

The Oral LD_{50} for male and female rats was placed at 11,300 milligrams per kilogram with lower and upper limits of 9940 to 12,880 milligrams per kilogram.

The compound was classed as practically non-toxic by oral ingestion in male and female rats.

Survival time was one to six days with most deaths occurring in three days.

Toxic symptoms included severe diarrhea, loss of appetite, lethargy, and increasing weakness.

At autopsy there was inflammation of the gastric mucosa, tissue edema, and liver congestion macroscopically.

GBRN002555

LEXOLDMON002175

TABLE II

THE MINIMUM LETHAL DOSE OF 'Aroclor 1262' BY SKIN ABSORPTION IN RABBITS

Sample Applied As A 50.0% Solution In Corn Oil

Animal No. - Sex	Weight Kg.	Dose Mg./Kg.	Weight Change 5 Days Later Kg.	Fate
1 - Female	2.2	794	- 0.1	Survived
2 - Female	2.7	1260	- 0.2	Survived
3 - Female	2.5	2000	- 0.3	Died — 7 Days
4 - Female	2.6	2000	- 0.2	Survived
5 - Female	2.4	3160	- 0.5	Died — 8 Days
6 - Female	2.7	5010	- 0.3	Died — 5 Days

DISCUSSION -

The Minimum Lethal Dose by Skin Absorption in female rabbits was found to be greater than 1260 milligrams per kilogram and less than 3160 milligrams per kilogram. There was one death and one survival at 2000 milligrams per kilogram.

The compound was classed as mildly toxic by skin absorption in female rabbits.

Survival time was five to eight days.

Toxic symptoms included poor appetite with resultant loss of weight, nervousness, increasing weakness, and sudden collapse.

At autopsy there was pulmonary hyperemia, liver discoloration, and renal congestion.

GBRN002556

LEXOLDMON002176

HAZLETON LABORATORIES
Biochemists... Pharmacologists... Analysts
PHONE TILDEN 6-2540

Certificate of Analysis

June 25th, 1962

SUBJECT -

Toxicological Investigation Of: Aroclor 1254

Monsanto Sample Number 38
Monsanto Project Number Y-62-36

STUDY CONDUCTED FOR -

Monsanto Chemical Company, St. Louis, Missouri

Y-62-36
6/25/62

EXPERIMENTAL PROCEDURE -

A) Oral LD_{50} (Rats)

The undiluted compound was fed by stomach tube to Sprague-Dawley strain albino rats.

After the approximate Median Lethal Dose was determined, groups of male and female rats were fed in increasing doses at increments of 0.1 fractional log intervals at four levels designed to blanket the toxicity range thereby supplying data for calculation of the LD_{50} which was done according to a modification of the method of L. J. de Beer.

Observations were made for toxic symptoms and the viscera of the animals that succumbed were examined macroscopically.

The data are shown in Table I.

B) Skin Absorption LD_{50} (Rabbits)

The undiluted compound was applied in increasing doses at increments of 0.2 fractional log intervals to the closely clipped, intact skin of New Zealand white female rabbits.

The treated areas were covered with plastic strips and the animals placed in metal stocks for periods up to twenty-four hours, after which time they were returned to individual cages.

Observations were made for toxic symptoms and the viscera of the animals that [illegible] were examined macroscopically.

The data are shown in Table II.

EXHIBIT
K-32

This confidential report is not to be used in any form of advertising without written permission

SUMMARY -

Aroclor 1254

A) Oral LD_{50} (Rats)

The Oral LD_{50} for male and female rats was placed at 11,900 milligrams per kilogram with lower and upper limits of 10,475 to 13,450 milligrams per kilogram.

The compound was classed as practically non-toxic by oral ingestion in male and female rats.

B) Skin Absorption MLD (Rabbits)

The Minimum Lethal Dose by Skin Absorption in female rabbits was found to be greater than 1260 milligrams per kilogram and less than 2000 milligrams per kilogram.

The compound was classed as mildly toxic by skin absorption in female rabbits.

YOUNGER LABORATORIES

Fred M. Younger

BY: FRED M. YOUNGER

The material in this report is to be used in development of the product and may be given to responsible sales contacts, but it is not to be used by them in advertising copy. The [illegible] of this material is not to be divulged until it appears in [illegible] publications. No exceptions to the established rule [illegible] without the approval of the Medical Department [illegible] St. Louis. Customers' inquiries regarding matters of [illegible] are to be referred as before to the Medical Department [illegible] St. Louis for reply.

— Monsanto Chemical Company

TABLE I

THE ORAL LD_{50} OF 'Aroclor 1254' FOR RATS

Sample Fed Undiluted

Animal No. - Sex	Weight Gm.	Dose Mg./Kg.	Fate
1- Female	210	7,940	Survived
2- Female	200	7,940	Survived
3- Female	215	7,940	Survived
4- Male	230	7,940	Survived
5- Male	225	7,940	Survived
6- Male	240	10,000	Died
7- Female	215	10,000	Survived
8- Female	220	10,000	Died
9- Female	230	10,000	Survived
10- Male	245	10,000	Survived
11- Male	250	12,600	Survived
12- Male	235	12,600	Died
13- Female	215	12,600	Died
14- Female	225	12,600	Survived
15- Female	230	12,600	Died
16- Male	250	15,800	Died
17- Male	245	15,800	Survived
18- Male	230	15,800	Died
19- Female	215	15,800	Died
20- Male	240	15,800	Died

DISCUSSION -

The Oral LD_{50} for male and female rats was placed at 11,900 milligrams per kilogram with lower and upper limits of 10,475 to 13,450 milligrams per kilogram.

The compound was classed as practically non-toxic by oral ingestion in male and female rats.

Survival time was two to six days with most deaths occurring in three days.

Toxic symptoms included severe diarrhea, loss of appetite, lethargy, salivation, and increasing weakness.

At autopsy there was inflammation of the gastric mucosa, pulmonary hyperemia, and liver discoloration macroscopically.

LEXOLDMON002179

TABLE II

THE MINIMUM LETHAL DOSE OF 'Aroclor 1254' BY SKIN ABSORPTION IN RABBITS

Sample Applied Undiluted

Animal No. - Sex	Weight Kg.	Dose Mg./Kg.	Weight Change 5 Days Later Kg.	Fate
1 - Female	2.4	501	0.0	Survived
2 - Female	2.5	794	0.0	Survived
3 - Female	3.0	1260	- 0.2	Survived
4 - Female	2.7	2000	- 0.4	Died — 8 Days
5 - Female	2.5	3160	- 0.5	Died — 7 Days
6 - Female	2.6	5010	———	Died — 4 Days

DISCUSSION -

The Minimum Lethal Dose by Skin Absorption in female rabbits was found to be greater than 1260 milligrams per kilogram and less than 2000 milligrams per kilogram.

The compound was classed as mildly toxic by skin absorption in female rabbits.

Survival time was four to eight days.

Toxic symptoms included poor appetite, lethargy after two to three days, marked weight loss and increasing weakness. Animals #1 and #2 were little affected while #3 became moderately weak but was not in danger of collapse.

At autopsy there was nasal congestion and liver discoloration (yellowish) macroscopically

GBRN002560

LEXOLDMON002180

YOUNGER LABORATORIES

Biochemists . . . Pharmacologists . . . Analysts

5620 TELEGRAPH ROAD
ST. LOUIS 29, MISSOURI

PHONE: TILDEN 6-2540

Certificate of Analysis

June 25th, 1962

C+25AA1

SUBJECT -

Toxicological Investigation Of: Aroclor 1260

Monsanto Sample Number 39
Monsanto Project Number Y-62-37

STUDY CONDUCTED FOR -

Monsanto Chemical Company, St. Louis, Missouri

COPY

EXPERIMENTAL PROCEDURE -

A) Oral LD_{50} (Rats)

The diluted compound was fed by stomach tube to Sprague-Dawley strain albino rats.

After the approximate Minimum Lethal Dose was determined, groups of male and female rats were fed in increasing doses at increments of 0.1 fractional log intervals at four levels designed to blanket the toxicity range thereby supplying data for calculation of the LD_{50} which was done according to a modification of the method of E. J. de Beer.

Observations were made for toxic symptoms and the viscera of the animals that succumbed were examined macroscopically.

The data, together with the dilution at which the compound was fed, are shown in Table I.

B) Skin Absorption MLD (Rabbits)

The diluted compound was applied in increasing doses at increments of 0.2 fractional log intervals to the closely clipped, intact skin of New Zealand white female rabbits.

The treated areas were covered with plastic strips and the animals placed in wooden stocks for periods up to twenty-four hours, after which time they were assigned to individual cages.

Observations were made for toxic symptoms and the viscera of the animals that succumbed were examined macroscopically.

The data, together with the dilution at which the compound was applied, are shown in Table II.

GBRN002561

This confidential report is not to be used in any form of c

EXHIBIT
K-33

LEXOLDMON002181

SUMMARY -

Aroclor 1260

A) Oral LD_{50} (Rats)

The Oral LD_{50} for male and female rats was placed at 10,000 milligrams per kilogram with lower and upper limits of 8,600 to 11,600 milligrams per kilogram.

The compound was classed as practically non-toxic by oral ingestion in male and female rats.

B) Skin Absorption MLD (Rabbits)

The Minimum Lethal Dose by Skin Absorption in female rabbits was found to be greater than 1260 milligrams per kilogram and less than 2000 milligrams per kilogram.

The compound was classed as mildly toxic by skin absorption in female rabbits.

YOUNGER LABORATORIES

BY: FRED M. YOUNGER

The material in this report is to be used in development of the product and may be given to responsible sales contacts, but it is not to be used by them in advertising copy. The source of this material is not to be divulged until it appears in formal publications. No exceptions to the established rule may be made without the approval of the Medical Department in St. Louis. Customers' inquiries regarding matters of toxicity are to be referred as before to the Medical Department in St. Louis for reply.

— Monsanto Chemical Company

GBRN002562

LEXOLDMON002182

TABLE I

THE ORAL LD_{50} OF 'Aroclor 1260' FOR RATS

Sample Fed As A 50.0% Solution In Corn Oil

Animal No. - Sex	Weight Gm.	Dose Mg./Kg.	Fate
1- Female	215	6,310	Survived
2- Female	230	6,310	Survived
3- Female	220	6,310	Survived
4- Male	245	6,310	Survived
5- Male	235	6,310	Survived
6- Male	250	7,940	Survived
7- Female	220	7,940	Survived
8- Female	210	7,940	Survived
9- Female	215	7,940	Died
10- Male	240	7,940	Survived
11- Male	225	10,000	Died
12- Male	235	10,000	Survived
13- Female	210	10,000	Survived
14- Female	225	10,000	Died
15- Female	210	10,000	Died
16- Male	230	10,000	Survived
17- Male	235	12,600	Died
18- Male	220	12,600	Died
19- Female	210	12,600	Survived
20- Female	215	12,600	Died
21- Male	240	12,600	Died

DISCUSSION -

The Oral LD_{50} for male and female rats was placed at 10,000 milligrams per kilogram with lower and upper limits of 8,600 to 11,600 milligrams per kilogram.

The compound was classed as practically non-toxic by oral ingestion in male and female rats.

Survival time was two to five days with most deaths occurring in three days.

Toxic symptoms included severe diarrhea, loss of appetite, and increasing lethargy and weakness.

At autopsy there was inflammation of the gastric mucosa and liver hyperemia macroscopically.

LEXOLDMON002183

T A B L E II

THE MINIMUM LETHAL DOSE OF 'Aroclor 1260' BY SKIN ABSORPTION IN RABBITS

Sample Applied As A 50.0% Solution In Corn Oil

Animal No. - Sex	Weight Kg.	Dose Mg./Kg.	Weight Change 5 Days Later Kg.	Fate
1 - Female	1.9	501	+ 0.1	Survived
2 - Female	2.2	794	0.0	Survived
3 - Female	2.4	1260	0.0	Survived
4 - Female	2.6	2000	- 0.4	Died — 8 Days
5 - Female	2.4	2000	- 0.4	Died — 7 Days
6 - Female	2.4	3160	- 0.5	Died — 7 Days

DISCUSSION -

The Minimum Lethal Dose by Skin Absorption in female rabbits was found to be greater than 1260 milligrams per kilogram and less than 2000 milligrams per kilogram.

The compound was classed as mildly toxic by skin absorption in female rabbits.

Survival time was seven to eight days.

Toxic symptoms included weight loss after two to three days with increasing weakness followed by collapse in six to seven days.

At autopsy there was evidence of liver dysfunction and renal hyperemia macroscopically.

GBRN002564

LEXOLDMON002184