

RECEIVED

MAR 04 1985

S. A. RASMUSSEN

#222-JN

The Activity of Environmental Samples in a Cell Culture Test for Asbestos Toxicity¹

BETTI REISS,* JAMES R. MILLETTE,† AND GARY M. WILLIAMS*

*Naylor Dana Institute for Disease Prevention, American Health Foundation, 1 Dana Road, Valhalla, New York 10595, and †Field Studies Division, Health Effects Research Laboratory, Cincinnati, Ohio 45268

Received January 15, 1980

The inhibition of colony-forming efficiency of cultured human embryonic intestine-derived epithelial (I-407) cells was utilized in order to assay the toxic potential of six coded samples of particulate matter provided by the United States Environmental Protection Agency (EPA). The results of the assay indicated that the most toxic of the EPA samples tested "blindly" was amosite which was equal in toxicity to the amosite used, by chance, as a positive control. The toxicities of the particulates from drinking water were approximately 100-fold less than the amosite; of these, the order of toxicity of the samples was San Francisco > Seattle > Duluth. The samples of attapulgite clay and taconite tailings displayed approximately half the toxicity of the drinking water particulates. These results indicate that this assay provides a sensitive and accurate method for screening asbestos and asbestiform contamination for potential toxicity.

INTRODUCTION

Occupational exposure to airborne asbestos fibers has been implicated in the production of pulmonary asbestosis, lung cancer, and pleural and peritoneal mesotheliomas (Selikoff, *et al.*, 1964, 1972; McDonald *et al.*, 1971; Enterline *et al.*, 1972). Data are also suggestive of a relationship between exposure to asbestos and the development of various forms of gastrointestinal cancer as a result of swallowing inhaled fibers or ingesting food and beverages containing asbestos fibers (Hammond *et al.*, 1965; Selikoff *et al.*, 1974). Since asbestos fibers penetrate the mucosal lining of the gastrointestinal tract, they may also lead to the production of tumors in remote organs (Westlake *et al.*, 1965; Storeygard and Brown, 1977; Cook and Olson, 1979).

These possibilities have caused concern about the presence of asbestos fibers in municipal water supplies in this country (Masson *et al.*, 1974; Levy *et al.*, 1976) and in Canada (Cunningham and Pontefract, 1971). The discovery in 1973 of asbestos fibers in Lake Superior, the source of municipal water for Duluth, Minnesota, prompted investigation of Duluth's drinking water (Cook *et al.*, 1976; Kramer, 1976). Studies revealed the presence of 600 million amphibole asbestiform fibers per liter in samples of Duluth tap water (Cook *et al.*, 1976). Contamination of the water supply with asbestos apparently began in 1955 as a result of the dumping of taconite mine tailings containing amphibole impurities into the lake

¹This work was supported by U.S. Environmental Protection Agency Grant R803998-01; parts of this study are available in Research Report EPA-600/1-79-023 (B. Reiss, J. H. Weisburger, and G. M. Williams, "Asbestos and Gastrointestinal Cancer: Cell Culture Studies").

A02507

(United States of America vs Reserve Mining Company, 5-22 Civil 19). Although, to date, surveys have demonstrated no consistent increase in cancer incidence (Masson *et al.*, 1974; Levy *et al.*, 1976) or mortality (Wigle, 1977) that can be specifically attributed to exposure to asbestos in drinking water, with the exception of one study which did indicate an increased incidence (McCabe and Millette, 1979), ongoing cancer surveillance in regions with high levels of asbestos fiber contamination in the drinking water has been initiated. Furthermore, it should be borne in mind that among persons occupationally exposed to asbestos dust, the increases in death rates from cancer of the stomach, colon, and rectum do not reach measurable proportions until many years after the onset of exposure. Thus, the lapse of time since Duluth water was first heavily polluted with mineral fibers is not yet sufficient to have produced a significant increase in death rates from these cancers, particularly if the risk for residents of Duluth is not as great as for workers occupationally exposed but still of important proportions. If increased risk exists, it will become apparent within another 5 to 15 years (National Research Council, 1977).

In order to study the cellular effects of asbestos, we have developed a cytotoxicity assay that measures the inhibition of colony-forming efficiency of cultured human embryonic intestine-derived (I-407), adult rat liver-derived epithelial (ARL-6), and mouse colon-derived epithelial-like (MCE-1) cells (Reiss *et al.*, 1980). Using this assay we were able to quantitate the cytotoxic effects of the chrysotile and amphibole (amosite and crocidolite) forms of asbestos. The I-407 line was found to be the most sensitive of these cell lines to the toxic effects of all forms of Union Internationale Contre Le Cancer (UICC) asbestos (Timbrell and Rendall, 1971-1972) that were tested. This cytotoxicity assay was utilized in the present investigation to determine "in a blind fashion" the toxicity of six coded samples of particulate matter provided by the EPA. Among these samples were particulates extracted by filtration from the drinking water of three large American cities, San Francisco, Seattle, and Duluth (the Duluth sample was obtained prior to the installation of a filtration plant), as well as taconite tailings, attapulgite clay, and the amosite form of asbestos.

The results of the assay indicated that the most toxic of the EPA samples tested "blindly" was the amosite which was equal in toxicity to the amosite used, by chance, as a positive control. The toxicities of the particulates from drinking water were approximately 100-fold less than the amosite control; of these, the order of toxicity of the samples was San Francisco > Seattle > Duluth. The toxicities of the attapulgite clay and the taconite tailings were approximately half that of the drinking water particulates. Thus, the test displayed a wide range in sensitivity to particulates of different composition, and, of these, the asbestos sample was the most toxic.

MATERIALS AND METHODS

Samples of Particulates

Six samples of particulates were provided by the EPA. At the time of testing, the identity of the samples was unknown; however, following the completion of the assays, the EPA supplied us with the following descriptions:

ASBI

- A. *Sample 1.* Prior to the amphibole c line fraction
- B. *Sample 2.* P. No filtration chrysotile a
- C. *Sample 3.* I water. The s analysis is s
- D. *Sample 4.* S which had b phibole fiber
- E. *Sample 5.* S eral which c chrysotile as from the mi checked by some water generally les bundles in di
- F. *Sample 8.* A ingested asbe tute of Envir Administrat

The UICC reference a 50% decrease in th 0.025 g/liter also pro cytotoxicity assay.

Before weighing, tl exposed to 130 erg/m contamination in a m chemical properties o inaccuracies in weig

ANALYSIS BY EI

Sample

Duluth
San Francisco
Seattle
Attapulgite

Single crystal shapes an Circular/1977 IC-8751.

A02508

UCC 017896

22 Civil 19). Although, se in cancer incidence (gle, 1977) that can be water, with the excep- (McCabe and Millette, evels of asbestos fiber rthermore, it should be d to asbestos dust, the on, and rectum do not set of exposure. Thus, ted with mineral fibers se in death rates from th is not as great as for oportions. If increased 15 years (National Re

we have developed a r-forming efficiency of dult rat liver-derived e (MCE-1) cells (Reiss he cytotoxic effects of s of asbestos. The 1-40th o the toxic effects of all asbestos (Timbrell and say was utilized in the e toxicity of six coded ng these samples were of three large American ple was obtained prior ilings, attapulgitic clay.

he EPA samples tested o the amosite used, by es from drinking water ; of these, the order of uluth. The toxicities of mately half that of the : range in sensitivity to bestos sample was the

At the time of testing, ving the completion of ptions:

- A. *Sample 1.* Particulates filtered from Duluth, Minnesota drinking water prior to the installation of the filtration plant at Duluth. Known to contain amphibole crystals, red clay, and biological fragments. From the crystalline fraction, the particle analysis is shown in Table 1.
- B. *Sample 2.* Particulates filtered from Seattle, Washington drinking water. No filtration is used in water treatment. The sample is known to contain chrysotile asbestos. The particle analysis is shown in Table 1.
- C. *Sample 3.* Particulates filtered from San Francisco, California drinking water. The sample is known to contain chrysotile asbestos. The particle analysis is shown in Table 1.
- D. *Sample 4.* Sample of the less than 2- μ m-size fraction of taconite tailings which had been prepared by a sedimentation separation procedure. Amphibole fibers were identified among the particulates in sample 4.
- E. *Sample 5.* Sample of attapulgitic (palygorskite) clay, a nonasbestos mineral which consisted of fibers of the same or smaller diameter range as chrysotile asbestos. The attapulgitic was obtained in relatively pure form from the mine in Attapulgis, Georgia. The purity of the sample was checked by X-ray diffraction. Attapulgitic fibers have been identified in some water supplies in Georgia and Florida. The attapulgitic fibers are generally less than 2 μ m in length and tend to clump and form small bundles in drinking water.
- F. *Sample 8.* Amosite fibers currently being used to study the effects of ingested asbestos on rats and hamsters sponsored by The National Institute of Environmental Health Sciences (NIEHS), The Food and Drug Administration (FDA), and the EPA (Moore, 1978).

The UICC reference sample of amosite at a CD_{50} (cytotoxic dose that produced a 50% decrease in the number of colonies formed in the cytotoxicity assay) of 0.025 g/liter also provided by the EPA was used as a positive control for the cytotoxicity assay.

Before weighing, the coded samples and the known sample of amosite were exposed to 130 erg/mm²/sec ultraviolet light for 2 hr in order to suppress viable contamination in a manner that would be least likely to alter the physical and chemical properties of the particulates (Reiss *et al.*, 1980). In order to eliminate inaccuracies in weighing caused by the electrostatic character of asbestos,

TABLE 1
ANALYSIS BY ELECTRON MICROSCOPY OF THE CRYSTALLINE FRACTION OF THE
SAMPLES OF PARTICULATES

Sample	Fibers* (%)	Equants (%)	Aggregates (%)	Acciculars (%)
Duluth	4.2	61.1	9.3	0.9
San Francisco	7.7	37.4	18.1	0.0
Seattle	2.0	80.9	1.5	0.0
Attapulgitic	100	Trace		

* Single crystal shapes and aggregate classification as described in the Bureau of Mines Information Circular/1977 IC-8751.

A02509

UCC 017897

aliquots of the coded samples or amosite were weighed in a known weight of water or culture medium immediately prior to use.

Cell Cultures

I-407 cells were used for the cytotoxicity assay. These cells were maintained at 37°C in Williams' Medium E (Williams and Gunn, 1974; Flow Laboratories, Rockville, Md.) supplemented with 10% fetal bovine serum (Flow Laboratories) and containing 50.0 units/ml mycostatin (Gibco, Grand Island, N.Y.) and 100 µg/ml gentamycin (Schering, Kenilworth, N.J.).

Cytotoxicity Assay

The cytotoxicity of the coded samples was quantified as described previously (Reiss *et al.*, 1980) by measuring the inhibition of I-407 epithelial cell colony formation following exposure. For this assay, 10 I-407 cells/cm² were inoculated into 25-cm² culture flasks. Twenty-four hours after the cells were seeded, the medium was replaced with medium containing an aliquot of the sample. Following 3 days of exposure, the cells were washed twice and reincubated in fresh medium. Between 1 and 2 weeks after the initiation of treatment, the cells were fixed in 10% formalin and stained with Giemsa for the determination of colony formation.

RESULTS

The dose-response for each unknown was determined over a range of concentrations from 0.001 to 5.0 g/liter. Testing was initiated by adding between 0.01 and 0.05 g/liter of each sample to the cultures. Depending upon the level of toxicity that the samples exhibited in these early experiments, a range of concentrations was chosen at quarter-log intervals in order to obtain several values of toxicity approximating a CD₅₀ for each sample.

The results of the cytotoxicity assay (Table 2) demonstrated that by assaying the inhibition of I-407 cell colony formation, the cytotoxic levels of samples coded only by reference number could be easily and reliably evaluated. The order of toxicity of these samples, as determined by the assay, was: sample no. 8 > 3 > 2 > 1 ≥ 5 ~ 4. The most toxic sample with a CD₅₀ of approximately 0.01 g/liter was sample 8, amosite. The similar toxicity of this amosite sample to the amosite control indicates the reliability of this assay.

The sample from San Francisco's drinking water (sample 3) was found to have a CD₅₀ between 0.75 and 1.0 g/liter. The CD₅₀ of the Seattle drinking water sample (sample 2) was between 1.0 and 2.5 g/liter and the CD₅₀ of the Duluth drinking water sample (sample 1), taken before installation of a filtration plant, was between 2.5 and 5.0 g/liter. The CD₅₀'s of both the samples of attapulgite clay (sample 5) and taconite tailings (sample 4) were at concentrations greater than 5.0 g/liter. Because of the difficulty of obtaining a homogeneous suspension at high concentrations, the assay was not performed at concentrations above this value.

DISCUSSION

The cytotoxicity assay described previously (Reiss *et al.*, 1980) and applied in this study is a rapid and efficient test which is shown to be useful for the screening

TABLE 2
TOXICITY TO THE I-407 CELL LINE OF SAMPLES OF PARTICULATES

Concentration of	Percentage inhibition of colony formation*				
	Duluth	Seattle	San Francisco	Taconite	Attapulgite

A02510

UCC 017898

known weight of water

cells were maintained at 74; Flow Laboratories, Inc (Flow Laboratories, Island, N.Y.) and 100

as described previously 37 epithelial cell colony cells/cm² were inoculated cells were seeded, the of the sample. Following incubated in fresh medium the cells were fixed in 10 of colony formation

d over a range of concentration adding between 0.01 and upon the level of toxicity a range of concentrations several values of toxicity

onstrated that by assaying ic levels of samples coded / evaluated. The order of was: sample no. 8 3 2 roximately 0.01 g/liter was the sample to the amosite

ple 3) was found to have a ttle drinking water sample D₅₀ of the Duluth drinking a filtration plant, was be amples of attapulgite clay concentrations greater than 50 eneous suspension at high trations above this value

et al., 1980) and applied to be useful for the screening

TABLE 2
TOXICITY TO THE I-407 CELL LINE OF SAMPLES OF PARTICULATES

Concentration of sample tested (g/liter)	Percentage inhibition of colony formation*					
	Amosite control	Duluth drinking water (1) ^a	Seattle drinking water (2)	San Francisco drinking water (3)	Taconite tailings (4)	Attapulgite clay (5)
0.001	—	—	—	—	—	18
0.0025	—	—	—	—	—	—
0.005	—	21	—	—	—	5
0.0075	—	—	—	—	—	—
0.01	—	—	10	—	—	—
0.025	59 ± 15	21	—	—	1	54 ± 13 ^c
0.05	—	0	—	—	0	72 ± 12
0.075	—	—	16	—	1	86
0.1	—	—	9	—	—	—
0.25	—	—	—	—	—	—
0.5	—	6	—	6	3	—
0.75	—	—	49	6	—	—
1.0	—	1	37 ^c	96	2	—
2.5	—	30	73	100	33	—
5.0	—	88 ^c	91	—	41	—

* These values represent the results of a total of seven experiments; many of these values are averages from duplicate experiments. Where more than two experiments are averaged, the standard deviation is included. Each value is the percentage inhibition of CFE of I-407 cells following exposure to water contamination samples. The average CFE for all experiments of I-407 cells not exposed to water contamination samples or amosite was 112.5 ± 41.2. An amosite standard at 0.025 g/liter was tested simultaneously with each experiment; the average percentage inhibition of CFE for all experiments of amosite-exposed cells was 59.2 ± 15.1. Chrysotile at 0.0025 g/liter exhibited approximately a 50% inhibition of CFE for all experiments of

^c Coded numbers used by the EPA to identify the six samples of particulates.

^d A result widely divergent from this value was omitted because it was inconsistent with the trend of increasing concentration of the sample.

AJ2511

UCC 017899

of samples of particulates from drinking water or other sources for the presence of asbestos. Although all of the samples tested were of similar densities, large differences in the toxicities of these samples were found using this assay. Furthermore, these toxicities correlated closely with the asbestos content described in the analysis of the samples provided by the EPA after the completion of the assays. The results of this investigation suggest that asbestos exerts its toxicity through specific physicochemical mechanisms and possesses a cytotoxic potential not found in nonasbestos samples composed primarily of attapulgite clay or taconite tailings.

This bioassay system developed for asbestos can be used as a rapid and accurate method for comparing the relative cytotoxicities of commercial fiber samples with fibers and particulates from relevant environmental samples. Among mineralogists there is considerable debate as to whether the amphibole fibers found in the water of Lake Superior should be called asbestos or asbestiform. Although the basic chemistries and crystal structure of the lake fibers are the same as a commercial form of asbestos, a number of mineralogists propose that there may be some critical differences which would change the hazard potential. Similar questions have been raised about the chrysotile associated with serpentinite quarry samples. The results of our assay suggest that the 4.2% fibers that are in Duluth drinking water (sample 1) have a toxic potential and behave like true asbestos; in contrast, the attapulgite clay containing 100% fibers displayed only minimal toxicity in this assay and is not asbestos-like. Thus, this *in vitro* assay, enables the screening of asbestos- and asbestiform-contaminated environmental samples for hazard potential, an undertaking that would be exceedingly difficult and require prolonged exposure *in vivo*.

ACKNOWLEDGMENTS

The authors wish to thank Mrs. Sondra Solomon for her skilled technical assistance and Mrs. Bette Meyer for preparation of the manuscript. The polygorskite (attapulgite) sample from Attapulgus, Georgia, was provided by Mr. Kenneth Wallingford, Field Investigations Unit, NIOSH, Cincinnati, Ohio. The sample of taconite tailings was provided by Dr. Philip Cook, EPA, Duluth, Minnesota.

REFERENCES

- Cook, P. M., Rubin, I. B., Maggiore, C. J., and Nicholson, W. J. (1976). X-ray diffraction and electron beam analysis of asbestiform minerals in Lake Superior waters. In "Proceedings of the International Conference on Environmental Sensing and Assessment, Las Vegas, Nev.," Vol. 34, pp. 1-9.
- Cook, P. M., and Olson, G. F. (1979). Ingested mineral fibers: elimination in human urine. *Science* 204, 195-198.
- Cunningham, H. M., and Pontefract, R. (1971). Asbestos fibers in beverages and drinking water. *Nature (London)* 232, 332-333.
- Enterline, P., DeCoufle, P., and Henderson, V. (1972). Mortality in relation to occupational exposure in the asbestos industry. *J. Occup. Med.* 14, 897-903.
- Hammond, E. C., Selikoff, I. J., and Churg, J. (1965). Neoplasia among insulation workers in the United States with special references to intra-abdominal neoplasia. *Ann. N.Y. Acad. Sci.* 132, 519-525.
- Kramer, J. R. (1976). Fibrous cummingtonite in Lake Superior. *Canad. Mineral* 14, 91-98.
- Levy, B. S., Sigurdson, E., Mandel, J., Laudon, E., and Pearson, J. (1976). Investigating possible effects of asbestos in city water: surveillance of gastrointestinal cancer incidence in Duluth, Minnesota. *Amer. J. Epidemiol.* 103, 362-368.
- Mason, T. J., McKay, F. (1978). Relation to cancer mortality. In "Proceedings of the International Conference on Environmental Sensing and Assessment, Las Vegas, Nev.," Vol. 34, pp. 1-9.
- McDonald, J. C., McDonald, J. C. (1978). Mortality in the chrysotile asbestos industry. In "Proceedings of the International Conference on Environmental Sensing and Assessment, Las Vegas, Nev.," Vol. 34, pp. 1-9.
- Reiss, B., Solomon, S., V. (1979). Different forms of asbestos. *Environ. Health* 25, 11.
- Selikoff, I. J., Churg, J., Hammond, E. C. (1978). Mortality in the asbestos industry. *Med. Assoc.* 188, 22-23.
- Selikoff, I. J., Hammond, E. C. (1978). Mortality in the asbestos industry. *Environ. Health* 25, 11.
- Selikoff, I. J., Hammond, E. C. (1978). Mortality in the asbestos industry. *Environ. Health* 25, 11.
- Storeygard, A. R., and Brechler, D. (1978). Mortality in the asbestos industry. *Mayo Clin. Proc.* 53, 1-2.
- Timbrell, V., and Rendall, J. (1978). Mortality in the asbestos industry. *Environ. Health* 25, 11.
- Westlake, G. E., Spjut, H. (1978). Mortality in the asbestos industry. *Environ. Health* 25, 11.
- Wiele, D. T. (1977). Cancer mortality in the asbestos industry. *Environ. Health* 25, 11.
- Williams, G. M., and Gunn, R. (1978). Mortality in the asbestos industry. *Environ. Health* 25, 11.

A02512

UCC 017900

for the presence of asbestos fibers in large quantities. Further, the described method of detection of the asbestos toxicity through the toxic potential of the clay or talc

is a rapid and efficient method for the detection of asbestos fibers in environmental samples. Asbestos fibers are the most common fibers found in the environment. It is proposed that the asbestos fibers have a potential for causing cancer in humans. The asbestos fibers are found in the environment with serpentine fibers that behave like asbestos fibers. The asbestos fibers displayed this *in vitro* and *in vivo* environmental toxicity exceeding that of asbestos

assistance and Mr. [unclear] sample from Atlanta, Ga. NIOSH, Cincinnati, Ohio. A. Duluth, Minn.

y diffraction and [unclear] proceedings of the [unclear]egas, Nev., [unclear]

in human urine

ages and drinking

to occupational

insulation workers
in. N.Y. Acad.

neral 14, 91-95
6). Investigating
cancer incidence in

- Masson, T. J., McKay, F. W., and Miller, R. W. (1974). Asbestos-like fibers in Duluth water supply: relation to cancer mortality. *J. Amer. Med. Assoc.* 228, 1019-1020.
- McCabe, L. J., and Millette, J. R. (1979). Health effects and prevalence of asbestos fibers in drinking water. In "Proceedings of the American Water Works Association Annual Conference, San Francisco." In press.
- McDonald, J. C., McDonald, A. D., Gibbs, G. W., Siemiatycki, J., and Rossiter, C. E. (1971). Mortality in the chrysotile asbestos mines and mills of Quebec. *Arch. Environ. Health* 22, 677-686.
- Moore, J. (1978). NIEHS Oral Asbestos Study. In "Workshop on Asbestos Definitions and Measurement Methods," NBS Special Publ. 506, 153-162.
- National Research Council. (1977). "Drinking Water and Health," Part I, pp. IV, 1-IV, 78. Natl. Acad. Sci., Washington, D.C.
- Reiss, B., Solomon, S., Weisburger, J. H., and Williams, G. M. (1980). Comparative toxicities of different forms of asbestos in a cell culture assay. *Environ. Res.* 22, 109-129.
- Schloff, I. H., Churg, J., and Hammond, E. C. (1964). Asbestos exposure and neoplasia. *J. Amer. Med. Assoc.* 188, 22-26.
- Schloff, I. J., Hammond, E. C., and Churg, J. (1972). Carcinogenicity of amosite asbestos. *Arch. Environ. Health* 25, 183-186.
- Schloff, I. J., Hammond, E. C., and Seidman, H. (1974). Cancer risk of insulation workers in the United States. *Insulation Hyg. Progr. Rep.* 6, 1-8.
- Shorey, A. R., and Brown, A. L. (1977). Penetration of the small intestinal mucosa by asbestos fibers. *Mayo Clin. Proc.* 52, 803-812.
- Tambrell, V., and Rendall, R. E. G. (1971-1972). Preparation of the U.I.C.C. standard reference samples of asbestos. *Powder Technol.* 5, 279-287.
- Westlake, G. E., Spjut, H. J., and Smith, M. N. (1965). Penetration of colonic mucosa by asbestos particles: An electron microscopic study in rats fed asbestos dust. *Lab. Invest.* 14, 2029-2033.
- Wigle, D. T. (1977). Cancer mortality in relation to asbestos in municipal water supplies. *Arch. Environ. Health* 34, 185-190.
- Williams, G. M., and Gunn, J. M. (1974). Long-term culture of adult rat liver epithelial cells. *Exp. Cell Res.* 89, 139-142.

A02513

