

METROPOLITAN LIFE INSURANCE COMPANY  
FREDERICK H. ECKER, PRESIDENT

NEW YORK CITY

ANTHONY J. LANZA, M.D.  
*Assistant Medical Director*

March 20, 1931

Mr. Allan Hardwall  
15 Broad Street  
New York City

Dear Mr. Hardwall:

Herewith is the report of the asbestos study. This is the summary report of the American plants studied. The Canadian summary report will be forthcoming shortly. In this report the plant of the Johns-Manville Company at Manville, N.J. is designated as Plant A. I am also sending a copy to Mr. Sayer and if you wish additional copies, please let me know. Needless to say, I will be very glad to discuss this report or amplify any portions of it if you so desire.

It is, of course, understood that this report is confidential and it will be given no publicity by us except with the consent of the firms concerned. After you have had a chance to go over it, I should like the opportunity of discussing with you and your associates the question of publication.

With best regards and assuring you of my sincere appreciation of your unfailing assistance which made this report possible, I am,

Sincerely yours,

A.J. Lanza, M.D.  
Assistant Medical Director.

Enc.

# INDUSTRIAL HYGIENE

F. G. PEDLEY, B.A., M.B., D.P.H.; J. G. CUNNINGHAM, B.A., M.B., D.P.H.

## ASBESTOSIS

FRANK G. PEDLEY, M.D., D.P.H.

WITHIN the past five or six years industrial hygienists have become interested in the effects of asbestos on the health of workers. In the Province of Quebec a very considerable part of the world's supply of asbestos is mined, and the industry has been in existence for forty years or more. To our knowledge, however, no case of pneumoconiosis has been reported from asbestos in Canada, and the general impression among miners is that the dust is not hazardous. It is from England that the first cases of pulmonary asbestosis have been reported. Dr. W. E. Cooke, in 1924, published the history of a case of pulmonary fibrosis in an asbestos worker. Since then a small literature has developed, which promises to grow very much larger as time goes on.

The cases of asbestosis so far reported—with the exception of four Rhodesian miners—have occurred in the manufacture of asbestos; but it is quite possible that the condition in mines may have gone unrecognized, for mines are usually situated in the country, whereas factories are apt to be in the city where facilities for autopsies are better. If, however, miners were seriously affected by asbestos the effects should be reflected in a high tuberculosis death rate, provided the mining population was fairly stationary and did not go to sanatoria or elsewhere to die. It is believed

that the population of Thetford Mines, P.Q., is quite stable. Since this town is given over entirely to the mining and milling of asbestos the following death rates per 100,000 from tuberculosis in Thetford Mines and the Province of Quebec during the years 1926, 1927 and 1928 will be of interest. (Quebec did not enter the registration area until 1926.)

|                    | 1926  | 1927  | 1928  |
|--------------------|-------|-------|-------|
| Thetford Mines     | 116.5 | 134.5 | 115.7 |
| Province of Quebec | 121.5 | 121.5 | 121.1 |

In view of the fact that the population of Thetford Mines is only about 12,000 it is to be expected that the death rates would fluctuate considerably from year to year. These figures, therefore, do not indicate an undue mortality from tuberculosis in Thetford Mines.

Canadian asbestos occurs as a silky, fibrous form of serpentine in seams of varying width. Its analysis shows it to consist largely of magnesium silicate, but there is usually some iron present. Mining is carried on in open quarries, and is not usually a very dusty job, but in the milling a great deal of fine dust is created, and men who work there are heavily exposed to whatever hazard may exist. As mentioned above, however, it is the general impression, both among miners and physicians, that asbestos dust is not particularly harmful. There can be little doubt, however, that the disease, asbestosis, is a clinical entity,

and it has  
lish autho-  
inclusion a  
(See: Jour  
28, 1930, p

Patholog  
dividual as  
a diffuse f  
in the gre  
ever, the a  
one thing  
morphologi  
dust can be  
In silicosis  
very difficu  
smaller. In  
ticles of d  
show yellow  
20 to 100  
dumb-bell  
mented. "bodies" an  
everywhere  
ported. Th  
but it is the  
gations of  
central spici

Unquestic  
bodies" have  
in the dist  
logists, whe  
east might  
the sight o  
From the  
however, it  
asbestosis w  
either from  
bidity or mo

The follow  
list of article

1. "Effects of  
on Men  
Factories"  
Labour

E

BA, M.B., D.P.H.

ation of Thetford quite stable. Since n over entirely to the ting of asbestos the rates per 100,000 in Thetford Mines and of Quebec during 1927 and 1928 will be Quebec did not enter area until 1926.)

|       |       |       |
|-------|-------|-------|
| 1926  | 1927  | 1928  |
| 176.1 | 224.5 | 212.7 |
| 231   | 221.5 | 221.1 |

e fact that the popula- d Mines is only about be expected that the old fluctuate consider- to year. These fig- do not indicate an from tuberculosis in

occurs as a silky, pentine in seams its analysis shows gely of magnesium re is usually some iron ng is carried on in open is not usually a very in the milling a great nt is created, and men re are heavily exposed hazard may exist. As we, however, it is the sion, both among min- ans, that asbestos dust arly harmful. There ht, however, that the mis, is a clinical entity,

## INDUSTRIAL HYGIENE

311

## INDUSTRIAL HYGIENE

and it has been recognized by the English authorities to the extent of its inclusion as a compensatable disease. (See: Jour. Amer. Med. Assn., June 28, 1930, p. 2078.)

Pathologically the lungs of an individual afflicted with asbestosis show a diffuse fibrosis, not unlike silicosis in the gross. Microscopically, however, the appearance is different. For one thing large particles of dust morphologically similar to asbestos dust can be found in the fibrotic areas. In silicosis the siliceous particles are very difficult to see and are very much smaller. In addition to the large particles of dust sections of the lung show yellowish, elongated bodies from 20 to 100 microns in length, usually dumb-bell shaped and often segmented. These are the "curious bodies" and have been described everywhere asbestosis has been reported. Their origin is not known, but it is thought that they are aggregations of protein material about a central spicule of asbestos.

Unquestionably these "curious bodies" have stimulated much interest in the disease, asbestosis. Pathologists, whose interest in such a disease might languish, become alive at the sight of the "curious bodies". From the public health standpoint, however, it seems hardly likely that asbestosis will become of importance either from the standpoint of morbidity or mortality.

The following is a fairly complete list of articles on the subject to date:

1. "Effects of Asbestos Dust on Workers' Health in Asbestos Mines and Factories".

Labour Gazette, Feb. 1912, 761.

2. "Fibrosis of the Lungs due to the Inhalation of Asbestos Dust". W. E. Cooke, Brit. Med. Jour., July 26, 1924, p. 147.
3. "Pulmonary Asbestosis in its Clinical Aspects". Thos. Oliver, Jour. Ind. Hyg., 9, 483, Nov., 1927.
4. "Pulmonary Asbestosis". W. E. Cooke, Brit. Med. Jour., (Dec. 3, 1927), 1024.
5. "Histology of Pulmonary Asbestosis". W. E. Cooke, Brit. Med. Jour., (Dec. 3, 1927), 1025.
6. "Asbestosis". McDonald Stuart, Brit. Med. Jour., 1927, ii 1025.
7. "Pulmonary Asbestosis in South Africa". F. W. Simson, Brit. Med. Jour., May 26, 1928, pp. 835-837.
8. "Asbestosis". M. J. Stewart, Brit. Med. Jour., 1928, ii 509.
9. "Asbestosis". H. E. Seiler, Brit. Med. Jour., ii 1928-1932.
10. "The Presence of Asbestos Fibre in the Lesions of Asbestos Workers". S. R. Gloyne, Tubercle, June, 1929, p. 405.
11. "A Case of Pulmonary Asbestosis". W. B. Wood & D. S. Page, Tubercle, July, 1929, p. 437.
12. "Asbestos Dust and the 'Curious Bodies' Found in Pulmonary Asbestosis". W. E. Cooke, Brit. Med. Jour., Sept. 28, 1929, 578.
13. "Clinical Aspects of Pulmonary Asbestosis". A. C. Haddon, Brit. Med. Jour., Sept. 29, 1929, 580.
14. "Further Observations on Pulmonary Asbestosis with Special Reference to Asbestos Dust and the 'Curious Bodies' Found in the Lungs". W. E. Cooke, Jour. Royal Microscopical Soc., 1930, vol. 50, p. 15.
15. "Pulmonary Asbestosis. A Socio-medical Study". Archiv für Gewerkepathologie und Geneserhygiene, Sonderabdruck aus 1. Band und 2. Heft, 1930.
16. "Asbestos". Occupation & Health, International Labour Office, Brochure No. 27.
17. "The Occurrence of Pulmonary Fibrosis and other Pulmonary Affections in Asbestos Workers". E. R. A. Mercuriether, Journ. Indust. Hygiene, 12, 196-222 and 239-257, May and June, 1930.

95  
5713729

# INDUSTRIAL HYGIENE

2. INDUSTRIAL HYGIENE

3. INDUSTRIAL HYGIENE

4. INDUSTRIAL HYGIENE

5. INDUSTRIAL HYGIENE

6. INDUSTRIAL HYGIENE

7. INDUSTRIAL HYGIENE

8. INDUSTRIAL HYGIENE

9. INDUSTRIAL HYGIENE

10. INDUSTRIAL HYGIENE

11. INDUSTRIAL HYGIENE

12. INDUSTRIAL HYGIENE

13. INDUSTRIAL HYGIENE

14. INDUSTRIAL HYGIENE

15. INDUSTRIAL HYGIENE

16. INDUSTRIAL HYGIENE

17. INDUSTRIAL HYGIENE

18. INDUSTRIAL HYGIENE

19. INDUSTRIAL HYGIENE

20. INDUSTRIAL HYGIENE

21. INDUSTRIAL HYGIENE

22. INDUSTRIAL HYGIENE

23. INDUSTRIAL HYGIENE

24. INDUSTRIAL HYGIENE

25. INDUSTRIAL HYGIENE

26. INDUSTRIAL HYGIENE

27. INDUSTRIAL HYGIENE

28. INDUSTRIAL HYGIENE

29. INDUSTRIAL HYGIENE

30. INDUSTRIAL HYGIENE

31. INDUSTRIAL HYGIENE

32. INDUSTRIAL HYGIENE

33. INDUSTRIAL HYGIENE

34. INDUSTRIAL HYGIENE

35. INDUSTRIAL HYGIENE

36. INDUSTRIAL HYGIENE

37. INDUSTRIAL HYGIENE

38. INDUSTRIAL HYGIENE

39. INDUSTRIAL HYGIENE

40. INDUSTRIAL HYGIENE

Bialy  
pml 115

