

PLAINTIFF'S
EXHIBIT
QA-105

1950

Report to
Quebec Asbestos Mining Association
re
Pulmonary Cancer

September 25, 1950.

PLAINTIFF'S
EXHIBIT

13058.0

216

Return to
General Electric Medical Equipment
70
Fairbury, Illinois

PHS Ex # 3 for rd
4/4/79

Report to
Canadian Asbestos Mining Association
on
Pulmonary Cancer

PROBLEM:

To ascertain if there is a relationship between the
inhalation of asbestos dust and pulmonary cancer or
has been stated publicly by certain British authorities that
this is a definite sign of wide exposure. As this
is necessary for us to be able to prove that this
statement is not correct or to accept it and the
sequence that goes with it. Our position in this
matter is the same as the Association and Industry
found itself in 1937 as regards asbestosis.

RECOMMENDATIONS:

It is recommended to the Association that they proceed
at once on two programs -

1. A Short Range Program
2. A Long Range Program

2. The Short Range Program

This would consist of an epidemiological study. This
statistical work or survey to be done under the aus-
pices of a statistician of repute and it is especially
recommended that this statistician should be, if
possible, a Canadian.

1. The Short Range Program - continued -

Suggestions for this job are:

Dr. A. D. Phillips of the National Cancer Institute of Canada, who are affiliated with the Canadian

Cancer Society. E. Royler Hammond was its

Director of Science, Secretary of the Cancer Prevention

Committee, likewise statistician of the Cancer Insti-

tute of the United States, Washington, D.C.; Mary Ross

of the School of Hygiene in Toronto could assist in this

work if it was done under the auspices of Dr. Hammond;

another person considered proper for this is Dr. A.

Valois, a statistician for the City of Montreal.

This survey should be undertaken under the auspices of
Saratoga.

2. The Long Range Program:

Under The Long Range Program it is recommended that the services of the Saratoga Laboratory of the Edward Tamm Foundation be utilized. They are presently setting up a cancer research dealing with aluminum, iron, carbon and silicon dust and could do this by putting in another dust rock and do our work paralleling these other elements.

of the sale of the property in Toronto could well be this

1. The first of these is the fact that the
2. second of these is the fact that the
3. third of these is the fact that the
4. fourth of these is the fact that the
5. fifth of these is the fact that the
6. sixth of these is the fact that the
7. seventh of these is the fact that the
8. eighth of these is the fact that the
9. ninth of these is the fact that the
10. tenth of these is the fact that the

[illegible]

SHORT RANGE PROGRAM:

The United States Public Health and Cancer Research, Dr. Macintosh, the National Cancer Institute have all published that there is a relationship between asbestos dust and pulmonary cancer. They have made these published statements only on statistical data, no research has been done, and these statistical data are based only on death statistics. For this reason the Short Range Program has been recommended to you since we are informed by our own medical officers that:

- a. This statistical information is not confirmed in their opinion is our workmen.
- b. The statistical survey is the most rapid, cheap, and more conclusive than any other method.
- c. We would be using the same methods as our antagonists.

Personnel Required:

The following personnel are considered necessary to make this survey:

- a. One statistician of repute or an actuary of the Insurance type. (A medical statistician is considered preferable.)
- b. X-ray consultant. Dr. Smith and Dr. Cartier could very well serve as consultants with Saranac reporting on border-line cases.

The estimated cost of making this report is \$4,000.

maximum and the required time to complete is four months.

THE FIRST PART OF THE
INVESTIGATION INTO THE
CAUSES OF THE

DISASTERS OF THE

NAVY OF THE

UNITED STATES

IN THE

YEAR 1897

BY

THE

NAVY DEPARTMENT

WASHINGTON

1898

THE

NAVY DEPARTMENT

WASHINGTON

1898

THE

NAVY DEPARTMENT

WASHINGTON

1898

THE

NAVY DEPARTMENT

WASHINGTON

1898

THE

NAVY DEPARTMENT

WASHINGTON

1898

THE

NAVY DEPARTMENT

WASHINGTON

1898

THE

NAVY DEPARTMENT

WASHINGTON

1898

LONG RANGE PROGRAM:

This program would consist of a research program and the following through on it for at least two years and probably three years. Two programs have been considered :

1. The transplanting of living tumors under the guidance of Dr. Leane and his group.
2. Utilizing the facilities as we have in the past of the Trudeau Foundation and subjecting animals to a high asbestos dust condition.

Long Range Program - 3:

The third phase:

1. It is the feeling of your committee that along with the Short Range Program and the Long Range Program a third phase should be developed. That phase would consist of more publicity coming from our Association than has in the past, rather than taking the defensive attitude that we have. We would point out that Dr. Paul Cartier has been published in France and Portugal regarding the incidence of heart disease and tuberculosis in the asbestos industry but there has been no mobilization, to our knowledge, dealing with this subject, on this continent. It is recommended that more of " " be published at least one yearly, dealing with the health aspects of our industry.
2. That we arrange to have our doctors lecture at Medical Schools yearly.
3. That we attempt to arrange to affiliate our doctors in some manner with Research Centres such as Saranac and in such organizations as the Industrial Hygiene Association and the Institute of Industrial Medicine as clinicians in the asbestos industry.

In this matter we feel that we would combat reports which we believe, with the information at hand, to be erroneous. We believe that publicity on the part of our Institute would combat erroneous reports such as Jean Kurokawa made in Thorford recently when he quoted from what he reported to be an official report of the Third International Conference on pneumoconiosis. This report has not as yet been issued.

Long Range Program - 2.

This would consist of an inhalation techniques with studies carried on at Saranac. It is recommended that if the Long Range Program is adopted, then this is the program that should be adopted:

Saranac is the central repository of industrial lung data. They have the inhalation chambers and technicians for doing this type of research and they have a number of years of research with us in asbestosis. Likewise Dr. Rhode who is connected with the Sloan-Kettering Cancer Research Foundation is in contact with Saranac.

This study would require a minimum of three years and probably five years. It is estimated to cost \$25,000. the first year and \$10,000. the second year. Each member company is in receipt of a copy of their proposal.

Dr. Rhode + Co

and the ... of ...

the ...

Take ...

...

...

...

...

...

...

...

...

...

...

...

Long Range Program - 1.

The proposal under this program consists of transplanting living tissue of an embryo mouse with tumor implanted into a living mouse to see if cancer can be produced. This proposal can only be considered as a screening and cannot be regarded as positive data, but only as a probability if the research doctors are able to reproduce pulmonary cancer. If there is no reproduction there is no proof, neither positive nor negative.

The estimated cost of this program would be \$10,000. for the first year with no further commitments on our part at the end of the first year, but it is not considered probable that any results would be forthcoming prior to two years.

Report to
Quebec Asbestos Mining Association
re
Pulmonary Cancer

PROBLEM:

To ascertain if there is a relationship between the inhalation of asbestos dust and pulmonary cancer. This has been stated publicly by certain Health Authorities and in certain magazines of wide coverage. It is necessary for us to be able to prove that this statement is not correct or to accept it and the consequences that goes with it. Our position in this matter is the same as the Association and Industry found itself in 1937 as regards asbestosis.

RECOMMENDATION:

It is recommended to the Association that they proceed at once on two programs -

1. A Short Range Program
2. A Long Range Program

1. The Short Range Program:

This would consist of an epidemiological study. This statistical work or survey to be done under the auspices of a statistician of repute and it is especially recommended that this statistician should be, if possible, a Canadian.

SHORT RANGE PROGRAM:

The United States Public Health and Cancer Research,
⑦ Dr. Marwether, the National Cancer Institute have
all published that there is a relationship between
asbestos dust and pulmonary cancer. They have made
these published statements only on statistical data,
no research has been done, and these statistical data
are based only on death statistics. For this reason
the Short Range Program has been recommended to you
since we are informed by our own medical officers that:

- a. This statistical information is not confirmed in their opinion in our workmen.
- b. The statistical survey is the most rapid, cheapest, and more conclusive than any other method.
- c. We would be using the same methods as our antagonists.

Personnel Required:

The following personnel are considered necessary to make this survey:

- a. One statistician of repute or an actuary of the insurance type. (A medical statistician is considered preferable.)
- b. X-ray consultant. Dr. Smith and Dr. Cartier could very well serve as consultants with Saranac reporting on border-line cases.

The estimated cost of making this report is \$4,000.

maximum and the required time to complete is four months.

When the presence of neoplasm in an individual is suggested by this technique, he should be further studied with the purpose of establishing or excluding this diagnosis with certainty. Other factors suspected of being carcinogenic, such as cigarette smoking, may constitute variables within the population to be studied. Information as to these variables must be obtained as well as information concerning exposure to asbestos. It appears that cigarette smoking is the most prominent variable factor in the epidemiological study proposed. Other suspected factors such as atmospheric hydrocarbons would be essentially constant for each group of individuals in the same geographic area where asbestos is being mined.

4. In view of these considerations, it is suggested that only certain phases of the epidemiological study be undertaken at this time pending the advice and recommendations of those with a wide experience in the conduct of such studies. Agencies with this experience are: the Medical Research Councils of Canada and the United States, the cancer societies of both countries, the respective tuberculosis societies, the National Cancer Institute of the U. S. Public Health Service, and the Sloan-Kettering Institute of the Cancer Memorial Hospital.

5. The phases of the epidemiological study which might be undertaken immediately include the clinical and statistical data already available for employees in industries such as those at Thorford and Asbestos, where routine physical examination and chest

Suggest advice & recommendations be first obtained then make

X-rays have been standard practice for years. In essence, this program should include scrutiny of all medical records, of chest roentgenograms of all employees, of autopsy protocols, and of death certificates for evidence of malignancy. It should be pointed out that statistical calculations relying on clinical records or on death certificates based on clinical diagnosis may contain a wide range of error in the fundamental material and may be unreliable as to the absolute numerical conclusions reached. Similarly, autopsy protocols may represent only a highly selective sample of the total mortality and may constitute, therefore, a weighted index of the incidence of cancer in subjects exposed to asbestos dust.

PART III

EXPERIMENTAL INVESTIGATIONS

6. The Baranec Laboratory with its background of study concerning chronic pulmonary disease, especially that resulting from the inhalation of industrial dusts including asbestos, is uniquely qualified to pursue certain phases dealing with the problem of asbestosis and pulmonary cancer.
7. For purpose of completeness and general information, certain brief comments seem justified for guidance in the design of the experimental approach which we propose.

'Susceptible' animals

A number of experimental methods may be employed to test the capacity of asbestos to produce cancer. It is important to keep in mind, however, certain facts concerning asbestos which affect the selection of the method:

(1) Asbestosis is a chronic pulmonary disease resulting from the deposition of asbestos fibers in the lung; therefore, the experimental method employed should direct emphasis on the lung.

(2) The weight of evidence is that asbestos fibers produce their irritation by a mechanical rather than by a chemical mode of action. The mechanical action is intimately related to the movement of the lung during respiration. Thus, an experimental method which neglects movement of the lung and thereby excludes the mechanical action in all probability would be highly inconclusive and subject to criticism.

(3) Asbestosis is a disease resulting from widespread dissemination of asbestos dust in the lung. Reliable conclusions in an experimental investigation with asbestos cannot be drawn solely from experiments with the intratracheal injection technique whereby a high concentration of dust becomes localized in a restricted portion of the lung and stimulates excessive focal irritation. When the asbestos dust is introduced into the animal in that manner, the injected dust may miss certain potentially cancer-producing sites in the lung which are remote from the

localized area of implantation of dust. It is further possible for the excessive focal irritation, which is not a characteristic feature of human asbestosis, to induce pulmonary cancer which would not have developed had the dust been uniformly distributed throughout the lung. Thus, the intratracheal injection technique alone might lead to unwarranted conclusions. The intratracheal method has been used for cancer studies by a number of investigators, probably because of the relative difficulty and tediousness of methods involving inhalation of the potential carcinogen. The inhalation method, however, has been employed by eminently recognized investigators of carcinogenic agents.

8. In passing, it might be said that lung tumors can be produced by numerous means in susceptible mice, i.e., mice possessing the tendency to develop tumors spontaneously. Some of the methods gathered from the literature are: repeated applications of coal tar to the skin; subcutaneous injection of carcinogenic hydrocarbons; intravenous injection of serum and charcoal dispersions of carcinogenic hydrocarbons; feeding large doses of dibenzanthracene; and introduction of carcinogenic hydrocarbons into the rectum, vagina, peritoneal cavity, or pleural cavity. For explanation of the occurrences of lung cancer following such methods, it has been suggested that the introduction of carcinogenic compounds into the body of the mouse alters the body state so that tumors arise at points of incidental irritation. The more plausible explanation advanced by certain investigators is that the lung tumors are pro-

duced because very small amounts of the carcinogenic agent leave the site of application and come in contact with extremely susceptible alveolar epithelium. If this is true, it is unlikely that such methods could be used for investigation of cancer and asbestos, a dust which remains within pulmonary tissue, which is not transported systemically as are soluble chemical agents, and which produces its irritating effect by a mechanical rather than by a chemical mode of action.

3. Reference has been made to inbred strains of mice in which the spontaneous occurrence of tumors of the lung is a characteristic feature. Strains are available in which the incidence of spontaneous lung tumors is accurately known. In some strains the incidence is low, so that an increase in the incidence after an experimental procedure is easily detectable. Other strains with high incidence of spontaneous tumors are suitable to determine whether the procedure has caused tumors to appear at an earlier age and whether the number of tumors per mouse has been increased.

The susceptibility of the pulmonary tissue in selected inbred strains of mice is such that they serve as excellent test animals for checking an alleged extrinsic carcinogen such as asbestos. A reasonably delicate assay of the potency of asbestos as a carcinogen is possible because of the unique multiplicity of tumors formed.

PART IV

PROPOSED EXPERIMENTAL INVESTIGATION

10. METHODS TO BE EMPLOYED

In view of the considerations mentioned in the preceding paragraphs, the Saranac Laboratory proposes to conduct an experimental investigation employing ⁽¹⁾ the inhalation method and also, for supportive evidence, ⁽²⁾ the intratracheal injection method. Two groups of pure inbred strain mice will be used, i.e., the low spontaneous lung tumor "strain C 57 black," and the high lung tumor "strain A" (obtained from the Roscoe B. Jackson Memorial Laboratory). The animals of each group will be divided into subgroups, equivalent as to sex, age, and weight at the time of inauguration of the actual experiments. After mixing, part of each subgroup will be exposed to atmospheric suspensions of long-fiber asbestos dust of controlled concentrations; part will be injected intratracheally with measured amounts of the dust suspended in sterile normal saline; part will be set aside in normal air for controls. Each animal will be numbered and a separate record will be kept.

Animals from each subgroup will be sacrificed at intervals, and the other members will be permitted to live out their natural life span. After death the lungs will be fixed by intratracheal injection of Zenker's formal or formalin solution. Sections passing transversely through both lungs and the mediastinal structures will be prepared for microscopic study.

11. EVALUATION OF EXPERIMENTAL RESULTS ✓

Carcinogenicity is to be evaluated by comparing the following five factors in all subgroups:

- ✓(1) The per cent of mice with lung tumors.
- ✓(2) The average number of tumors per mouse.
- ✓(3) The number of tumors arising in the asbestotic areas versus the non-asbestotic areas of tissue in the exposed mice.
- ✓(4) The induction time for tumors.
- ✓(5) Evidence for direct origin of tumor from asbestos, i.e., the histological appearance of tumor will be compared with the non-cancerous epithelial hyperplasias that may be present.

12. COMPARATIVE ANALYSIS WITH OTHER INVESTIGATIONS UNDERWAY

The studies concerning asbestos will be integrated with other investigations of the Saranac Laboratory. These investigations concern the carcinogenic properties of inhaled quartz and of non-irritating dusts, such as carbon and iron oxide. The data obtained will form a basis for statistical study of the occurrence of pulmonary cancer in mice exposed to dusts other than asbestos.

13. ESTIMATED DURATION OF THE EXPERIMENTAL INVESTIGATIONS

The investigations proposed involve disease processes which develop very slowly. This applies to asbestosis as well as to pulmonary cancer. With respect to asbestosis, previous studies indicate that at least sixteen months of continuous exposure to dust are necessary in order to produce definite fibrosis in a highly susceptible

animal such as the guinea pig. In contrast, the mouse is more resistant and develops his reaction even more slowly under similar conditions of exposure. The development of asbestosis in mice may be speeded up considerably by increasing the concentration of the dust to which they are exposed. Thus, it is anticipated that rather definite results can be obtained in not less than two years by the inhalation method, and in not less than one year in mice receiving an intratracheal injection of the dust. Regarding the latter method, it must be stressed that although the injection experiment is of shorter duration, the results are inconclusive. An inhalation study of relatively longer duration should be performed to obtain reliable results.

With respect to pulmonary cancer, there are many complex problems inherent in any study of the subject. The very nature of the disease and the experience of other investigators dictate that at least a two-year, and preferably a five-year program, should be envisaged. This should not discourage the sponsorship of the program by the companies, as unfortunately there is no other alternative if reliable results are to be obtained.

14. REPORTS OF EXPERIMENTS

Progress reports will be issued from time to time during the experiment and a comprehensive report will be submitted at the conclusion of the investigation. These reports will be forwarded to the proper authorities identified by the sponsoring agencies.

18. PUBLICATIONS OF RESULTS OF INVESTIGATIONS

The Bureau Laboratory reserves the right to publish the results of its investigations in the recognized manner commonly employed by the scientific fraternity and in accordance with established practices, whereby the manuscript, prior to its publication, is submitted to the sponsoring company for its review and approval.

Look 6

19. COST ESTIMATE

The combined project including both the inhalation and the intratracheal techniques requires the services of a number of staff members, the use of an exposure chamber or room and special apparatus needed for control, the purchase of a large number of mice of a pure inbred strain, the use of special mouse cages, and a large quantity of expendable materials such as chemicals, glassware, microscopic slides, and cover slips.

The cost for this comprehensive study will approximate \$18,000.00 for the first year and \$10,000.00 for the second year.

ATV:bw

Cost of 5 yr. Program?

1st.	16,000
2nd -	10,000
3rd -	5,000
4th -	5,000
5th -	5,000
}	
41,000	