

PRELIMINARY DRAFT
LEAD PAINT INGESTION STUDY

SEP 10 1973

R.A. Moore

*original
was Revised*

FINAL REPORT
14 September 1973

Contract No. 62-W-62GC&NPC
MRI Project No. 3729-B

For

National Paint and Coatings Association
1500 Rhode Island Avenue, N.W.
Washington, D.C. 20005

Attn: Mr. Royal A. Brown

0007-SWP-036910

N22050

PRELIMINARY DRAFT
LEAD PAINT INGESTION STUDY

FINAL REPORT
14 September 1973
Contract No. 62-W-62GCSNPC
MRI Project No. 3729-B

For
National Paint and Coatings Association
1500 Rhode Island Avenue, N.W.
Washington, D.C. 20005

Attn: Mr. Royal A. Brown

0007-SWP-036911

0007-SWP-000112898

PREFACE

This report was prepared at Midwest Research Institute, 425 Volker Boulevard, Kansas City, Missouri 64110, under Contract No. 62-W-62GC&NPC with the National Institutes of Health, Department of Health, Education and Welfare, MRI Project No. 3729-B, "Lead Paint Ingestion Study." The research was sponsored by the National Paint and Coatings Association, 1500 Rhode Island Avenue, N.W., Washington, D.C. 20005. Dr. John P. Frawley, Chief Toxicologist, Hercules, Inc., Wilmington, Delaware 19899, was the project monitor.

The research was conducted in the Biological Sciences Division, under the direction of Dr. W. B. House from 1 December 1972 through 31 July 1973. Dr. Thomas R. Castles, Principal Pharmacologist, was the principal investigator, assisted by Dr. Jaime L. Sanyer, Associate Pathologist, and Mrs. Jane Hoch, Biology Research Assistant. Dr. James L. Spigarelli, Senior Chemist supervised the lead analysis with the assistance of Mrs. Hope M. Miller, Assistant Chemist.

Approved for:

MIDWEST RESEARCH INSTITUTE

W. B. House, Director
Biological Sciences Division

14 September 1973

TABLE OF CONTENTS

	<u>Page</u>
I. Introduction.	1
II. Method	2
A. Preparation of <u>Paint</u> Films Containing Different Concentrations of Lead.	2
B. Preparation of Diets.	3
C. Experimental Procedure.	3
D. Analyses	4
E. Pathology	8
F. Statistical Analyses.	9
III. Results	9
A. Concentrations of Lead in Paint Films	9
B. Lead Analyses of Diets Containing Different Paint Films	9
C. Preliminary Feeding Studies	10
D. Gross Observations, Feed Consumptions and Body Weights	10
E. Weekly and Total Lead Consumption	11
F. Hematology of Rats Fed Paint Films Containing Different Concentrations of Lead.	12
G. Tissue and Fluid Chemistry of Rats Fed Paint Films Containing Different Concentrations of Lead	12
H. Lead Content in Tissues of Rats Fed Paint Films Containing Different Concentrations of Lead	13
I. Pathology of Rats Fed Paint Films Containing Different Concentrations of Lead	15
IV. Discussion	17
V. Conclusions	19
References	20
Figures	22
Tables	26

I. INTRODUCTION

Few persons will disagree that old lead-based paint presents a dangerous and unnecessary health hazard to young children. As a result of reports by Kehoe,^{1/} Chisholm,^{2/} and King^{3/} medical researchers, paint manufacturers and legislators have recognized the seriousness of this problem and are acting to establish safe concentrations of lead in paint. To do this, they have voluminous reports on clinical observations and animal research from which to draw. Unfortunately, most of this information is based upon the old lead-based paints or soluble lead salts, which do not reflect the ~~enamel~~ ^{modern} paint formulations on today's market.

Before safe concentrations of lead in paint can be intelligently established, we must evaluate the toxicity of the lead as it exists currently in modern enamel paints. These studies must be as complete as possible and must be performed upon several animal species, since a complete model for man has not yet been found.

It was the purpose of this study to evaluate the toxicity in rats produced by modern enamel paints containing lead octoate, lead chromate or lead carbonate. The results are presented in the following pages.

II. METHODS

A. Preparation of Paint Films Containing Different Concentrations of Lead

Paint films containing no lead or lead in the form of lead octoate or lead chromate were supplied by the Sherman-Williams Paint Company. Enamels which contained either lead octoate or lead chromate (medium yellow) were prepared using a flat alkyd ~~enamel~~ ^{Paint} formula based upon a TT-R-266D Type I Class B alkyd. The flat alkyd ~~enamel~~ ^{Paint} without ~~either~~ ^{added} lead compound was used for control. The percent of lead in each sample was calculated on the basis of the nonvolatile material. Dispersion was accomplished with a Cowles Dissolver and the ~~painting~~ ^{Fig. Vol. Conc.} was held at one level. The paints were placed in pans and allowed to air dry. This was followed by force drying at 120°F to volatilize any remaining solvent. Paint films were ground, sieved and chips ranging from 0.5 to 1.0 mm in size used for this study. *This chips used.*

The concentrations of lead in the different paint films were determined independently by the DeSoto ~~Paint Company~~ ^{Inc.}, Sherman-Williams ~~Paint Company~~ ^{WPA} and Midwest Research Institute, using Atomic Absorption Spectrophotometry.

Samples of old lead-based paint were obtained from the U.S. Department of Commerce, National Bureau of Standards, Washington, D.C. This paint had been pulverized and sieved through 325 mesh screen and contained 11.87% lead by weight. This sample will be called "NBS lead paint" for this report.

B. Preparation of Diets

Diets were prepared by mixing paint films with Purina Rat Chow mash in the concentration of 0.1% (w/w). This was accomplished by first preparing a 10% concentrate (paint film/feed, w/w) in a high-speed twin shell mixer. This concentrate was then added to the appropriate amount of Purina Rat Chow mash to give a final concentration of 0.1% (paint film/feed, w/w) and mixed for 10 min in a bulk mixer.

Each diet was prepared three times during the experiment and samples of each of these preparations were assayed for lead content.

C. Experimental Procedure

Two hundred weanling (40-60 gm) Charles River rats (100 males and 100 female rats) were used for this study. Upon arrival rats were housed individually in air conditioned quarters in polycarbonate cages containing hardwood bedding and filter tops. After 5-7 days of equilibration, all rats were assigned one of the following diets.

1. Rat chow plus paint film without lead
2. Rat chow plus paint film containing 0.08% lead as lead octoate
3. Rat chow plus paint film containing 0.53% lead as lead octoate
4. Rat chow plus paint film containing 2.05% lead as lead octoate
5. Rat chow plus paint film containing 0.42% lead as lead chromate
6. Rat chow plus paint film containing 1.95% lead as lead chromate
7. Rat chow plus paint film containing 12.43% lead as lead chromate
8. Rat chow plus paint film containing 66.05% lead as lead carbonate
9. Rat chow plus NBS lead paint containing 11.92% lead in an unknown form

Twenty male and 20 female rats were assigned to the control diet, and 10 males and 10 female rats were assigned to each of the remaining diets. Each diet was fed in a 9 oz wide-mouth glass jar which was secured to the cage. Tap water was available ad. libitum during the entire experiment.

Feed consumption was measured daily and body weights were measured weekly. All rats were observed twice or more each week for toxic signs.

After 4, 8, and 13 weeks a few male and female rats were placed in metabolism cages for the collection of 24-hr urine samples. A portion of the collected urine was used for urinalysis and the remainder frozen for assay of delta aminolevulinic acid and coproporphyrin. After urine collection each rat was anesthetized with ether, exsanguinated via the abdominal aorta and bone marrow smears were prepared. Blood samples were heparinized, cooled and used immediately for hematology and enzyme assays. Aliquots of the remaining blood were taken for protoporphyrin and lead assays. After urine and blood samples were obtained, each rat was necropsied and tissue taken for lead analysis or microscopic examination.

D. Analyses

1. Lead analyses of paint films and diets containing paint film:

a. Paint films: Ten milligram samples of paint films containing no lead or 0.08% lead octoate were charred with 3 ml of concentrated

nitric acid and dry-ashed at 500°C for 2 hr. The ash was dissolved in 1 ml of aqua regia, diluted to 5 ml with distilled water and this final dilution measured for lead content by atomic absorption.

All other paint films (10 mg samples) were digested in 15 ml of 2:1 concentrated nitric acid: 37% perchloric acid, evaporated and the remaining perchloric acid solution diluted to 25 ml with distilled water. This final dilution was used for atomic absorption spectrophotometry.

b. Diets: One-gram samples of feed were digested in 2:1 concentrated nitric acid: 37% perchloric acid and their final dilutions adjusted to make their lead concentrations within the detection limits of the atomic absorption technique.

2. Analyses performed upon each rat:

a. Hematology:

(1) Hematocrit: Hematocrit was determined in capillary tubes using a microcapillary centrifuge (International Equipment Company, Model MB).

(2) Hemoglobin: Hemoglobin was measured as cyanomethemoglobin.^{4/}

(3) Erythrocyte and leukocyte counts: Total erythrocyte and leukocyte were counted using a Coulter Electronic Particle Counter with 100 μ aperture.^{5/}

(4) Reticulocytes: Reticulocytes were counted by the methylene blue method using the Miller disc.^{6/}

(5) Differential leukocyte counts: Wright's stain was used to stain the leukocytes for examination.

(6) Erythrocyte osmotic fragility: Osmotic fragility of erythrocytes was quantitatively determined by subjecting heparinized whole blood to sodium chloride solutions of different osmolarities.^{7/} The concentration of sodium chloride which hemolyzed 50% of the erythrocytes was obtained from a plot of percent hemolysis versus sodium chloride concentration.

b. Body fluid and tissue chemistry:

(1) Plasma protein electrophoretic patterns: Total plasma protein were determined using the Hycal Biuret Reagent (Hycal, Inc., Houston, Texas). The quantity of each plasma protein was determined electrophoretically on cellulose acetate and expressed as a percentage of the total plasma protein.

(2) Erythrocyte- δ -aminolevulinic acid dehydrase (ALAD): δ -Aminolevulinic acid dehydrase activity was determined by the method of Lichtman and Feldman.^{8/} This procedure was performed immediately after each blood sample was taken.

(3) Erythrocyte protoporphyrin: Protoporphyrin in erythrocytes was measured by the method of Heller, et al.^{9/} This analysis was performed the day after blood was withdrawn.

(4) Urinary δ -aminolevulinic acid (ALA): Urinary δ -aminolevulinic acid was measured according to the Davis and Andelman^{10/} modification of Mauzerall's and Cranick's Method.^{11/} These analyses were performed in subdued light.

(5) Urinary coproporphyrin: Urinary coproporphyrins were determined by the method of Schlenker and Kitchell.¹² These analyses were performed in subdued light.

c. Urinalysis:

(1) Urinary protein (albumin): Urinary protein was measured with "Uristix" reagents strips (Ames Company, Elkhart, Indiana).

(2) Microscopic examination of urine: Urine samples were centrifuged, the residues resuspended, and examined microscopically for the presence of erythrocytes and leukocytes under high power field and for casts under low power field.

d. Tissue lead:

(1) Blood: One milliliter of blood was mixed with 1 ml of 3:1 v/v 5% trichloroacetic acid:37% perchloric acid. The sample was centrifuged and the supernate filtered through an AAWP 0.8 μ Millipore filter. The supernate was analyzed for lead with an atomic absorption spectrophotometer using standards prepared in control rat blood.

(2) Other tissues: Bone, liver, kidney, and brain were digested in concentrated nitric acid, evaporated to dryness and reconstituted to the lowest possible volume with 20% nitric acid. These solutions were analyzed for lead by atomic absorption spectrophotometry.

(3) Atomic absorption analysis: Lead concentrations were measured using a Varian-Techtron AA-5 atomic absorption spectrophotometer. Sample solutions were aspirated into an air-acetylene flame and the absorbance was measured at 2,833 nm. Background interference was determined with the

use of a hydrogen continuum lamp at 2,833 nm and subtracted from the absorbance obtained at 2,833 nm with the Pb hollow cathode lamp.

E. Pathology

1. Gross pathology: At necropsy, rats were examined for gross abnormalities and their livers, kidneys, spleens, hearts, gonads, thyroids, brains, and adrenals weighed and the relative organ weights calculated. After weighing, a portion of the liver, kidney, and brain were taken for microscopic examination and the rest of each organ was measured for lead content. A femur was removed to be used for the measurement of lead in bone.

2. Microscopic pathology: Bone marrow smears were prepared for a myeloid/erythroid cell count. The following tissues were fixed in buffered neutral 10% formalin: brain, liver, spleen, stomach, small intestine (duodenum, jejunum, ileum), colon, pancreas, kidneys, urinary bladder, adrenals, thymus, gonad, thyroid (with parathyroid attached), mesenteric salivary gland, lymph nodes, heart, lungs, diaphragm, skeletal muscle, and prostate or uterus.

Tissues from all 13 week rats fed the control, 2.05% lead octoate, 12.43% lead chromate, 66.05% lead carbonate, and the 11.92% NBS lead paint diets were embedded in paraffin, sectioned to 6 μ thick, stained with hematoxylin and eosin and examined for histopathology.

F. Statistical Analyses

Control and treatment values were compared statistically using Dunnett's multiple-comparison test with $P < 0.05$ as the criteria of significance.

III. RESULTS

A. Concentrations of Lead in Paint Films

The concentrations of lead in the paint films supplied by the Sherman-Williams Paint Company were analyzed for lead content by the De Soto ~~Paint Company~~, Sherman-Williams Paint Company, and Midwest Research Institute (MRI). The results of these analyses are shown in Table 1. The control enamel which did not contain lead octoate or chromate was found to contain approximately 0.01% lead. The agreement between the different analyses was good, so an average ppm of lead was computed and used for the actual lead concentration of each paint film. Lead concentrations of the lead paint obtained from the National Bureau of Standards were averaged in a similar manner.

B. Lead Analyses of Diets Containing Different Paint Films

Table 2 shows the lead content of each diet. Purina Rat Chow mash was found to contain approximately 4 $\mu\text{g}/\text{gm}$ of lead (assuming 0.1 $\mu\text{g}/\text{gm}$ of lead was contributed by the control enamel). The μg of lead/ gm of feed which was attributable to each added paint film was reasonably close to its respective theoretical value. The average total μg of lead/ gm of feed for each diet was used for the calculation of weekly and total lead consumption for each rat.

C. Preliminary Feeding Studies

Before beginning this study, a pilot experiment was performed to see if the rats were consuming the lead paint film along with their feed. Briefly, three rats were placed on the control diet and three rats on the 66.05% lead carbonate diet. After 48 hr, feces were collected and their lead content measured. Feces from rats on the control diet contained 11.5 µg of lead/gm feces while feces from rats fed the 66.05% lead carbonate diet contained 3,683 µg of lead/gm of feces. Thus, rats did consume the paint film along with their feed.

D. Gross Observations, Feed Consumptions and Body Weights

During the entire study only two rats had remarkable changes. One female rat on 2.05% lead octoate developed an abscess on her left shoulder during the ninth treatment week and one female on 66.05% lead carbonate developed abscessed hind feet in the 10th week and chew her toes off.

The feed consumption of male and female rats are shown in Figures 1 and 2, respectively. Except for the rats fed the diet containing 11.92% lead as NBS lead paint, all groups ate at the same rate as the control group throughout the entire experiment. The reason the 11.92% NBS lead paint group ate significantly less initially, was that they were started a month after the other rats and were initially smaller. Since this group was consuming feed at the same rate as the control group by 4 weeks, we consider their initial food consumption normal also (for their size).

The body weight gains for each group of rats are shown in Figures 3 and 4. As mentioned above, the rats on the 11.92% NBS lead paint diet were small initially but became similar to controls by either the fourth or eighth week. Throughout the entire period the male rats in the other treatment groups gained weight at a normal rate. The female rats did show a change in body weights during the first eight weeks of feeding which appears related to the concentration of lead octoate. After 4 weeks of feeding, rats in the 2.05% lead octoate group weighed significantly less than control. The pattern was the same at eight weeks. After 12 weeks their average body weights still exhibited the same pattern, but the 2.05% lead octoate group was not significantly different from control. This was due to fewer animals and a slight increase in variability.

E. Weekly and Total Lead Consumption

The weekly and total lead consumption during the experiment were calculated on the basis of the total lead/diet and amount of each diet eaten by each rat. These calculations are shown in Tables 3 and 4. During the 13th week period, the control group consumed approximately 105 ug of lead/day, while rats eating 2.05% lead octoate, 12.43% lead chromate, 66.05% lead carbonate, and 11.92% NBS lead paint consumed an average of 786.9, 3,591.1, 13,435.8, and 3,184.3 ug of lead/day, respectively. Using the average body weights of rats surviving 13 days, the rats in the control, 2.05% lead octoate, 12.43% lead chromate, 66.05% and 11.92% NBS lead paint groups consumed 245.2, 1,957.5, 8,469.6, 34,430.8, and 7,804.7 ug of lead/day/kg of body weight, respectively.

F. Hematology of Rats Fed Paint Films Containing Different Concentrations of Lead

The results of the hematology of rats fed paint films containing different concentration of lead are shown in Tables 5-13. No differences were observed in the erythrocyte count, reticulocyte count, hematocrit, leukocyte count, differential leukocyte count, and the erythrocyte osmotic fragility.

G. Tissue and Fluid Chemistry of Rats Fed Paint Films Containing Different Concentrations of Lead

Tables 14-22 show the tissue and fluid chemistry of rats fed paint films containing different concentrations of lead. The serum electrophoresis, erythrocyte protoporphyrin, urinary coproporphyrin, and urinary delta-aminolevulinic acid were not altered by any of the diets.

The activity of the erythrocyte enzyme, delta-aminolevulinic acid dehydrase (ALAD), in the control, lead octoate (0.08%, 0.53%, and 2.05%) and lead chromate (0.42%, 1.95% and 12.43%) groups did not differ throughout the experiment (Tables 14-22). ALAD activity was depressed in rats fed 66.05% lead carbonate and 11.92% NBS lead paint. At 4 weeks there was a depression of 56% and 59% in these two groups, respectively. At 8 weeks the ALAD activity was depressed 44% in the 66.05% lead carbonate group, but the ALAD activity in the NBS lead paint group was not significantly different from control. At this time a sex difference was observed in both of these groups. The ALAD activity in the male rats remained depressed at the 4-week levels while the ALAD activities of the female rats returned to control levels. This sex difference may be linked to

fact that these female rats had matured and were probably beginning their menstrual cycles.

By 13 weeks the sex difference had disappeared and the ALAD activity in 66.05% lead carbonate group and the 11.92% NBS lead paint group were depressed 56% and 53%, respectively.

H. Lead Content in Tissues of Rats Fed Paint Films Containing Different Concentrations of Lead

Tables 23-27 show the lead content of blood, brain, liver, kidney, and bone from rats fed paint films containing different concentrations of lead. The values for the lead concentrations in blood at 4 and 8 weeks were omitted from Table 23 because it was discovered that these samples had been contaminated by their storage containers. At 13 weeks the blood lead levels in the rats fed lead octoate and the rats fed 0.42% and 1.95% lead chromate were not different from control. In the rats fed 12.43% lead chromate, 66.05% lead carbonate, and 11.92% NBS lead paint, blood levels were found to be elevated significantly.

At 4 weeks, tissue lead levels in brain, liver, and kidney were below the detection limits of our assay. The results were the same at 8 weeks with the exception of the NBS lead paint group. All four rats in this group showed detectable lead in their kidneys.

At 13 weeks, detectable levels of lead in the brain appeared only in the rats fed 11.92% NBS lead paint. In the liver detectable levels of lead were found in two rats fed 66.05% lead carbonate and nine rats fed 11.92% NBS lead paint. No lead was detected in the kidney of the

control group at 13 weeks. In contrast, lead was detected in the kidneys of 1-2 rats in the lead octoate groups and the 0.42% and 1.95% lead chromate groups. Lead was detected in the kidneys of three rats, eight rats, and 12 rats in the 12.43% lead chromate, 66.05% lead carbonate and 11.95% NBS lead paint groups, respectively.

The femurs of rats fed the control diet, lead octoate or lead chromate exhibited very little lead at 4 weeks. However, the femurs of rats fed 66.05% lead carbonate or 11.92% NBS lead paint contained 13.0 and 14.9 ug of lead/gm of bone.

At 8 weeks, the femurs of rats fed the control, lead octoate or lead chromate diets contained similar quantities of lead, ranging from 1.8-4.2 ug lead/gm bone. Femur lead in the 66.05% lead carbonate and 11.92% NBS groups were now 16.0 and 16.4 ug of lead/gm of bone. Both of these values were significantly above control.

After 13 weeks, the rats fed lead octoate or lead chromate had no more lead in their femurs than at 8 weeks and these levels were no higher than those in the control group. The 66.05% lead octoate group exhibited even higher levels of lead in their femurs (21.7 ug lead/gm bone) than at 8 weeks. The 11.92% NBS lead paint group had less lead in their femurs (10.9 ug lead/gm bone) than at 8 weeks. The lead levels in both of these groups was still significantly greater than those observed in the control rats.

I. Pathology of Rats Fed Paint Films Containing Different Concentrations of Lead

The pathology of rats fed the control, 2.05% lead octoate, 12.43% lead chromate 66.05% lead carbonate or 11.92% NBS lead paint diets for 13 weeks is shown in Tables 29-33. The control rats had a few naturally occurring lesions. Fourteen of these rats had mild to moderate lymphoid hyperplasia in the lungs, characteristic of early murine pneumonia. One of these rats had a moderate pneumonia and another one a mild emphysema.

Other lesions included an unspecific myocarditis in two rats; foci of subacute inflammation in the liver of two rats; a cross-section of a parasite (roundworm) in the colon of one rat; and foci of mononuclear cell infiltration in the kidneys of two rats. The bone marrow myeloid/erythroid cell ratios (M/E) of all rats were within normal limits.

The lesions observed in rats fed the 2.05% lead octoate diet were similar to the lesions seen in the control rats. Seven out of the 12 rats had mild to moderate lymphoid hyperplasia in the lungs and one had mild pneumonia. Four rats had mild foci of subacute inflammation in the liver characterized by a loss of several hepatic cord cells which were replaced by macrophages and lymphocytes. Another rat had a foci of inflammation in the interstitial tissue of the pancreas. The M/E ratios of these rats were normal.

The rats fed the 12.43% lead chromate diet had lesions similar to the lesions seen in the control rats. Seven of 12 rats had mild to moderate lymphoid hyperplasia, and one had mild pneumonia. Only one rat had a moderate unspecific myocarditis. The M/E ratios were normal.

The lesions seen in the rats fed the 11.92% NBS lead paint diet were similar to those lesions seen in the control rats. Eight of 12 rats had different degrees of lymphoid hyperplasia in the lungs. In one rat pneumonia was evident. Unspecific myocarditis occurred in 2 rats while foci of inflammation in the liver were observed in only one rat. The M/E ratios were normal.

In conclusion, the pathology seen in the control and treated rats were characteristic of those in a normal population of rats. No evidence of lead induced lesions were found.

Scientific Am
Interim

IV. DISCUSSION

The levels of lead fed to rats in the study were sufficient to produce significant elevation of body lead burden of rats and dogs if the lead is in the form of lead nitrate^{13/} or lead acetate.^{14/} Although it is questionable whether these levels are toxic to the rat,^{15/} they are quite toxic to the dog.^{13/} If we equate the amount of lead consumed by the average rat fed the 2.05% lead octoate, 12.43% lead (carbonate) and 11.92% NBS lead paint diets to a 14.6 kg 3-year old child, we find that the lead consumption would equal 28.6, 123.7, 503.0, and 113.9 mg of lead/day, respectively. On the basis of a 2% lead paint, the total paint intake would be 1.5, 6.2, 25.0, and 3.7 gm/day. In humans, these daily intakes can produce lead poisoning.^{13/} *at the level of a soluble form?*

There were no changes in hematology, urinalysis or serum proteins throughout the entire study. However, there was one effect on porphyrin metabolism. As early as 4 weeks, rats fed either the 66.05% lead carbonate diet or the 11.92% NBS lead paint diet exhibited 50% depression of erythrocyte ALAD activity. This effect appeared to be inversely proportional to the concentration of blood lead as has been reported by others.^{16,17/} However, there was one exception, blood lead in the 12.43% lead chromate group became elevated by the 13th week, but there was no depression of ALAD activity.

This discrepancy could be related to the duration of elevated blood lead or the age of the animal when blood lead becomes elevated whatever the explanation, the fact remains although erythrocyte ALAD activity was inhibited 50%, no changes in erythrocyte protoporphyrin urinary delta-aminolevulinic acid or urinary coproporphyrin levels were observed. Thus the effects on paraphyrin metabolism must have been mild.

The body lead burden of rats fed paint containing lead octoate or lead chromate were no different from control until 13 weeks. At this time lead was found in the kidneys of 1-3 rats on each diet and the blood lead levels in the 12.43% lead chromate group became significantly elevated. Since rats fed lead nitrate show elevated lead levels in blood, kidneys, bone and liver within 2 weeks,^{13/} we conclude that the lead octoate and lead chromate were not either readily absorbed or available for absorption.

Pathology was performed only on the rats that were fed the control, 2.05% lead octoate, 12.43% lead chromate, and 11.92% NBS lead paint diets for 13 weeks. All of these rats were relatively free of lesions, and those lesions which were observed, occur naturally in rat colonies.

V. CONCLUSIONS

This study shows that rats can consume large quantities of enamel paint containing as much as 2.05% lead as lead octoate or 12.43% lead as lead chromate for 13 weeks without any toxic effects. The lack of any marked change in the body lead-burden in these rats suggests that very little lead was absorbed from the gastrointestinal tract. Since there is considerable evidence that the rat can absorb lead in several forms, one must conclude that the lead in the enamel paints was not readily available for absorption. In conclusion, the results reported herein support the contention that lead is not as readily available in the newer enamel paints as in the older lead paints.

OK
not
determined
by
this
study
has
not
been
determined

Lead compound in
1.05% paint analyzed
As determined!

REFERENCES

1. Kehoe, R. A., The Metabolism of Lead in Man in Health and Disease;
Lecture II. J. Roy. Inst. Public Health Hyg., 24:101 (1961).
2. Chisholm, J. J., Ped. Clin., North Amer., 17:591 (1970).
3. King, B. G., Maximum Daily Intake of Lead Without Excessive Body
Lead-Burden in Children, Amer. J. Dis. Child., 122:337 (1971).
4. Selegson, D., Standard Methods of Clinical Chemistry, Academic Press,
Inc., New York, Vol. 2, p. 52 (1958).
5. Brecher, G., M. Schneiderman, and G. Z. William, Am. J. Clin. Path.,
26:1439 (1956).
6. Brecher, G., and M. Schneiderman, Am. J. Clin. Path., 20:1079
(1950).
7. Faulkner, W. R., and J. W. King, Eds., Manual of Clinical Laboratory
Procedures, 2nd Ed., p. 178 (1970).
8. Lichtman, H. C., and F. Feldman, J. Clin. Invest., 42:830 (1963).
9. Heller, S. R., Labbe, R. F., and Nutter, J., Clin. Chem., 17:525
(1971).
10. Davis, F. R., and S. L. Andelman, Arch. Envir. Health, 15:53 (1967).
11. Mauzerall, D., and S. Granick, J. Biol. Chem., 219:435 (1956).
12. Schlenker, F. S., and C. L. Kitchell, Am. J. Clin. Pathol.,
29:593 (1958).

13. Gage, J. C., and Litchfield, M. H., J. Oil. Col. Chem. Assoc.,
52:236 (1969).
14. Stowe, H. D., Goyer, R. A., Krigman, M. M., Wilson, M., and Cates,
M., Arch. Pathol., 95:106 (1973).
15. Hammond, P. B., In: Essays in Toxicology, Vol. 1, F. R. Blood
(Ed.), Academic Press, N.Y., 1969, p. 150.
16. Goldwater, L. J., Indust. Med., 41:13 (1972).
17. Millar, J. A., Battistini, V., Cumming, R. L. C., Carswell, F.,
and Goldberg, A., Lancet, Oct. 2:695 (1970).

Figure 1
Feed Consumption of Male Rats Fed Different
Concentrations of Lead in Faint Film





Figure 2
Feed Consumption of Female Rats Fed Different Concentrations of Lead in Paint Film







Figure 4
Bodyweights of female Rats Fed Different
Concentrations of Lead in Palm Oil



0007-SWP-036938

TABLE 1

CONCENTRATION OF LEAD IN PAINT FILM

<u>Paint Film</u>	<u>Theoretical Lead</u>	<u>Measured Lead Concentration</u>		
	<u>Concentration</u>	<u>Paint Companies</u>	<u>MRI</u>	<u>Average^{b/}</u>
	<u>(%)</u>	<u>(%)</u>	<u>(%)</u>	<u>(%)</u>
Control enamel	0	0.01	< 0.02	0.01
Lead octoate	0.06	0.08	0.07	0.08
Lead octoate	0.50	0.50	0.56	0.53
Lead octoate	2.00	1.90	2.20	2.05
Lead chromate	0.50	0.42	0.41	0.42
Lead chromate	2.00	1.80	2.10	1.95
Lead chromate	15.00	12.57	12.30	12.43
Lead carbonate	64.12	63.20 ^{c/}	68.90	66.05
NBS lead paint	-	11.87 ^{c/}	11.98	11.92 ^{c/}

^{a/} Measured by U.S. Department of Commerce, National Bureau of Standards, Washington, D.C.

^{b/} Average of paint companies and MRI values.

^{c/} Average of Bureau of Standards and MRI values.

TABLE 2

LEAD ANALYSES OF DIETS CONTAINING DIFFERENT PAINT FILMS

<u>Diet</u>	<u>Theoretical^{a/}</u> <u>(ug/gm)</u>	<u>Measured</u>	
		<u>Total^{b/}</u> <u>(ug/gm)</u>	<u>Paint Lead^{c/}</u> <u>(ug/gm)</u>
Control enamel plus rat chow	0.1	4.1 (6)	-
Lead octoate (0.08%) plus rat chow	0.8	5.2 (6)	1.1 (6)
Lead octoate (0.53%) plus rat chow	5.3	12.0 (6)	7.9 (6)
Lead octoate (2.05%) plus rat chow	20.5	27.2 (6)	23.1 (6)
Lead chromate (0.42%) plus rat chow	4.2	11.0 (6)	6.9 (6)
Lead chromate (1.95%) plus rat chow	19.5	23.5 (6)	19.4 (6)
Lead chromate (12.43%) plus rat chow	124.3	140.2 (6)	136.1 (6)
Lead carbonate (66.05%) plus rat chow	660.5	516.0 (5)	511.9 (5)
NBS lead based paint (11.92%) plus rat chow	119.2	124.0 (3)	119.9 (3)

a/ Theoretical concentration of lead in feed contributed by indicated paint film.

b/ Mean ug of lead/gm feed for number of samples shown in parenthesis.

c/ Mean ug of lead from paint films/gm feed for number of samples shown in parenthesis.

TABLE 3
MEAL AND TOTAL LEAD CONSUMPTION BY MALE RATS PER SAJIST FILM

	1	Treatment Week												Total/ meal
		2	3	4	5	6	7	8	9	10	11	12	13	
	meal	meal	meal	meal	meal	meal	meal	meal	meal	meal	meal	meal	meal	meal
Control	470.32/ (20)	572.0 (20)	638.1 (20)	731.0 (20)	749.8 (16)	769.2 (16)	806.3 (16)	778.9 (16)	751.9 (12)	786.4 (12)	838.0 (12)	879.4 (12)	838.0 (12)	9.0
Lead acetate 0.02%	597.0 (10)	715.3 (10)	882.2 (10)	920.9 (10)	932.3 (8)	948.2 (8)	979.2 (8)	964.6 (8)	997.4 (8)	1,011.9 (8)	1,044.7 (8)	1,019.2 (8)	1,042.9 (8)	12.2
0.5%	1,110.8 (10)	1,206.0 (10)	1,804.0 (10)	2,097.6 (10)	2,382.0 (8)	2,158.0 (8)	2,444.5 (8)	2,301.6 (8)	2,304.0 (8)	2,306.0 (8)	2,309.2 (8)	2,332.0 (8)	2,340.0 (8)	27.0
2.0%	3,179.7 (10)	6,017.4 (10)	4,549.6 (10)	6,500.3 (10)	5,100.0 (8)	5,004.6 (8)	5,064.6 (8)	5,407.4 (8)	5,305.3 (8)	5,369.3 (8)	5,276.4 (8)	5,404.3 (8)	5,331.6 (8)	64.3
Lead chromate 0.5%	1,501.3 (10)	1,107.1 (10)	1,006.3 (10)	3,017.4 (10)	2,163.7 (8)	2,140.3 (8)	2,449.6 (8)	2,156.0 (8)	2,032.9 (8)	2,363.9 (8)	2,237.0 (8)	2,310.0 (8)	2,502.3 (8)	21.3
1.0%	2,019.4 (10)	3,267.1 (10)	3,931.6 (10)	6,456.0 (10)	6,210.2 (8)	6,567.3 (8)	6,440.3 (8)	6,441.3 (8)	5,566.0 (8)	5,394.0 (8)	5,971.6 (8)	5,236.6 (8)	5,442.4 (8)	61.0
12.5%	14,771.0 (10)	20,413.1 (10)	23,944.0 (10)	25,010.0 (10)	27,262.9 (8)	26,040.0 (8)	27,945.9 (8)	23,553.0 (8)	23,001.5 (8)	24,044.3 (8)	20,727.1 (8)	20,931.1 (8)	20,717.0 (8)	314.0
Lead acetate 50.0%	37,436.8 (10)	73,179.6 (10)	83,363.2 (10)	94,636.4 (10)	90,607.6 (8)	97,005.2 (8)	101,497.2 (8)	95,356.0 (8)	102,219.6 (8)	102,942.0 (8)	104,025.6 (8)	102,942.0 (8)	104,748.0 (8)	1,222.7
MMS lead paint 11.5%	10,542.6 (10)	14,440.4 (10)	19,356.4 (10)	22,134.0 (10)	27,654.0 (8)	23,436.0 (8)	26,770.0 (8)	22,404.4 (8)	22,020.4 (8)	24,990.4 (8)	20,344.4 (8)	20,474.0 (8)	27,333.6 (8)	104.0

a/ Average ug of lead consumed/rat/week for the number of rats shown in parentheses.
b/ Average total mg of lead consumed/rat for the rats surviving 13 weeks.

TABLE 4
DAILY AND TOTAL LEAD CONSUMPTION BY TESTED DATE FOR PAINT FILMS

	Treatment Week													Total (lb)
	1	2	3	4	5	6	7	8	9	10	11	12	13	
Control	477.6 ^{1/} (20)	502.3 (20)	511.6 (20)	559.3 (20)	608.6 (16)	591.1 (16)	505.4 (16)	508.3 (16)	516.9 (12)	502.6 (10)	519.7 (11)	545.3 (11)	597.0 (11)	7.2 (11)
Lead acetate 0.001	342.4 (10)	655.2 (10)	644.3 (10)	610.8 (10)	707.5 (8)	695.2 (8)	691.6 (8)	704.7 (8)	606.3 (6)	717.1 (6)	755.5 (6)	604.3 (6)	702.5 (6)	8.8 (6)
0.33%	1,164.4 (10)	1,352.4 (10)	1,470.4 (10)	1,579.2 (10)	1,579.2 (8)	1,671.6 (8)	1,621.2 (8)	1,582.6 (8)	1,530.8 (8)	1,530.8 (8)	1,510.8 (6)	1,522.0 (6)	1,700.0 (6)	20.4 (6)
1.01%	3,179.7 (10)	4,017.4 (10)	4,545.6 (10)	4,900.5 (10)	5,140.0 (8)	5,064.6 (8)	5,064.6 (8)	5,407.4 (8)	5,306.2 (8)	5,269.3 (8)	5,216.1 (6)	5,446.3 (6)	5,321.6 (6)	64.4 (6)
Lead chromate 0.47%	1,185.8 (10)	1,316.7 (10)	1,401.4 (10)	1,447.6 (10)	1,406.1 (8)	1,506.2 (8)	1,510.5 (8)	1,509.2 (8)	1,540.8 (8)	1,540.8 (8)	1,479.3 (6)	1,401.6 (6)	1,340.8 (6)	19.7 (6)
1.53%	2,500.6 (10)	2,941.0 (10)	3,043.3 (10)	3,042.0 (10)	3,119.4 (8)	3,108.7 (8)	3,109.4 (8)	3,207.0 (8)	3,139.4 (8)	3,421.6 (6)	3,413.6 (6)	3,400.4 (6)	3,190.0 (6)	41.0 (6)
11.43%	16,320.5 (10)	17,176.5 (10)	16,470.2 (10)	17,255.3 (10)	21,002.0 (8)	19,922.4 (8)	19,630.0 (8)	20,110.7 (8)	18,446.6 (8)	18,450.3 (8)	18,042.9 (6)	17,959.6 (6)	23,040.0 (6)	254.7 (6)
Lead carbonate	31,200.4 (10)	39,950.2 (10)	41,446.0 (10)	43,912.4 (10)	44,670.0 (8)	46,099.6 (8)	47,905.6 (8)	44,654.0 (8)	46,812.0 (8)	43,912.4 (8)	45,790.4 (6)	49,310.4 (6)	46,059.6 (6)	811.8 (6)
MSB lead paint 11.93%	9,982.0 (10)	14,110.4 (10)	14,570.8 (10)	16,144.0 (10)	-3/ ^{2/} 10,480.4 (10)	-3/ ^{2/} 10,480.4 (8)	19,096.0 (8)	17,773.2 (8)	15,650.4 (8)	17,967.6 (8)	17,620.4 (6)	18,230.0 (6)	16,445.6 (6)	311.0 ^{2/} (6)

2/ Average mg of lead consumed/rat/week for the number of rats shown in parenthesis.

3/ Average total mg of lead consumed/rat for the rats surviving 13 weeks.

4/ Food consumption not measured.

5/ Total includes average Pb intake (17,316.6 mg/rat/week) for weeks 4 and 5 as an estimate for the week 3 intake.

TABLE 5

HEMATOLOGY OF RATS FED PAINT FILM
CONTAINING NO LEAD

Analyses	Treatment Week		
	4 (N=8) ^{a/}	8 (N=8)	13 (N=24)
Erythrocytes ($\times 10^6/\text{mm}^3$)	6.17 $\pm$ 0.14 ^{b/}	6.33 $\pm$ 0.19	6.42 $\pm$ 0.29
Reticulocytes, %	1.5 $\pm$ 0.7	0.8 $\pm$ 0.3	1.4 $\pm$ 0.1
Hematocrit, vol., %	46.1 $\pm$ 0.8	44.6 $\pm$ 1.2	44.6 $\pm$ 0.3
Hemoglobin, gm, %	14.0 $\pm$ 0.3	14.6 $\pm$ 0.3	16.0 $\pm$ 0.1
Leukocytes ($\times 10^3/\text{mm}^3$)	5.0 $\pm$ 0.5	7.6 $\pm$ 1.3	8.0 $\pm$ 0.6
Neutrophils, %	11.8 $\pm$ 2.9	21.7 $\pm$ 5.6	13.1 $\pm$ 1.5
Lymphocytes, %	86.3 $\pm$ 3.3	75.4 $\pm$ 5.3	84.1 $\pm$ 1.6
Bands, %	0.3 $\pm$ 0.2	0	0.1 $\pm$ 0.1
Eosinophils, %	0.9 $\pm$ 0.4	1.3 $\pm$ 0.4	0.5 $\pm$ 0.2
Basophils, %	0	0	0
Monocytes, %	1.1 $\pm$ 0.6	1.6 $\pm$ 0.6	2.1 $\pm$ 0.4
Atypical, %	0	0	0
Nucleated RBC, %	0	0	0.1 $\pm$ 0.1
Erythrocyte osmotic fragility [NaCl] ^{c/}	0.381 $\pm$ 0.004	0.391 $\pm$ 0.008	0.400 $\pm$ 0.004

^{a/} Number of rats per period.

^{b/} Mean $\pm$ standard error.

^{c/} Concentration of NaCl that hemolyzed 50% of the erythrocytes.

TABLE 6

HEMATOLOGY OF RATS FED PAINT FILM
CONTAINING 0.08% LEAD AS LEAD OCTOATE

Analyses	Treatment Week		
	4 (N=4) ^{a/}	8 (N=4)	13 (N=12)
Erythrocytes ($\times 10^6/\text{mm}^3$)	$6.14 \pm 0.32^{\text{b/}}$	5.33 ± 0.09	5.92 ± 0.39
Reticulocytes, %	0.9 ± 0.5	1.0 ± 0.5	1.4 ± 0.2
Hematocrit, vol., %	41.3 ± 0.9	46.0 ± 0.7	43.0 ± 0.5
Hemoglobin, gm, %	14.1 ± 0.4	15.4 ± 0.2	14.7 ± 0.8
Leukocytes ($\times 10^3/\text{mm}^3$)	5.1 ± 0.8	5.5 ± 0.7	7.7 ± 0.9
Neutrophils, %	24.5 ± 1.6	20.3 ± 4.7	16.3 ± 1.7
Lymphocytes, %	74.3 ± 1.8	76.3 ± 4.7	80.9 ± 1.8
Bands, %	0	0	0.3 ± 0.2
Eosinophils, %	0.8 ± 0.3	1.5 ± 0.6	0.7 ± 0.3
Basophils, %	0	0	0
Monocytes, %	0.5 ± 0.5	2.0 ± 0.8	1.9 ± 0.5
Atypical, %	0	0	0
Nucleated RBC, %	0	0	0
Erythrocyte osmotic fragility [NaCl] ^{c/}	0.381 ± 0.005	0.410 ± 0.007	0.395 ± 0.005

^{a/} Number of rats per period.

^{b/} Mean $\pm$ standard error.

^{c/} Concentration of NaCl that hemolyzed 50% of the erythrocytes.

TABLE 7

HEMATOLOGY OF RATS FED PAINT FILMS
CONTAINING 0.53% LEAD AS LEAD OCTOATE

Analyses	Treatment Week		
	4 (N=4) ^{a/}	8 (N=4)	13 (N=12)
Erythrocytes ($\times 10^6/\text{mm}^3$)	6.27 $\pm$ 0.2 ^{b/}	6.04 $\pm$ 0.31	5.6 $\pm$ 0.4
Reticulocytes, %	0.9 $\pm$ 0.7	0.9 $\pm$ 0.4	1.6 $\pm$ 0.2
Hematocrit, vol., %	42.3 $\pm$ 0.7	43.8 $\pm$ 0.5	43 $\pm$ 0.7
Hemoglobin, gm, %	14.2 $\pm$ 0.2	14.9 $\pm$ 0.3	15.3 $\pm$ 0.3
Leukocytes ($\times 10^3/\text{mm}^3$)	5.0 $\pm$ 1.4	7.5 $\pm$ 2.8	8.3 $\pm$ 1.0
Neutrophils, %	15.8 $\pm$ 3.8	19.3 $\pm$ 3.8	16.2 $\pm$ 1.9
Lymphocytes, %	83.0 $\pm$ 4.4	77.5 $\pm$ 3.8	81.4 $\pm$ 1.8
Bands, %	0	0	0.1 $\pm$ 0.1
Eosinophils, %	0.8 $\pm$ 0.3	1.0 $\pm$ 0.4	0.8 $\pm$ 0.3
Basophils, %	0	0	0
Monocytes, %	0.5 $\pm$ 0.5	2.3 $\pm$ 0.5	1.6 $\pm$ 0.3
Atypical, %	0	0	0
Nucleated RBC, %	0	0	0
Erythrocyte osmotic fragility [NaCl] ^{c/}	0.383 $\pm$ 0.006	0.388 $\pm$ 0.012	0.401 $\pm$ 0.006

a/ Number of rats per period.

b/ Mean $\pm$ standard error.

c/ Concentration of NaCl that hemolyzed 50% of the erythrocytes.

TABLE 8

HEMATOLOGY OF RATS FED PAINT FILM
CONTAINING 2.05% LEAD AS LEAD OCTOATE

Analyses	Treatment Week		
	4 (N=4) ^{a/}	8 (N=4)	13 (N=12)
Erythrocytes ($\times 10^6/\text{mm}^3$)	$6.30 \pm 0.2^b/$	6.46 ± 0.08	6.72 ± 0.42
Reticulocytes, %	1.1 ± 0.6	0.7 ± 0.1	2.0 ± 0.6
Hematocrit, vol., %	41.8 ± 0.5	45.0 ± 0.4	43.4 ± 0.8
Hemoglobin, gm, %	14.4 ± 0.2	15.5 ± 0.2	15.6 ± 0.3
Leukocytes ($\times 10^3/\text{mm}^3$)	5.1 ± 1.1	6.7 ± 1.7	8.1 ± 0.7
Neutrophils, %	13.5 ± 3.0	13.0 ± 2.1	14.1 ± 2.6
Lymphocytes, %	85.8 ± 3.2	85.0 ± 2.3	82.8 ± 2.8
Bands, %	0	0	0
Eosinophils, %	0.8 ± 0.5	0.3 ± 0.3	0.9 ± 0.3
Basophils, %	0	0	0
Monocytes, %	0	1.8 ± 0.9	2.2 ± 0.7
Atypical, %	0	0	0
Nucleated RBC, %	0	0	0
Erythrocyte osmotic fragility [NaCl] ^{c/}	0.381 ± 0.006	0.399 ± 0.008	0.393 ± 0.006

^{a/} Number of rats per period.

^{b/} Mean $\pm$ standard error.

^{c/} Concentration of NaCl that hemolyzed 50% of the erythrocytes.

TABLE 9

HEMATOLOGY OF RATS FED PAINT FILM
CONTAINING 0.42% LEAD AS LEAD CHROMATE

Analyses	Treatment Weeks		
	4 (N=6) ^{a/}	8 (N=6)	13 (N=12)
Erythrocytes ($\times 10^6/\text{mm}^3$)	$6.71 \pm 0.16^{\text{b/}}$	6.18 ± 0.16	6.6 ± 0.3
Reticulocytes, %	0.9 ± 0.8	0.8 ± 0.2	1.6 ± 0.2
Hematocrit, vol., %	43.3 ± 0.6	43.0 ± 1.1	43.6 ± 0.7
Hemoglobin, gm, %	14.7 ± 0.4	15.1 ± 0.3	15.6 ± 0.3
Leukocytes ($\times 10^3/\text{mm}^3$)	5.1 ± 0.6	5.1 ± 0.5	6.9 ± 0.7
Neutrophils, %	8.5 ± 1.7	19.0 ± 2.0	14.3 ± 2.1
Lymphocytes, %	90.3 ± 2.0	79.5 ± 1.3	83.4 ± 2.0
Bands, %	0	0	0
Eosinophils, %	0	0.8 ± 0.8	1.2 ± 0.4
Basophils, %	0	0	0
Monocytes, %	1.3 ± 0.8	0.8 ± 0.8	1.2 ± 0.4
Atypical, %	0	0	0
Nucleated RBC, %	0	0.3 ± 0.3	0
Erythrocyte osmotic fragility, [NaCl] ^{c/}	0.383 ± 0.003	0.395 ± 0.12	0.392 ± 0.007

^{a/} Number of rats per period, except where indicated otherwise.

^{b/} Mean $\pm$ standard error.

^{c/} Concentration of NaCl that hemolyzed 50% of the erythrocytes.

TABLE 10

HEMATOLOGY OF RATS FED PAINT FILM
CONTAINING 1.95% LEAD AS LEAD CHROMATE

Analyses	Treatment Week		
	4 (N=4) ^{a/}	8 (N=4)	13 (N=12)
Erythrocytes ($\times 10^6/\text{mm}^3$)	6.17 $\pm$ 0.05 ^{b/}	6.21 $\pm$ 0.13	6.37 $\pm$ 0.52
Reticulocytes, %	0.8 $\pm$ 0.4	0.8 $\pm$ 0.3	1.2 $\pm$ 0.2
Hematocrit, vol. %	40.8 $\pm$ 0.6	44.3 $\pm$ 0.3	44.1 $\pm$ 0.7
Hemoglobin, gm, %	13.9 $\pm$ 0.2	15.3 $\pm$ 0.3	16.2 $\pm$ 0.4
Leukocytes ($\times 10^3/\text{mm}^3$)	4.9 $\pm$ 0.8	6.1 $\pm$ 1.2	8.2 $\pm$ 1.0
Neutrophils, %	10.0 $\pm$ 2.4	18.5 $\pm$ 6.6	11.4 $\pm$ 2.3
Lymphocytes, %	92.0 $\pm$ 1.8	80.5 $\pm$ 6.6	85.9 $\pm$ 2.2
Bands, %	0	0	0
Eosinophils, %	0.5 $\pm$ 0.3	0.3 $\pm$ 0.3	1.5 $\pm$ 0.5
Basophils, %	0	0	0
Monocytes, %	0	0.8 0.3	1.2 $\pm$ 0.3
Atypical, %	0	0	0
Nucleated RBC, %	0	0	0
Erythrocyte osmotic fragility, [NaCl] ^{c/}	0.383 $\pm$ 0.007	0.385 $\pm$ 0.007	0.404 $\pm$ 0.005

^{a/} Number of rats per period.

^{b/} Mean $\pm$ standard error.

^{c/} Concentration of NaCl that hemolyzed 50% of the erythrocytes.

TABLE 11

HEMATOLOGY OF RATS FED PAINT FILM
CONTAINING 12.43% LEAD AS LEAD CHROMATE

Analyses	Treatment Week		
	4 (N=4) ^{a/}	8 (N=4)	13 (N=12)
Erythrocytes ($\times 10^6/\text{mm}^3$)	6.32 $\pm$ 0.14 ^{b/}	6.23 $\pm$ 0.18	6.61 $\pm$ 0.38
Reticulocytes, %	1.95 $\pm$ 1.28	0.9 $\pm$ 0.1	1.6 $\pm$ 0.2
Hematocrit, vol., %	41.3 $\pm$ 1.1	43.8 $\pm$ 0.5	44.1 $\pm$ 0.7
Hemoglobin, gm, %	13.9 $\pm$ 0.4	15.2 $\pm$ 0.1	15.8 $\pm$ 0.2
Leukocytes ($\times 10^3/\text{mm}^3$)	4.8 $\pm$ 0.5	5.3 $\pm$ 1.8	7.6 $\pm$ 0.6
Neutrophils, %	21.5 $\pm$ 6.3	15.0 $\pm$ 2.1	12.8 $\pm$ 1.6
Lymphocytes, %	78.0 $\pm$ 6.0	82.8 $\pm$ 2.0	84.8 $\pm$ 1.7
Bands, %	0	0	0.1 $\pm$ 0.1
Eosinophils, %	0.5 $\pm$ 0.5	1.0 $\pm$ 0	1.0 $\pm$ 0.4
Basophils, %	0	0	0
Monocytes, %	0	1.3 $\pm$ 0.3	1.3 $\pm$ 0.3
Atypical, %	0	0	0
Nucleated RBC, %	0	0.3 $\pm$ 0.3	0
Erythrocyte osmotic fragility [NaCl] ^{c/}	0.384 $\pm$ 0.007	0.385 $\pm$ 0.010	0.396 $\pm$ 0.006

^{a/} Number of rats per period.

^{b/} Mean $\pm$ standard error.

^{c/} Concentration of NaCl that hemolyzed 50% of the erythrocytes.

TABLE 12

HEMATOLOGY OF RATS FED PAINT FILM
CONTAINING 66.05% LEAD AS LEAD CARBONATE

Analyses	Treatment Weeks		
	4 (N=3) ^{a/}	8 (N=4)	13 (N=12)
Erythrocytes ($\times 10^6/\text{mm}^3$)	$5.94 \pm 0.56^{\text{b/}}$	6.13 ± 0.06	6.60 ± 0.37
Reticulocytes, %	0.4 ± 0.2	1.1 ± 0.3	2.9 ± 1.3
Hematocrit, vol., %	41.3 ± 0.9	44.0 ± 1.5	44.7 ± 1.2
Hemoglobin, gm, %	13.9 ± 0.5	15.0 ± 0.2	15.6 ± 0.6
Leukocytes ($\times 10^3/\text{mm}^3$)	4.3 ± 1.2	5.0 ± 0.9	7.7 ± 0.5
Neutrophils, %	14.3 ± 4.1	15.0 ± 4.1	15.2 ± 2.6
Lymphocytes, %	82.0 ± 5.6	84.0 ± 4.2	82.7 ± 2.7
Bands, %	0	0	0.2 ± 0.1
Eosinophils, %	0.7 ± 0.7	0.5 ± 0.5	0.9 ± 0.3
Basophils, %	0	0	0
Monocytes, %	3.0 ± 2.0	0.5 ± 0.3	1.0 ± 0.3
Atypical, %	0	0	0
Nucleated RBC, %	0	0	0.1 ± 0.1
Erythrocyte osmotic fragility [NaCl] ^{c/}	0.388 ± 0.009	0.394 ± 0.012	0.397 ± 0.005

^{a/} Number of rats per period.

^{b/} Mean $\pm$ standard error.

^{c/} Concentration of NaCl that hemolyzed 50% of the erythrocytes.

TABLE 13

HEMATOLOGY OF RATS FED LEAD BASED
PAINT CONTAINING 11.92% LEAD

Analyses	Treatment Week		
	4 (N=4) ^{a/}	8 (N=4)	13 (N=12)
Erythrocytes ($\times 10^6/\text{mm}^3$)	5.72 ± 0.15 ^{b/}	4.49 ± 0.83	5.63 ± 0.34
Reticulocytes, %	2.5 ± 0.3	1.0 ± 0.4	1.3 ± 0.1
Hematocrit, vol., %	39.7 ± 0.8	42.8 ± 1.1	42.9 ± 0.4
Hemoglobin, gm, %	13.6 ± 0.2 ^{c/}	14.7 ± 0.3	14.7 ± 0.1
Leukocytes ($\times 10^3/\text{mm}^3$)	9.7 ± 1.4	6.7 ± 1.2	7.0 ± 0.5
Neutrophils, %	9.5 ± 2.1	7.0 ± 1.7	14.2 ± 2.1
Lymphocytes, %	90.2 ± 2.3	92.0 ± 2.1	83.6 ± 2.2
Bands, %	0.3 ± 0.3	0	0
Eosinophils, %	0	0	1.4 ± 0.4
Basophils, %	0	0	0
Monocytes, %	0	1.0 ± 0.7	0.8 ± 0.3
Atypical, %	0	0	0
Nucleated RBC, %	0	0	0.1 ± 0.1
Erythrocyte osmotic fragility [NaCl] ^{d/}	0.389 ± 0.006	0.412 ± 0.006	0.408 ± 0.005

a/ Number of rats per period, except where indicated otherwise.

b/ Mean $\pm$ standard error.

c/ Two rats.

d/ Concentration of NaCl that hemolyzes 50% of the erythrocytes.

TABLE 14

TISSUE AND FLUID CHEMISTRY OF RATS FED PAINT FILM
CONTAINING NO LEAD

Analyses	Treatment Week		
	4 (N=8) ^{a/}	8 (N=8)	13 (N=24)
Serum electrophoresis			
Albumin, %	43 ± 1 ^{b/}	43 ± 2	48 ± 1
Alpha 1 globulin, %	28 ± 1	23 ± 1	21 ± 1
Alpha 2 globulin, %	5 ± 1	5 ± 1	5 ± 1
Beta globulin, %	20 ± 1	20 ± 1	21 ± 1
Gamma globulin, %	6 ± 1	10 ± 1	6 ± 1
Total protein, mg %	5.4 ± 0.1	6.2 ± 0.2	6.3 ± 0.1
Albumin/globulin ratio	0.76 ± 0.03	0.77 ± 0.07	0.93 ± 0.04
Erythrocyte			
ALAD, μ mol. PBG/	Males: 23.2 ± 2.3 ^{c/}	15.1 ± 2.3 ^{c/}	14.5 ± 1.4 ^{d/}
100 ml RBC/hr	Females: 16.6 ± 3.1 ^{c/}	16.3 ± 2.2 ^{c/}	13.4 ± 0.7 ^{d/}
	Total: 19.9 ± 2.2	15.8 ± 1.5	15.0 ± 1.0
Protoporphyrin,	19.6 ± 1.6	23.5 ± 1.9	29.2 ± 2.0
μ g/100 ml RBC			
Urine			
Coproporphyrin,	2.5 ± 1.2	3.3 ± 7.4	6.0 ± 1.9
μ g/24 hr			
ALA,	78.0 ± 7.4	83.3 ± 11.0	43.4 ± 5.0
μ g/24 hr			

^{a/} Number of rats per period, except where indicated otherwise.

^{b/} Mean ± standard error.

^{c/} Four rats.

^{d/} Twelve rats.

TABLE 15

HEMISTRY OF RATS FED PAINT FILM
1% LEAD AS LEAD OCTOATE

Treatment Week		
4	8	13
<u>(N=4)</u> ^{a/}	<u>(N=4)</u>	<u>(N=12)</u>
0.5 ± 1 ^{b/}	44 ± 3	48 ± 1
5 ± 1	20 ± 2	20 ± 1
5 ± 1	4 ± 1	5 ± 1
2 ± 1	20 ± 1	22 ± 1
3 ± 1	12 ± 1	7 ± 1
0.9 ± 0.1	6.3 ± 0.2	6.2 ± 0.1
0.6 ± 0.02	0.79 ± 0.08	0.93 ± 0.04
13.3 ± 7.7 ^{e/}	14.9 ± 1.3 ^{e/}	12.0 ± 2.2 ^{d/}
15.5 ± 4.4 ^{e/}	18.7 ± 0.3 ^{e/}	16.1 ± 1.3 ^{d/}
12.2 ± 3.9	16.8 ± 1.2	14.0 ± 1.4
11.4 ± 2.4	24.2 ± 1.6	22.8 ± 2.5
0.2 ± 0.2	4.8 ± 2.4	2.9 ± 1.4
14 ± 16.6	107.6 ± 34.1	35.0 ± 6.7

except where indicated otherwise.

TABLE 16

TISSUE AND BODY FLUID CHEMISTRY OF RATS FED PAINT
FILM CONTAINING 0.53% LEAD AS LEAD OCTOATE

Analyses	Treatment Week		
	4 (N=4) ^{a/}	8 (N=4)	13 (N=12)
Serum electrophoresis			
Albumin, %	41 ± 2 ^{b/}	41 ± 4	50 ± 1
Alpha 1 globulin, %	26 ± 1	23 ± 3	20 ± 1
Alpha 2 globulin, %	5 ± 1	6 ± 2	4 ± 1
Beta globulin, %	21 ± 1	20 ± 1	20 ± 1
Gamma globulin, %	8 ± 1	11 ± 2	7 ± 1
Total protein, mg %	5.7 ± 0.1	6.3 ± 0.1	6.0 ± 0.1
Albumin/globulin ratio	0.68 ± 0.04	0.70 ± 0.13	1.00 ± 0.07
Erythrocytes			
ALAD, μ mol. PBG/ Males: 21.4 ± 6.3 ^{c/}		10.3 ± 0.8 ^{c/}	12.5 ± 0.7 ^{d/}
100 ml RBC/hr Females: 15.5 ± 2.7 ^{c/}		15.3 ± 0.2 ^{c/}	17.0 ± 1.2 ^{d/}
Total: 18.4 ± 3.3		12.8 ± 1.5	14.7 ± 0.9
Protoporphyrin, 23.1 ± 1.0		21.7 ± 1.1	27.7 ± 3.3
ug/100 ml RBC			
Urine			
Coproporphyrin, 10.7 ± 5.9		1.1 ± 0.6	4.9 ± 2.4
ug/24 hr			
ALA, 97.4 ± 19.2		101.4 ± 30.1	50.5 ± 6.6
ug/24 hr			

^{a/} Number of rats per group, except where indicated otherwise.

^{b/} Mean ± standard error.

^{c/} Two rats.

^{d/} Six rats.

TABLE 17

TISSUE AND BODY FLUID CHEMISTRY OF RATS FED PAINT
FILM CONTAINING 2.05% LEAD AS LEAD OCTOATE

Analyses	Treatment Week		
	4 (N=6) ^{a/}	8 (N=6)	13 (N=12)
Serum electrophoresis			
Albumin, %	41 ± 3 ^{b/}	46 ± 2	48 ± 2
Alpha 1 globulin, %	27 ± 2	22 ± 1	20 ± 1
Alpha 2 globulin, %	4 ± 1	4 ± 1	6 ± 1
Beta globulin, %	20 ± 1	19 ± 1	21 ± 1
Gamma globulin, %	9 ± 1	9 ± 1	6 ± 1
Total protein, mg %	5.7 ± 0.2	6.0 ± 0.1	6.2 ± 0.1
Albumin/globulin ratio	0.70 ± 0.07	0.85 ± 0.07	0.96 ± 0.07
Erythrocytes			
ALAD, μ mol. PBG/ 100 ml RBC/hr	Males: 24.1 ± 4.0 ^{c/} Females: 18.7 ± 1.0 ^{c/} Total: 21.4 ± 2.3	12.1 ± 1.7 ^{c/} 11.3 ± 0.2 ^{c/} 11.7 ± 0.7	11.4 ± 0.8 ^{d/} 16.3 ± 3.9 ^{d/} 13.8 ± 2.0
Protoporphyrin, μ g/100 ml RBC	29.5 ± 2.8	30.8 ± 2.5	24.0 ± 2.2
Urine			
Coproporphyrin, μ g/24 hr	1.6 ± 0.6	3.0 ± 0.8	7.7 ± 2.9
ALA, μ g/24 hr	80.7 ± 17.3	82.7 ± 10.4	64.5 ± 8.8

^{a/} Number of rats per period, except where indicated otherwise.

^{b/} Mean ± standard error.

^{c/} Two rats.

^{d/} Six rats.

TABLE 18

TISSUE AND BODY FLUID CHEMISTRY OF RATS FED PAINT
FILM CONTAINING 0.3% LEAD AS LEAD CHROMATE

Analyses	Treatment Week		
	4 (N=4) ^{a/}	8 (N=4)	13 (N=12)
Serum electrophoresis			
Albumin, %	42 ± 2 ^{b/}	44 ± 2	47 ± 1
Alpha 1 globulin, %	28 ± 1	20 ± 2	22 ± 1
Alpha 2 globulin, %	4 ± 1	4 ± 1	5 ± 1
Beta globulin, %	19 ± 1	19 ± 1	21 ± 1
Gamma globulin, %	8 ± 3	12 ± 2	6 ± 1
Total protein, mg %	5.8 ± 0.2	6.0 ± 0.2	6.1 ± 0.1
Albumin/globulin ratio	0.71 ± 0.06	0.80 ± 0.08	0.90 ± 0.03
Erythrocyte			
ALAD, μ mol. PBG/	Males: 23.4 ± 3.5 ^{c/}	14.1 ± 0.6 ^{c/}	15.7 ± 1.5 ^{d/}
100 ml RBC/hr	Females: 13.4 ± 0.9 ^{c/}	17.4 ± 0.1 ^{c/}	16.5 ± 1.4 ^{d/}
	Total: 18.4 ± 3.3	15.7 ± 1.0	16.1 ± 1.0
Protoporphyrin,	27.3 ± 2.8	28.3 ± 0.6	27.9 ± 2.1
μ g/100 ml TBC			
Urine			
Coproporphyrin,	8.2 ± 6.7	1.5 ± 0.7	6.3 ± 2.2
μ g/24 hr			
ALA,	69.3 ± 6.6	65.2 ± 17.4	85.3 ± 31.4
μ g/24 hr			

^{a/} Number of rats per period, except where indicated otherwise.

^{b/} Mean ± standard error.

^{c/} Two rats.

^{d/} Six rats.

TABLE 19

TISSUE AND BODY FLUID CHEMISTRY OF RATS FEED PAINT
FILM CONTAINING 1.95% LEAD AS LEAD CHROMATE

Analyses	Treatment Week		
	4 (N=6) ^{a/}	8 (N=6)	13 (N=12)
Serum electrophoresis			
Albumin, %	41 ± 2 ^{b/}	45 ± 1	47 ± 2
Alpha 1 globulin, %	25 ± 1	19 ± 1	19 ± 2
Alpha 2 globulin, %	4 ± 1	4 ± 1	5 ± 1
Beta globulin, %	20 ± 1	20 ± 1	23 ± 1
Gamma globulin, %	9 ± 1	11 ± 1	6 ± 1
Total protein, mg %	5.7 ± 0.1	6.20 ± 0.04	6.4 ± 0.1
Albumin/globulin ratio	0.73 ± 0.06	0.81 ± 0.04	0.94 ± 0.08
Erythrocyte			
ALAD, umol. PBG/ Males: 29.5 ± 3.1 ^{c/}		12.3 ± 1.0 ^{c/}	13.3 ± 1.4 ^{d/}
100 ml RBC/hr Females: 20.4 ± 4.3 ^{c/}		14.7 ± 0.6 ^{c/}	12.9 ± 2.3 ^{d/}
Total: 24.9 ± 3.4		13.5 ± 0.8	13.1 ± 1.3
Protoporphyrin, ug/100 ml RBC	26.5 ± 4.7	26.8 ± 1.8	27.4 ± 1.5
Urine			
Coproporphyrin, ug/24 hr	0.9 ± 0.1	7.0 ± 6.4	4.4 ± 1.9
ALA, ug/24 hr	80.8 ± 7.2	67.7 ± 6.8	66.5 ± 17.5

^{a/} Number of rats per period, except where indicated otherwise.

^{b/} Mean ± standard error.

^{c/} Two rats.

^{d/} Six rats.

TABLE 20

TISSUE AND BODY FLUID CHEMISTRY OF RATS FED PAINT
FILM CONTAINING 12.43% LEAD AS LEAD CHROMATE

Analyses	Treatment Week		
	4 (N=4) ^{a/}	8 (N=4)	13 (N=12)
Serum electrophoresis			
Albumin, %	40 ± 2 ^{b/}	47 ± 3	46 ± 2
Alpha 1 globulin, %	27 ± 1	21 ± 1	21 ± 1
Alpha 2 globulin, %	3 ± 1	4 ± 1	5 ± 1
Beta globulin, %	22 ± 1	18 ± 1	22 ± 1
Gamma globulin, %	8 ± 1	9 ± 2	6 ± 1
Total protein, mg %	5.6 ± 0.1	6.1 ± 0.1	6.4 ± 0.1
Albumin/globulin ratio	0.68 ± 0.05	0.90 ± 0.12	0.88 ± 0.06
Erythrocyte			
ALAD, umol. PBG/ Males:	26.4 ± 7.4 ^{c/}	16.5 ± 1.4 ^{c/}	12.6 ± 1.6 ^{d/}
100 ml RBC/hr Females:	14.3 ± 3.2 ^{c/}	13.4 ± 3.8 ^{c/}	13.7 ± 1.1 ^{d/}
Total:	20.3 ± 4.9	15.0 ± 1.9	13.1 ± 1.0
Protoporphyrin, ug/100 ml RBC	25.5 ± 1.6	24.2 ± 2.6	24.6 ± 2.0
Urine			
Coproporphyrin, ug/24 hr	2.6 ± 1.8	0.9 ± 0.1	2.6 ± 0.8
ALA, ug/24 hr	89.7 ± 19.3	78.4 ± 6.8	49.7 ± 12.5

^{a/} Number of rats per period, except where indicated otherwise.

^{b/} Mean ± standard error.

^{c/} Two rats.

^{d/} Six rats.

TABLE 21

TISSUE AND BODY FLUID CHEMISTRY OF RATS FED PAINT
FILM CONTAINING 66.05% LEAD AS LEAD CARBONATE

Analyses	Treatment Week		
	4 (N=3) ^{a/}	8 (N=4)	13 (N=12)
Serum electrophoresis			
Albumin, %	41 ± 6	45 ± 1	47 ± 2
Alpha 1 globulin, %	27 ± 1	21 ± 2	22 ± 1
Alpha 2 globulin, %	3 ± 1	4 ± 1	5 ± 1
Beta globulin, %	21 ± 2	19 ± 1	20 ± 1
Gamma globulin, %	8 ± 2	10 ± 1	7 ± 1
Total protein, mg %	5.4 ± 0.2	6.0 ± 0.06	6.3 ± 0.1
Albumin/globulin ratio	0.73 ± 0.17	0.82 ± 0.04	0.86 ± 0.06
Erythrocyte			
ALAD, μ mol. PBG/ Males:	12.3 ^{e/}	4.4 ± 0.1 ^{d/}	5.4 = 1.0 ^{e/}
100 ml RBC/hr Females:	9.1 ± 2.1 ^{d/}	12.4 ± 0.4 ^{d/}	7.4 = 0.8 ^{e/}
	10.2 ± 1.6 ^{f/}	8.4 ± 2.3 ^{f/}	6.4 = 0.7 ^{f/}
Protoporphyrin, μg/100 ml RBC	23.2 ± 0.9	20.4 ± 2.1	27.2 ± 6.8
Urine (N=4)			
Coproporphyrin, μg/24 hr	2.0 ± 0.7	0.8 ± 0.3	6.7 ± 3.2
ALA, μg/24 hr	90.3 ± 27.7	72.3 ± 13.4	42.2 ± 8.8

a/ Number of rats per period, except where indicated otherwise.

b/ Mean ± standard error.

c/ One rat.

d/ Two rats.

e/ Six rats.

f/ Significantly different from control (P < 0.05) as shown by Dunnett's multiple comparison test following an analysis of variance.

TABLE 22

TISSUE AND BODY FLUID CHEMISTRY OF RATS FED LEAD
BASED PAINT CONTAINING 11.9% LEAD

Analysis	Treatment Week		
	4 (N=6) ^{a/}	8 (N=6)	13 (N=12)
Serum electrophoresis			
Albumin, %	52 ± 1 ^{b/}	44 ± 2	44 ± 2
Alpha 1 globulin, %	21 ± 1	21 ± 2	24 ± 1
Alpha 2 globulin, %	4 ± 1	8 ± 5	7 ± 1
Beta globulin, %	20 ± 1	21 ± 3	21 ± 1
Gamma globulin, %	3 ± 1	6 ± 1	5 ± 1
Total protein, mg %	5.4 ± 0.2	6.0 ± 0.2	6.1 ± 0.1
Albumin/globulin ratio	1.1 ± 0.03	0.78 ± 0.06	0.82 ± 0.06
Erythrocyte			
ALAD, μ mol. PBC/ Males:	8.5 ± 3.0 ^{c/}	5.0 ± 0.3 ^{c/}	4.9 ± 0.6 ^{d/}
100 ml RBC/hr Females:	10.1 ± 0.1 ^{c/}	15.0 ± 5.1 ^{c/}	8.8 ± 1.0 ^{d/}
Total:	9.3 ± 2.1 ^{c/}	10.0 ± 3.6	6.9 ± 0.8 ^{e/}
Protoporphyrin, μ g/100 ml RBC	14.8 ± 4.3	24.5 ± 1.8	26.3 ± 1.2
Urine (N=4)			
Coproporphyrin, μ g/24 hr	2.0 ± 0.8	6.8 ± 4.9	4.6 ± 2.1
ALA, μ g/24 hr	110.1 ± 21.5	72.0 ± 13.6	51.3 ± 3.6

^{a/} Number of rats per period, except where indicated otherwise.

^{b/} Mean ± standard error.

^{c/} Two rats.

^{d/} Six rats.

^{e/} Significantly different from control ($P < 0.05$) as indicated by Dunnnett's multiple comparison test following an analysis of variance.

TABLE 23

LEAD CONTENT OF WHOLE BLOOD

<u>Diet</u>	<u>Treatment Week</u>		
	<u>4^a/</u>	<u>8^a/</u>	<u>13</u>
Control			13.1 ± 0.9 (16) ^{b/}
0.08% Lead Octoate			12.8 ± 0.8 (8)
0.53% Lead Octoate			16.4 ± 2.4 (7)
2.05% Lead Octoate			14.5 ± 1.5 (8)
0.42% Lead Chromate			13.2 ± 1.0 (8)
1.95% Lead Chromate			12.0 ± 1.4 (8)
12.43% Lead Chromate			19.4 ± 2.2 (8) ^{c/}
66.05% Lead Carbonate			24.9 ± 2.2 (8) ^{c/}
11.92% Lead Paint			23.9 ± 1.3 (12) ^{c/}

^{a/} Values deleted due to contamination.

^{b/} Average μ g of lead/100 ml blood $\pm$ Standard Error of number of rats in parenthesis.

^{c/} Significantly different from control ($P < .05$) as shown by Dunnett's multiple comparison test following an analysis of variance.

TABLE 24
LEAD CONTENT OF BRAIN

Diet	Treatment Week		
	4	8	13
Control	n.d. ^{a/}	n.d.	n.d.
0.08% Lead Octoate	n.d.	n.d.	n.d.
0.53% Lead Octoate	n.d.	n.d.	n.d.
2.05% Lead Octoate	n.d.	n.d.	n.d.
0.42% Lead Chromate	n.d.	n.d.	n.d.
1.95% Lead Chromate	n.d.	n.d.	n.d.
12.43% Lead Chromate	n.d.	n.d.	n.d.
66.05% Lead Carbonate	n.d.	n.d.	n.d.
11.92% Lead Pain	n.d.	n.d.	0.2 ± 0.1 ^{b/} (7)

^{a/} Not detectable. Below sensitivity of 0.07 ug/gm of brain.

^{b/} Average ug of lead/gm brain ± standard error number of rats shown in parenthesis.

TABLE 25

LEAD CONTENT OF LIVER

<u>Diet</u>	<u>Treatment Week</u>		
	<u>4</u>	<u>8</u>	<u>13</u>
Control	n.d. ^{a/}	n.d.	n.d.
0.08% Lead Octoate	n.d.	n.d.	n.d.
0.53% Lead Octoate	n.d.	n.d.	n.d.
2.05% Lead Octoate	n.d.	n.d.	n.d.
0.42% Lead Chromate	n.d.	n.d.	n.d.
1.95% Lead Chromate	n.d.	n.d.	n.d.
12.43% Lead Chromate	n.d.	n.d.	n.d.
66.05% Lead Carbonate	n.d.	n.d.	1.1 ± 0.6 ^{b/} (2)
11.92% Lead Paint	n.d.	n.d.	0.4 ± 0.1 (9)

^{a/} Not detectable. Below sensitivity of 0.07 µg/gm liver.

^{b/} Values are average µg of lead/gm liver ± standard error of number of rats shown in parenthesis.

TABLE 26

LEAD CONTENT OF KIDNEY

<u>Diet</u>	<u>Treatment Week</u>		
	<u>4</u>	<u>8</u>	<u>12</u>
Control	n.d. ^{a/}	n.d.	n.d.
0.08% Lead Octoate	n.d.	n.d.	0.33 ^{a/} (1)
0.33% Lead Octoate	n.d.	n.d.	0.32 (1)
2.05% Lead Octoate	n.d.	n.d.	0.37 ± 0.1 (2)
0.42% Lead Chromate	n.d.	n.d.	0.52 (1)
1.95% Lead Chromate	n.d.	n.d.	4.2 (1)
12.43% Lead Chromate	n.d.	n.d.	0.4 ± 0.0 (3)
66.05% Lead Carbonate	n.d.	n.d.	2.3 ± 0.2 (8)
11.92% Lead Paint	n.d.	1.4 ± 0.1 (4)	1.86 ± 0.1 (12)

^{a/} Not detectable. Below sensitivity of 0.07 µg/gm kidney.

^{b/} Values are µg of lead/gm kidney or average µg of lead/gm kidney ± standard error of number of rats in parenthesis.

TABLE 27

LEAD CONTENT OF BONE (FEMUR)

<u>Diet</u>	<u>Treatment Week</u>		
	<u>4</u>	<u>8</u>	<u>13</u>
Control	1.6 ^a (1)	3.9 ± 0.4 (8)	3.2 ± 0.2 (24)
0.08% Lead Octoate	3.1 (1)	4.2 ± 0.6 (4)	3.3 ± 0.3 (12)
0.53% Lead Octoate	n.d. ^b	3.9 ± 0.2 (4)	3.1 ± 0.2 (12)
2.05% Lead Octoate	n.d.	3.0 ± 0.7 (4)	3.9 ± 0.5 (12)
0.42% Lead Chromate	2.9 (1)	2.6 ± 0.9 (4)	3.8 ± 0.6 (12)
1.95% Lead Chromate	0.9 (1)	1.8 ± 0.2 (4)	4.1 ± 0.5 (12)
12.43% Lead Chromate	3.3 (1)	3.1 ± 1.1 (4)	3.5 ± 0.5 (10)
66.05% Lead Carbonate	13.0 ± 4.1 (4)	16.0 ± 2.1 (4) ^c	21.7 ± 2.4 (12) ^c
11.92% Lead Paint	14.9 ± 1.5 (4)	16.4 ± 0.4 (4) ^c	10.9 ± 0.9 (12) ^c

^a/ Values are µg of lead/gm bone or average µg of lead/gm bone ± standard error of number of rats in parenthesis.

^b/ Not detectable. Below sensitivity of 0.4 µg/gm of bone.

^c/ Significantly different from control (P<0.5) as shown by Dunnett's multiple comparison test following an analysis of variance.

TABLE 26

RELATIVE ORGAN WEIGHTS OF RATS FED DIETS CONTAINING DIFFERENT CONCENTRATIONS OF LEAD IN PAINT
FILM FOR 13 WEEKS

Diet	(N)	Thyroid mg/100gm Body Weight	Spleen gm/100gm Body Weight	Heart gm/100gm Body Weight	Kidneys gm/100gm Body Weight	Liver gm/100gm Body Weight	Organs with sex differences					
							Adrenals		Brain		Conduits	
							mg/100gm Body Weight	gm/100gm Body Weight	gm/100gm Body Weight	gm/100gm Body Weight	Male	Female
Control	25	4.15 ± 0.12 ^{a/}	0.16 ± 0.01	0.28 ± 0.01	0.66 ± 0.01	3.08 ± 0.06	12.05 ± 0.79	28.75 ± 1.17	0.53 ± 0.01	0.70 ± 0.02	0.60 ± 0.04	0.04 ± 0.01
0.001% lead	12	6.28 ± 0.33	0.17 ± 0.01	0.70 ± 0.01	0.65 ± 0.01	2.97 ± 0.10	12.99 ± 0.49	27.76 ± 1.00	0.46 ± 0.01	0.70 ± 0.02	0.62 ± 0.03	0.03 ± 0.01
0.01% lead	12	6.31 ± 0.44	0.17 ± 0.01	0.70 ± 0.01	0.67 ± 0.02	3.12 ± 0.21	12.79 ± 0.91	27.54 ± 1.00	0.53 ± 0.03	0.71 ± 0.02	0.67 ± 0.04	0.04 ± 0.01
0.05% lead	12	6.73 ± 0.67	0.17 ± 0.02	0.70 ± 0.01	0.66 ± 0.01	3.12 ± 0.09	12.30 ± 0.93	26.44 ± 1.66	0.44 ± 0.01	0.76 ± 0.02	0.64 ± 0.02	0.03 ± 0.01
0.1% lead	12	6.10 ± 0.38	0.16 ± 0.01	0.70 ± 0.01	0.66 ± 0.02	3.11 ± 0.09	11.53 ± 0.96	27.76 ± 1.76	0.41 ± 0.02	0.71 ± 0.01	0.55 ± 0.03	0.05 ± 0.01
0.2% lead	12	5.57 ± 0.53	0.16 ± 0.01	0.70 ± 0.01	0.65 ± 0.02	3.14 ± 0.09	12.86 ± 1.03	25.77 ± 1.09	0.47 ± 0.01	0.66 ± 0.03	0.61 ± 0.02	0.03 ± 0.01
0.4% lead	12	6.18 ± 0.45 ^{b/}	0.16 ± 0.01	0.70 ± 0.01	0.64 ± 0.01	3.20 ± 0.07	11.26 ± 0.37	25.46 ± 1.00	0.53 ± 0.02	0.69 ± 0.02	0.56 ± 0.03	0.03 ± 0.01
0.8% lead	12	7.79 ± 0.67	0.21 ± 0.03	0.70 ± 0.01	0.70 ± 0.02	3.12 ± 0.12	16.95 ± 1.55	30.33 ± 1.66	0.45 ± 0.02	0.72 ± 0.04	0.65 ± 0.03	0.05 ± 0.01
11.9% lead	12	6.20 ± 0.46	0.18 ± 0.01	0.77 ± 0.01	0.65 ± 0.01	3.26 ± 0.03	12.77 ± 0.66	25.31 ± 1.23	0.46 ± 0.02	0.68 ± 0.02	0.58 ± 0.03	0.05 ± 0.01
Paint												

a/ Average ± standard error of number (N) of rats indicated.

b/ Eleven rats.

PATHOLOGY OF RATS FED FOR 18 WEEKS WITH PAINT FILM CONTAINING NO LEAD

Est. No.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	
<u>Lungs</u>																																									
Lymphoid hyperplasia	25/	2	1	1				1				2	1											1	1	1	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2
Pneumonia																								2																	
Emphysema																																									
<u>Heart</u>																																									
Myocarditis																																									
<u>Liver</u>																																									
Foci of inflammation														1																											
<u>Gall</u>																																									
Parasites (round worm)																																									
<u>Spleen</u>																																									
Mononuclear cell infiltration																																									
<u>Brain</u>																																									
N/G cells	1, 2	0, 2	1, 3	1, 2	1, 4	1, 4	1, 4	1, 4	1, 4	1, 4	1, 4	1, 4	1, 4	1, 4	1, 4	1, 4	1, 4	1, 4	1, 4	1, 4	1, 4	1, 4	1, 4	1, 4	1, 4	1, 4	1, 4	1, 4	1, 4	1, 4	1, 4	1, 4	1, 4	1, 4	1, 4	1, 4	1, 4	1, 4	1, 4	1, 4	

Viscera not listed were normal.

1/ Severity of lesions; 1-mild; 2-moderate; 3-severe; 4-very severe; 5-quotable

James Earl Ray

aggravated by the fact that the Government has been unable to secure adequate information regarding the activities of the various groups.

TABLE 30

PATHOLOGY OF RATS FED FOR 13 WEEKS PAINT FILM CONTAINING
2.05% LEAD AS LEAD OCTOATE

	<u>83</u>	<u>86</u>	<u>87</u>	<u>88</u>	<u>89</u>	<u>90</u>	<u>93</u>	<u>96</u>	<u>97</u>	<u>98</u>	<u>99</u>	<u>100</u>
<u>Lungs</u>												
Lymphoid hyperplasia					1 ^{a/}	2	1	1		1	2	1
Pneumonia					1							
<u>Liver</u>												
Foci of inflammation	1			1				1				1
<u>Pancreas</u>												
Foci of inflammation		1										
<u>Bone Marrow</u>												
M/E ratio	1.2	1.0	1.7	1.0	1.2	1.1	1.1	1.1	1.4	1.2	1.3	1.8

Tissues not listed were normal.

^{a/} Severity of lesions: 1-minimal; 2-moderate; 3-severe; 4-very severe; = questionable

TABLE 31

PATHOLOGY OF RATS FED FOR 13 WEEKS WITH PAINT FILM CONTAINING
12.43% LEAD AS LEAD CHROMATE

<u>Rat No.</u>	<u>143</u>	<u>146</u>	<u>147</u>	<u>148</u>	<u>149</u>	<u>150</u>	<u>153</u>	<u>156</u>	<u>157</u>	<u>158</u>	<u>159</u>	<u>160</u>
<u>Lungs</u>												
Lymphoid hyperplasia	2 ²	2		1		1				2	2	1
Pneumonia	1											
<u>Heart</u>												
Myocarditis		2										
<u>Bone Marrow</u>												
M/E ratio	1.2	1.0	1.0	1.2	1.1	1.4	1.1	1.4	1.5	1.4	1.2	1.5

Tissues not listed were normal.

g/ Severity of lesions: 1-minimal; 2-moderate; 3-severe; 4-very severe; ± questionable.

TABLE 32

PATHOLOGY OF RATS FED FOR 13 WEEKS WITH PAINT FILM
CONTAINING 66.05% LEAD AS LEAD CARBONATE

<u>Rat No.</u>	<u>163</u>	<u>166</u>	<u>167</u>	<u>168</u>	<u>169</u>	<u>170</u>	<u>173</u>	<u>176</u>	<u>177</u>	<u>178</u>	<u>179</u>	<u>180</u>
<u>Lungs</u>												
Lymphoid hyperplasia		1 ^{a/}	1	1		1			2		1	1
Pneumonia						1						
<u>Heart</u>												
Myocarditis		1	2		1							
<u>Liver</u>												
Foci of inflammation	1			1								
<u>Skeletal muscle</u>												
Miositis					1							
<u>Bone Marrow</u>												
M/E ratio	1.2	1.2	1.2	1.0	1.1	1.3	1.2	1.0	1.4	1.3	1.1	1.4

Tissues not listed were normal.

a/ Severity of lesions: 1-minimal; 2-moderate; 3-severe; 4-very severe; = questionable

TABLE 33

PATHOLOGY OF RATS FED FOR 13 WEEKS WITH NBS LEAD PAINT
CONTAINING 11.92% LEAD

<u>Rat No.</u>	<u>183</u>	<u>184</u>	<u>185</u>	<u>188</u>	<u>189</u>	<u>190</u>	<u>193</u>	<u>194</u>	<u>195</u>	<u>198</u>	<u>199</u>	<u>200</u>
<u>Lungs</u>												
Lymphoid hyperplasia	2 ^{a/}	2		2		2			1	2	1	2
Pneumonia				2								
<u>Heart</u>												
Myocarditis			2		1							
<u>Liver</u>												
Foci of inflammation									1			
<u>Bone Marrow</u>												
M/E ratio	1.2	1.4	1.3	1.2	1.5	1.2	1.2	1.4	1.5	1.3	1.2	1.3

Tissues not listed were normal.

a/ Severity of lesions: 1-minimal; 2-moderate; 3-severe; 4-very severe; = questionable