

Pat,

Here are the two papers you requested.
They are from the "Documentation of the
Threshold Limit Values", Fourth Edition, 1980, by
American Conference of Governmental Industrial
Hygienists, Inc. Let me know if you
need anything else.

Linda Ballard
9694

R&S 040589

RECEIVED

FEB 25 1981

VCM TECH CTR

VCM and EDC.

Note: This is ICGI recommended limit, not a statute.
Some states have passed laws that are more strict.

VINYL CHLORIDE

Chloroethene

$\text{CH}_2 = \text{CHCl}$

TLV, 5 ppm ($\approx 10 \text{ mg/m}^3$) — Appendix A1a —
Recognized Carcinogen

A colorless, highly flammable gas with an ethereal odor.
Vinyl chloride has a molecular weight of 62.50. It boils at

-13.9°C and freezes at -159.7°C . Vinyl chloride is usually
handled as a liquid under pressure, and containing a po-
lymerization inhibitor (phenol). It is slightly soluble in wa-
ter, but dissolved by alcohol and ether.

The chief use of vinyl chloride is as a raw material for
the manufacture of polyvinyl chloride resins. It is also em-
ployed in organic syntheses.

Since vinyl chloride is a gas at room temperature and
pressure, the common route of toxic exposure is by inhala-
tion. As with many liquified gases, contact of the skin or

eyes with escaping compressed vinyl chloride can produce freezing frostbite.⁽¹⁾

Vinyl chloride has long been considered to be very low in toxicity by acute inhalation. Lehmann and Flury⁽²⁾ summarized the literature and reported work by Schauman who considered vinyl chloride to be a candidate surgical anesthetic. Schauman reported little pathological change even after repeated exposure to anesthetic concentrations. Further work on the anesthetic potential of vinyl chloride indicated that vinyl chloride was unsafe for use as a surgical anesthetic in dogs and that because of its flammability, poor efficacy and its ability to cause cardiac irregularities at anesthetic concentrations vinyl chloride was not suitable for use as an anesthetic in humans.

Despite the early reports ascribing low toxicity to vinyl chloride, injury during the production of polyvinyl chloride (PVC) resins was reported as early as 1949. Significantly this report came from Europe where production of PVC in Europe preceded U.S. production by several years and today the quantity produced in Europe still exceeds U.S. production by about two fold. In 1949, Tribuhk *et al*⁽³⁾ reported numerous effects in PVC workers in what, by today's standards, must be considered as primitive production facilities. These authors found a "considerable number of cases of hepatitis among workers" but were more concerned with other hepatotoxic chemicals such as chlorinated diphenyl and chlorinated naphthylene (Holowax [sic]) than they were with vinyl chloride.

As a result of two deaths in Canada, the acute inhalation toxicity of vinyl chloride was studied by Mastromatteo *et al*⁽⁴⁾ who reported that exposure of mice, rats and guinea pigs to 10, 20 and 30 volume percent vinyl chloride caused the following mortality:

NUMBER OF DEATHS IN DIFFERENT GROUPS OF FIVE MICE, RATS AND GUINEA PIGS EXPOSED FOR THIRTY MINUTES TO VARYING CONCENTRATIONS OF VINYL CHLORIDE IN AIR

Vinyl Chloride concentration (percent by volume in air)	Laboratory animal			TOTAL
	Mice	Rats	Guinea pigs	
10	0/5	0/5	0/5	0/15
20	1/5	0/5	0/5	1/15
30	5/5	5/5	1/5*	11/15
40	—	—	2/5*	2/5

*A delayed death occurred within 24 hours following exposure.

Some pulmonary hyperemia and engorgement was observed by these investigators, but liver and kidney injury were remarkably low. Deaths were due to narcosis.

The first report of studies to determine the effect of long-term repeated exposure (6 months) were summarized by Torkelson, Owen and Rowe⁽⁵⁾ as follows:

"Repeated exposures of laboratory animals at several concentrations of vinyl chloride in air were conducted to determine the chronic toxicity of this material towards animals in order to assess the hazard to humans. Vinyl chloride was found to have a slight capacity to cause liver and kidney injury on repeated exposures. Male and female rats showed micropathological changes after repeated daily 7-hour exposures

at 500 ppm for 4.5 months. Repeated 7-hour exposures at 200 ppm for six months resulted in micropathological changes in the livers of rabbits and statistically significant increases in the average weight of the livers of male and female rats, but no detectable changes in dogs and guinea pigs. Repeated 7-hour exposures at 100 ppm resulted in slight increases in the average weight of rat livers, the other species were not affected. All species studied tolerated repeated daily 7-hour exposures at 50 ppm for six months with no detectable injury.

Repeated daily 1-hour exposures at 200 and 100 ppm of vinyl chloride were without effect, longer exposures caused a slight increase in liver weight.

The standard for evaluating regular daily 7- or 8-hour exposures may be defined as the concentration below which practically all analytical results must fall. The value of 100 ppm is suggested as this standard for vinyl chloride, with a time-weighted average for all exposures not to exceed 50 ppm."

Lester, Greenberg and Adams⁽⁵⁾ took exception to the conclusion of Torkelson *et al* (1961) that 50 ppm should be a maximum time-weighted average exposure for workers. On the basis of 3 months exposure of rats to 2 volume percent and 19 days to 5 volume percent, they concluded that 500 ppm was acceptable as a TLV despite minor changes which they observed in rat livers and which they considered "were within the normal range and were not pathologic in nature."

Since 1949 numerous articles describing conditions and problems in PVC production plants have appeared particularly in the Eastern European literature. Filatova and Gronsberg,⁽⁶⁾ Gabor *et al*,⁽⁷⁾ Suciu *et al*,⁽⁸⁾ Gabor *et al*,⁽⁹⁾ Grigorescu and Toba,⁽¹⁰⁾ Antonyuzhenko,⁽¹¹⁾ and Kudryavtseva⁽¹²⁾ have all described the effects of apparent gross chronic exposure. These papers and abstracts are difficult to interpret since there are generally inadequate descriptions of the exposure conditions and analysis of the workroom air, so no dose-response relationship can be determined. The injuries and effects described by the authors are not consistent with the levels of exposures claimed by the authors nor are the levels of exposure consistent with past or even present-day chemical technology. Furthermore, mixtures of chemicals are involved making it possible to ascribe the effect to any one of them.

For example, Suciu *et al*⁽⁹⁾ (through translation) described nervous disorders including euphoria with whistling and laughing, incoordination and dizziness similar to alcohol intoxication. However, Suciu *et al* ascribes these results to exposure of the order of 5.5 mg/m³ (2 ppm v/v) which is not consistent with other publications which indicate these effects will be apparent only if concentrations greatly exceed 10,000 to 20,000 ppm v/v. Therefore, the following conclusions by the authors can be construed as being the result of massive and apparently repeated exposures:

1. Vinyl chloride and the vinyl monomers possess a narcotic action and produce, depending upon concentration, in addition to characteristic neurologic manifestation, a state of euphoria (12%), followed by a state of inebriation similar to that of alcohol intoxication. In certain cases narcosis can appear.

After leaving the working environment, a state of somnolence (45%) persists, with hypersomnia. Vinyl chloride acts on the skin and produces a sensation of formication and of heat.

2. After repeated exposure, a neurologic asthenia sets in which somnolence predominates.
3. After a variable period of time, dyspeptic disturbances are added to the neurologic manifestations; these are at first not characteristic; they are in the form of epigastric pains (16%), swelling, discomfort, feeling of heaviness in the right hypochondrium (7%) or the left (5%) with anorexia, particularly for fats. In 30.2% of the cases, congestive hepatomegaly appears, which may mimic toxic hepatitis without jaundice; some cases may become chronic. In 6% of the cases, the hepatomegaly is accompanied by splenomegaly. The proteinogram and the aldolases are the most sensitive tests and show changes similar to those of acute hepatitis: increase in α -globulins and of the β - and γ -globulins; and thymol test, Greenstedt's reaction and the zinc sulfate test are positive only in few of the cases.
4. After 3 years of exposure in 9% of the cases a syndrome typical of ulcer without radiologic changes becomes manifest.
5. In 6% of the cases the Raynaud syndrome has appeared, particularly among the young men. Plethysmography shows in half of the cases an inhibition of the vasomotor centers.
6. In addition, allergic dermatitis in 4.4% of the cases, and scleroderma in 3.6%, has been observed.
7. The clinical and laboratory findings are of great importance in occupational pathology because in numerous cases diseases appear in man that cannot be reproduced in the animal (Raynaud's syndrome and scleroderma).

The sudden and frequent appearance of these manifestations in the PVC division of several plants, and in certain divisions in normal individuals who are still relatively young, and their disappearance in the majority of the cases after the institution of protective measures and change of work, have shown us decisively that vinyl chloride and the vinyl monomers have played a part in the production of these manifestations. (End of author's summary).

In 1967, reports appeared in the literature describing a condition known as acroosteolysis in workmen engaged in polymerization of vinyl chloride to polyvinyl chloride. Harris and Adams⁽¹²⁾ reported on two cases in Europe. Wilson et al.⁽¹³⁾ reported on 37 cases in the B. F. Goodrich Company. Juhe et al.⁽¹⁴⁾ described a syndrome consisting of (arranged in decreasing order of occurrence) thrombopenia, splenomegaly, liver damage, obstruction of ventilation, circulation, obstruction, and skin and bone alteration.

As a result of this problem, the University of Michigan in 1967 was retained by the Manufacturing Chemists Association to investigate acroosteolysis in sponsoring American companies. The results of a large scale epidemiological study of workers then currently employed in vinyl chloride plants in the United States were reported in three publications by this group Dinman et al.⁽¹⁵⁾ Cook et al.⁽¹⁶⁾ and Dodson et al.⁽¹⁷⁾

Dinman et al.⁽¹⁵⁾ summarized the study as follows:

"An epidemiological study was performed covering 5,011 employees with 27,510 man-years experience in various phases of vinyl chloride (VC) and polyvinyl chloride (PVC) manufacturing in 32 plants throughout the United States and Canada. The total number of definitive cases of acroosteolysis (AOL) was 25; 16 other individuals were under suspicion. This condition is clearly associated with the hand cleaning of polymerizers. Workers engaged in other phases of VC or PVC manufacturing do not appear to be at risk of developing AOL. The importance of Raynaud's phenomenon as a concomitant of AOL is emphasized. Several statistical approaches for rapid medical survey are suggested. Acroosteolysis appears to be a systemic rather than local disease. Presently, neither the etiological agent nor its portal of entry is known."

Cook et al.⁽¹⁶⁾ describes the polyvinyl chloride production process in considerable detail. They concluded that although no etiological agent could be identified, "There appeared to be a correlation between the extent of degassing prior to entry into the reactor" and the incidence of acroosteolysis.

Mutchler and Kramer⁽¹⁸⁾ presented a paper at the 1968 Gordon Research Conference which was subsequently published (1972), which reported on *The Correlation of Clinical and Environmental Measurements for Workers Exposed to Vinyl Chloride*. The authors drew the following conclusion:

"Our findings suggest that repeated exposure to vinyl chloride at TWA levels of 300 ppm or above for working lifetime together with a very low level of vinylidene chloride may result in slight changes certain physiologic and clinical laboratory parameters. The possibility of some impairment in liver function tests must be considered, even though overt clinical disease was evident in any of the individuals studied. We shall continue our study, suggest that similar studies to help clarify the effect of this material be performed for other worker populations exposed to vinyl chloride alone."

P. L. Viola, in an attempt to produce acroosteolysis in animals, exposed rats 4 hours per day, 5 days per week to 30,000 ppm (3%) vinyl chloride vapor. In his first report on the results of 12 months exposure, he described metaplastic changes in the bones which he considered similar to the human disease acroosteolysis. He made no mention of having observed cancer in these animals until the Tenth International Cancer Congress in May 1970. In the abstracts of this meeting, and subsequently in May 1971, Viola, Bigotti and Caputo⁽¹⁹⁾ reported tumors of the skin, lungs and bones occurring first after 10 months of exposure. The authors summarized this work as follows:

"Rats (Ar. RF Wistar strain) exposed for 12 months to vapors of vinyl chloride developed tumors of the skin, lungs, and bones. The cutaneous tumors, which always appeared in the area in which submaxillary and parotid glands are located, have been histologically recognized as epidermoid carcinomas, papillomas, and mucocutaneous carcinomas. The histological characteristics of lung tumors, which

R&S 040591

occurred in a lower percentage, were mainly of the adenocarcinoma type, with the exception of a single epidermoid tumor originating from the epithelial covering cells. In a minor number of rats, a large proliferation of cartilaginous tissue diagnosed as osteochondroma developed in the metacarpal and metatarsal regions of the four limbs."

The report by Viola *et al.*⁽²¹⁾ is apparently the earliest publication in which carcinogenic activity has been ascribed to vinyl chloride in man or animals. Although there were obvious deficiencies in Viola's study, such as his very impure sample, the presence of food and bedding in the exposure chamber, the excessive exposure concentration as well as in the statistical evaluation and interpretation of the lesions, the report was of serious concern and resulted in additional animal and epidemiological studies which are currently underway in Italy by Maltoni,⁽²²⁾ Maltoni and Lefemine^(23,24) and in the U.S., Keplinger *et al.*⁽²⁵⁾

On January 22-23, 1974, the B. F. Goodrich Company notified its employees, NIOSH, the Kentucky State Department of Labor, and the public, that three workers had died of angiosarcomas of the liver. The case reports of the first subject has been published by Creech and Johnson.⁽²⁶⁾ The subject, a 36 year old male, was hospitalized January 5, 1970 and subsequently succumbed September 27, 1971. He had worked in PVC production from November 1955 until his illness. The history, clinical course and pathologic findings are consistent with the others who died of angiosarcoma.

The work of Maltoni and Lefemine^(23,24) has been reported publicly at the OSHA hearing, Washington, D.C., February 15, 1974, and included in the 1974 publication of the Second International Symposium on Cancer Detection and Prevention, Bologna, Italy, April 9-12, 1973. In these studies groups of rats as well as mice and hamsters have been exposed at concentrations of 10,000 to 50 ppm vinyl chloride vapor. Maltoni and Lefemine (1974) reported carcinomas of the Zymbal glands, nephroblastoma and angiosarcomas of the livers of rats at concentrations of 250 ppm to 10,000 ppm but not at 50 ppm. Subsequent unpublished information (August 31, 1974) reported, "1 liver angiosarcoma, 1 extrahepatic angiosarcoma and 1 nephroblastoma, in three animals of the first experiment, exposed at 50 ppm of VC for 1 year, and surviving 135 weeks from the beginning of the treatment." The authors conclude that,

"a dose-response relationship clearly emerges, as far as angiosarcomas and nephroblastomas are concerned, in the lower dose ranges: from 500 ppm to 50 ppm for angiosarcomas, and from 250 ppm to 50 ppm for nephroblastomas. A comparison of the results available at the present moment in rats exposed for 12 months and 4 months (BT1 and BT3 experiments) shows that the neoplastic response, as far as angiosarcomas and nephroblastomas are concerned, is affected by the length of exposure to VC."

In their experiment BT3 Maltoni and Lefemine reported possible interproduction of angiosarcomas in offspring of pregnant rats exposed at 10,000 and 6,000 ppm.

Keplinger *et al.*⁽²⁵⁾ in a study sponsored by several companies, have confirmed the findings of Maltoni and Lefemine. In this study, groups of 100 rats, mice and hamsters of both sex are being exposed seven hours per day, five days per week to either 2,500, 200 to 500 ppm vinyl chloride monomer. After seven months of exposure an-

giosarcoma and lung adeaomas have been observed in mice at all exposure levels. Although the data are preliminary in nature and require confirmation, angiosarcomas were apparently also observed in rats at 2,500 and 200 ppm and in a single hamster at 2,500 ppm. This study is still in progress and will not be completed until 1976 or 1977.

Epidemiological studies on U.S. workers have been conducted by Tabershaw-Cooper Associates for the Manufacturing Chemists Association.⁽²⁷⁾ The summary of this study is as follows:

This historical prospective mortality study of 8384 men who had at least one year of occupational exposure to vinyl chloride before December 31, 1972, demonstrated that cancers of the digestive system (primarily angiosarcoma), respiratory system, brain, and cancers of unknown site, as well as lymphomas, occurred more often than expected in those members of the study population with the greatest estimated exposure. The mortality from other cancers was lower than that of the general male population, with the exception of cancers of the buccal cavity and pharynx. The explanation for the latter finding is not apparent.

The other major findings of the study are: (1) The overall mortality of the study population was approximately 75% of what would be expected in a comparable population of U.S. males; (2) No cause of death showed a statistically significant excess over what would be expected in a comparable U.S. male population; and (3) No deaths identified as angiosarcoma of the liver were found other than those previously identified.

This is the first epidemiological study which suggests that in humans vinyl chloride may also be associated with cancer of multiple sites.

It is difficult to derive a reasonable TLV from the data presented in the literature summarized above. There is indirect evidence that intermittent exposures to vinyl chloride of the order of thousands of ppm, in this country as well as Russia, have not been infrequent in PVC plants. No data on the past (or even present) concentrations of vinyl chloride in plants where angiosarcoma cases have occurred, or not occurred, appear to be available. One report⁽²⁸⁾ indicates that 21 of 26 of the early cases of angiosarcoma occurred in former reactor cleaners. Cleaning of reactors was apparently responsible for most of the acroosteolysis cases investigated by Cook and associates,⁽¹⁷⁾ and has resulted in death from acute poisoning by vinyl chloride.⁽³⁶⁾

It is the sense of the Committee that, if the average exposure to vinyl chloride does not exceed 5 ppm, there will be no increase in the incidence of cancer, specifically of angiosarcoma of the liver. It is probable that the cancers reported and attributed to vinyl chloride among PVC workers resulted from exposures many times this level.

NIOSH recommended a limit of 1 ppm as a TWA, with a ceiling of 5 ppm. The 1 ppm value was apparently based on the erroneous belief that this was the lowest concentration that could be readily measured.

Gehring and co-workers,⁽³⁰⁾ using a probit model, and based on studies with rats, found the predicted incidence of hepatic angiosarcomas from 8 hour/day, 5 days/week 35 year exposure at 1 ppm to be 1.5 times 10^{-8} .

Recent papers have included a report of 4 cases of respiratory cancer among vinyl chloride workers, but no dose-response relationship.⁽³¹⁾

On the other hand, Fox and Collier,⁽³²⁾ in a study of 7000 men exposed to vinyl chloride in PVC manufacture between 1940 and 1974, found no evidence of cancers due to vinyl chloride at sites other than the liver. There are four liver cancers, two of them angiosarcomas.

Delorme and Theriault⁽³³⁾ described 10 cases of liver angiosarcoma among workers in vinyl chloride polymerizing plant in Quebec, which were accompanied by fibrosis of the liver. Details of 64 cases were presented by Sputas and Kaminski.⁽³⁴⁾

Mutufugi, in Japan, noted that in contrast to western countries, in which 70 angiosarcomas cases (associated with vinyl chloride exposure) had been reported, no cancer, but many poisoning cases, have been reported in the USSR.⁽³⁵⁾

Based on the above data, an A1a classification as a confirmed carcinogen is given vinyl chloride and a TLV of 5 ppm as a time-weighted average is suggested. If this value is not exceeded, there should be no increase in the incidence of cancer, especially angiosarcoma of the liver.

Limits adopted in other countries, subsequent to the surfacing of vinyl chloride exposure associated cancers, as are follows, according to a 1977 summary: Australia (1973) 25 ppm; Finland (1975), Holland (1973), Poland (1976), Switzerland (1976) and USSR (1977) about 10 ppm; Italy (1975) 5 ppm; Japan (1975) and Sweden (1978) 1 ppm.

References:


1. Torkelson, T.R. et al: *Am. Ind. Hyg. Assoc. J.* 22(5):354 (1961).
2. Lehman, K.B., Flury, F.: *Toxicology & Hygiene of Industrial Solvents* (1938).
3. Tribukh, et al: *Arg. Saint.* 10:38 (1949).
4. Mastromatteo, E. et al: *Am. Ind. Hyg. Assoc. J.* 21(5):394 (1961).
5. Lester, D. et al: *Ibid.* 24:265 (1963).
6. Filatova, V.A., Gronsberg, E.S.: *Gigiena i. Sanit.* 22(1):33, abstract (1957).
7. Gabor, S. et al: *Prom. Toksikol. i. Klinika Prof. Zaholevanii Khim. Etiol. Sb.* 221, abstract (1962).
8. Suciu, I. et al: *Medicina Interna (Bucharest)* XV(8):967 (1963).
9. Gabor, S. et al: *Ingiena Bucharest* 13(5):409 (1964).
10. Grigorescu, I., Toba, G.: *Rev. Chim.* 17(8):499, abstract (1966).
11. Antonyuzhenko, V.A.: *Gig. Tr. Prof. Zabol.* 12(3):50, abstract (1968).
12. Kudryavtseva, O.F.: *Ibid.* 14(8):54, abstract (1970).
13. Harris, D.K., Adams, W.G.F.: *Brit. Med. J.* 5567:712, abstract (1967).
14. Wilson, R.H. et al: *JAMA* 207(8):577 (1967).
15. Juhe, S. et al: *Dtsch. Med. Wschr.* 98:2034 (1973).
16. Dinman, B.D. et al: *Arch. Env. Health* 22:61 (1971).
17. Cook, W.A. et al: *Ibid.*, p. 74.
18. Dodson, V.N. et al: *Ibid.*, p. 83.
19. Kramer, C.G., Mutchler, J.E.: *Am. Ind. Hyg. Assoc. J.* 33(1):19 (1971).
20. Viola, P.L.: *Medicina del Lavoro* 61(3) (March 1970).
21. Viola, P.L. et al: *Cancer Research* 31:516 (1971).
22. Maltoni, C.: *Proc. 2nd Intl. Symp. on Cancer Detection & Prevention*, Bologna, 1973, Excerpta Medica, Amsterdam (1974).
23. Maltoni, C., Lefemine, G.: *Lincei-Rendiconto Della Classe di Science, Tesiche, Mathmatische* 56:1 (1974).
24. Eidem: *Carcinogenic Bioassays of Vinyl Chloride*, unpublished data (1974).
25. Keplinger, J.L. et al: *Annals of NY Acad. of Sciences Working Group* (May 10, 1974).
26. Creech, J.L., Johnson, M.N.: *JOM* 16(3):150 (1974).
27. Tabershaw, I.R., Gaffey, W.R.: *Ibid.*, p. 508.
28. Patty, F.A. et al: *U.S. Pub. Health Reports* 45, 1963, abstract (1930).
29. Current Intelligence, *J. Occup. Med.* 16:809 (1974).
30. Gehring, P.T. et al: *Tox. Appl. Pharm.* 49:15 (1979). IHD abstract 753/79.
31. Buffler, P.A. et al: *J. Occup. Med.* 21:195 (1979). *Ibid.* 54/79.
32. Fox, A.J., Collier, P.F.: *Brit. J. Ind. Med.* 34:1 (1977). *Ibid.* 501/77.
33. Delorme, F., Theriault, G.: *J. Occup. Med.* 20:338 (1978). *Ibid.* 688/78.
34. Sputas, R., Kaminski, R.: *Ibid.*, p. 427. *Ibid.* 828/78.
35. Matufugi, H.: *J. Science Labor* 54 585 (1978). *Ibid.* 54/79.
36. Massachusetts Div. of Occup. Hygiene: Unpublished report.

References:

1. Lucas, G.H.W.: *J. Pharm.* 34:223 (1928).
2. Von Oettingen, W.F.: *The Halogenated Hydrocarbons, Toxicity and Potential Dangers*, US Public Health Service Publ. No. 414, p. 152, Washington, DC (1955).
3. Kochmann, M.: *Muench. Med. Wochenschr.* 75:1334 (1928) Cited in ref. 2, p. 153.
4. Olmstead, E.V.: *Arch. Ind. Health* 21:525 (1960).
5. Rowe, V.K., Spencer, H.C., McCollister, D.D., Hollingsworth, R.L., Adams, E.M.: *Arch. Ind. Hyg. & Occup. Med.* 6:158 (1952).
6. Rowe, V.K., Hollingsworth, R.L., McCollister, D.D.: *Agric. & Food Chem.* 2:1318 (1954).
7. McCollister, D.D., Hollingsworth, R.L., Oyen, F., Rowe, V.K.: *Arch. Ind. Health* 13:1 (1956).
8. NIOSH: *Criteria for a Recommended Standard — Occupational Exposure to Ethylene Dibromide*, DHEW (NIOSH) Pub. No. 77-221 (1977).
9. Kochmann, M.: *Muench. Med. Wochenschr.* 75:1334 (1928) Cited in ref. 8.
10. Olson, W.A. et al: *J. Natl. Cancer Inst.* 51:993 (1973). Ibid.
11. Powers, M.B. et al: *Tox. Appl. Pharm.* 33:171 (1975). Ibid.
12. Ward, J.M., Haberman, R.T.: *Lab. Invest.* 30:392 (1974). Ibid.
13. Idem: *Bull. Soc. Pharmacol. Exp. Path.* 2:10 (1974). Ibid.
14. Ott, M.G., Scharnweber, H.C., Langner, R.R.: Unpublished report submitted to NIOSH by the Dow Chemical Co. (1977). Ibid.
15. Stanford Research Institute: Report submitted to NIOSH under contract No. DCD-99-74-31 (1976). Ibid.
16. Amir, D., Volcani, R.: *Nature* 206:99 (1965). Ibid.
17. Alumot, E. et al: *Poult. Sci.* 47:1979 (1968). Ibid.
18. Short, R.D. Jr. et al: Report No. EPA-560/6-76-018, US Environmental Protection Agency, Office of Toxic Substances (1967) Ibid.
19. NIOSH: *Toxic Substances List*, H.E. Christensen ed., p. 4 (1973).

ETHYLENE DICHLORIDE

1,2-Dichloroethane

TLV, 10 ppm ($\approx 40 \text{ mg/m}^3$)STEL, 15 ppm ($\approx 60 \text{ mg/m}^3$)

Ethylene dichloride is a colorless liquid with the odor typical of chlorinated hydrocarbons. It has a molecular weight of 98.96 and, at 20° C, a specific gravity of 1.2569 and a vapor pressure of 62 mm Hg. The boiling point is 83.5° C, freezing point is -35.5° C and closed cup flash point is 55° F. It is soluble in about 120 parts water and miscible with alcohol, chloroform and ether.

It has been used as an intermediate in the manufacture of vinyl chloride, a scavenger in leaded gasoline, a degreaser, fumigant, solvent, and in paint removers, wetting and penetrating agents, ore flotation and soaps and scouring compounds.

Animal studies have uniformly indicated liver and kidney injury from ethylene dichloride, although to a lesser extent than from comparable exposures to carbon tetrachloride. Smyth et al⁽¹⁾ reported an oral LD₅₀ in rats to be 770 mg/kg.

Heppel and associates⁽²⁾ on the basis of animal studies, concluded that ethylene dichloride is one of the more toxic of the common chlorinated hydrocarbons. They found increased mortality in 5 species of animals exposed 7 hr/d, 5 d/wk, at 200 ppm. Loss in weight, pulmonary congestion and liver changes were noted. At 100 ppm rats bred successfully and autopsy revealed no abnormal findings.

Smyth and co-workers⁽³⁾ found loss in weight and increased liver weight in guinea pigs similarly exposed for 220 days at 100 ppm. Various species with similar effects died by 200 ppm, and high mortality occurred at 400 ppm.

Hofmann et al⁽⁴⁾ found high mortality in various species exposed to 400 ppm for 17 weeks at 500 ppm. At 100 ppm rats showed a decreased growth rate, but other species were unaffected.

Opacity of the cornea has been observed in dogs and foxes following administration of ethylene dichloride, according to von Oettingen.⁽⁵⁾

A number of deaths from the accidental ingestion of ethylene dichloride have been reported. In addition, Browning⁽⁶⁾ noted three fatal cases from inhalation; liver and kidney injury were found in all, pulmonary edema in one. Several other episodes of occupational intoxication, with nausea and vomiting the predominating symptoms, have been reported.^(5,7,8)

Experience in one plant indicated that concentrations in the range of 25 to 50 ppm were safe for prolonged exposure.⁽⁹⁾ Bardodej⁽¹⁰⁾ however, considered it a most undesirable substance, comparable in toxicity to carbon tetrachloride.

In its criteria document,⁽¹¹⁾ NIOSH cited French,⁽¹²⁾ Italian,^(13,14) German⁽¹⁵⁾ and Russian⁽¹⁶⁾ papers which indicated fatalities from inhalation of ethylene dichloride in concentrations insufficient to cause anesthesia. A Polish paper⁽¹⁷⁾ reported symptoms such as nausea, vomiting and dizziness from exposures between 10 and 37 ppm in one of six workers, and physical examinations revealed abnormalities, notably blood changes, in 8 of 16 workers.

The NIOSH recommendation for a workplace air standard at this time was 5 ppm as TWA, with a 15-minute ceiling of 15 ppm. This recommendation appears to have been based largely on a paper by Kozik⁽¹⁸⁾ in Russia. Workers with exposures generally averaging below 16 ppm, but in a few instances between 30 and 50 ppm, experienced numerous adverse nervous system and liver effects.

Another paper⁽¹⁹⁾ reported no changes in blood or internal organs among 100 Russian workers exposed up to five years at concentrations of ethylene dichloride not exceeding 25 ppm, but various complaints and nervous system effects were noted.

NIOSH has since issued a recommended limit of 1 ppm or 1 ppm, with a 2 ppm ceiling.⁽¹¹⁾ Cancer is given as one of the health effects considered.

Ethylene dichloride clearly belongs in the group of hepatotoxic halogenated hydrocarbons, as opposed to chloroform with primarily narcotic effects. A reduction of the TLV from

50 to 10 ppm has been adopted and a STEL of 15 ppm is recommended.

Other recommendations from a 1977 listing: West Germany, Sweden, Switzerland, 20 ppm; Czechoslovakia, East Germany, Romania, Yugoslavia, 12 ppm; Poland, U.S.S.R., 2.5 ppm; others, including Japan and the Council of Europe retained the 50 ppm value. In 1978 Sweden reduced its TWA to 5 ppm, and in 1979 West Germany added the designation of potential carcinogen, but retained the 20 ppm MAK.

References:

1. Smyth, H.F. Jr. et al: *Am. Ind. Hyg. Assoc. J.* 30:470 (1969).
2. Heppel, L.A., Neal, P.A., Perrin, T.L., Porterfield, V.T.: *J. Ind. Hyg. & Tox.* 28:113 (1946).
3. Spencer, H.C., Rowe, V.K., Adams, E.M., McCollister, D.D., Irish, D.D.: *Arch. Ind. Hyg. & Occup. Med.* 4:482 (1951).
4. Hofmann, H.T., Birnstiel, H., Jobst, P.: *Arch. Toxikol.* 27:248 (1971).
5. von Oettingen, W.F.: *The Halogenated Hydrocarbons, Toxicity and Potential Dangers*, US Public Health Service, Pub. No. 414, p. 144, Washington, DC (1955).
6. Browning, E.: *Toxicity & Metabolism of Organic Solvents*, p. 252, Elsevier Publishing Co., Amsterdam (1965).
7. Wirtschafter, Z.T., Schwartz, E.D.: *J. Ind. Hyg. & Tox.* 21:126 (1939).
8. McNally, W.D., Fostvedt, G.: *Ind. Med.* 10:373 (1941).
9. Fassett, D.: Private communication to TLV Committee (1964).
10. Bardodej, Z.: in *Documentation of MAC in Czechoslovakia*, p. 63, Prague (1969).
11. NIOSH: *Criteria for a Recommended Standard — Occupational Exposure Ethylene Dichloride*, HEW Pub. No. (NIOSH) 76-139 (1976).
12. Olivier, H. et al: *Soc. Med. Leg.* 34:261 (1954). Cited by NIOSH in ref. 10.
13. Guarino, A., Koia, N.: *Folia Med. (Napoli)* 41:676 (1958). *Ibid.*
14. Triosi, F.M., Cavallazzi, D.: *Med. Lav.* 52:612 (1961). *Ibid.*
15. Menschick, H.: *Arch. Dewerbepath. Gewerbehyg.* 15:241 (1957). *Ibid.*
16. Rosenbaum, N.D.: *Gig. Sanit.* 12(2):17 (1947). *Ibid.*
17. Brzozowski, J.: *Med. Pracy* 5:89 (1954). *Ibid.*
18. Kozik, I.: *Gig. Tr. Prof. Zabol.* 1:32 (1957). *Ibid.*
19. NIOSH: *Special Hazard Review—Ethylene Dichloride* (1978).

R&S 040595

RECEIVED

FEB 20 1981

VCM TECH CTR