

TABLE 1

PREDICTED INCIDENCE OF ANGIOSARCOMA IN RATS EXPOSED TO VINYL CHLORIDE USING VARIOUS MODELS VERSUS DOSE ( $v$ ,  $\mu\text{g}$  VC METABOLIZED PER DAY) COMPARED TO EXPERIMENTAL RESULTS

| Exposure <sup>a</sup><br>(ppm) | Dose <sup>b</sup><br>( $v$ , $\mu\text{g/day}$ ) | Experimental <sup>c</sup><br>(%) | Predicted percentage (Models) <sup>c</sup> |      |      |      |
|--------------------------------|--------------------------------------------------|----------------------------------|--------------------------------------------|------|------|------|
|                                |                                                  |                                  | A                                          | B    | C    | D    |
| 10,000                         | 5521                                             | 14.8 (9/61)                      | 19.8                                       | 20.0 | 19.7 | 18.9 |
| 6,000                          | 5403                                             | 21.7 (13/60)                     | 19.3                                       | 20.5 | 19.3 | 18.6 |
| 2,500                          | 5030                                             | 22.0 (13/59)                     | 17.9                                       | 18.1 | 17.9 | 17.4 |
| 500                            | 3413                                             | 11.9 (7/59)                      | 12.1                                       | 11.8 | 12.2 | 12.2 |
| 250                            | 2435                                             | 6.8 (4/59)                       | 8.1                                        | 8.0  | 8.7  | 8.8  |
| 50                             | 739                                              | 1.7 (1/59)                       | 1.4                                        | 1.4  | 2.6  | 2.8  |

<sup>a</sup> From Maltoni and Lefemine (1975).

<sup>b</sup> Calculated from  $v = [5706 (\mu\text{g VC}/4 \text{ hr}) \cdot S(\mu\text{g/liter})] / [860 (\mu\text{g/liter}) + S(\mu\text{g/liter})]$ , where  $S = 2.56 (\mu\text{g/ppm/liter}) \times \text{exposure (ppm)}$  (Gehring *et al.*, 1978).

<sup>c</sup> Model A, Probit % =  $-1.625 + 1.543 \text{ Log (dose)}$ ; Model B, % =  $-1.48 + 0.389 \times 10^{-2} (\text{dose})$ ; Model C, % =  $0.3565 \times 10^{-2} (\text{dose})$ ; Model D, % =  $[1 - e^{-\beta(\text{dose})}] 100$ , where  $\beta = 0.38 \times 10^{-4}$ .

## RESULTS AND DISCUSSION

The incidence of hepatic angiosarcoma in subgroups of 9677 workmen with respect to their duration of exposure are given in Table 2. These workers include 95.1% of the 10,173 in the cohort study; the remainder could not be traced.

The atmospheric concentrations of vinyl chloride to which these workers had been exposed were not quantitated, except subjectively into high, medium, and low categories. Since the time-weighted-average (TWA) exposure recommended by the American Conference of Governmental Industrial Hygienists (ACGIH) prior to 1972 was 500 ppm and subsequently 200 ppm until adoption of the Occupational Safety and Health Act standard of less than 1 ppm in 1974, it is assumed that even those exposures subjectively deemed low were high. Indeed, it is reasonable to expect the TWA exposures of 200 ppm and greater for 8 hr were common rather than the exceptions.

Table 3 depicts the average theoretical rat equivalent daily doses of biotransformation products of VC received by workers exposed to 200 or 500 ppm for 8 hr daily,

5 days/week adjusted for the fraction of their working life during which exposure occurred. The doses,  $v$ , are expressed as micrograms per 8 hr and were calculated from Eq. 3 (given under Introduction) multiplied by the fraction of an assumed 35-year working life during which exposure occurred.

Having estimated the rat equivalent daily doses of biotransformation products of VC received by the various subgroups of 9677 workmen exposed to a TWA of 200 or 500 ppm VC, the predicted incidence of hepatic angiosarcoma for each subgroup was calculated (Table 4). Examination of the data in this table indicates that Models C and D overestimate the incidence of hepatic angiosarcoma experienced by these workers while Model B underestimates the incidences. Models C and D may overpredict because they do not address either the enzymatic biotransformation of VC or repair of the lesion leading to development of angiosarcoma, both of which are likely to be distributed normally in human or animal populations.

It may be argued that Models C and D overpredict because the recommended TWA exposure values of 200 or 500 ppm VC are

TABLE 2  
INCIDENCE OF ANGIOSARCOMA IN 9677 WORKMEN EXPOSED TO VINYL CHLORIDE<sup>a</sup>

| Population               | Duration of exposure (years) | Mean exposure duration (years) <sup>b</sup> | Fraction of working life <sup>c</sup> | Observed angiosarcomas    |
|--------------------------|------------------------------|---------------------------------------------|---------------------------------------|---------------------------|
| 4384 (0.31) <sup>d</sup> | <4                           | 2                                           | 0.06                                  | 1 (18 years) <sup>e</sup> |
| 2339 (0.27)              | 5-9                          | 7                                           | 0.20                                  | 0                         |
| 946 (0.58)               | 10-14                        | 12                                          | 0.34                                  | 2 (15 and 23 years)       |
| 1007 (1.0)               | 15-19                        | 17                                          | 0.49                                  | 2 (18 and 19 years)       |
| 677 (1.0)                | 20-24                        | 22                                          | 0.63                                  | 0                         |
| 324 (1.0)                | 25+                          | 27                                          | 0.77                                  | 0                         |
| Total<br>9677            |                              |                                             |                                       |                           |

<sup>a</sup> From "Epidemiological Study of Vinyl Chloride Workers", Final Report January 1978, prepared by Equitable Environmental Health, Inc., Rockville, Md., for Manufacturing Chemists' Association, Washington, D.C.

<sup>b</sup> Assumed mean duration of exposure.

<sup>c</sup> Mean duration of exposure divided by 35 years.

<sup>d</sup> Fraction of population incurring first exposure 15 or more years prior to study.

<sup>e</sup> Duration between first exposure and diagnosis.

higher than those actually incurred by workers included in the epidemiological study. Undoubtedly, some were exposed to smaller levels. Many were reported exposed to much higher levels; in some cases at or above the odor threshold of approximately 3500 ppm. Unfortunately, in the absence of

analytically determined exposure levels, approximations of the actual TWA values must be used. For a TWA exposure of 100 ppm VC, the predicted number of workers developing angiosarcoma in this epidemiological cohort are 4, 0, 30, and 32 for mathematical Models A, B, C, and D, respectively. For a 50-ppm TWA, the corresponding numbers are 1, 0, 17, and 18. Thus, utilizing values for TWA exposures which are thought to underestimate the actual exposures appear to overpredict still the number of workers developing angiosarcoma of the liver when mathematical Models C and D are used, albeit Model A apparently underpredicts when a TWA exposure of 50 ppm is used.

When a TWA of 200 ppm is utilized to represent the atmospheric concentration to which these workers were exposed, the predicted incidence of angiosarcoma of 10 cases using Model A compares favorably with that experienced, five cases. For a significant number of workers, the time elapsed since initial exposure has yet to reach 15 years, the shortest time needed for development of angiosarcoma in 1 of 5 of the afflicted workers (Table 2). Thus, it may be

TABLE 3

AVERAGE THEORETICAL RAT EQUIVALENT DAILY DOSES OF BIOTRANSFORMATION PRODUCTS RECEIVED BY A 70-kg HUMAN EXPOSED TO 200 OR 500 ppm VINYL CHLORIDE FOR VARIOUS FRACTIONS OF AN ASSUMED 35-YEAR WORKING LIFE

| Mean exposure duration (years) | Fraction of working life | D, micrograms per 8 hr averaged over 35 hr <sup>a</sup> |     |
|--------------------------------|--------------------------|---------------------------------------------------------|-----|
|                                |                          | 500 <sup>b</sup>                                        | 200 |
| 2                              | 0.06                     | 60                                                      | 38  |
| 7                              | 0.20                     | 200                                                     | 125 |
| 12                             | 0.34                     | 341                                                     | 213 |
| 17                             | 0.49                     | 491                                                     | 306 |
| 22                             | 0.63                     | 631                                                     | 394 |
| 27                             | 0.77                     | 771                                                     | 481 |

<sup>a</sup>  $D = [1675 \mu\text{g}/8 \text{ hr} \cdot S \mu\text{g}/\text{liter}] / [860 \mu\text{g}/\text{liter} + S \mu\text{g}/\text{liter}] \times (\text{fraction of working life})$ .

<sup>b</sup> Parts per million.

PREDICT  
8 HR/DA  
OBS

Population

4384  
2339  
946  
1007  
677  
324

Total  
9677

<sup>a</sup> Observed  
Report Janua  
Chemists Ass  
<sup>b</sup> Model A,  
% = 0.3565 ×  
models were c  
chloride (Geh  
<sup>c</sup> Parts per r

anticipated  
hepatic ang  
population.  
of cases, as  
not to be an

As indicat  
the previous  
experiments  
of hepatic  
workers app  
served incide  
biotransform  
by humans a  
extrapolation

The foreg  
percentage i  
revealed that  
rats may be  
hepatic angio  
able accurac  
model will b  
tion of the r.

UCC  
002871

TABLE 4

PREDICTED NUMBER OF ANGIOSARCOMAS IN WORKERS EXPOSED TO 500 OR 200 PPM VINYL CHLORIDE 8 HR/DAY, 5 DAYS/WEEK FOR VARIOUS FRACTIONS OF A 35-YEAR WORKING LIFE VERSUS THE NUMBERS OBSERVED USING BIOTRANSFORMATION AND ANGIOSARCOMA INCIDENCE DATA FROM RATS AND FOUR MODELS FOR EXTRAPOLATION\*

| Population | Angiosarcomas observed | Angiosarcomas predicted (Models) <sup>a</sup> |     |     |     |      |      |      |      |
|------------|------------------------|-----------------------------------------------|-----|-----|-----|------|------|------|------|
|            |                        | A                                             |     | B   |     | C    |      | D    |      |
|            |                        | 500 <sup>b</sup>                              | 200 | 500 | 200 | 500  | 200  | 500  | 200  |
| 4384       | 1                      | 0.2                                           | 0.0 | 0   | 0   | 9.4  | 5.9  | 10.0 | 6.3  |
| 2339       | 0                      | 2.4                                           | 0.8 | 0   | 0   | 16.7 | 10.4 | 17.7 | 11.1 |
| 946        | 2                      | 3.0                                           | 1.1 | 0   | 0   | 11.5 | 7.2  | 12.1 | 7.6  |
| 1007       | 2                      | 6.8                                           | 2.7 | 4.0 | 0   | 17.6 | 11.0 | 18.6 | 11.7 |
| 677        | 0                      | 7.3                                           | 3.0 | 6.6 | 0.3 | 15.2 | 9.5  | 16.0 | 10.1 |
| 324        | 0                      | 4.9                                           | 2.1 | 4.9 | 1.3 | 8.9  | 5.6  | 9.4  | 5.9  |
| Total      |                        |                                               |     |     |     |      |      |      |      |
| 9677       | 5                      | 25                                            | 10  | 16  | 2   | 79   | 50   | 84   | 53   |

\* Observed number of angiosarcomas from "Epidemiological Study of Vinyl Chloride Workers," Final Report January 1978, prepared by Equitable Environmental Health Inc., Rockville, Md., for Manufacturing Chemists Association, Washington, D.C.

<sup>a</sup> Model A, Probit % =  $-1.625 + 1.543 \text{ Log (dose)}$ ; Model B, % =  $-1.48 + 0.389 \times 10^{-2} (\text{dose})$ ; Model C, % =  $0.3565 \times 10^{-2} (\text{dose})$ ; Model D, % =  $[1 - e^{-\beta (\text{dose})}] 100$  where  $\beta = 0.38 \times 10^{-4}$ . The constants in these models were obtained using the biotransformation and angiosarcoma incidence data for rats exposed to vinyl chloride (Gehring *et al.*, 1978).

<sup>b</sup> Parts per million.

anticipated that a few additional cases of hepatic angiosarcoma will occur in this population. It is comforting that an epidemic of cases, as has been envisioned by some, is not to be anticipated.

As indicated in the beginning of the paper, the previous attempt to extrapolate data from experiments in rats to predict the occurrence of hepatic angiosarcoma in vinyl chloride workers appeared to overpredict the observed incidence even when differences in the biotransformation of VC to active products by humans and rats were incorporated in the extrapolation.

The foregoing analysis using a probit percentage incidence model (Model A) has revealed that data collected in experiments on rats may be used to predict the incidence of hepatic angiosarcoma in humans with reasonable accuracy. Further verification of this model will be assisted greatly by determination of the rate of biotransformation of VC

by humans exposed to sufficient concentrations of VC to allow such measurement. Currently, this must be estimated by the rate of biotransformation in rats and its assumed relationship to body surface area (Gehring *et al.*, 1978). Nonetheless, the correlation between predicted and observed incidence of hepatic angiosarcoma in vinyl chloride workers seems to justify both the assumptions and the model (Model A, Probit %) used to make the predictions.

Since the predicted incidence of hepatic angiosarcoma in humans using results from studies in rats appears justifiable, it is of interest to use the same procedures to estimate the incidence in workers exposed to 1 ppm 8 hr/day for a 35-year working life, the upper limit of the current Occupational Safety and Health Act standard. These predictions are shown in Table 5 for all four models.

Models C and D predict an incidence

TABLE 5

PREDICTED INCIDENCE OF ANGIOSARCOMA IN WORKERS EXPOSED TO 1 ppm VINYL CHLORIDE 8 HR/DAY, 5 DAYS/WEEK FOR 35 YEARS USING BIOTRANSFORMATION AND ANGIOSARCOMA INCIDENCE DATA FROM STUDIES IN RATS\*

- (A) Probit % or logarithm probability  
 $1.5 \times 10^{-6}\%$  or 1.5/100,000,000
- (B) Linear not forced through origin  
 None predicted below exposures of 98.7 ppm.
- (C) Linear forced through origin  
 $1.7 \times 10^{-2}\%$  or 1.7/10,000 or 177/1,000,000
- (D) One-hit model (CAG)  
 $1.89 \times 10^{-2}\%$  or 1.9/10,000 or 189/1,000,000

\* Rat equivalent daily dose of biotransformation products received by workmen exposed for 8 hr to 1 ppm vinyl chloride is:

$$V = [1675 \mu\text{g/hr} \cdot 2.56 \mu\text{g/liter}] / [860 \mu\text{g/liter} + 2.56 \mu\text{g/liter}] = 4.97 \mu\text{g/8 hr.}$$

comparable to that experienced by workers exposed to much higher levels, again revealing their inadequacy because of overprediction. Model B, although a linear model, predicts zero response at 98.7 ppm VC. Thus, exposures to levels below this are not predicted to induce hepatic angiosarcoma. Using what appears to be the most reliable model (Model A, Probit %), the predicted incidence of hepatic angiosarcoma is  $1.5 \times 10^{-6}\%$  or 1.5 cases in 100,000,000 workers. It is emphasized that use of Models A, C, and D to predict the incidence of hepatic angiosarcoma in workers exposed to 1 ppm VC assumes no threshold or even an inflection point in the dose-response curve (even though a threshold might occur).

In summary, the use of animal data for the assessment of risk in humans exposed to a chemical is not as mystical as frequently assumed. Dose-response data for the induction of angiosarcoma in rats exposed to various levels of VC together with attendant biotransformation data have been used to

estimate, reliably, the risk of developing angiosarcoma in persons exposed to VC. There is every reason to believe that similar data can and should be used in the assessment of risk of exposure to specified levels of other substances.

## REFERENCES

- BARBIN, A., BRÉSIL, H., CROISY, A., JACQUIGNON, P., MALAVIELLE, C., MONTESANO, R., AND BARTSCH, H. (1975). Liver-microsome-mediated formation of alkylating agents from vinyl bromide and vinyl chloride. *Biochem. Biophys. Res. Commun.* 67, 596-603.
- BARTSCH, H., MALAVIELLE, C., AND MONTESANO, R. (1975). Human, rat, and mouse liver mediated mutagenicity of vinyl chloride in *Salmonella typhimurium* strains. *Int. J. Cancer* 15, 429-437.
- BOLT, H. M., KAPPUS, H., KAUFMANN, R., APPEL, K. E., BUCHTER, A., AND BOLT, W. (1975). Metabolism of  $^{14}\text{C}$ -vinyl chloride *in vitro* and *in vivo*. *Inserm* 52, 151-164.
- BOLT, H. M., KAPPUS, H., BUCHTER, A., AND BOLT, W. (1976). Disposition of (1,2- $^{14}\text{C}$ )vinyl chloride in the rats. *Arch. Toxicol.* 35, 153-162.
- CREECH, J. L., AND JOHNSON, M. N. (1974). Angiosarcoma of liver in the manufacture of polyvinyl chloride. *J. Occup. Med.* 16, 150-151.
- FOX, A. L., AND COLLIER, P. F. (1977). Mortality experience of workers exposed to vinyl chloride monomer in the manufacture of polyvinyl chloride in Great Britain. *Brit. J. Ind. Med.* 34, 1-10.
- GEHRING, P. J., WATANABE, P. G., AND PARK, C. N. (1978). Resolution of dose-response toxicity data for chemicals requiring metabolic activation: Example—vinyl chloride. *Toxicol. Appl. Pharmacol.* 44, 581-591.
- KAPPUS, H., BOLT, H. M., BUCHTER, A., AND BOLT, W. (1976). Liver microsomal uptake of [ $^{14}\text{C}$ ]vinyl chloride and transformation to protein alkylating metabolites *in vitro*. *Toxicol. Appl. Pharmacol.* 37, 461-471.
- MAKK, L., DELMORE, F., CREECH, J. L., OGDEN, L. L., FADELL, E. H., SONGSTER, C. L., CLANTON, J., JOHNSON, M. N., AND CHRISTOPHERSON, W. H. (1976). Clinical and morphologic effects of hepatic angiosarcoma in vinyl chloride workers. *Cancer* 37, 149-163.
- MALAVIELLE, C., BARTSCH, H., BARBIN, A., CAMUS, A. M., AND MONTESANO, R. (1975). Mutagenicity of vinyl chloride chloroethyleneoxide, chloroacetaldehyde, and chloroethanol. *Biochem. Biophys. Res. Commun.* 63, 363-370.

MALTONI, C.  
 genicity as  
*Ann. N.Y.*  
 TABERSHAW,  
 Mortality  
 vinyl chlo  
 16, 509-51  
 WATANABE,  
 E. O., A  
 [ $^{14}\text{C}$ ]vinyl  
 in rats. *To*  
 WATANABE, I

UCC  
 002873

veloping  
o VC.  
similar  
assess-  
levels of

SON, P.,  
ARTSCH,  
lation of  
nd vinyl  
67, 596-

SANO, R.  
mediated  
lmonella  
437.

APPEL,  
Meta-  
in vivo.

D BOLT,  
loride in

Angio-  
polyvinyl

ality ex-  
chloride  
chloride  
10.

JK. C. N.  
city data  
ivation:  
armacol.

D BOLT,  
C]vinyl  
alkylating  
acol. 37,

OGDEN,  
LANTON,  
W. H.  
f hepatic  
Cancer

CAMUS,  
agenicity  
chloro-  
em. Bio-

MALTONI, C., AND LEFEMINE, G. (1975). Carcino-  
genicity assays of vinyl chloride: Current results.  
*Ann. N.Y. Acad. Sci.* 246, 195-224.

TABERSHAW, I. R., AND GAFFEY, W. R. (1974).  
Mortality study of workers in the manufacture of  
vinyl chloride and its polymers. *J. Occup. Med.*  
16, 509-518.

WATANABE, P. G., MCGOWAN, G. R., MADRID,  
E. O., AND GEHRING, P. J. (1976a). Fate of  
[<sup>14</sup>C]vinyl chloride following inhalation exposure  
in rats. *Toxicol. Appl. Pharmacol.* 37, 49-59.

WATANABE, P. G., MCGOWAN, G. R., AND GEHRING,

P. J. (1976b). Fate of [<sup>14</sup>C]vinyl chloride after  
single oral administration in rats. *Toxicol. Appl.*  
*Pharmacol.* 36, 339-352.

WATANABE, P. G., HEFNER, R. E., JR., AND GEHRING,  
P. J. (1976c). Vinyl chloride induced depression of  
hepatic nonprotein sulfhydryl content and effects  
on bromosulphthalein (BSP) clearance in rats.  
*Toxicology* 6, 1-8.

WATANABE, P. G., ZEMPEL, J. H., PIGO, D. G., AND  
GEHRING, P. J. (1978). Hepatic macromolecular  
binding following exposure to vinyl chloride. *Toxi-*  
*col. Appl. Pharmacol.* 44, 571-579.