

Incidence of cancer among vinyl chloride and polyvinyl chloride workers

S STORETVEDT, HILDAAS, S L LANGAARD, AND A ANDERSEN

From the Research Center,^a Norsk Hydro a.s. Porsgrunn Fabrikker, 3900 Porsgrunn, Telemark Sentralsjukhus,^b Department of Occupational Medicine, 3900 Porsgrunn, and Cancer Registry of Norway,^c Montebello, Oslo 3, Norway.

ABSTRACT The results of a follow up study of the incidence of cancer and the mortality in a cohort of 454 male workers producing vinyl chloride and polyvinyl chloride are presented. The study population was restricted to employees with more than one year's work experience in the study plant between 1950 and 1969 and the cohort was followed up from 1953 to the end of 1979. Twenty three new cases of cancer were observed compared with 20.2 expected; one case of liver angiosarcoma was found. Five cases of lung cancer were found (2.8 expected) and four cases of malignant melanoma of the skin were observed (0.8 expected). The possibility of a causal relationship between exposure to vinyl chloride and the development of malignant melanomas is discussed.

Since 1974 when Creech and Johnson presented their report,¹ human exposure to vinyl chloride (VCM) has been associated with the occurrence of angiosarcoma of the liver. Several studies have confirmed these initial results as reviewed by Spirtas and Kaminski.² In animal studies where VCM was given to mice, rats, and hamsters by different routes of administration neoplasms have also been induced in other organs.³

Whether VCM has the potency to induce cancer in man in organs other than the liver has not yet been confirmed. Lung cancer,⁴⁻⁷ brain tumours,^{8,9} and cancer in the digestive system, primarily angiosarcoma of the liver,¹⁰⁻¹² have been suggested to be in excess in workers producing VCM and polyvinyl chloride (PVC), and the prime purpose of the present investigation was to study the incidence of cancer in workers exposed to VCM, with special emphasis on cancers other than liver tumours.

Material and method

PLANT DESCRIPTION

The study plant is located in the county of Telemark in south east Norway. In addition to VCM and PVC the company produces fertilisers, magnesium, and chlorine in adjacent plant complexes. The produc-

tion of VCM and PVC was started in 1950. VCM was produced from acetylene in the room where the polymerisation reactors were located. In 1967 the VCM production was partly altered, and cracking of ethylenedichloride (EDC) was introduced as one of two production methods. The production of VCM was discontinued in 1971 after which time the plant was operated as a polymerisation unit. From 1955 onwards VCM and PVC were produced in separate rooms; table 1 shows the volume of VCM and PVC production and the number of employees over time.

STUDY POPULATION

A list containing the names of employees who had started work before the end of 1974 and whose period of employment had exceeded one month was constructed from the personnel register provided by the plant. The health department of the company has kept the health records of all present and previous workers employed from 1940 and onwards. These records also contain some information on the place of work of each employee. By combining these

Table 1 Annual vinyl chloride and polyvinyl chloride production and number of employees in the study plant

Work period	VCM (tons)	PVC (tons)	No of employees
1950-55	1000-2500	1000-2500	45-50
1956-60	2500-5000	2500-5000	45-90
1961-70	5000-30000	5000-30000	90-130
>1971	None	30000-60000	120-130

Received 20 August 1982

Accepted 14 October 1982

two sources of information a list of names was constructed which was presumed to be complete. The individual records contained the following information on each worker: full name, date of birth, identity number, last known address, dates for beginning and terminating employment, departments in the plant at which each person had worked, and the duration of work in each department. Data were obtained for 1233 workers.

It was decided to restrict the analyses to male workers who had been employed for at least one year and who had first been employed at the plant before 1970. These limitations reduced the number of subjects to 454.

ESTIMATION OF EXPOSURE

As a consequence of the technical design of the plant and the process operation, the level of exposure to VCM in the past must have been high but no industrial hygiene survey had been performed before 1974. Estimates of the average atmospheric concentration in the past were based on sporadic measurements that had been carried out with an "explosion-meter" which was graded from 0% to 100% of the estimated lower explosion limit; 1%, equivalent to 400 ppm of VCM. Two sources of supplementary information for estimating the VCM concentration were used: interviews with workers, some of whom had been employed in the plant since the start of production, and a suggested odour threshold of about 500 ppm VCM. Based on this information, we have assumed that the VCM concentration was about 2000 ppm from 1950 to 1954, about 1000 ppm from 1955 to 1959, about 500 ppm from 1960 to 1967, and about 100 ppm from 1968 to 1974. During autoclave cleaning the exposure level may have been as high as 3000 ppm.¹⁴ Both emulsion PVC and suspension PVC were produced. The surplus monomer in the PVC leaving the man-

ufacturing plant before 1970 may have contained 500 ppm VCM or even higher concentrations.¹⁴

The plant was closed for reconstruction in October 1974 and since 1975 the VCM in the working atmosphere has been monitored continuously with automatic measuring devices, including alarm systems that warn the workers during even minor leakages. Since 1975 the VCM exposure level at the production units has been below 1 ppm most of the time.

JOB CLASSIFICATIONS

Jobs considered to have been associated with high VCM exposure include production, autoclave cleaning, maintenance, and tapping. Owing to the great amounts of surplus monomer in the PVC, one may also assume that the exposure level in the packing unit may have been high until 1970. Those working in the development laboratory which contained small autoclaves in small rooms, in the control laboratory, and at processing in the customers' service laboratory, may also have been heavily exposed, but no information about VCM levels in these working places is available. Nine different exposure categories were defined (table 2), a number of the workers having been employed in more than one exposure category. Membership of an exposure category was defined as that category in which the longest time had been spent. If an individual had been a production worker for some years and subsequently spent a greater number of years at a job that clearly entailed minimal exposure, he would by definition belong to a low exposure category.

As the exposure level was high in the early 1950s, an attempt was also made to classify the jobs as determined by the exposure level, using the number of "years of 500 ppm" exposure. The procedure was as follows: one year of production work in 1951,

Table 2. Observed (O) and expected (E) deaths from all causes, and all new cases of cancer in the study population except non-melanoma skin cancer

Exposure categories	No. of workers	All deaths			All cancers			Person-years 1953-79
		O	E	O/E	O	E	O/E	
01 Vinyl chloride (acetylene)	17	1	2.41	NC*	0	0.81	NC	368.5
02 Vinyl chloride (1-DX)	8	0	0.50	NC	1	0.20	NC	110.0
03 Research laboratory	48	2	4.32	0.46	0	1.69	NC	912.0
04 Polyvinyl chloride production	117	19	16.42	1.16	9	5.10	1.76	2308.5
05 Autoclave cleaning	17	5	3.04	1.64	2	1.03	1.94	309.5
06 Maintenance	43	4	7.64	0.52	1	2.87	NC	888.0
07 Packing/drying	95	5	8.44	0.59	5	2.95	1.69	1742.5
08 Customers service laboratory	59	7	8.77	0.80	3	3.06	0.98	1065.5
09 Various jobs	50	7	7.80	0.98	2	2.44	0.82	971.5
Total	454	50	59.34	0.84	23	20.16	1.14	8676.0

*NC = Not calculated.

ad Andersen

the working atmosphere contained about 2000 ppm VCM on average, was classified as four "500 ppm-years." The same type of work in 1961, when the atmosphere was about 500 ppm, would give one 500 ppm-year. This additional categorization of exposure could be given for VCM production, PVC production, autoclave cleaning, maintenance work, and PVC packing. Owing to the surplus monomer in the resin, one year of packing PVC would give about one 500 ppm-year for each year up to 1970, and it is even possible that this high level continued until 1974. In this way high exposure for short periods in workers categorised by jobs with low exposure level could be accounted for.

All workers were placed in an exposure category before we had any knowledge of their state of health or diagnoses.

FOLLOW UP

The main element of the method is to identify those individuals who meet the criteria for membership of the study population and to compare this population with a constructed Norwegian population whose age distribution was identical with that of the study group. (The method has been described in more detail by Langård *et al.*¹²)

Total mortality and cancer incidence from 1953 to the end of 1979 have been determined for the study population. The Cancer Registry in Norway has records of all new cases of cancer since 1953 and has access to information on all causes of death provided by the Central Bureau of Statistics. This study is based on a comparison of observed and expected total mortality and incidence of cancer for the period 1953-79. To estimate the expected number of cases of cancer, the national five year age specific incidence rates were used. According to the data of the Cancer Registry, the rates of cancer incidence in the Telemark county are between 0.90 and 0.95 of the national figures.¹³ Consequently, the more robust national rates could be used for reference purposes and the expected number of deaths are based on the national rates.

Results

Results

Deaths from all causes and the number of new cases of cancer, excluding non-melanoma skin cancer, are presented in table 2. The number of person-years at risk is also shown together with a tabulation of the nine different exposure categories. Among 454 men there were 50 deaths from all causes (expected 59.3).

During the 27 years of follow up, 23 new cases of cancer were found against 20.2 expected. The nine exposure categories have been combined into three groups reflecting high, medium, and low exposure (table 3); the increased incidence of cancer is accounted for almost entirely by the high exposure group.

There was only one liver angiosarcoma, recruited from exposure category (H), and this case has been included in a previous survey.⁷

Table 3 presents the observed and expected figures for some malignant neoplasms of interest. Five cases of lung cancer were observed in the whole study population compared with 2.8 expected. The mean time between first exposure and the time of diagnosis (latent period) in these five cases was 17.2 years (range 3-26) (not shown in the table). Four of the five cases were in the high exposure group (1.82 expected).

Four malignant melanomas of the skin were identified in the study population whereas only 0.8 was expected; three of the cases were observed in the high exposure group (0.5 expected). The tumours were located as follows: two on the trunk, one in the face-neck area, and one on a foot. After the observation period one more case has been diagnosed in the medium exposure group; this was in the face-neck area. We are aware of one case of incipient malignant melanoma located on the trunk

in the early 1950s. To classify the jobs as level, using the number of years spent. If an individual worked for some years, the number of years at minimal exposure, he was in the low exposure category.

Study population except

DIE	Person-years 1953-79
NC	368.5
NC	110.0
NC	912.0
1-76	2308.5
1-94	309.5
NC	888.0
1-69	1742.5
0-98	1065.5
0-82	971.5
1-14	8676.0

Table 3 Broad categories of work indicating exposure level. Observed (O) and expected (E) deaths from all causes, cases of cancer (all sites), colon cancer (ICD 153), bronchial cancer (ICD 162/163), malignant melanoma of the skin (ICD 190), and cancer of the thyroid gland (ICD 194). For work label, see table 2

Exposure level ^a	No. of workers	All cancers		ICD 153		ICD 162/163		ICD 190		ICD 194		Person-years 1953-79	
		O	E	O	E	O	E	O	E	O	E	O	E
Group 1	297	18	12.96	1.4	3	0.92	3.3	4	1.82	2.2	3	0.51	5.9
Group 2	107	3	4.75	0.6	0	0.34	NC	0	0.67	NC	1	0.18	NC
Group 3	50	2	2.44	0.8	0	0.18	NC	1	0.35	NC	0	0.10	NC
Total	454	23	20.16	1.1	3	1.44	2.1	5	2.84	1.8	4	0.79	5.9

^aGroup 1 (high vinyl chloride exposure) = Work labels 01, 02, 04, 05, 06, 07.

Group 2 (medium vinyl chloride exposure) = Work labels 03, 08.

Group 3 (low vinyl chloride exposure) = Work label 09.

NC = Not calculated.

Table 4. Observed (O) and expected (E) cases of cancer (all sites), lung cancer (ICD 162/163), and malignant melanoma of the skin (ICD 190) in relation to time of first employment in the plant

Years of first employment	No. of workers	All cancers			ICD 162/163			ICD 190			Person-years 1953-79
		O	E	O/E	O	E	O/E	O	E	O/E	
1950-4	105	6	7.21	0.8	2	1.01	2.0	1	0.23	NC*	2619.0
1955-9	128	8	6.51	1.2	2	0.92	2.1	1	0.25	NC*	2717.0
1960-4	123	6	4.33	1.4	0	0.59	NC*	2	0.19	11.1	2093.5
1965-9	98	3	2.12	1.4	1	0.29	NC*	0	0.12	NC*	1246.5
Total	454	23	20.16	0.8	5	2.82	1.8	4	0.79	5.1	8676.0

NC* = Not calculated.

Table 5. Observed (O) and expected (E) cases of cancer, all sites, of the colon (ICD 153), of the lung (ICD 162/163), malignant melanoma of the skin (ICD 190), and cancer of the thyroid gland (ICD 194), by 500 ppm-year exposure

No. of 500 ppm-years*	No. of workers	All cancers		ICD 153			ICD 162/163			ICD 190			ICD 194			Person-yr 1953-79	
		O	E	O/E	O	E	O/E	O	E	O/E	O	E	O/E				
<1	169	6	7.06	0.8	0	0.51	NC*	2	1.01	2.0	1	0.29	NC*	0	0.06	NC	3072.0
1-5	124	3	3.10	1.0	1	0.21	NC*	0	0.40	NC*	0	0.16	NC*	0	0.03	NC	2088.5
>5	161	14	9.95	1.4	2	0.74	2.7	3	1.44	2.1	3	0.33	9.1	2	0.06	33.5	3540.5
Total	454	23	20.16	1.1	3	1.44	2.1	5	2.84	1.8	4	0.79	5.1	2	0.16	12.5	8676.0

NC* = Not calculated.

*For classification of exposure for these cases see material and methods.

Table 6. Distribution of exposure categories among workers in whom cancer has been diagnosed (all cases of cancer). Figures give duration of exposure under different categories in years

Age when cancer was diagnosed	Time of first employment	Year of diagnosis	ICD No.*	Exposure categories†							
				01	02	03	04	05	06	07	08
63	1962	1970	151							2	
67	1964	1967	151								3
62	1956	1972	153					5			
67	1962	1973	153				11				
55	1968	1977	153							6	
56	1950	1971	155				22				
63	1950	1963	162								
67	1952	1978	162/177				3				
58	1955	1977	162/194				7			6	10
54	1958	1977	162					11		10	
56	1968	1971	162							2	
70	1950	1973	177				2				15
26	1961	1967	178						18		
45	1952	1979	181								
38	1952	1957	190				4				
45†	1953	1981	190				2				
40	1958	1980	190					4			
41	1961	1974	190								13
41‡	1961	1977	190								15
66	1964	1979	190				14				
62	1969	1972	193								
33	1957	1960	194			11					
55	1955	1966	199						2		

*ICD 151 (stomach cancer), ICD 153 (colon cancer), ICD 155 (cancer of the biliary passages and liver), ICD 162/163 (lung and cancer), ICD 177 (cancer of the prostate), ICD 178 (testis cancer), ICD 181 (bladder cancer), ICD 190 (malignant melanoma of the skin), ICD 194 (cancer of the thyroid gland).

†Exposure categories corresponding with the numbers in table 2.

‡This case occurred after the end of observation period.

§Incipient case.

pations have been associated with the development of these tumours. Tear gas (*o*-chloroacetophenone) and polychlorinated biphenyls²⁰ are chemicals that have been suggested as possible inducers. In our study population one more case of melanoma has been diagnosed after the closure of the study and before the increased incidence of tumour was known to us. This additional case considerably strengthens the association between exposure to VCM and the development of malignant melanomas. As the medical services in the Telemark county have been of a high quality and easily accessible, we do not believe that there has been any bias due to the presence of the occupational health service in the company. In fact, none of the four cases was diagnosed by the occupational physicians. One of the five subjects with skin cancer had survived until June 1982. The incidence of malignant melanomas in the rural areas of Telemark county is slightly higher than the national rate, while the incidence in urban areas is fairly equal to the national rate.²¹ All subjects with skin cancer had lived in the urban area.

We observed two cancers of the thyroid gland (0-16 expected); both cases occurred in the high exposure group and were of the same histological type. We are not aware of other studies indicating an excess of this type of cancer but as only two cases have been observed, no conclusions can be drawn.

In a Canadian study on workers exposed to VCM for five years or more Theriault and Allard showed an excess of cancer in the digestive system,² but the authors concluded that the excess is accounted for exclusively by cancer of the liver. In the present study we found a slight excess of cancer of the colon. Many of the other studies referred to include cancer of the liver in digestive cancers, and they conclude that the excess risk is due to this tumour. There are few reports on occupationally induced cancer of the colon and rectum although it has been indicated that asbestos workers, textile workers, and steel workers have an increased risk from these diseases.²² Watanabe *et al.* in their kinetic study showed some faecal elimination of ¹⁴C after the inhalation of ¹⁴C-VCM, but they discuss only urinary and pulmonary elimination in their paper.¹⁰

The results presented in the present study, considered with other epidemiological studies, seem to support the view that VCM may act as a multicarcinogen in man. This is in accordance with observations from animal studies.⁴ Some of our findings differ from those presented by other workers, and will be followed up later. The observation period is short for some of the members of the cohort, and the study group is small. Nevertheless, the fact that the excess of different cancers was seen in the group with the highest estimated exposure level streng-

thens the suggested association between VCM exposure and these types of cancers, in particular the malignant melanomas.

We thank the Plastic Division of Norsk Hydro for cooperation throughout the study. O. C. Hockman's critical comments were of great help. We thank Mr P. A. H. for linguistic help and Mrs U. Danielsen for typing the manuscript.

References

- Cressch JJ, Johnson MN. Angiosarcoma of liver in the manufacture of polyvinyl chloride. *JAMA* 1974;16:150-1.
- Spiras R, Kaminski R. Angiosarcoma of the liver in vinyl chloride polymerized chloride workers. 1977 update of the NIOSH register. *JAMA* 1978;239:427-9.
- Althaus C, Telemann G, Gilibert A, Conti G, Carrozzini D. Carcinogenicity bioassays of vinyl chloride monomer: a model of risk assessment on an experimental basis. *Environ Health Perspect* 1981;41:1-29.
- Althaus C, Peters JM, Johnson MN. Proportional mortality among vinylchloride workers. *Lancet* 1974;ii:397-8.
- Hehlhaus H, Gailley WK. Mortality study of workers in the manufacture of vinyl chloride and its polymers. *JMA* 1974;16:509-18.
- Wassweiler RJ, Strasser W, Wagoner JK, Jones J, Falk H, Can U. Mesoplastic risk among workers exposed to vinyl chloride. *Am J Hyg* 1976;211:40-8.
- Hallberg PA, Wood S, Eiler C, Sauer L, Kilian DJ. Mortality experience of workers in a vinyl chloride monomer production plant. *JAMA* 1979;241:195-202.
- Toy AJ, Collier PL. Mortality experience of workers exposed vinyl chloride monomer in the manufacture of polyvinyl chloride in Great Britain. *Br J Ind Med* 1977;34:1-10.
- Theriault G, Allard P. Cancer mortality of a group of Canadian workers exposed to vinyl chloride monomer. *JAMA* 1981;245:1-6.
- Harries AW. Vinyl chloride and the production of PVC. *Proceedings of the Royal Society of Medicine* 1976;69:277-81.
- Karsch K. PVC: health implications and production in. *Environ Health Perspect* 1976;17:107-15.
- Langård S, Andersen AA, Gylseth B. Incidence of cancer or leukaemia among and leukaemia workers. *Br J Ind Med* 1980;37:114-20.
- Cancer Registry of Norway. Incidence of cancer in 1972-1976. Oslo: The Norwegian Cancer Society, 1977.
- Kjelle G, Johansen AA, Langekreft (Borge) en oversikt hos Kreftregisteret materiale. *Tidsskr Nor Læge* 1982;102:480-4.
- Wegman DH, Peters JM. Cell lung cancer in selected workers: a case-control study. *JAMA* 1978;239:793-6.
- Axelsson O. Aspects on confounding in occupational epidemiology. *Scand J Work Environ Health* 1978;4:9.
- Dupont P, Fabry JP, Gradiski D, Magadur JL. An approach to industrial poisoning: blood kinetics and elimination of ¹⁴C-vinylchloride monomer (VCM). *Acta Pharmacol* 1977;31, suppl 1:142-5.
- Watanabe PG, McCosman GR, Madrid ED, Gehring PJ. ¹⁴C-vinyl chloride following inhalation exposure in rats. *Appl Pharmacol* 1976;37:49-59.
- Magnus K. Habits of sun exposure and risk of melanoma: an analysis of incidence rates in Norway 1977 by cohort, sex, age and primary tumour site. *1981* 48:2329-35.
- Sklar AJ, Fitzpatrick TB. Genetic and environmental factors in malignant melanoma in man. *Pigment Cell* 1979;3:8.
- Jansson H. Colorectal cancer—an occupational disease. *Scand J Work Environ Health* 1978;4:321-3.