Table 3 Pertinent Physical Findings by Exposure Subgroup

Characteristic	Unexposed		Ever PCP		Ever PCP Chloracne		Only PCP		Only PCP Chloracne	
	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
Hyperpigmentation	27	(8.9)	28	(7.7)	7	(10.8)	22	(8.5)	5	(16.1)
Hypertrichosis[1]	1	(0.3)	4	(1.1)	3	(4.6)*	2	(0.8)	2	(6.5)*
Abnormal liver consistency[2]	1	(0.3)	4	(1.1)	2	(3.1)	4	(1.5)	2	(6.5)*
Absent right ankle reflex[2,3]	25	(8.3)	48	(13.1)	6	(9.2)	39	(15.0)	3	(9.7)
Absent left ankle reflex[2,3]	23	(7.6)	50	(13.7)	8	(12.3)	39	(15.0)	3	(9.7)
Abnormal vibration [2,3]	125	(41.4)	194	(53.0)	30	(46.2)	140	(53.9)	16	(51.6)
Stocking loss of pinprick[2,3]	23	(7.6)	50	(13.7)	12	(18.5)	35	(13.5)*	7	(22.6)
Tremor, positional, upper[4] extremity	23	(8.2)	44	(13.2)	13	(21.3)	35	(14.7)	9	(30.0)***

[1] *Unadjusted p*
Dichotomous exposure
* $.01 < p \leq .05$

[2] P value adjusted for age and alcohol
Dichotomous exposure
* $.01 < p \leq .05$
** $.001 < p \leq .01$
*** $p \leq .001$

[3] P value adjusted for age, height, and alcohol and excluding participants with diabetes, renal failure, and neurotoxic medications (chloramphenicol, nitrofurantoin, vincristine, and hydralazine)

[4] P value adjusted for age, alcohol, mercury exposure, and current smoking and excluding participants on theophylline and epinephrine

STLCOPCB4043948