

the synthetic fibers of glass wool apparently are too inflexible to pass easily through the nose, pharynx, trachea and bronchi and seldom reach the terminal bronchioles and alveoli.

Inhaled particulate matter comes to rest throughout the terminal air spaces (alveolar duct, atria, alveoli) in all parts of the lungs; inhaled asbestos fibers are first retained in the respiratory bronchioles. These very small tubes are immediately distal to bronchioles lined by ciliated epithelium. Their own essential lining is a low cuboidal type of epithelium but, as their name implies, they actually function in respiration through lateral alveoli given off as pouches along their walls. Either these pouches, or the abrupt change in the character of the lining epithelium, or the decrease in diameter of the tube, or perhaps the combination of all three factors is responsible for local retention of the inhaled fiber. Only after asbestosis is well established are appreciable numbers of fibers carried into the more peripheral air spaces.

6. Rate of Tissue Reaction to Asbestos Fibers

The rate of tissue reaction to asbestos is much more rapid than to an active dust like quartz. Evidences of tissue response appear as soon as fibers have localized in sufficient concentration in specific areas. In rats receiving asbestos fibers by intratracheal injection this evidence is visible as early as two weeks after injection; for quartz dust the latent period might be two months or more.

The behavior of the tissue reaction to inhaled dust after the termination of exposure is not the same in silicosis as in asbestosis. In silicosis, the young nodules become larger; in asbestosis, young scar tissue, that may have formed, contracts and becomes more dense but the area of involvement decreases in size. If exposure to asbestos dust is terminated after a brief period, the recently-inhaled