

FILE NAME: Oil Industry and American Petroleum Institute (API)


DATE: 1959

DOC#: API112

DOCUMENT DESCRIPTION: Book Excerpt - Occupational Cancers & Industrial Carcinogens

Editor in Chief: ANTHONY J. LANZA, M.D.


Consulting Editor: RICHARD H. ORR, M.D.



Industrial Carcinogens

By R. E. ECKARDT, M.D., Ph.D., F.A.C.P.

Director, Medical Research Division, Esso Research and Engineering Company, Linden, N. J.; Associate Clinical Professor of Industrial Medicine, New York University, Postgraduate Medical School; Instructor in Medicine, Cornell University, Medical School



GRUNE & STRATTON

1959

NEW YORK AND LONDON

to the crude saltpeter. At best it can probably be said at this time that this occupation requires further intensive study.

In 1929, Martland reported on bone sarcomas among radium watch-dial painters. These cases have received so much publicity and attention, that it suffices here only to mention them in their historical perspective.

In 1932, Grenfell reported the occurrence of lung and sinus cancer cases among workers in the Mond nickel works. It is believed by most people today that these cases resulted from long continued exposures to low concentrations of nickel carbonyl, a highly toxic gas in higher concentrations; but this is not definitely established as yet.

In 1935, Lynch and Smith reported lung cancers among asbestos workers who had developed asbestosis. Although there are still those who do not agree thoroughly that asbestos was the cause in these cases, most authorities believe that the incidence of lung cancer among asbestosis cases is higher than would be anticipated.

Finally in 1952, Weil et al. reported nasal sinus cancers among workers in isopropyl alcohol manufacture. Although a waste product called isopropyl oil is suspect as the specific agent, considerable doubt exists as to the exact etiologic agent.

In reviewing the historical aspects of occupational cancer, it is essential to realize that cancer of the scrotum occupies a unique position. This is so because cancer of the scrotum is such a rare disease that it is easy to associate it with a given occupation if more than one or two cases are observed. Only about 15 cases of scrotal cancer occur per 100,000,000 males per year. Thus, whenever cancer of the scrotum is observed, it should be considered occupational in origin until it is proven otherwise. Even in cases reported as having occurred among occupations not usually associated with exposure to carcinogens* (such as commercial, finance and insurance occupations) a complete and careful occupational history from childhood to the development of the disease seems warranted. Such a history, if carefully taken, may disclose earlier occupational contacts which really were responsible for the development of the disease. Thus, many individuals who enter more sedentary occupations as an adult, may have engaged in strenuous and dirty occupations in adolescence. Thus, a few years ago, a prominent New York newspaper featured a picture of the son of the Governor of the State of New York engaged in a summer occupation of road-building. He was pictured working, bare-

* See page 37 of Reference 3.

backed, with a shovel. The being listed as "day laborer code. The sole purpose of this occupation of an individual present occupation, whereas, a disease such as occupational cancer that it may have been founded of it by careful quest

Although it is relatively c of the scrotum with occupati cancers of other sites. Thus, to incidence of about 30 pe times greater incidence than of cancer of the scrotum ma the occupational origin of th cancer of the skin, or cancer between the occupation and sound to attribute a given ty be established with statistical of cancer among employees i in an employee group compa occupation. To do so requir cancers, but also of the pop for the occurrence of cancer of occupational cancer mus meaningful epidemiological been raised by observations cautioned not to make prem dence has a high statistical the field of occupational ca admonition and hence the lit pational cancers that cannot evidence for their occupation

The history of occupation

Examination of this table important to point out in or can be kept in proper perspec

First, over 75% of the o Of these, about 60%, or ab

n the bronchial epithelium
jority of cases. However,
ese cancers is available.
y the end of 1950. Early
ubtly will be the most

n of dust respirators may
se cases. Thus, the Chief
cancers, and only 2 lung
yment started after 1924,
In addition, no cases have
out both in Canada and

se cases has not yet been
gest that it is nickel car-
nature of this compound,
difficult to devise experi-
re of animals without pro-
ated finely divided nickel,
ds with the production of
y decomposes on contact
monoxide, it has always
es, such as the lung and
divided nickel. Although
te theory that nickel car-
be accepted as conclusive
ere not involved. In view
various commercial chem-
rseed oil to make edible
er alcohols from hydro-
nite answers about nickel
adily formed from nickel
tions of temperature and
used to flush out units
0% of carbon monoxide.
l carbonyl have occurred
s in which nickel was the
nickel carbonyl may be
ving if nickel carbonyl is

responsible for the cancers observed in the nickel refining industry in England—and if so—in what amounts.*

Asbestos and Asbestosis

Clinical Features—The lung cancers seen in cases of asbestosis are generally squamous cell carcinomas arising from the bronchial epithelium. In addition to the cancers observed, squamous metaplasia of the epithelium in various adjacent and distant regions of the bronchial epithelium is observed. The most complete analysis of a large series of these cases is contained in the Annual Report of the Chief Inspector of Factories of England for the year 1954.³ In this report, a total of 344 asbestosis deaths was analyzed, of which 205 were men and 139 women. Cancer of the lung was observed in 55, or 16%. Among the men, 41 lung cancers, or 20%, were observed while 14 were seen in women, or 10.1%. This makes the male:female ratio for lung cancer approximately 2:1, which compares with a male:female ratio in the general population of 5:1 or greater. The mean age at death of men with lung cancer and asbestosis was 55 years, while the mean age in women with lung cancer and asbestosis was 45.6 years. This compares with a mean age of 53.7 years for men with asbestosis without lung cancer and 42.8 for women with asbestosis without lung cancer. The mean age at death for asbestosis complicated by tuberculosis was 45.2 years in men and 35.1 years in women. These data suggest that asbestosis, uncomplicated by lung cancer poses a greater threat to longevity than when lung cancer is present. Thus, the control of asbestosis should eliminate this occupational cancer.

No case of lung cancer developed in an individual, male or female, who had less than 2 years exposure, although 7 cases, all in women, of asbestosis without lung cancer or tuberculosis, and 5 cases, again all in women, of asbestosis with tuberculosis developed with exposures of less than 2 years duration. The first cases of lung cancer in women developed when they were between 30–39 years of age, whereas in men the first cases appeared in the 40–49 year age group.

The diagnosis of occupational cancer in asbestos workers is considerably simplified in comparison with other occupational cancers because

* In a recent private communication from Dr. J. Gwyne Morgan, Chief Medical Officer of the Clydach works of the Moud Nickel Company, Ltd., calcined dust, rather than nickel carbonyl was proposed as the etiologic agent. Dr. Morgan has prepared a complete manuscript of his observations which is now in press.

of the requirement of the pre-existence of asbestosis. This chronic fibrosing disease of the lung could hardly be missed at the autopsy table. Another factor which supports the occupational origin of this type of cancer is the presence of asbestos bodies. These bodies have been well described by Lynch,¹⁷ but a review of their characteristics may be helpful. They vary from 10 to more than 100 microns in length, and from about 1 to 12 microns in thickness. They have a central translucent fiber, presumably asbestos, surrounded by a shiny, golden or brownish substance. They may have knobs at one or both ends. The shaft may frequently be beaded. Spherical forms of various sizes may also be seen, although these may be the longer-knobbed-end structures seen on end under the microscope, or broken off from the shaft. Characteristically, they give a Prussian Blue reaction for iron. In cases of asbestosis, they are seen usually in the bronchial tract, the adjacent alveoli and the peribronchial connective tissues and lymph nodes. In the alveoli, they may be enclosed by mono- or multi-nuclear phagocytes or foreign body giant cells. In the connective tissue they may cause some surrounding fibrosis. They may be found in the sputum, but their presence there is not essential, since they are not found in the sputum of some advanced cases of asbestosis. It would seem that a case of lung cancer in an asbestos worker should not be diagnosed as occupational in origin unless asbestosis and/or asbestos bodies have been found. It should be remembered, however, that asbestos bodies may be absent from advanced cases of asbestosis or present in individuals with no evidence of asbestosis. Their presence is only an indication of exposure to asbestos. Their absence has no significance.

Lung cancer in asbestosis cases seems not to vary from lung cancer in general. The prognosis and treatment are identical, although the embarrassment of pulmonary reserve by the fibrosis in the lungs may limit the cases in which pneumonectomy can be performed.

Etiologic Agent—It is not clear at present whether these cancers result from the asbestos per se, or whether they arise as a result of some other mineral impurities contained in certain forms of asbestos. Although this may seem academic, it is of importance for two reasons. First, it may help us understand carcinogenic mechanisms, and second, if some forms of asbestos do not lead to lung cancer formation, this may be extremely important from a practical and medico-legal standpoint. A recent study by Braun and Truan⁵ denies the existence of a lung cancer hazard in the Canadian asbestos industry, even though asbestosis may be observed.

This suggests that the from that for lung cancer impurities present in squamous metaplasia is that asbestos only serves simply as a suit the malignant changes carcinogen was present may be unimportant, promoted by asbestos w

Isopropyl Alcohol M

In 1952, Weil, Smyt one case of lung cancer loma of the vocal cord facture of isopropyl a another manufacturer and two cancers of the experience of these two sinus and intrinsic lary lung cancer case obser occupation, since it has should serve to alert the developments in lung c

Etiologic Agent—De what the causative age volved, evidence was c began their work in the suggests that whatever removed by alterations date. The most signifi about this time was a r In brief, the process acid to form isoprop water to form isoprop alcohol is distilled off trated for re-use. Whe high, a small amount o oil. This is a complex

s. This chronic fibrosis at the autopsy table. origin of this type of nodules have been well characteristics may be helpful in length, and from central translucent fiber, dense or brownish substance. The shaft may frequently also be seen. Structures seen on end of. Characteristically, in cases of asbestosis, they in the alveoli and the periphery of the alveoli, they may be foreign body giant cells surrounding fibrosis. Since there is not essentially advanced cases of asbestosis in an asbestos worker unless asbestosis is remembered, however, advanced cases of asbestosis. Their absence has

ity from lung cancer is not identical, although the similarity in the lungs may be formed.

These cancers result from the result of some other asbestos. Although this is not the case, First, it may be noted, if some forms of asbestosis may be extremely rare. A recent study of lung cancer hazard in asbestos may be observed.

This suggests that the causative agent of asbestosis may be different from that for lung cancer. This lends some support to the concept of impurities present in some asbestos but not in others. Finally, since squamous metaplasia is also observed in practically all cases, it may be that asbestos only serves to produce the squamous metaplasia, which serves simply as a suitable field for some other carcinogen to produce the malignant changes. Cancer, then, would only develop where this carcinogen was present. From a medico-legal standpoint, of course, this may be unimportant, since whether lung cancer is produced or only promoted by asbestos would have little significance.

Isopropyl Alcohol Manufacture

In 1952, Weil, Smyth and Nale²⁶ reported 4 cases of sinus cancer, one case of lung cancer, one carcinoma of the vocal cord and one papilloma of the vocal cord among a group of men involved in the manufacture of isopropyl alcohol. This experience has been confirmed by another manufacturer of isopropyl alcohol, where three sinus cancers and two cancers of the intrinsic larynx were found. From the combined experience of these two plants, it seems possible to conclude that the sinus and intrinsic larynx cases were related to occupation. The single lung cancer case observed can hardly be said to have any relation to occupation, since it has no statistical significance. Its presence, however, should serve to alert the profession to continue careful review of future developments in lung cancer incidence in this operation.

Etiologic Agent—Despite extensive animal testing, it is not clear today what the causative agent in these cases was. In each of the plants involved, evidence was developed that no cancers appeared if employees began their work in the plant after a particular date. This observation suggests that whatever the causative agent may have been, it was removed by alterations of the manufacturing process subsequent to that date. The most significant change in the process, which occurred at about this time was a reduction of the strength of the sulfuric acid used. In brief, the process consists of bubbling propylene through sulfuric acid to form isopropyl sulfates. This is subsequently hydrolysed by water to form isopropyl alcohol and weak sulfuric acid. The isopropyl alcohol is distilled off the weak sulfuric acid, and the acid reconcentrated for re-use. When the concentration of the sulfuric acid used was high, a small amount of impurities was formed which is called isopropyl oil. This is a complex, variable mixture of isopropyl ethers, isopropyl

of the prolonged inhalation of the control of the dustiness with the provision of exhaust dustiness. Use of wet processes of good housekeeping routine. Elimination of all unnecessary dust may settle will also help cannot be controlled by any should be provided. The use of one for work clothes, seems washing or showering facilities in the working environment are increasing, the reasons for

Individuals selected for the examination. This should include respiratory abnormalities, or an A pre-placement chest X-ray examination. Any pulmonary abnormalities before the individual is examined should be re-examined. Sputum examinations for asbestos bodies from the sputum should be present in the sputum should be of excessive exposure for asbestos bodies include asbestos hydroxide, followed by asbestos bodies are spread over several examined. The electron microscope bodies more carefully, but fracture.

It should be remembered that there is a wide variety of other occupations and activities that dust can be present in. It is not permitted to be prolonged, repeated exposure to a hazard exists. Each such exposure has the potential hazard. There is a variation in the susceptibility to dust while others may have no effect, even after considerable exposure.

Asbestos Handling

The solution of the problem of lung cancer resulting from asbestos lies in the control of the problem of asbestosis. The disease is the result

f control measures and would
onal personal or unit control

e sinus and lung cancer cases
sure in this case would be to a
itated by more precise knowl-
nowledge has been developed,
are, designed to contain within
inhalable materials, gaseous or
he best, and wherever gaseous
ned or ventilated, use of respi-
ickel carbonyl decomposes on
e principle might be taken in design-
control equipment.

am in this industry should be
and lung cancers. A thorough
nasopharynx, performed prior
t periodic intervals, should be
thoroughly familiar with the
s, and preferably the periodic
done by the same individual in
rays of the sinuses and chest
nation and interpreted by some-
a. Any changes observed should
ative diagnosis has been made.
pt to correlate urine or blood
atmospheric nickel concentra-
nal check of exposures. When-
other means, the men should be
l instructed adequately in the
the men in the importance of
the time of the periodic re-

cancer resulting from asbestos
stosis. The disease is the result

of the prolonged inhalation of asbestos dust, so that its control lies in the control of the dustiness of the operation. Enclosure of equipment with the provision of exhaust ventilation should eliminate much of the dustiness. Use of wet processes wherever possible, and the establishment of good housekeeping routines to clean up accumulated dust will help. Elimination of all unnecessary ledges, nooks and areas where asbestos dust may settle will also help control this operation. Wherever dustiness cannot be controlled by any of these means, adequate dust respirators should be provided. The use of double lockers, one for street clothes and one for work clothes, seems warranted in this industry, but no special washing or showering facilities seem necessary. Periodic dust measurements in the working environments should be undertaken and if levels are increasing, the reasons for this should be sought.

Individuals selected for this work should have a careful pre-placement examination. This should be designed specifically to detect any pulmonary abnormalities, or any encroachment on the pulmonary reserve. A pre-placement chest X-ray should be included as a part of this examination. Any pulmonary abnormalities found should be carefully evaluated before the individual is permitted to work in this occupation. These examinations should be repeated at yearly intervals, at which time sputum examinations for asbestos bodies should be done. The absence of such bodies from the sputum has no special significance, but their presence in the sputum should serve to alert the physician to the possibility of excessive exposures. The technique of examining the sputum for asbestos bodies includes digestion of the sputum by warm sodium hydroxide, followed by centrifugation and decanting. The decanted residues are spread over several slides, mounted in Canada balsam and examined. The electron microscope can also be used for examining these bodies more carefully, but great care must be exercised or they may fracture.

It should be remembered that asbestos is widely used in a large variety of other occupations and industries. Whenever it is used in such a way that dust can be produced, and the work is of such a nature as to permit prolonged, repeated inhalation of the dust, a potential asbestosis hazard exists. Each such use must be carefully evaluated in terms of the potential hazard. There seems to be a considerable individual variation in the susceptibility to asbestosis, some people developing it rapidly, while others may have comparable exposures with no signs of the disease, even after considerable periods of exposure.