

Using Urinary Biomarkers to Elucidate Dose-related Patterns of Human Benzene Metabolism

Sungkyoon Kim¹, Roel Vermeulen², Suramya Waidyanatha¹, Brent A. Johnson¹, Qing Lan², Nathaniel Rothman², Martyn T. Smith³, Luoping Zhang³, Guilan Li⁴, Min Shen², Songnian Yin⁴, and Stephen M. Rappaport^{1*}

¹ School of Public Health, University of North Carolina, Chapel Hill, NC 27599, USA.

² National Cancer Institute (NCI), National Institutes of Health (NIH), Department of Health and Human Services (DHHS), Bethesda, MD 20892, USA.

³ School of Public Health, University of California, Berkeley, CA 94720, USA.

⁴ Chinese Center for Disease Control and Prevention, Beijing, China.

* To whom requests for reprints should be addressed, at CB 7431, School of Public Health, University of North Carolina, Chapel Hill, NC 27599. Phone: (919) 966-5017; Fax: (919) 966-0521; E-mail: stephen_rappaport@unc.edu.

Grant support:

NIEHS through grant P42ES05948 and P30ES10126 (S. M. R.) and RO1ES06721, P42ES04705, and P30ES01896 (M.T.S.)

ABSTRACT

Although the toxicity of benzene has been linked to its metabolism, the dose-related production of metabolites is not well understood in humans, particularly at low levels of exposure. We investigated unmetabolized benzene in urine (UBz) and all major urinary metabolites [phenol (PH), *E,E*-muconic acid (MA), hydroquinone (HQ), and catechol (CA)] as well as the minor metabolite, *S*-phenylmercapturic acid (SPMA), in 250 benzene-exposed workers and 139 control workers in Tianjin, China. Median levels of benzene exposure were about 1.2 ppm for exposed workers (interquartile range: 0.53–3.34 ppm) and 0.005 ppm for control workers (interquartile range: 0.002 – 0.007 ppm). (Exposures of control workers to benzene were predicted from levels of benzene in their urine). Metabolite production was investigated among groups of 30 workers aggregated by their benzene exposures. We found that the urine concentration of each metabolite was consistently elevated when the group's median benzene exposure was at or above the following air concentrations: 0.2 ppm for MA and SPMA, 0.5 ppm for PH and HQ, and 2 ppm for CA. Dose-related production of the 4 major metabolites and total metabolites ($\mu\text{mol/l}$ per ppm benzene) declined between 2.5 and 26 fold as group median benzene exposures increased between 0.027 and 15.4 ppm. Reductions in metabolite production were most pronounced for CA and PH below one ppm, indicating that metabolism favored production of the toxic metabolites, HQ and MA, at low exposures.

Key words: benzene, metabolism, urinary biomarkers, dose

Abbreviations

BO, benzene oxide; CA, catechol; CV, coefficient of variation; EI, electron ionization; GC-MS, gas chromatography-mass spectrometry; GM, geometric mean; HQ, hydroquinone; LOD, limit of detection; MA, *E,E*-muconic acid; PH, phenol; SPMA, *S*-phenylmercapturic acid; TMS-, trimethylsilyl derivative; UBz, urinary benzene.

INTRODUCTION

Benzene is an important industrial chemical that is also emitted into the air from gasoline, engine exhausts, and combustion of organic materials (including cigarette smoke) [1,2].

Occupational exposures to benzene at air levels greater than about 10 ppm, have long been linked to hematotoxicity and to acute myelogenous leukemia [3-5]. A recent report of hematotoxic effects in workers exposed to benzene below one ppm [6] has raised additional concerns regarding the health consequences of low exposures to this contaminant.

While it is generally accepted that benzene causes toxic effects via metabolism, the particular toxic metabolite(s) remains elusive [7]. The major metabolic pathways are shown in Figure 1. Benzene is metabolized by CYP enzymes (primarily CYP2E1) to benzene oxide (BO, which is in equilibrium with its tautomer, oxepin), an electrophile that binds to macromolecules [8-11] and is the source of all other metabolites. Spontaneous rearrangement of BO produces phenol (PH), which can undergo another CYP oxidation to give hydroquinone (HQ). Hydrolysis of BO via epoxide hydrolase produces benzene dihydrodiol which can be converted to catechol (CA), via dihydrodiol dehydrogenases, or to benzene diolepoxides via CYP oxidation. Hydroquinone and CA can be oxidized to 1,4-benzoquinone and 1,2-benzoquinone, respectively, which also bind to macromolecules [11-13]. A second CYP oxidation of oxepin, followed by ring opening, produces the muconaldehydes. These reactive species are also capable of binding to macromolecules [14-17] and are ultimately converted to *E,E*-muconic acid (MA) [18]. The major phenolic metabolites, i.e., PH, HQ, and CA are easily conjugated and excreted in urine [19-21]. Minor pathways of benzene metabolism include *S*-phenylmercapturic acid (SPMA) following reaction of BO with glutathione [16], and 1,2,4-trihydroxybenzene via CYP oxidation of HQ. Of all the benzene metabolites, 1,4-benzoquinone (derived from HQ) has most often been linked to the spectrum of toxic effects observed in humans and animals [7,22-24].

Since risks of hematotoxicity and leukemia are thought to arise from benzene metabolism, it is important that the dose-related production of benzene metabolites be understood. Two recent studies have reported levels of urinary metabolites in both benzene-exposed workers and control workers in China [25-28]. Metabolite profiles from those studies suggested shifts among the competing metabolic pathways at increasing levels of benzene exposure. The Chinese studies also pointed to relatively high levels of the phenolic metabolites, PH, CA and HQ in control workers, presumably from dietary and endogenous sources of these compounds. These background levels of metabolites tend to obscure the amounts of PH, CA and HQ derived from ambient benzene exposure and from cigarette consumption. Also, because benzene air concentrations levels were not reported for control subjects in previous Chinese studies, they could not be used to determine the utility of the various metabolites as biomarkers of exposure to benzene at environmental levels.

In the present study we report levels of benzene in air and urine, as well as urinary levels of PH, CA, HQ, MA and SPMA, among 389 subjects (250 exposed and 139 controls) in a Chinese population that has shown evidence of hematotoxic effects [6]. Because some personal measurements of airborne benzene were below the limits of detection (LOD) or were missing, we predict benzene exposures in these (mostly control) subjects from the levels of unmetabolized benzene in urine (UBz). Then, we use the data to characterize relationships between metabolite levels and benzene exposures in groups of subjects stratified by exposure level. This analysis identifies useful ranges of the various metabolites as biomarkers of benzene exposure. Finally, we estimate the dose-related production of each metabolite ($\mu\text{mol/l}$ urine per ppm benzene) after adjusting for background levels.

MATERIALS AND METHODS

Chemicals

Benzene, CA (99%) and HQ (>99%) were obtained from Fluka Chemical Co. (Switzerland). NaCl was purchased from Fisher Scientific (Pittsburg, PA) and was heated at 120°C overnight prior to use. PH (99%+, redistilled), MA (98%), and [$^2\text{H}_6$]benzene (99.86%) were obtained from Aldrich Chemical (Milwaukee, WI). [$^{13}\text{C}_6$]PH (99%) and [$^2\text{H}_6$]CA (98%) were purchased from Cambridge Isotope Laboratories (Woburn, MA). Phenylmercapturic acid, Tri-Sil reagent, and concentrated hydrochloric acid (Optima grade) were purchased from TCI America (Portland, OR), Pierce (Rockford, IL), and Fisher Scientific (Pittsburgh, PA), respectively. Ethyl acetate (analytical reagent grade) and hexane (nanograde) were obtained from Mallinckrodt Baker (Paris, KT). Anhydrous Na_2SO_4 was purchased from J.T. Baker (Phillipsburg, NJ). 2,5- $^{13}\text{C}_2$]MA, [$^2\text{H}_5$]SPMA and [$^{13}\text{C}_6$]HQ were kindly provided by Drs. Avram Gold, Ramiah Sangaiah and Alistair Henderson from the University of North Carolina at Chapel Hill. All headspace solid phase microextraction (HS-SPME) supplies were obtained from Supelco (Bellefonte, PA).

Subject Recruitment and Sample Collection

Subjects ($n = 390$) were recruited with informed consent from five factories in Tianjin, China. Exposed subjects ($n = 250$) worked in two shoe-making factories, and control subjects ($n = 140$) worked in three clothes-manufacturing factories as previously described [6,29]. The eligibility and exclusion criteria for participation were the same as in previous reports, and one control subject who did not provide a urine sample was removed. Exposed and control subjects were frequency matched by region and gender. As shown in Table 1, demographic characteristics were similar among exposed and control workers in terms of gender, age and

smoking status. Smoking status was highly correlated with gender. Personal protective equipment was not used.

Full-shift exposures to benzene were monitored with passive samplers (Organic Vapor Monitors, 3M Corp., St. Paul, MN), and urine was collected from each participant at the end of each working shift for which air had been monitored. Air samples were analyzed for benzene by gas chromatography with flame-ionization detection as described previously [29]. The nominal LOD was 0.20 ppm of benzene in air.

Urine samples were collected repeatedly (up to 4 times) in 139 subjects at intervals of about 42 days (median value) from June 2000 to June 2001. The other 250 subjects provided a single urine specimen. Urine samples were aliquoted immediately after collection, and stored at -80°C for 1.6 – 4.0 y (for UBz) and 2.2 – 3.8 y (for urinary metabolites) prior to analysis. Samples were identified by randomly assigned numbers. Information about exposure levels and demographic factors were released after all assays had been completed and results reported to collaborators. A total of 620 urine specimens were analyzed from 389 subjects. Urinary creatinine was quantified with a commercial kit (Monarch Instrument, Amherst, NH) as described previously [25].

For quality control purposes, positive controls were assayed with each batch of urine samples. They were prepared from a single urine specimen obtained from an unexposed volunteer to which had been added sufficient quantities of standard analytes to achieve the following concentrations: UBz – 24.5 nmol/l, MA – 32.3 µmol/l, SPMA – 2.65 µmol/l, and PH – 92.8 µmol/l, CA – 94.1 µmol/l and HQ – 69.7 µmol/l. Batches of positive controls were aliquoted into 2-ml vials and stored at -80°C prior to analysis.

This study was approved by the Institutional Review Boards of the University of North Carolina, the University of California, Berkeley, the U.S. National Cancer Institute and the Chinese Academy of Preventive Medicine.

Measurement of Urinary Benzene

Levels of UBz were determined according to the method of Waidyanatha *et al.* [26], with minor modifications. Briefly, to 0.5 ml of urine in a 2-ml vial containing ca. 0.5 g of NaCl was added one μl of 59.4 $\mu\text{mol/l}$ [$^2\text{H}_6$]benzene in methanol (internal standard). The vial was immediately sealed with a PTFE/silicone septum and benzene was extracted from the headspace with a 30- μm polydimethylsiloxane fiber (Supelco, PA) at 40°C for 15 min. Samples were analyzed by gas chromatography-mass spectrometry (GC-MS) in electron ionization (EI) mode using a HP 5980 Series II gas chromatograph coupled to a HP 5971-A mass selective detector. Sample injections were performed with a Varian Model 8200 autosampler (Walnut Creek, CA) equipped with a carousel (Strumenti Scientific, Padova, Italy). The injector, MS-transfer line and ion source temperatures were 200, 280, and 168°C, respectively. A DB-1 fused silica capillary column (60 m, 0.25-mm i.d., 0.25- μm film thickness) was used with He as the carrier gas at a flow rate of 1 ml/min. The GC oven was held at 45°C for 9 min, and was then increased at 10 °C/min to 100 °C. Late eluting compounds were removed by rapidly raising the oven temperature to 250°C. The mass spectrometer was operated with an electron energy of 70 eV, and ions were monitored at m/z 78 and 84 for benzene and [$^2\text{H}_6$]benzene, respectively. The retention times were 8.56 and 8.50 min for benzene and [$^2\text{H}_6$]benzene, respectively. Quantification was based on peak areas relative to the isotopically labeled internal standard.

Measurement of Benzene Metabolites

Levels of CA, HQ, MA, PH and SPMA were determined according to the method of Waidyanatha *et al.*, with minor modifications [30]. Briefly, to 0.5 ml of urine in a 4-ml vial was added 10 μ l of a mixture (in methanol) of 1.73 mmol/l [$^{13}\text{C}_2$]MA, 0.205 mmol/l [$^2\text{H}_5$]SPMA, 6.24 mmol/l [$^{13}\text{C}_6$]PH, 2.19 mmol/l [$^2\text{H}_4$]CA, and 2.15 mmol/l [$^{13}\text{C}_6$]HQ (internal standards). After adding 50 μ l of concentrated HCl, the mixture was extracted with 1.5 ml of ethyl acetate and the organic phase, containing MA, SPMA, and free PH, CA, and HQ, was transferred to another 4-ml vial. Then, a 10- μ l aliquot of a mixture containing the above quantities of internal standards of PH, CA, and HQ was added to the aqueous phase, which was heated at 100°C for one h to hydrolyze the conjugates. After cooling to room temperature, the aqueous phase was extracted with an additional 1.5 ml of ethyl acetate. The two organic extracts were combined, dried with anhydrous Na_2SO_4 , reduced under N_2 , transferred to a 500- μ l flat-bottomed insert, and brought to dryness under N_2 . The residue was dissolved in 100 μ l hexane and derivatized with 100 μ l Tri-Sil reagent at 70°C for one h.

Samples were analyzed by GC-MS in EI mode with the same instrumentation described above using automated liquid injection. A 2- μ l injection was analyzed in splitless mode for SPMA; here, the GC oven was maintained at 75°C for 4 min, and was then increased at 8°C/min to 245°C, where it was held for 10 min. A 1- μ l portion was analyzed in (1:20) split mode for PH, CA, HQ, and MA; here, the GC oven was maintained at 75 °C for 4 min, and was then increased at 10 °C/min to 230°C and held for 4 min. In both cases, late eluting compounds were removed by rapidly raising the oven temperature to 270°C and holding this temperature for 15 min. The following ions were monitored for trimethylsilyl (TMS)-derivatives of the analytes: TMS-PH [m/z 166, M^+], TMS-[$^{13}\text{C}_6$]PH [m/z 172, M^+], TMS-CA [m/z 254, M^+], [$^2\text{H}_4$]CA [m/z 258, M^+],

TMS-HQ [m/z 254, M^+], TMS- $^{13}\text{C}_6$]HQ [m/z 260, M^+], TMS-MA [m/z 271, $(M-15)^+$], TMS- $^{13}\text{C}_2$]MA [m/z 273, $(M-15)^+$], TMS-SPMA [m/z 252, $(M-\text{NH}_2\text{COCH}_3)^+$], and TMS- $^2\text{H}_5$]SPMA [m/z 257, $M-\text{NH}_2\text{COCH}_3)^+$]. The respective retention times were 12.66, 12.66, 17.65, 17.62, 18.83, 18.83, 21.91, and 21.91 min for TMS-PH, TMS- $^{13}\text{C}_6$]PH, TMS-CA, TMS- $^2\text{H}_4$]CA, TMS-HQ, TMS- $^{13}\text{C}_6$]HQ, TMS-MA, and TMS- $^{13}\text{C}_2$]MA in split mode, and 32.65 and 32.59 for TMS-SPMA and TMS- $^2\text{H}_5$]SPMA in splitless mode. Quantification was based on peak areas relative to the corresponding isotopically labeled internal standards.

Creatinine Adjustment for Benzene Metabolites

It is customary to adjust urinary metabolite levels for urinary creatinine to control for urine volume at the time of collection. However, such adjustments can introduce biases and reduce precision, due to variation of creatinine levels with gender, age, body mass index, and other physiological factors [31-33]. In preliminary regression models, we observed lower adjusted- R^2 values using creatinine-adjusted metabolite levels compared to unadjusted values. We also found significant effects of age, gender and body mass index upon creatinine levels in our samples of workers. Thus, for all statistical analyses we used metabolite levels without adjustment for urinary creatinine. The following median levels of urinary creatinine were observed in our samples of workers: all subjects – 11.6 mmol/l; female controls – 10.4 mmol/l; male controls – 11.5 mmol/l; female exposed – 11.4 mmol/l; and male exposed – 13.9 mmol/l.

Predicting Low-level Exposures from Urinary Benzene

All air samples from control subjects were below the LOD ($n = 160$) and some measurements from exposed subjects were either below the LOD ($n = 70$) or were missing ($n = 23$). In order to predict the levels of benzene exposure in these samples, we used the log-scale linear regression of UBz on air benzene (228 data pairs) shown in Figure 2A. For subjects

having multiple air and urine samples, the mean values of the natural logarithms of urine and air levels were used. The following calibration curve was derived from the parameters of the simple linear regression: $\text{benzene (ppm)} = \exp\left(\frac{\ln(\text{UBZ, nmol/l}) - 5.42}{0.886}\right)$. Note that these predicted values of benzene exposure include contributions from smoking. Uncertainties in this relationship were evaluated via a bootstrapping technique as described below.

Investigating Exposure-Metabolite Relationships

The relationships between levels of metabolites and exposure were explored using median values and interquartile ranges. Based upon 13 groups of 30 subjects stratified by exposure the following median air concentrations (ppm) were observed: 0.001, 0.002, 0.004, 0.007, 0.027, 0.273, 0.508, 0.689, 1.02, 1.57, 2.41, 5.45, and 15.4. Multiple comparison tests were utilized to generate step-functions showing statistically significant increases in geometric mean (GM) metabolite levels from group to group. These step functions were used to define levels of benzene exposure at which metabolite concentrations were statistically distinguishable from background values.

A similar grouping scheme (30 subjects stratified by exposure level) was used to investigate the dose-related production of each benzene metabolite, after adjustment for background metabolite levels. Background levels were estimated as median metabolite concentrations in 60 subjects with the lowest benzene exposures (median predicted benzene concentration = 0.002 ppm). Adjustment involved subtracting background metabolite levels from the corresponding urine concentrations observed in each subject. After grouping by benzene exposure ($n = 30$ per group) the median metabolite level in each group was divided by the median air concentration of benzene to estimate dose-related production of each metabolite ($\mu\text{mol/l urine per ppm benzene}$) for the 9 groups with estimated median benzene exposures

between 0.027 and 15.4 ppm. Uncertainties in estimated median values of benzene exposure and background-adjusted metabolite levels were evaluated via a bootstrapping technique as described below.

Statistical Analyses

All statistical analyses were performed using SAS software for Windows v. 9.12 (SAS Institute, Cary, NC). Between one and four concurrent air and urine samples from each subject were available for statistical analysis (median = 2/subject). For subjects with multiple samples, estimated individual GM levels were used for all analyses. Non-detectable levels of urinary analytes were replaced by $LOD/\sqrt{2}$ [UBz: $n = 8$ (1.29%); SPMA: $n = 30$ (4.84%)] [34]. The precision of the assay for each urinary analyte was estimated as the coefficient of variation (CV) from 84 pairs of duplicate urine samples (from exposed and control workers), which had been aliquoted and assigned random identification numbers prior to shipment to the laboratory. The CV was estimated as $CV = \sqrt{\exp(s_e^2) - 1}$, where s_e^2 is the estimated error variance obtained from a one-way analysis of variance of the log-transformed levels of each analyte (Proc NESTED of SAS).

To evaluate possible effects of sample storage at -80°C upon the levels of UBz and the benzene metabolites, the logged level of each urinary analyte was regressed upon the corresponding logged air concentration of benzene and the time of storage using a mixed-effects model, which included a random-subject effect to account for multiple urine specimens being collected from some persons (Proc MIXED of SAS).

Median levels of benzene exposure and those of the urinary metabolites were compared by exposure status, gender, and smoking status using the Wilcoxon two-sample test (two-sided tests with a significance level of 0.05, Proc NPAR1WAY of SAS). Trends of (median)

metabolite levels versus benzene exposure were tested with Spearman correlation coefficients, (Proc CORR of SAS). Differences in mean values of logged metabolite levels among groups of 30 subjects aggregated by benzene exposure were tested using Tukey's multiple comparisons test (Proc GLM of SAS). These tests were used to determine levels of benzene exposure where metabolite levels could be differentiated from background values.

We predicted air exposures of control subjects from their levels of urinary benzene and used these predicted exposures to group subjects for aggregated analyses. Thus, to estimate dose-related profiles of metabolite production, and the associated uncertainties, bootstrapping was performed based upon median values of background-adjusted metabolite levels and the corresponding levels of benzene exposure. First, resampling was conducted (with replacement) 1000 times to predict nondetected air exposures from urinary benzene levels, using the 228 data pairs shown in Figure 2. For each of the 1000 samples, predicted air concentrations were combined with observed air concentrations, median background levels from the lowest 60 exposed persons were subtracted from individual metabolite levels, and data were aggregated for groups of 30 subjects. Then, 5th, 50th and 95th percentiles of the distributions of estimated median values of air and metabolite concentrations were obtained. Differences in dose-related production of each metabolite between adjacent exposure groups were evaluated with Kruskal-Wallis tests for overall group effects and trends (Proc FREQ of SAS).

RESULTS

Precision, Sensitivity and Storage Stability

Estimates of *CVs* for the urinary analytes were as follows: UBz – 21.1%, MA – 6.15%, SPMA – 46.2%, PH – 4.97%, CA – 6.50%, and HQ – 4.81%. Based on a volume of 0.5 ml urine and a signal-to-noise ratio of 3:1, the following LODs were estimated: UBz, 0.128 nmol/l, MA,

5.66 nmol/l (one μ l injected with 1:20 split), and SPMA, 0.836 nmol/l (2 μ l with splitless injection). Values of LOD were not estimated for PH, CA, and HQ due to the presence of background levels of these compounds in all urine specimens. The LOD for measurements of airborne benzene had previously been assigned a nominal value of 0.2 ppm [29]. When the period of storage was included as a fixed effect in mixed-effects models of the logged levels of UBz and the benzene metabolites on the logged benzene air concentrations, the observed regression coefficients were very small relative to the intercepts and showed no evidence of statistical significance ($p > 0.2$). This indicates that all analytes were stable during long periods of storage at -80°C in this study.

Summary Statistics

Estimates of median levels, ranges and interquartile ranges of air exposures to benzene and all urinary analytes are summarized in Table II, after stratification by exposure status, gender, and smoking status. Among occupationally exposed subjects, air levels of benzene ranged from 0.017 to 88.9 ppm (interquartile range: 0.529 – 3.34 ppm), with median values of about 1.2 ppm in all categories of subjects. Among control subjects, median exposures (predicted from UBz levels) were between 3 and 6 parts per billion (ppb) depending upon gender and smoking status. Although air and urinary concentrations of benzene were comparable in all categories of exposed workers, females had significantly higher median levels of SPMA ($p = 0.048$) and PH ($p = 0.038$) than males. No significant differences were observed between smokers and nonsmokers in the exposed group. Control subjects who were smokers had higher median levels of all urinary analytes than nonsmokers, notably UBz ($p = 0.003$), CA ($p = 0.034$) and HQ ($p = 0.001$), indicating the likely presence of these analytes in cigarette smoke. Although there were no significant effects of gender on metabolite levels in control subjects, when nonsmokers were

further stratified by gender, females ($n = 84$) had significantly higher median levels of PH (63.3 vs. 46.4 $\mu\text{mol/l}$, $p = 0.037$) and CA (13.2 vs. 7.99 $\mu\text{mol/l}$, $p = 0.010$) than males ($n = 16$).

Distinguishing Benzene Metabolites from Background Sources

Figure 3 shows median levels and interquartile ranges for concentrations of each metabolite among groups of 30 subjects aggregated by benzene exposure. As exposure increased, the levels of metabolites also increased (estimated Spearman coefficients of median values for PH: 0.973, HQ: 0.945, CA: 0.934, MA: 0.973, and SPMA: 0.973, all with $p < 0.001$). The step functions signify nominal exposure concentrations at which statistically significant increases in metabolite levels were observed. As exposure increased, the contribution of benzene to each metabolite could consistently be discerned at some point (Tukey's multiple comparisons test), as indicated by an arrow in Figure 3. Although there were marginal increases in median levels of MA, PH, CA, and HQ at median benzene exposures as low as 0.02 ppm, elevations were not unambiguous below air concentrations of 0.2 ppm for MA and SPMA, 0.5 ppm for PH and HQ and 2 ppm for CA. From these results, it appears that none of the metabolites would be useful for biomonitoring of benzene at environmental levels (generally below 0.01 ppm).

Dose-Related Production of Benzene Metabolites

Median urine concentrations of all metabolites per ppm of benzene exposure are presented in Figure 4 for the 9 groups of workers having exposures above 0.01 ppm (after adjustment for background values). Each point in the figure and the corresponding error bars represent median values and 5th and 95th percentiles of distributions of median values, as determined by bootstrapping. These curves point to the dose-related production of benzene metabolites according to the pathways shown in Figure 1. Over the range of exposures investigated (median benzene concentrations between 0.027 and 15.4 ppm) significant

downward trends in metabolite production were observed ($p < 0.001$, Kruskal-Wallis tests) for all major metabolites (PH, CA, HQ and MA) and for total metabolites ($p < 0.001$), while a significant upward trend was observed for SPMA ($p < 0.001$). The downward trend of total benzene metabolites is consistent with partial saturation of CYP metabolism of benzene between 0.027 and 15.4 ppm. The transitions were particularly apparent between the first and second exposure groups (median benzene concentrations between 0.027 and 0.274 ppm for all metabolites).

DISCUSSION

The connections between benzene metabolism and toxicity have been extensively debated [15,22,23,35,36]. To gain a better understanding of human benzene metabolism, we measured the key metabolites (PH, CA, HQ, MA and SPMA) and UBz in 250 benzene exposed workers and 139 control workers from a Chinese population. Of the studies devoted to urinary biomarkers of benzene in Asian populations, none compared levels of all key metabolites with air levels representing both environmental and occupational exposures [25,27,28,30,37,38]. Our goals were to determine the usefulness of the urinary metabolites as biomarkers of benzene exposure in exposed and control subjects and to elucidate the dose-related patterns of benzene metabolism along the prominent pathways shown in Figure 1.

Since control workers in our study were exposed to benzene in environmental air and tobacco smoke, we predicted their exposures from the corresponding UBz concentrations, using a calibration curve obtained from exposed subjects (Figure 2). Such predictions should be reasonably unbiased, given previous findings of strong linear trends between levels of benzene in urine and environmental air [39-42]. This conjecture is reinforced by Figure 2B which plots levels of UBz versus air benzene from the current study (open circles) as well as those reported

by Ghittori *et al.* [39] for 63 nonsmoking workers exposed to measured benzene concentrations between 0.01 and 4 ppm (closed circles). The two sets of measurements point to a linear log-scale relationship down to the lowest measured air level of about 0.01 ppm. Also, the median benzene air concentration predicted from UBz levels for our control subjects (3.55 ppb, $n = 139$) is similar to air concentrations estimated from personal monitoring of adult populations in the U.K. (3.82 ppb daytime mean value, $n = 50$) [43], in Germany (3.44 ppb median value, $n = 113$) [44], in the U.S. (2.29 – 7.01 ppb range of median values for 5 cities, $n = 421$) [45], and Italy (2.82 and 1.88 ppb for control subjects in two cities, $n = 107$) [46].

Comparing levels of urinary analytes between smoking and nonsmoking controls, the median values of UBz, CA, HQ, MA and SPMA were all higher in smokers, while the median level of PH was marginally lower in smokers (Table I). The apparent anomaly for PH is probably related to gender because male controls had lower PH levels than females and 92% of the smokers were male. [Among male controls, smokers had a higher median level of PH (58.7 $\mu\text{mol/l}$, $n = 36$) than male nonsmokers (46.4 $\mu\text{mol/l}$, $n = 16$)]. Other investigators have reported that non-occupationally exposed smokers had significantly higher urinary levels of UBz [39-42], MA [30,40,47-50], and SPMA [30,40,51] than nonsmokers.

To estimate the equivalent air level of benzene from cigarette consumption, we regressed the (logged) predicted air concentration of benzene on the number of cigarettes smoked per day among control subjects. This led to the following relationship:

$\ln(\text{benzene, ppm}) = -5.81 + 0.050(\text{cigarettes/d})$ ($n = 134$, p -value for slope = 0.003), from which we predict that smoking 20 cigarettes/d would be equivalent to an occupational benzene exposure of 8.2 ppb (26 $\mu\text{g}/\text{m}^3$). This estimated benzene exposure of 8.2 ppb is the same as that predicted from British data [52] for an urban smoker consuming 20 cigarettes/d.

In order to distinguish benzene-derived metabolites from background levels of the same compounds, we investigated median metabolite levels for groups of 30 subjects, aggregated by benzene exposure (Figure 3). Using statistical significance to discern exposure-related increases in urinary levels, we detected marginal elevations at benzene exposures between 0.02 and 0.1 ppm for all metabolites except SPMA. However, when we considered exposure levels which produced unambiguous increases in metabolite concentrations, MA and SPMA were the most sensitive biomarkers of exposure to benzene (about 0.2 ppm), followed by PH and HQ (about 0.5 ppm) and CA (about 2 ppm).

Using aggregated data, we also observed changes in dose-related metabolism of benzene (Figure 4) for median benzene exposures between 0.027 and 15.4 ppm. We anticipated that saturable metabolism would be detected in this range, given results from other Chinese studies [25-27,30], and because benzene, oxepin and PH are all thought to compete for the same enzymes (CYP2E1) [53-55]. And, in fact, downward trends were apparent for dose-related production of PH (9.9-fold), CA (26-fold), HQ (3.4-fold), MA (2.5-fold), and total metabolites (14-fold) (Figure 4). The 2.5-fold decrease observed for MA is comparable to a 3.3-fold reduction that we estimated from grouped data reported by Melikian *et al.* [27] (using differences between post-work and pre-work measurements of MA) for Chinese workers exposed to benzene concentrations between 0.34 and 22.6 ppm (estimated mean values).

Surprisingly, SPMA displayed an upward trend of production, with a 5.8-fold increase between 0.027 and 15.4 ppm (Figure 4). This might reflect the action of glutathione-S-transferase or be the result of mechanisms [16] or sites of formation of SPMA which differ fundamentally from those of the other metabolites.

It is noteworthy that the most dramatic reductions in dose-related metabolism occurred between the first two groups in Figure 4, with benzene exposures of 0.027 and 0.274 ppm, respectively. In this range, production of CA dropped by 16-fold and that of PH by 4.4-fold. Yet, only marginal reductions were observed in this exposure range for production of MA (44% reduction) and HQ (36% reduction). Thus, it appears that metabolism shifted away from CA and PH at low doses in favor of MA and HQ, the only major benzene metabolites requiring two CYP oxidations (Figure 1). When exposures exceeded about one ppm, the curves in Figure 4, representing major metabolites (PH, MA, HQ, and CA), became quasi-parallel, suggesting that metabolism of benzene to BO had become rate-limiting.

If our conjecture is correct that exposures to benzene below one ppm favor production of HQ and MA, there could be important implications for risk assessment. Certainly, HQ is the precursor of 1,4-benzoquinone, which is generally regarded as most hematotoxic of the benzene metabolites [18,22,24,36,56], and MA is derived from the extremely reactive and toxic muconaldehydes [14,15,17]. We are currently applying various nonlinear models to these data to more fully examine the dose-related metabolism of benzene and to estimate effects of physiological and genetic factors upon benzene metabolites.

Acknowledgements

This research was supported by the National Institute for Environmental Health Sciences through grants P42ES05948 and P30ES10126 (S. M. R.) and RO1ES06721, P42ES04705, and P30ES01896 (M.T.S.) and by intramural funds of the National Cancer Institute. M.T.S. has received consulting and expert testimony fees from law firms representing both plaintiffs and

defendants in cases involving exposure to benzene. G.L. has received funds from the American Petroleum Institute for consulting on benzene-related health research.

References

1. IARC (1987) *IARC monographs on the evaluation of carcinogenic risks to humans. Overall evaluations of carcinogenicity: An updating of IARC Monographs*. World Health Organization, International Agency for Research on Cancer, Lyon, France.
2. Wallace, L. (1996) Environmental exposure to benzene: an update. *Environ Health Perspect*, 104 Suppl 6, 1129-36.
3. IARC (1982) *IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans. Some Industrial Chemicals and Dyestuffs*. World Health Organization International Agency for Research on Cancer, Lyon, France.
4. Hayes, R.B., Yin, S.N., Dosemeci, M., Li, G.L., Wacholder, S., Travis, L.B., Li, C.Y., Rothman, N., Hoover, R.N. and Linet, M.S. (1997) Benzene and the dose-related incidence of hematologic neoplasms in China. Chinese Academy of Preventive Medicine--National Cancer Institute Benzene Study Group. *J Natl Cancer Inst*, 89, 1065-71.
5. Savitz, D.A. and Andrews, K.W. (1997) Review of epidemiologic evidence on benzene and lymphatic and hematopoietic cancers. *Am J Ind Med*, 31, 287-95.
6. Lan, Q., Zhang, L., Li, G., Vermeulen, R., Weinberg, R.S., Dosemeci, M., Rappaport, S.M., Shen, M., Alter, B.P., Wu, Y., Kopp, W., Waidyanatha, S., Rabkin, C., Guo, W., Chanock, S., Hayes, R.B., Linet, M., Kim, S., Yin, S., Rothman, N. and Smith, M.T. (2004) Hematotoxicity in workers exposed to low levels of benzene. *Science*, 306, 1774-6.
7. Snyder, R. (2000) Recent developments in the understanding of benzene toxicity and leukemogenesis. *Drug Chem Toxicol*, 23, 13-25.
8. Lindstrom, A.B., Yeowell-O'Connell, K., Waidyanatha, S., McDonald, T.A., Golding, B.T. and Rappaport, S.M. (1998) Formation of hemoglobin and albumin adducts of benzene oxide in mouse, rat, and human blood. *Chem Res Toxicol*, 11, 302-10.
9. Lindstrom, A.B., Yeowell-O'Connell, K., Waidyanatha, S., McDonald, T.A. and Rappaport, S.M. (1999) Investigation of benzene oxide in bone marrow and other tissues of F344 rats following metabolism of benzene in vitro and in vivo. *Chem Biol Interact*, 122, 41-58.
10. Rappaport, S.M., Waidyanatha, S., Qu, Q., Shore, R., Jin, X., Cohen, B., Chen, L.C., Melikian, A.A., Li, G., Yin, S., Yan, H., Xu, B., Mu, R., Li, Y., Zhang, X. and Li, K. (2002) Albumin adducts of benzene oxide and 1,4-benzoquinone as measures of human benzene metabolism. *Cancer Res*, 62, 1330-7.
11. Yeowell-O'Connell, K., Rothman, N., Waidyanatha, S., Smith, M.T., Hayes, R.B., Li, G., Bechtold, W.E., Dosemeci, M., Zhang, L., Yin, S. and Rappaport, S.M. (2001) Protein adducts of 1,4-benzoquinone and benzene oxide among smokers and nonsmokers exposed to benzene in China. *Cancer Epidemiol Biomarkers Prev*, 10, 831-8.
12. McDonald, T.A., Waidyanatha, S. and Rappaport, S.M. (1993) Production of benzoquinone adducts with hemoglobin and bone-marrow proteins following administration of [¹³C₆]benzene to rats. *Carcinogenesis*, 14, 1921-5.
13. Waidyanatha, S., Yeowell-O'Connell, K. and Rappaport, S.M. (1998) A new assay for albumin and hemoglobin adducts of 1,2- and 1,4-benzoquinones. *Chem Biol Interact*, 115, 117-39.

14. Amin, R.P. and Witz, G. (2001) DNA-protein crosslink and DNA strand break formation in HL-60 cells treated with trans,trans-muconaldehyde, hydroquinone and their mixtures. *Int J Toxicol*, 20, 69-80.
15. Golding, B.T. and Watson, W.P. (1999) Possible mechanisms of carcinogenesis after exposure to benzene. *IARC Sci Publ*, 75-88.
16. Henderson, A.P., Barnes, M.L., Bleasdale, C., Cameron, R., Clegg, W., Heath, S.L., Lindstrom, A.B., Rappaport, S.M., Waidyanatha, S., Watson, W.P. and Golding, B.T. (2005) Reactions of Benzene Oxide with Thiols Including Glutathione. *Chem Res Toxicol*, 18, 265-270.
17. Oshiro, Y., Balwierz, P.S. and Witz, G. (2001) Micronucleus formation in mouse bone marrow cells in vivo in response to trans, trans-muconaldehyde. *Toxicol Lett*, 121, 159-66.
18. Snyder, R. and Hedli, C.C. (1996) An overview of benzene metabolism. *Environ Health Perspect*, 104 Suppl 6, 1165-71.
19. Sabourin, P.J., Bechtold, W.E., Griffith, W.C., Birnbaum, L.S., Lucier, G. and Henderson, R.F. (1989) Effect of exposure concentration, exposure rate, and route of administration on metabolism of benzene by F344 rats and B6C3F1 mice. *Toxicol Appl Pharmacol*, 99, 421-44.
20. Seaton, M.J., Schlosser, P. and Medinsky, M.A. (1995) In vitro conjugation of benzene metabolites by human liver: potential influence of interindividual variability on benzene toxicity. *Carcinogenesis*, 16, 1519-27.
21. Kenyon, E.M., Seeley, M.E., Janszen, D. and Medinsky, M.A. (1995) Dose-, route-, and sex-dependent urinary excretion of phenol metabolites in B6C3F1 mice. *J Toxicol Environ Health*, 44, 219-33.
22. Smith, M.T. (1996) The mechanism of benzene-induced leukemia: a hypothesis and speculations on the causes of leukemia. *Environ Health Perspect*, 104 Suppl 6, 1219-25.
23. Snyder, R. (2000) Overview of the toxicology of benzene. *J Toxicol Environ Health A*, 61, 339-46.
24. Inayat-Hussain, S.H. and Ross, D. (2005) Intrinsic pathway of hydroquinone induced apoptosis occurs via both caspase-dependent and caspase-independent mechanisms. *Chem Res Toxicol*, 18, 420-7.
25. Rothman, N., Bechtold, W.E., Yin, S.N., Dosemeci, M., Li, G.L., Wang, Y.Z., Griffith, W.C., Smith, M.T. and Hayes, R.B. (1998) Urinary excretion of phenol, catechol, hydroquinone, and muconic acid by workers occupationally exposed to benzene. *Occup Environ Med*, 55, 705-11.
26. Waidyanatha, S., Rothman, N., Fustinoni, S., Smith, M.T., Hayes, R.B., Bechtold, W., Dosemeci, M., Guilan, L., Yin, S. and Rappaport, S.M. (2001) Urinary benzene as a biomarker of exposure among occupationally exposed and unexposed subjects. *Carcinogenesis*, 22, 279-86.
27. Melikian, A.A., Qu, Q., Shore, R., Li, G., Li, H., Jin, X., Cohen, B., Chen, L., Li, Y., Yin, S., Mu, R., Zhang, X. and Wang, Y. (2002) Personal exposure to different levels of benzene and its relationships to the urinary metabolites S-phenylmercapturic acid and trans,trans-muconic acid. *J Chromatogr B Analyt Technol Biomed Life Sci*, 778, 211-21.
28. Qu, Q., Melikian, A.A., Li, G., Shore, R., Chen, L., Cohen, B., Yin, S., Kagan, M.R., Li, H., Meng, M., Jin, X., Winnik, W., Li, Y., Mu, R. and Li, K. (2000) Validation of biomarkers in humans exposed to benzene: urine metabolites. *Am J Ind Med*, 37, 522-31.

29. Vermeulen, R., Li, G., Lan, Q., Dosemeci, M., Rappaport, S.M., Bohong, X., Smith, M.T., Zhang, L., Hayes, R.B., Linet, M., Mu, R., Wang, L., Xu, J., Yin, S. and Rothman, N. (2004) Detailed exposure assessment for a molecular epidemiology study of benzene in two shoe factories in China. *Ann Occup Hyg*, 48, 105-16.
30. Waidyanatha, S., Rothman, N., Li, G., Smith, M.T., Yin, S. and Rappaport, S.M. (2004) Rapid determination of six urinary benzene metabolites in occupationally exposed and unexposed subjects. *Anal Biochem*, 327, 184-99.
31. Zacur, H., Kaufman, S.C., Smith, B., Westhoff, C., Helbig, D., Lee, Y.J. and Gentile, G. (1997) Does creatinine adjustment of urinary pregnanediol glucuronide reduce or introduce measurement error? *Gynecol Endocrinol*, 11, 29-33.
32. Hall Moran, V., Leathard, H.L. and Coley, J. (2001) Urinary hormone levels during the natural menstrual cycle: the effect of age. *J Endocrinol*, 170, 157-64.
33. Ikeda, M., Ezaki, T., Tsukahara, T., Moriguchi, J., Furuki, K., Fukui, Y., Okamoto, S., Ukai, H. and Sakurai, H. (2003) Bias induced by the use of creatinine-corrected values in evaluation of beta2-microglobulin levels. *Toxicol Lett*, 145, 197-207.
34. Hornung, R.W. and Reed, L.D. (1990) Estimation of average concentration in the presence of nondetectable values. *Appl Occup Environ Hyg*, 5, 46-51.
35. Lovern, M.R., Cole, C.E. and Schlosser, P.M. (2001) A review of quantitative studies of benzene metabolism. *Crit Rev Toxicol*, 31, 285-311.
36. Snyder, R. (2004) Xenobiotic metabolism and the mechanism(s) of benzene toxicity. *Drug Metab Rev*, 36, 531-47.
37. Ong, C.N., Kok, P.W., Lee, B.L., Shi, C.Y., Ong, H.Y., Chia, K.S., Lee, C.S. and Luo, X.W. (1995) Evaluation of biomarkers for occupational exposure to benzene. *Occup Environ Med*, 52, 528-33.
38. Ong, C.N., Kok, P.W., Ong, H.Y., Shi, C.Y., Lee, B.L., Phoon, W.H. and Tan, K.T. (1996) Biomarkers of exposure to low concentrations of benzene: a field assessment. *Occup Environ Med*, 53, 328-33.
39. Ghittori, S., Fiorentino, M.L., Maestri, L., Cordioli, G. and Imbriani, M. (1993) Urinary excretion of unmetabolized benzene as an indicator of benzene exposure. *J Toxicol Environ Health*, 38, 233-43.
40. Ghittori, S., Maestri, L., Fiorentino, M.L. and Imbriani, M. (1995) Evaluation of occupational exposure to benzene by urinalysis. *Int Arch Occup Environ Health*, 67, 195-200.
41. Fustinoni, S., Giampiccolo, R., Pulvirenti, S., Buratti, M. and Colombi, A. (1999) Headspace solid-phase microextraction for the determination of benzene, toluene, ethylbenzene and xylenes in urine. *J Chromatogr B Biomed Sci Appl*, 723, 105-15.
42. Fustinoni, S., Buratti, M., Campo, L., Colombi, A., Consonni, D., Pesatori, A.C., Bonzini, M., Farmer, P., Garte, S., Valerio, F., Merlo, D.F. and Bertazzi, P.A. (2005) Urinary t,t-muconic acid, S-phenylmercapturic acid and benzene as biomarkers of low benzene exposure. *Chem Biol Interact*, 153-154, 253-6.
43. Leung, P.L. and Harrison, R.M. (1998) Evaluation of personal exposure to monoaromatic hydrocarbons. *Occup Environ Med*, 55, 249-57.
44. Hoffmann, K., Krause, C., Seifert, B. and Ullrich, D. (2000) The German Environmental Survey 1990/92 (GerES II): sources of personal exposure to volatile organic compounds. *J Expo Anal Environ Epidemiol*, 10, 115-25.

45. Rappaport, S.M. and Kupper, L.L. (2004) Variability of environmental exposures to volatile organic compounds. *J Expo Anal Environ Epidemiol*, 14, 92-107.
46. Fustinoni, S., Consonni, D., Campo, L., Buratti, M., Colombi, A., Pesatori, A.C., Bonzini, M., Bertazzi, P.A., Foa, V., Garte, S., Farmer, P.B., Levy, L.S., Pala, M., Valerio, F., Fontana, V., Desideri, A. and Merlo, D.F. (2005) Monitoring low benzene exposure: comparative evaluation of urinary biomarkers, influence of cigarette smoking, and genetic polymorphisms. *Cancer Epidemiol Biomarkers Prev*, 14, 2237-44.
47. Melikian, A.A., O'Connor, R., Prahalad, A.K., Hu, P., Li, H., Kagan, M. and Thompson, S. (1999) Determination of the urinary benzene metabolites S-phenylmercapturic acid and trans,trans-muconic acid by liquid chromatography-tandem mass spectrometry. *Carcinogenesis*, 20, 719-26.
48. Scherer, G., Renner, T. and Meger, M. (1998) Analysis and evaluation of trans,trans-muconic acid as a biomarker for benzene exposure. *J Chromatogr B Biomed Sci Appl*, 717, 179-99.
49. Melikian, A.A., Prahalad, A.K. and Secker-Walker, R.H. (1994) Comparison of the levels of the urinary benzene metabolite trans,trans-muconic acid in smokers and nonsmokers, and the effects of pregnancy. *Cancer Epidemiol Biomarkers Prev*, 3, 239-44.
50. Taniguchi, S., Niitsuya, M., Inoue, Y., Katagiri, H., Kadowaki, T. and Aizawa, Y. (1999) Evaluation of passive smoking by measuring urinary trans, trans-muconic acid and exhaled carbon monoxide levels. *Ind Health*, 37, 88-94.
51. Maestri, L., Ghittori, S. and Imbriani, M. (1997) Determination of specific mercapturic acids as an index of exposure to environmental benzene, toluene, and styrene. *Ind Health*, 35, 489-501.
52. Duarte-Davidson, R., Courage, C., Rushton, L. and Levy, L. (2001) Benzene in the environment: an assessment of the potential risks to the health of the population. *Occup Environ Med*, 58, 2-13.
53. Gilmour, S.K., Kalf, G.F. and Snyder, R. (1986) Comparison of the metabolism of benzene and its metabolite phenol in rat liver microsomes. *Adv Exp Med Biol*, 197, 223-35.
54. Snyder, R., Chepiga, T., Yang, C.S., Thomas, H., Platt, K. and Oesch, F. (1993) Benzene metabolism by reconstituted cytochromes P450 2B1 and 2E1 and its modulation by cytochrome b5, microsomal epoxide hydrolase, and glutathione transferases: evidence for an important role of microsomal epoxide hydrolase in the formation of hydroquinone. *Toxicol Appl Pharmacol*, 122, 172-81.
55. ATSDR (1997) Toxicological profile for benzene. In DHHS, U. (ed.), *PB/98/101157/AS*. Agency for Toxic Substances and Disease Registry.
56. Smith, M.T., Skibola, C.F., Allan, J.M. and Morgan, G.J. (2004) Causal models of leukaemia and lymphoma. *IARC Sci Publ*, 373-92.

Table I. Demographic characteristics of the study population.

Exposure Status	Gender	<i>n</i> (%)	Age median (range)	Current smokers <i>n</i> (%)	Smoking intensity^a median (range)
Control	Male	52 (37.4)	28 (18–51)	36 (69.2)	10 (1–40)
	Female	87 (62.5)	28 (18–51)	3 (3.45)	NR ^b
	All	139 (100)	28 (18–51)	39 (28.1)	10 (1–40)
Exposed	Male	86 (34.4)	23 (18–44)	47 (54.7)	10 (1–30)
	Female	164 (65.6)	33 (18–52)	5 (3.05)	4.5 (2–10)
	All	250 (100)	29 (18–52)	52 (20.8)	7 (1–30)

^a Average number of cigarettes/day.

^b Not reported.

Table II. Summary statistics for exposure to benzene and urinary analytes.

Variable	Control				Exposed			
	Female (<i>n</i> = 87)	Male (<i>n</i> = 52)	Nonsmokers (<i>n</i> = 100)	Smokers (<i>n</i> = 39)	Female (<i>n</i> = 164)	Male (<i>n</i> = 86)	Nonsmokers (<i>n</i> = 198)	Smokers (<i>n</i> = 52)
Air benzene [†]								
Median	3.40 ppb	3.71 ppb	3.09 ppb	6.07 ppb	1.28 ppm	1.05 ppm	1.18 ppm	1.18 ppm
Range	0.146 – 21.2 ppb	0.146– 553 ppb	0.146 – 21.2 ppb	0.146 – 553 ppb	0.017 – 88.9 ppm	0.122 – 50.2 ppm	0.017 – 88.9 ppm	0.122 – 40.1 ppm
IQ range	1.70 – 6.80 ppb	2.13 – 9.71 ppb	1.70 – 5.91 ppb	2.56 – 17.6 ppb	0.523 – 4.21 ppm	0.572– 2.36 ppm	0.520– 3.34 ppm	0.588 – 3.02 ppm
<i>p-value</i>	0.273		0.003		0.588		0.889	
UBz (nmol/l)								
Median	1.48	1.59	1.36	2.47	283	216	267	197
Range	0.091–7.47	0.091–130	0.091–7.47	0.091–130	6.21–53900	19.4–42600	6.21–53900	19.4–41600
IQ range	0.799–2.73	0.973–3.74	0.797–2.41	1.15–6.32	87.3–1020	96.3–643	96.3 – 810	85.3–654
<i>p-value</i>	0.273		0.003		0.635		0.702	
MA (μmol/l)								
Median	1.06	1.09	1.06	1.11	13.5	10.3	12.3	12.4
Range	0.152–6.17	0.132–5.78	0.152–6.17	0.132–4.96	0.644–426	1.50–370	0.644–426	2.15–347
IQ range	0.549–1.75	0.774–1.72	0.543–1.70	0.803–1.74	6.33–34.5	5.39–22.4	6.13–28.5	6.22–28.4
<i>p-value</i>	0.565		0.384		0.272		0.473	
SPMA (nmol/l)								
Median	1.94	3.24	1.96	3.17	262	137	228	171
Range	0.591–86.4	0.591–68.1	0.591–86.4	0.591–68.1	1.50–29400	3.68–33000	1.50–29400	4.06–33000
IQ range	1.17–3.62	1.04–6.93	1.13–4.25	1.09–7.37	50.6–788	48.9–353	50.1–631	47.9 – 412
<i>p-value</i>	0.479		0.493		0.048		0.638	
PH (μmol/l)								
Median	61.4	53.2	61.3	56.3	171	134	158	135
Range	9.48–208	7.58–336	9.48–208	7.58–336	15.9–3740	41.8–4140	15.9–4140	41.8–3380
IQ range	45.0–103	28.8–82.0	43.7–89.6	29.0–98.5	102–310	86.6–190	100–261	84.6–313
<i>p-value</i>	0.117		0.415		0.038		0.391	
CA (μmol/l)								
Median	13.3	12.7	12.1	15.7	21.8	21.0	20.5	23.7
Range	2.30–61.1	2.45–46.6	2.30–61.1	2.45–46.6	3.50–420	4.20–438	3.50–438	7.08–348
IQ range	8.06–17.0	7.07–18.9	7.50–15.8	9.90–19.8	13.3–33.0	12.6–29.4	12.6–32.1	13.8–35.5
<i>p-value</i>	0.786		0.034		0.411		0.153	
HQ (μmol/l)								
Median	5.95	7.35	5.93	8.52	17.7	18.5	17.1	19.6
Range	1.54–39.1	1.32–41.7	1.40–39.1	1.32–41.7	2.29–427	3.20–341	2.29–427	5.76–375
IQ range	4.44–8.88	4.48–11.7	4.11–8.86	5.65–13.9	11.2–36.8	11.5–27.8	10.4–29.1	14.9–35.9
<i>p-value</i>	0.106		0.001		0.925		0.084	

Legend: IQ range - interquartile range, benzene - benzene in air, UBz - urinary benzene, MA - *E,E*- muconic acid, SPMA - *S*-phenylmercapturic acid, PH - phenol, CA - catechol, HQ - hydroquinone.

[†] Exposure to benzene for control subjects was predicted from the relationship between UBz and air exposure.

Figure legends

Fig. 1. Simplified metabolic scheme for benzene showing major pathways.

Fig. 2. Calibration curve for predicting air concentrations of benzene from the corresponding urinary levels. (A) Points represent observed air and urine concentrations and the line represents the least-squares regression equation: $\ln(\text{urinary benzene, nmol/l}) = 5.42 + 0.886 \times \ln(\text{air benzene, ppm})$ ($n = 228$, adjusted $R^2 = 0.428$; 95 confidence of intercept, 5.24 – 5.61; 95 confidence interval of slope, 0.752 – 1.02). The curves represent 95% confidence intervals and 95% prediction intervals. (B) Scatter plot of urinary benzene versus air benzene from the current study (open circles) and data from Ghittori *et al.* [39] obtained from measurements from 63 nonsmoking subjects exposed to benzene (closed circles).

Figure 3. Levels of urinary metabolites versus air exposures to benzene. Each vertical box depicts median, 25th and 75th percentile values of a given metabolite at the median level of benzene exposure for data grouped by exposure levels (30 subjects per group). Horizontal lines represent groups with equivalent metabolite levels (Tukey's multiple comparison tests). Arrows represent air exposure levels where it is possible to distinguish benzene metabolites from background sources. Legend: MA - *E,E*-muconic acid, SPMA - *S*-phenylmercapturic acid, PH – phenol, CA - catechol, and HQ– hydroquinone.

Figure 4. Dose-related production of urinary metabolites versus air exposure to benzene. Each point and the corresponding error bars represent median values and 5th and 95th percentiles of distributions of estimated median values of air benzene and dose-related metabolite levels (background-adjusted metabolite concentration/air benzene concentration) for groups of 30 subjects, determined by bootstrapping (1000 iterations). Legend: MA - *E,E*-muconic acid, SPMA - *S*-phenylmercapturic acid, PH – phenol, CA - catechol, and HQ– hydroquinone Total – total metabolites.

Fig. 1.




Fig. 2.




Fig. 3.




Fig. 4.

