

PCFB
→

A REVIEW OF CHLORINATED BIPIHENYL CONTAMINATION IN NATURAL WATERS

GILMAN D. VLIETH and G. FRED LIE

Water Chemistry Laboratory
University of Wisconsin, Madison, Wis. 53706, U.S.A.

(Received 10 November 1969)

Abstract—A review of the present state of knowledge concerning environmental contamination by organochlorine compounds indicates that the chlorinated biphenyls may be one of the more widespread contaminants. The significance of this contamination has not yet been evaluated due, in part, to the lack of systematic analytical procedures for the quantitative and qualitative determinations of the components of chlorinated biphenyl mixtures. The possibility of the chlorinated biphenyls producing serious errors in chlorinated pesticide analyses is emphasized.

INTRODUCTION

CHEMICAL determinations of the types and concentrations of chlorinated pesticides in natural waters have become increasingly important to fish management, wildlife ecology, and water pollution control. Based on gas chromatographic (GLC) analyses of environmental samples, pp'DDE is generally considered to be one of the more widespread contaminants. However, implicit in every GLC analysis is the possibility that unresolved and/or unidentified components which have specific retention volumes similar to that of the desired substance may introduce serious errors in the interpretation of the chromatogram. Increasing numbers of investigators have reported the presence of organochlorine compounds in environmental samples which cannot be identified as the common chlorinated pesticides. Although the unidentified compounds have been attributed from presumptive evidence to the metabolites of common pesticides, plasticizers, or chlorinated biphenyls, proper chemical identifications have not been reported. Therefore, when studying natural water systems which receive industrial wastes, it may be more correct to consider unidentified compounds as chlorinated organic contaminants until confirmation techniques are developed.

Within the last three years, the chlorinated biphenyls (or polychlorinated biphenyls-PCB) have received considerable study. The preliminary evidence indicates that this group of organochlorine compounds may be widespread in the environment. For example, WIDMARK (1967) reported that mass-spectrum analyses of compounds eluting from GLC of fish and bird extracts indicated that unknown organochlorine compounds were present which behaved as chlorinated biphenyls of higher chlorine content. More recently, mass numbers and chlorine numbers per molecule for each peak in the chromatograph of commercial chlorinated biphenyl were reported (JENSEN *et al.*, 1968). However, to our knowledge, no spectral data or other confirmatory data for positive structural identification of these compounds have ever been published.

With this improper basis of identification, a number of other publications have followed which identify unknown compounds as chlorinated biphenyls. Since mixtures of chlorinated biphenyls may contain greater than 20 compounds which can only be

Cla

is of the
5 (CITCE).

Board:

idgo, England
1 BOCKRIS
ylvania, USA

ute of
Switzerland
RYSELBERGHE
USA
'ENSI

ing papers on
ach your,
imica Acta thus
ut the rapidly

imica Acta:

osses

electrodes

ont valency states

ectrodes in

tion copy



NEY

11/69

WATER 4/4-3

265

resolved by GLC with great difficulty, assignment of unidentified peaks to one of the chlorinated biphenyls has been accomplished by comparison of a chlorinated biphenyl with the nearest relative retention time. For example, HOLDEN and MARSDEN (1967), who had earlier reported the presence of unknown compounds in fish (HOLDEN, 1966), reported the presence of chlorinated biphenyls in seals with no supporting data other than relative retention data from GLC columns.

ROBINSON *et al.* (1967) reported compounds of unknown identity in marine organisms which gave GLC retention times longer than pp'DDE and microcoulometric response. The authors indicated that the chromatographic characteristics of the unknowns suggested they were not metabolites of the "cyclodiene-type" pesticides as has been proposed by ROHURN (1965).

HOLMIS *et al.* (1967) attributed unknown peaks from GLC to the chlorinated biphenyls and noted that the substances seemed to accumulate in organisms associated with natural waters (seals, fish and birds) to a greater extent than in domestic animals. This work reported relative retention data with no supporting data.

RISERBROUGH *et al.* (1968 a) and RISERBROUGH *et al.* (1968 b) reported organochlorine compounds in fish, birds and bird eggs which were attributed to the chlorinated biphenyls. With no supporting data and despite the absence of structural confirmations for the chlorinated biphenyls, the investigators reported concentrations of the unknowns (as chlorinated biphenyls) as well as ratios of DDT to chlorinated biphenyls. From these ratios, it was concluded that a chlorinated biphenyl fallout pattern existed around industrial centers and that air transport of the contaminants was significant.

KORMAN *et al.* (1969) presented GLC chromatograms of extracts from environmental samples as well as mass numbers and numbers of chlorine atoms per molecule for each peak of the chromatogram. Although no spectral data were presented, the mass number data strongly suggested that chlorinated biphenyls were present. This paper also presented the results of the analyses for each unknown as being "present" or "absent" in a particular sample, rather than attempting to quantitate the amount of material present.

THE CHLORINATED BIPHENYLS

Since contamination of the environment by chlorinated biphenyls has been implicated in the brief review presented, the need for the environmental scientist to become familiar with the present state of knowledge concerning these molecules is apparent. Mixtures of chlorinated biphenyls have been manufactured since 1929 and present registered trademarks for commercial brands include: Arochlor (Monsanto), Chlorxetol (Allis-Chalmers), Dykanol (Cornell-Dubilier Division, Federal Pacific Electric), Intercen (Westinghouse), Noflamol (Wagner Electric), Pyramol (G. E.), Therminol (Monsanto) in the U.S.; and Clophen (I. G. Farbenindustrie A. G., Germany), Fencolor (Caffaro, Italy), Kannechlor (Kanebuchi Chemical Co., Japan), Pyralene (Prodelec, France), and Sovat (Russia) (HUBBARD, 1964). The chlorinated biphenyls have been used as plasticizers for adhesives, polyvinyl acetates, acrylic resins, in sealing compounds, pipe joint compounds, printing ink, rubber manufacturing, high pressure-high temperature lubricants, fungicidal insulations, anti-corrosion, nail coatings, water proofings, diffusion pump fluid, stabilization of polymers,

dielectrics, preventing (MONSANTO *et al.*, 1955)

The toxic has not be indicate the "safe" is 1. highly chle with chlori reported re biphenyls effects of / cited) to 1 5460 to be

More re biphenyls t tion, and t factor of 1 dieldrin an

Howeve with organ possibility biological review of other met enzyme in great as py facts make cules have concentrat

Some e studies of biphenyl. analyses. have been make the of chemis

For exa step whiel residues (was treat It was cor nitration :

dielectrics, heat transfer media, coax-cable insulations, paints and varnishes, ice-preventing coatings, and is an evaporation retardant for pesticide applications (MONSANTO O/PL-306; HUBBARD, 1964; HOPKINS and HOFFMAN, 1955; SULLIVAN *et al.*, 1955).

The toxicity and sublethal effects of the chlorinated biphenyls to biological systems has not been thoroughly evaluated (REICHEL, 1969). Inhalation tests on animals indicate that maximum concentrations of chlorinated biphenyls as vapor which are "safe" is 1.0 mg/m³ for the lower chlorinated biphenyls and 0.5 mg/m³ for the more highly chlorinated biphenyls (HUBBARD, 1964). Continuous or repeated skin contact with chlorinated biphenyl leads to a condition called "chloracne" which has been reported repeatedly with industrial workers handling products containing chlorinated biphenyls (MUGS *et al.*, 1953). SULLIVAN and HORNSTEIN (1953) found no toxic effects of Arochlor 5460 on flies, but the chlorinated terphenyls were toxic (no levels cited) to the elm leaf beetle (DUDA, 1957). DEONIER *et al.* (1946) found Arochlor 5460 to be non-toxic at 0.1 ppm (24 hr) to the larvae of *Anopheles quadrimaculatus*.

More recently, LICHTENSTEIN *et al.* (1969) reported the toxicity of the chlorinated biphenyls to houseflies (*D. melanogaster*) decreased with increasing degree of chlorination, and that the chlorinated biphenyls appear to be less toxic than dieldrin by a factor of 1000. The chlorinated biphenyls were found to increase the toxic effect of dieldrin and DDT on houseflies.

However, in addition to investigating the sources of environmental contamination with organochlorine compounds, the environmental scientist is concerned with the possibility that these organochlorine molecules may exhibit detrimental effects on biological communities at sublethal levels. KUPFER (1967) has presented an extensive review of the effects of chlorinated pesticides on steroid hydroxylase induction and other metabolic systems. Also, RISENBROUGH *et al.* (1968 b) have demonstrated that enzyme induction by the chlorinated biphenyls (Arochlor 1262) was five times as great as pp'DDE or technical DDT as determined via oestradiol degradation. These facts make tolerance limits set by bioassay less significant since organochlorine molecules have the capacity to produce physiological effects in organisms at sublethal concentrations.

ANALYTICAL PROCEDURES

Some controversy has been generated over the analytical procedures utilized in studies of organochlorine contaminants which have been attributed to the chlorinated biphenyl. In general, the procedures are similar to typical chlorinated pesticide analyses. However, chemical reactions such as dehydrohalogenation and nitration have been introduced prior to electron capture gas chromatography in attempts to make the procedures more selective for the chlorinated biphenyls. It is the addition of chemical treatment steps which has caused disagreement in the literature.

For example, after extract clean up, JENSEN and WIDMARK (1967) used a nitration step which is similar to that used in removing DDT and metabolites from toxaphene residues (ERRO *et al.*, 1967; KAWANO *et al.*, 1969). Briefly, the concentrated extract was treated with a mixture of H₂SO₄ and HNO₃ (conc. 1:1, v/v) at 0°C for 5 min. It was concluded incorrectly that only PCB and BHC isomers are unaffected by the nitration step. RISENBROUGH *et al.* (1969) have reported the use of similar procedures.

To the contrary, REYNOLDS (1969) has been unable to repeat the nitration step reproducibly for a number of reasons. First, the reaction is not specific for the PCB and BHC isomers, since heptachlor epoxide remains unaffected by nitration. Also, the work of Bevenue (EARO, 1967; KAWANO, 1969) has demonstrated that toxaphene, the reaction product of the chlorination of camphene which produces in excess of 20 unresolved peaks on GLC, is apparently unaltered by nitration. Second, Reynolds found some loss of the earlier eluting chlorinated biphenyls during nitration. Third, and most significant, peaks with longer retention volumes appeared in the chromatograms after nitration. These peaks resulted from the nitration of other aromatic molecules in the mixture because nitro groups reduce the vapor pressures in the gas phase (GLC) and the aromatic nitro grouping has a high affinity for the secondary electrons in the electron capture detector.

Results of studies of this laboratory (VEITH and LEE, 1969) have substantiated earlier reports (DEGURSE, 1969) that fish and natural waters in several areas of Wisconsin contain unidentified organochlorine contaminants. Using essentially the same procedures discussed by REYNOLDS (1969), the resulting chromatograms contain many peaks which have retention volumes similar to those of components of several commercial chlorinated biphenyl mixtures. However, components with corresponding retention volumes were also observed in chromatograms of technical chlordane, heptachlor epoxide, benzene hexachloride, and toxaphene, as well as analytical standards of a number of the DDT family of pesticides. Accordingly, qualitative and quantitative interpretations of such chromatograms from Wisconsin fish and water await more detailed and systematic investigations of the extracts.

CONCLUSIONS

The preliminary evidence of the past several years, which has been reviewed in this communication, indicates that the environment associated with natural waters is becoming contaminated with unidentified organochlorine compounds from unknown sources. The facts that chlorinated biphenyls have been strongly implicated as contaminants, that the chlorinated biphenyls have been widely used in industry and have been prepared commercially since 1929, and that the mixtures produce GLC peaks with retention volumes similar to common chlorinated pesticides create a need for basic studies concerning analytical procedures for chlorinated hydrocarbons. Also, it is apparent that comprehensive and systematic investigations into the chemistry and the physiological effects of the contaminants as well as the degree and sources of the contamination are essential before control measures can be considered.

REFERENCES

- DEGURSE P. (1969) Personal communication from the Wisconsin Department of Natural Resources, Madison.
 DEONIER C. C., JONES H. A. and INCHEO H. H. (1946) Organic compounds effective against larvae of *Anopheles quadrimaculatus*. *J. econ. Ent.* 39, 459-462.
 DUDA J. (1957) The use of chlorinated polyphenyls to increase the effective insecticidal life of lindane. *J. econ. Ent.* 50, 218-219.
 EARO P., BEVENUE A. and BECKMAN H. (1967) A method for the determination of toxaphene in the presence of DDT. *Bull. Environ. Contam. Toxicol.* 2, 372-379.
 HOLDEN A. V. (1966) Organochlorine insecticide residues in Salmonid fish. *J. appl. Econ.* 3, 45-53.

HOLDEN A.
 216, 12
 HOLMES D.
 Nature
 HOPKINS T.
 again
 HUBBARD J.
 Chemi
 JENSEN S.
 Decem
 JENSEN S.
 Septer
 KAWANO Y.
 nitric
 Chem
 KORMAN J.
 fish, p
 211, 1
 KUPFER D.
 bolism
 LICHSTEIN
 betwe
 MILES J. V.
 J. Am
 MONSANTO
 RICHIE L.
 Wildt
 REYNOLDS
 analy
 RISEBROOK
 alism
 RISEBROOK
 chlor
 RISEBROOK
 poly
 ROBINSON
 chlor
 ROBINSON
 ions.
 and s
 SULLIVAN
 flies.
 SULLIVAN
 ould
 VEITH G.
 Rep
 WEDMARD

nitration step
for the PCB
ion. Also, the
xaphene, the
excess of 20
nd, Reynolds
ation. Third,
he chromat-
omatic mole-
the gas phase
ary electrons

stiated earlier
if Wisconsin
e same prob-
contain many
several com-
corresponding
l chlordane,
s analytical
sensitive and
h and water

reviewed in
al waters is
n unknown
ted as con-
y and have
GLC peaks
a need for
ons. Also,
mistry and
rces of the

- HOLDEN A. V. and MARSDEN K. (1967) Organochlorine pesticides in seals and porpoises. *Nature, Lond.* 216, 1274-1276.
- HOLMES D. C., SINKOVIS J. H. and TATTON J. O'G. (1967) Chlorinated hydrocarbons in British wildlife. *Nature, Lond.* 216, 227-229.
- HOPKINS T. L. and HOFFMAN R. A. (1955) Effectiveness of dilon and certain candidate synergists against DDT resistant house flies. *J. econ. Ent.* 48, 146-147.
- HURNARD H. L. (1964) Chlorinated biphenyl and related compounds. *Kirk-Othmer Encyclopedia of Chemical Technology*, 2nd edn, Vol. 5, 289-297.
- JENSEN S., NUCCI B. and WIDMARK G. (1968) *The OECD Study of Analysis of PCB*. Stockholm, December, mimeo.
- JENSEN S. and WIDMARK G. (1967) *Organochlorine Residues, OECD Preliminary Study*. Stockholm, September, mimeo.
- KAWANO Y., BEVINCE A., BUCKMAN H. and ERKO F. (1969) Studies on the effect of sulfuric-fuming nitric acid treatment on the analytical characteristics of toxaphene. *J. Ass. Official Analyt. Chem.* 52, 167-172.
- KOLMAN J. H., DE BRAUW M. C., TEN NOUVER and DE VOS R. H. (1969) Chlorinated biphenyls in fish, mussels and birds from the Rhine River and the Netherlands coastal area. *Nature, Lond.* 221, 1126-9.
- KUPFER D. (1967) Effects of some pesticides and related compounds on steroid function and metabolism. *Residue Rev.* 19, 11-30.
- LICHTENSTEIN F. P., SCHULZ K. R., FUHRMANN T. W. and LIANG T. T. (1969) Biological interaction between plasticizers and insecticides. *J. econ. Ent.* 62, 761-765.
- MILES J. W., ALBOM J. J. and KORTIN B. L. (1952) Chlordane from an unusual exposure to Arochlor. *J. Am. med. Ass.* 154, 1417-1418.
- MONSANTO COMPANY, *Technical Bulletin O/P1*-306, St. Louis, Missouri.
- RICHIE W. L. (1969) Chief, section of Chemistry. Personal communication from the Patuxent Wildlife Research Center, Laurel, Maryland.
- REYNOLDS L. M. (1969) Polychlorobiphenyls (PCB's) and their interference with pesticide residue analysis. *Bull. Env. Contam. Toxicol.* 4, 128-143.
- RISHBROUGH R. W., HUGGITT R. J., GRIFFIN J. J. and GOLDBERG E. D. (1968 a) Pesticides: Transatlantic movements in the Northeast Trades. *Science* 159, 1233-1234.
- RISHBROUGH R. W., RICHIE P., HERMAN S. G., IFARALL D. B. and KIRVIN M. N. (1968 b) Polychlorinated biphenyls in the global ecosystem. *Nature, Lond.* 220, 1098-1102.
- RISHBROUGH R. W., RICHIE P. and OLCOFF H. S. (1969) Current progress in the determination of the polychlorinated biphenyls. *Bull. Env. Contam. Toxicol.* 4, 192-201.
- ROBINSON J., REHARDSON A., CHASTRIE A. N., COULSON J. C. and POTTIS G. R. (1967) Organochlorine residues in marine organisms. *Nature, Lond.* 214, 1307-1311.
- ROBURN J. (1965) A simple concentration cell technique for determining small amounts of halide ions and its use in the determination of residues of organochlorine pesticides in extracts of animal and plant tissue. *Analyst* 90, 467-475.
- SULLIVAN W. N. and HORNSTEIN I. (1953) A comparison of evaporation rates and toxicity to house flies of lindane and lindane-chlorinated polyphenyl deposits. *J. econ. Ent.* 46, 882-894.
- SULLIVAN W. N., HORNSTEIN I. and YEMANS A. H. (1955) Improved deposits for controlling insects outdoors. *J. econ. Ent.* 48, 153-155.
- VEITH G. D. and LEE G. F. (1969) *Chlorinated Organic Contaminants in the Milwaukee River*. Progress Report to the Wisconsin Department of Natural Resources, Madison, October, mimeo.
- WIDMARK G. (1967) Pesticide Residues. *J. Ass. Official Analyt. Chem.* 50, 1069.

Resources,
nt larvae of
of lindane,
here in the
s, 45-53.