



INORGANIC FIBROUS MATERIALS AS SUBSTITUTES FOR ASBESTOS

MATERIAL : ATTAPULGITE

KNOWN BIOLOGICAL EFFECTS

- Membranolytic (1).
 - Cytotoxic to pulmonary macrophages (2).
 - Causes mesothelioma in experimental animals (3).
 - Adsorbs polycyclic aromatic hydrocarbons (4).
 - Lung fibrosis (1).
-

REFERENCES:

- 1- Bignon, J. et al. (1980) *in* Biological Effects of Mineral Fibres, Vol. 1:163-181, Ed. J.C. Wagner, IARC Pub. No. 30.**
- 2- AIRD: unpublished data.**
- 3- Pott, F. et al. (1976) Ann. Anat. Pathol. 21: 237-246.**
- 4- Harvey et al. (1984) Brit. J. Ind. Med. 41: 396-400.**

A01048

INORGANIC FIBROUS MATERIALS AS SUBSTITUTES FOR ASBESTOS

MATERIAL : REFRACTORY (CERAMIC) FIBERS

KNOWN BIOLOGICAL EFFECTS :

- Alumino-silicate (ceramic) fibers are both fibrogenic and cancerogenic, after administration in experimental animals by implantation and by inhalation (1) .
 - Intraperitoneal administration (25 mg) of man-made refractory fibers (mean diameter 1.8 μ) and of UICC crocidolite resulted in an incidence of 86% and 80% respectively of abdominal tumours (2) .
 - Under normal service conditions, alumino-silicate (ceramic) fibers can be transformed partly into free silica, cristobalite and mullite (3). These crystalline forms of silica have been reported to be even more fibrogenic than quartz (4).
-

REFERENCES :

- 1- Davis, J.M.G. et al. (1983) // Biological Effects of Man-Made Mineral Fibres, WHO/IARC, Euro Reports and Studies 81 : p. 124
- 2- Schultz, D.L. (April 22, 1985) Sohio Engineered Materials Co. Letter to EPA, Washington, Re : FYI-OTS-0184-0288
- 3- Vine, G. et al. (1985) Ann. Occup. Hyg.28 : 356-359
- 4- Stöber, W. (1968) Arch. Environ. Health 16 : 706-707,

A01049

INORGANIC FIBROUS MATERIALS AS SUBSTITUTES FOR ASBESTOS

MATERIAL : GLASS FIBERS

KNOWN BIOLOGICAL EFFECTS :

- Causes mesothelioma in experimental animals (1 , 2) .
 - Effects in glass-dusted animals parallel those observed in asbestos-dusted animals (3) .
 - Small diameter glass fibers are as cytotoxic in cell systems as comparable size asbestos fibers (4) .
 - Small diameter glass fibers induce metaplasia in organ cultures of trachea, a condition predisposing to malignant transformation by carcinogens found in tobacco smoke (5) .
 - Glass fibers of respirable size can have an effect on the (human) lung similar to that of asbestos (6) .
 - Studies in fibrous-glass production workers are consistent with a growing number of animal studies which demonstrate a carcinogenic response to glass fibers less than 1 μ in diameter (7) .
-

REFERENCES :

- 1- Pott F. et al. (1974) Environ. Health Perspect. 9 : 313-315.
- 2- Mohr, U. et al. (1984) Exper. Pathol. 26 : 179-183.
- 3- Miller, K. (1980) *in* Biological Effects of Mineral Fibers, Vol. 1 : 459-465, Ed. J.C. Wagner, IARC Pub. No. 30.
- 4- Tilkes, F. and Beck, E. G. (1983) Environ. Health, Perspect. 51 : 275-279
- 5- Woodworth, C. D. et al. (1983) Cancer Res. 43 : 4906-4912)
- 6- Chiappino, G. et al. (1981) Med. del Lavoro 72 : 96-101.
- 7- Bayliss, D.L. et al. (1976) U.S. Dept. of Commerce, NTIS, No. PB-257 784.

A01050

ORGANIC FIBROUS MATERIALS AS SUBSTITUTES FOR ASBESTOS

MATERIAL : ARAMID FIBERS

KNOWN BIOLOGICAL EFFECTS :

- Cytotoxic to pulmonary macrophages (1).
 - Exposure of test animals to respirable fraction has caused slight fibrosis, and a low incidence of lung tumors (2).
-

REFERENCES :

- 1- Dunnigan, J. et al. (1984) Toxicology Letters 20 : 277-282
- 2- Reinhardt, C.F. (1985) E.I. du Pont de Nemours & Co. Letter to EPA, Washington, Re : 8EHQ-0485-0550

A01051