

8076 NT598 P 2078
EMME
J. CANCER RES CLIN
Clinical Oncology

RECEIVED Springer-Verlag 1981

FEB 9 1982

J. T. BARR

Guest Editorial*

Malignant Tumors After Chronic Exposure to Vinyl Chloride

K. H. Emmerich¹ and K. Norpoth²

¹ Institute of Pathology (Director: Prof. E. Grundmann, MD, University of Münster, Domagkstr. 17, D-4400 Münster, Federal Republic of Germany

² Institute of Industrial Medicine and Silicosis Research

Summary. Correlations between exposure to vinyl chloride and the development of malignant tumors in the liver have been known since 1974 and have been confirmed by many an experimental investigation.

Based on the evaluation of mortality statistics from nine different countries an increased incidence of malignant tumors of the lung, the gastrointestinal tract, and the central nervous system (CNS), and of malignant lymphomas is documented in connection with exposure to vinyl chloride. Statistically significant increases, however, are only found in the incidence of malignant liver tumors. Metabolism and toxicology of vinyl chloride are discussed in detail.

Key words: Human carcinogenesis - Vinyl chloride - Toxicology - Liver tumors - Metabolism

Introduction

The industrial production of monomeric vinyl chloride (VCM) started some 30 years ago. In 1976, the USA produced 2.58, Western Europe 3.925, and Japan 1.281 million tons. About 95% of this output went into polymerization, the remaining 5% into the production of other chemicals, such as methylchloroform (IARC 1978).

• The "Journal of Cancer Research and Clinical Oncology" publishes in loose succession "Editorials" and "Guest Editorials" on current and/or controversial problems in experimental and clinical oncology. These contributions represent exclusively the personal opinion of the author. The Editors Die Zeitschrift "Journal of Cancer Research and Clinical Oncology" bringt in zwangloser Folge "Editorials" und "Guest Editorials" zu aktuellen und/oder kontroversen Problemen der experimentellen und klinischen Onkologie. Diese Beiträge geben ausschließlich die persönliche Meinung des Autors wieder.
Die Herausgeber

Offprint requests to: K. H. Emmerich, MD (address see above)

COPYRIGHT
SPRINGER VERLAG 81
NEW YORK NY

0171-5216/81/0102/0001/\$2.20

The very nature of the polymerization process allows for various ways of VCM release into the environment. Workers were exposed to particularly high concentrations of VCM during manual autoclave cleansing. Up to the 1960s the health risk of vinyl chloride was thought to be negligible (Lefaux 1966). Concentrations of 1,000 ppm at the individual working place were not uncommon in certain plants. First recommendations for limiting the exposure to 500 ppm were issued in 1955 in the United States. The average concentration levels of that period can be estimated approximately in retrospect:

1945-1955: $\pm$ 1,000 ppm
1955-1960: 400-500 ppm
1960-1970: 300-400 ppm
1973: $\pm$ 140 ppm
1975: $\pm$ 5 ppm (Barnes 1976).

In the Federal Republic of Germany the statutory MAK level¹ was fixed at 500 ppm until 1970, then reduced to 100 ppm until 1974. Today the technical limit concentration at the work place in the FRG is below 5 ppm for older and currently working plants, under 2 ppm for new plants under construction. The management soon realized that VCM concentration could be effectively reduced to the statutory levels by technical innovation and financial investments. One Swedish firm was able to reduce mean VCM concentrations from 12.2 ppm (2nd quarter of 1974) to 0.6 ppm (4th quarter of 1975) (Englund and Holmerg 1976).

Due to the physicochemical binding of VCM in PVC products, workers in PVC-processing or -manufacturing industries are equally exposed to a certain concentration of VCM, although at much lower levels than during the process of polymerization. In a 1974 survey of PVC-manufacturing plants levels were below 1 ppm in more than 60% of all firms checked.

The first reports about the potential risk of VCM had been published in 1949. Since 1957 reports had appeared here and there about a certain disease occurring in VCM industrial workers, manifested in scleroderma-like skin changes, Raynaud syndrome, and acro-osteolysis (Juehe and Lange 1972). Some years later, further symptoms of the disease were described as liver damage, splenomegaly, and thrombocytopenia (K. T. Müller et al. 1976). Detailed investigations were carried out on liver changes since the first cases of hemangiosarcoma had been found among VCM workers. These liver changes were graded histologically in five steps, correlating the length of exposure, and the severity of recorded changes (Gedigk et al. 1975; R. Müller et al. 1975). Even in VCM-exposed workers showing no clinical symptoms of liver damage, sequential scintigraphy would reveal RHS changes in liver and spleen (Biersack et al. 1975).

In addition to liver changes a considerable number of workers showed an impairment of pulmonary functions, and roentgenologic changes up to pulmonary fibrosis (Miller et al. 1975; Miller 1975; Lilis et al. 1976). These findings equally reflected a signification correlation between the length of exposure and the severity of changes.

The first reports about tumors induced by VCM were published in 1974, recording the increased incidence of a very rare neoplasm, liver hemangiosarcoma,

¹ MAK = maximale Arbeitsplatz-Konzentration (maximum concentration at work)




Fig. 1. Major metabolic pathways of vinyl chloride. The main urinary metabolites are underlined

among workers of a PVC-producing firm in the USA (Creech and Johnson 1974). More cases of this uncommon tumor, all among VCM workers, were published in the following years (Lloyd 1974, 1975; Lange et al. 1974, 1975; Byrén and Holmberg 1975). Based on 13 cases of liver hemangiosarcoma reported in the USA, the risk of VCM workers for this rare tumor was determined to be 400 times higher than that of the normal population (Heath et al. 1975). Histological studies of Thomas and Popper (1975) suggested the possibility of multicentric sarcoma development. Until 1978 the cases of liver hemangiosarcoma among VCM workers amounted worldwide to a total of 90 (Stafford 1980).

Sixteen cases of death of liver hemangiosarcoma had been recorded in the FRG until 1978 (Reinl et al. 1979). The cases subjected to detailed investigation until then revealed a longer latency period between first exposure and diagnosis, and a higher age at first diagnosis of hemangiosarcoma. A possible cause for this pattern was seen in the higher initial doses to which workers had been exposed during the first years of VCM production (Spirtas and Kaminski 1978). Environmental studies conducted in the immediate neighborhood of VCM-producing plants failed to assess an increased incidence of hemangiosarcoma in the general population of this area (Saric et al. 1976).

VCM Metabolism and Pharmacokinetics

The metabolism of monomeric vinyl chloride was studied by Green and Hathaway (1975), Watanabe et al. (1976), G. Müller et al. (1976), and many others. Figure 1 presents the major metabolic steps according to current concepts. In the mammalian organism monomeric vinyl chloride undergoes oxidative biotransforma-

tion involving the cytochrome P 450 system. Chloroethylenoxide, an alkylating carcinogen, is the resulting primary metabolite (Malaveille et al. 1975; Greim et al. 1975; Kappus et al. 1975) which is subsequently rearranged to chloroacetic aldehyde. Via several, partly unknown metabolic steps, this substance is transformed to S-carboxymethyl-cystein and thiodiacetic acid (Yllner 1971). The latter, a rather stable terminal metabolite, may be excreted via the urine. Quantitative demonstration of thiodiacetic acid is done by GC/MS technique after methylation (G. Müller et al. 1979). Another metabolic pathway, coupling chloroethylenoxide to glutathione, leads to β -hydroxyethyl mercapturic acid (Watanabe et al. 1976). The urinary level of this equally stable end product may be demonstrated by exchange chromatography after hydrolysis, or by the GC/MS test after derivation (G. Müller et al. 1980). It must be emphasized that β -hydroxyethyl mercapturic acid is the main urinary metabolite in the rat (Watanabe et al. 1976), but not in man (G. Müller et al. 1980).

The concept of VCM carcinogenicity being essentially mediated by its metabolite chloroethylenoxide has been substantially verified in recent years. In a mutagenicity test with *Salmonella typhimurium* TA 1535 the effect of this strongly reacting alkylating agent was 450 times stronger than that of the metabolite chloroacetaldehyde (Rannug et al. 1976). Tumor induction in rats with chloroethylenoxide was 100% successful, whereas chloroacetaldehyde failed to show any cancerogenic effect in a parallel assay (Zajdela et al. 1980). The role of intracellular chloroacetaldehyde, i.e., whether or not it might contribute essentially to VCM carcinogenesis (Guengerich et al. 1979), is still unclear.

VCM pharmacokinetics in rats and in man were extensively studied by Hefner et al. (1975), Whitey (1976), Bolt et al. (1976, 1977), Bolt (1978), Buchter et al. (1978), and Filser and Bolt (1979). According to them, man reacts to VCM levels around 2 ppm by establishing within a few minutes an equilibrium between environmental (air) and intracorporeal concentration. The constants of this equilibrium depend on the individual constitution, being approximately 0.8 in persons of normal weight. Overweight individuals will absorb a higher amount in proportion to their weight, and so the constant rises above 1.0, possibly due to the higher proportion of fatty tissue (Buchter et al. 1978). Experimental studies with rats have shown that the speed constant of VCM uptake and clearance after stopping the exposure ($k = 0.216 \text{ min}^{-1}$) is always much higher than the constant of metabolization ($k = 0.0037 \cdot \text{min}^{-1}$) and of urinary metabolite excretion ($k = 0.0032 \cdot \text{min}^{-1}$). Assuming human conditions to be similar, we may deduce that physiologic variations of ventilation levels will have either no or only minor influence on the VCM uptake by inhalation.

From data on the dose dependency of VCM biotransformation (G. Müller et al. 1976; Gehring et al. 1978, 1979) we may infer that reactive cancerogenic VCM metabolites will not increase in linear correlation with exposure, but rather follow the Michaelis-Menten function. Tumor incidence in rats (Gehring et al. 1978) and even in man (Gehring et al. 1979) is most adequately demonstrated in a probate model of frequency percentage correlated with exposure (Gehring et al. 1979). Functional correlation may be complicated if experimental exposure is under 50 ppm, by the glutathione-mediated detoxication of chloroethylenoxide, which is an effective protection mechanism at lower levels of exposure. Gehring et al. (1979) had even dis-

cussed the possibility of a veritable threshold dose of VCM, below which the tumor latency period would be longer than the normal life expectancy of the exposed person.

VCM Genotoxicity and Embryotoxicity

The mutagenicity of monomeric vinyl chloride on salmonella typhimurium strain in vitro was studied by Rannug et al. (1974), Bartsch et al. (1975), Greim et al. (1975), McCann et al. (1975), Andrews et al. (1976), and Garro et al. (1976). In a recent paper De Meester et al. (1980), though confirming the often cited mutagenic effect, reported that its action was considerably enhanced by the addition of an oxygenase-enriched preparation of mammalian liver tissue (S9 Mix). Hubermann et al. (1975) described the mutagenic effect of VCM on cultures mammalian cells. In studies of *Drosophila melanogaster* Magnusson and Ramel (1978) recorded an increase of lethal dominants under VCM exposure. Pretreatment with phenobarbital for 24 h would enhance the mutagenicity of VCM, probably via the induction of mixed-function oxidases which enhance the biotransformation of VCM to chloroethylenoxide.

In contrast to in vitro evidence of VCM mutagenicity, Short et al. (1977) and Peter and Ungvary (1980) were unable to register increased lethal dominants after VCM exposure in rats or mice. On the other hand, Basler and Röhrborn (1980) reported a time-dependent increase in sister-chromatid exchanges and structural chromosome aberrations in the bone marrow of Chinese hamsters after forced aspiration of high dose VCM. Fleig and Thies (1978) had already registered an increasing number of chromosomal aberrations in humans and animals with VCM disease, and an increased proportion of chromosomal aberrations was also found by Ducatman et al. (1975) among VCM-exposed workers in the USA, and in Sweden by Funes-Cravioto et al. (1975). Szentesi et al. (1976) recorded a rise of chromatid aberrations and instable chromosomal aberrations among VCM-exposed workers as compared to non-exposed controls. In his cytogenetic studies repeated at 30-month intervals, Hansteen et al. (1978) found a high proportion of chromosome fractures in lymphocyte cultures from workers exposed to high levels of VCM, which would return to lower values when doses were reduced. Purchase et al. (1978) in similar studies also registered a significant rise of chromosomal anomalies in VCM-exposed workers as compared to non-exposed control collectives. Heath et al. (1977) pointed out that cytogenetic damage might be due to other agents besides VCM. Finally, Infante et al. (1976) reported an increased rate of miscarriages among the wives of workers exposed to high VCM levels. According to them, an increasing rate of malformations was found even among the neonates of the general population of towns with VCM-processing factories.

Experimental Studies of VCM Carcinogenicity

The first experimental investigations of VCM-induced carcinogenesis were performed by Viola et al. (1971) who exposed Wistar rats to high doses of VCM (30,000 ppm) for 20 h each week over a whole year. Under this extreme exposure the animals developed tumors of skin, lungs, and bone. Subsequent animal studies

Table 1. Standardized mortality rates for tumor disease among VCM workers

Authors	Period	No.	SMR
Byrén et al. 1976	1958-1974	771	-
Chiazze et al. 1977	1964-1973	65,000	1.19 m 1.31 f
Duck et al. 1975	1955-1975	2,120	0.96
Fox and Collier 1977	1940-1974	7,717	0.75
Monson et al. 1974	1947-1973	(161 deaths)	1.50
Nicholson et al. 1975	1946-1974	255	3.90
Reinl et al. 1979	1946-1974	7,021	1.03
Tabershaw and Gaffey 1974	1946-1972	8,384	1.10
Waxweiler et al. 1976	1941-1973	1,294	1.49

revealed that the cancerogenic effect of low doses (50 ppm) would depend on the duration of the exposure. Extensive serial studies recorded VCM-induced tumors in all rodent species, such as rats, mice, and hamsters. Besides angiosarcoma mice were found to develop mammary carcinomas, pulmonary adenomas, and skin tumors. Apart from angiosarcomas rats would develop malignant neoplasms of the skin and zymbal gland, nephroblastomas, heptomas, angiomas, and neuroblastomas. In hamsters, VCM induced liver hemangiosarcomas, trichoepithelomas, and lymphomas (Maltoni and Lefemine 1975). Suzuki (1978) induced pulmonary tumors in mice by exposing them to VCM doses of 2,500 and 6,000 ppm for 5-6 months.

On the whole, these experimental results suggest that VCM exposure is likely to induce neoplastic changes also in other organs than the liver.

Epidemiological Data

Reports about angiosarcoma incidence among VCM workers, and also the results of experimental studies, have stimulated a survey of mortality statistics and proportional VCM mortality among former and present workers in VCM-processing factories. These studies attempted to determine the particular risk of VCM workers to acquire certain diseases, compared with the average normal population. Risks were assessed quantitatively by comparison with the standardized mortality (SMR), but the studies are based on a wide variety of collectives and observation periods (Table 1).

The only Swedish VCM factory registered an increased incidence of liver tumors (SMR = 4.5) and an accumulation of cardiovascular diseases among 58 workers dying between 1958 and 1974. The problem of a general increase in the overall SMR of tumors was not discussed by the authors of this report (Byrén et al. 1976).

An extensive study was conducted between 1965 and 1975 in some 65,000 people working in the American VCM production (Chiazze et al. 1977). Cancer mortality was generally increased (SMR in males = 1.19, females = 1.31). Men showed an increased incidence of gastrointestinal tumors (SMR = 1.34), liver tumors (SMR = 1.43), pulmonary tumors (SMR = 1.17) tumors of the CNS (= 1.15) and of the lymphatic system (= 1.30). Women showed a particular increase in the

Table 2. Standardized mortality rates of malignancies among VCM workers according to different organ sites

Authors	Gastroint.	Liver	Lung	CNS	Lymph.	Others
Byrén et al. 1976	-	4.13	1.68	6.12	-	-
Chiazze et al. 1977	M 1.34	1.43	1.17	1.15	1.30	1.09
	F 1.66	0.0	1.07	1.14	1.18	2.42
Duck et al. 1975	0.99	-	1.03	-	-	-
Fox and Collier 1977	0.91	3.22	0.89	0.54	0.99	-
Monson et al. 1974	-	11.00	1.60	4.20	1.50	-
Reini et al. 1979	1.30	15.23	0.96	1.62	2.14	-
Tabershaw and Gaffey 1974	0.94	-	1.12	-	1.06	1.89
Waxweiler et al. 1976	>1	11.55	1.56	3.29	1.59	1.55

incidence of urogenital tumors (SMR = 2.42); tumors of the gastrointestinal tract had a SMR of 1.66, that of pulmonary tumors was only slightly elevated (= 1.07) and so were the SMR of CNS (= 1.14) and lymphatic tumors (= 1.18).

An age-standardized survey of 2,100 workers of a VCM-processing factory in England failed to reveal an increased SMR for tumors, with the exception of cerebrovascular disease (SMR = 1.41) (Duck et al. 1975). Fox and Collier (1977) had been unable to detect an increasing SMR for tumors in general among 7,717 workers of PVC-producing industries in Great Britain, except for malignant liver disease with a distinctly elevated SMR of 3.22. Nevertheless, the general tumor SMR was relatively higher than that of cardiovascular disease. The possibility of cancerogenic action on organs other than the liver was refused by these authors.

Another study analyzed 161 deaths recorded among the workers of two PVC-producing factories in the USA between 1954 and 1973. SMR for tumors in general was 1.5 in this period; a striking increase was found in liver tumors (= 11.0) and in tumors of the CNS (= 4.2). A minor rise was seen in pulmonary tumors (= 1.6) and malignant lymphatic disease (= 1.5) (Monson et al. 1974). A smaller study of 255 VCM-exposed workers (Nicholson et al. 1975) recorded a generally increased mortality risk (= 1.27) and a marked increase in tumor SMR (= 3.9).

A recent epidemiologic survey of 7,021 VCM-exposed workers in the Federal Republic of Germany (Reini et al. 1979) revealed no significant overall increase in malignant tumors (SMR = 1.03), but the rates for individual organ tumors were markedly elevated: gastrointestinal tumor = 1.30, liver tumors = 15.23(1), CNS tumors = 1.62, tumors of the lymphatic system = 2.14. Tabershaw and Gaffey (1974) has assessed standardized mortality rates in 8,384 VCM-exposed workers employed in 33 different factories in the USA. The rate for tumors in general was only slightly elevated (1.10), but organ tumor rates were more distinct: 1.89 for tumors of the oral cavity and pharynx, 1.55 for cancer with unknown primary. There was also the case report of a 22-year-old man who, after 14 years of habitually chewing PVC-coated cables, developed a keratinizing squamous carcinoma of the buccal mucosa (Casterline et al. 1977).

A survey of 1,294 VCM-exposed workers from 1941 to 1973) Waxweiler et al. (1976) recorded an increase in the SMR of tumors in general (1.49) with a distinct

Table 3. Increased standard mortality rates for non-neoplastic disease in VCM workers

Authors	SMR	Disease
Byrén et al. 1976	1.52	Cardiovascular
Chiazze et al. 1977	1.05	Cardiovascular (males)
Duck et al. 1975	1.41	Cerebrovascular
Reini et al. 1979	1.27	Cardiovascular
Waxweiler et al. 1976	1.04	Cardiovascular
	1.76	Pulmonary

rise in liver tumors (11.55) and CNS tumors (3.29), and lesser elevations in pulmonary tumors (1.56), lymphatic malignancies (1.59); tumors of the gastrointestinal tract are registered with the rate "greater than one".

Increasing standardized mortality rates for other diseases are not considered in the same way by all these authors. Some of them did mention rising SM rates for certain diseases, but the latter are not confirmed by other authors. Table 3 compares the rates registered by different authors for nonneoplastic disease occurring in different organs.

Conclusions

Various surveys of standardized mortality rates in workers under chronic exposure to VCM have assessed the risk of developing fatal tumor disease to be distinctly higher in these workers than in the normal population, with the exception of several studies conducted in Great Britain. Besides a very striking elevation in the risk of liver tumors, nearly all these studies revealed an increasing incidence of gastrointestinal, pulmonary, CNS, and lymphatic malignancies. Although the SMR for malignant neoplasms in general was only slightly higher than 1, exposed workers were found to have an increased risk of dying of these tumors. This may be due to the high VCM concentrations registered in factories during the first years of VCM production and processing to PVC, i.e., at levels that were subsequently found to be positively cancerogenic in experimental studies.

Recent years brought a worldwide endeavour to reduce VCM concentrations in the factories to minimize the risk for the exposed workers. Although the levels have been effectively lowered, the potential latency period of 20 years for the induction of VCM tumors in man is a strong point in favor of keeping a close watch on occupational tumor development in the next decades. Ten or fifteen years are a period of risk especially in VCM or PVC workers, because the dangerously high concentrations had been recorded in some factories until 1974. The effectivity of such closely knit surveillance systems in the early detection of VCM tumors in the exposed group is hard to predict. It is equally undecided whether CEA tests (proposed by Anderson et al. 1978) or alkaline phosphatase tests (Lilis et al. 1975) might help to detect liver damage at an early stage. A practicable way may be found in coordinating biological monitoring of individual exposure, with the analytical monitoring of concentration limits in the occupational environment.

References

- Anderson HA, Snyder J, Lewinson T, Woo C, Liliis R, Selikoff IJ (1978) Levels of CEA among vinyl chloride and polyvinyl chloride workers. *Cancer* 42 [Suppl 3]:1560-1567
- Andrews AW, Zanistowski ES, Valentine CR (1976) A comparison of the mutagenic properties of vinyl chloride and methyl chloride. *Mutat Res* 40:273-276
- Barnes AW (1976) Vinyl chloride and the production of PVC. *Proc R Soc Med* 69:277-281
- Bartsch H, Malaveille C, Montesano R (1975) Human, rat and mouse liver-mediated mutagenicity of vinyl chloride in *S. typhimurium* strains. *Int J Cancer* 15:429-437
- Basler A, Röhrborn G (1980) Die Induktion von strukturellen Chromosomenaberrationen und Schwester-Chromatid-Austauschen (SCE) im Knochenmark des Chinesischen Hamsters nach Begasung mit Vinylchlorid. Statusseminar der Gesellschaft für Strahlen- und Umweltforschung mbH, München-Neuherberg, 8.-9. 11. 1979, München
- Biersack HJ, Lange CE, Ebinger H, Marsteller HJ, Leibach WK, Veltman G, Winkler C (1975) Sequenzintigraphische Untersuchungen von Leber und Milz bei Patienten mit Vinylchloridkrankheit. *Dtsch Med Wochenschr* 100:615-617
- Bolt HM (1978) Pharmacokinetics of vinyl chloride. *Gen Pharmacol* 9:91-95
- Bolt HM, Kappus H, Buchter A, Bolt W (1976) Disposition of (1,2-¹⁴C)-vinyl chloride in the rat. *Arch Toxicol* 35:153-162
- Bolt HM, Laib RJ, Kappus H, Buchter A (1977) Pharmacokinetics of vinyl chloride in the rat. *Toxicology* 7:179-188
- Buchter A, Bolt HM, Filser J, Georgens HW, Laib RJ, Bolt W (1978) Pharmakokinetik und Karzinogenese von Vinylchlorid. *Verh Dtsch Ges Arbeitsmed* 18:11-124
- Byrén D, Holmberg B (1975) Two possible cases of angiosarcoma of the liver in a group of Swedish vinyl chloride-polyvinyl chloride workers. *Ann N Y Acad Sci* 246:249-250
- Byrén D, Engholm G, Englund A, Westerholm P (1976) Mortality and cancer morbidity in a group of Swedish VCM and PVC production workers. *Environ Health Perspect* 17:167-170
- Casterline CL, Casterline PF, Javques DA (1977) Squamous cell carcinoma of the buccal mucosa associated with chronic oral polyvinyl chloride exposure: Report of a case. *Cancer* 39:1686-1688
- Chiazze L Jr, Nichols WE, Wong O (1977) Mortality among employees of PVC fabricators. *J Occupation Med* 19:623-628
- Creech JLIR, Johnson MN (1974) Angiosarcoma of the liver in the manufacture of polyvinyl chloride. *J Occup Med* 16:150-151
- De Meester C, Duverger-van Bogaert M, Lambotte-van de Paer M, Roberfroid M, Poncelet F, Mercier M (1980) Mutagenicity of vinyl chloride in the Ames test. Possible artifacts related to experimental conditions. *Mutat Res* 77:175-179
- Ducatman A, Hirschhorn K, Selikoff IJ (1975) Vinyl chloride exposure and human chromosome aberrations. *Mutat Res* 31:163-168
- Duck BW, Carter JT, Coombes EJ (1975) Mortality studies of workers in a polyvinyl production plant. *Lancet* II:1197-1199
- Englund A, Holmberg B (1976) Recent achievements and research initiated in the Swedish plastics and rubber industries. *Environ Health Perspect* 17:237-239
- Filser JG, Bolt HM (1979) Pharmacokinetics of halogenated ethylenes in rats. *Arch Toxicol* 42:123-136
- Fleig I, Thies AM (1978) External chromosome studies on persons and animals with VC illness; (Abstract). *Mutat Res* 53:187
- Fox AJ, Collier PF (1977) Mortality experience of workers exposed to vinyl chloride monomer in the manufacture of polyvinyl chloride in Great Britain. *Br J Ind Med* 34:1-10
- Funes-Cravioto F, Lambert B, Lindstein J, Ehrenberg L, Natarjan AT, Osterman-Golkar S (1975) Chromosome aberrations in workers exposed to vinyl chloride. *Lancet* I:459
- Garro AJ, Guttenplan JB, Milvy P (1976) Vinyl chloride-dependent mutagenesis: effects of liver extracts and free radicals. *Mutat Res* 38:81-88
- Gedigk P, Müller R, Bechtelsheimer H (1975) Morphology of liver damage among polyvinyl chloride workers. A report on 51 cases. *Ann N Y Acad Sci* 246:278-285
- Gehring PJ, Watanabe PG, Park CN (1978) Resolution of dose-response toxicity data for chemicals requiring metabolic activation: Example: Vinyl chloride. *Toxicol Appl Pharmacol* 44:581-591
- Gehring PJ, Watanabe PG, Park CN (1979) Risk of angiosarcoma in workers exposed to vinyl chloride as predicted from studies in rats. *Toxicol Appl Pharmacol* 49:15-21

- Green T, Hathaway DE (1975) The biological fate in rats of vinyl chloride in relation to its oncogenicity. *Chem Biol Interact* 11:545-562
- Greim H, Bonse G, Radwan Z, Reichert D, Henschler D (1975) Mutagenicity in vitro and potential carcinogenicity of chlorinated ethylenes as a function of metabolic oxirane formation. *Biochem Pharmacol* 24:2013-2017
- Guengerich FP, Crawford WM, Watanabe PG (1979) Activation of vinyl chloride to covalently bound metabolites: Role of 2-chloroethylene oxide and 2-chloroacetaldehyde. *Biochemistry* 18:5177-5182
- Hansteen JL, Hillestad L, This-Evensen E, Heldaas SS (1978) Effects of vinyl chloride in man. A cytogenetic follow-up study. *Mutat Res* 51:271-278
- Heath CW, Falk H, Creech JL (1975) Characteristics of cases of angiosarcoma of the liver among vinyl chloride workers in the US. *Ann N Y Acad Sci* 246:231-236
- Heath CW, Dumont CR, Gamble J, Waxweiler RJ (1977) Chromosomal damage in men occupationally exposed to vinyl chloride monomer and other chemicals. *Environ Res* 14:58-72
- Hefner RE, Watanabe PG, Gehring PJ (1975) Preliminary studies of the fate of inhaled vinyl chloride monomer in rats. *Ann N Y Acad Sci* 246:135-148
- Hubermann E, Bartsch H, Sachs L (1975) Mutation induction in Chinese hamster V 79 cells by two vinyl chloride metabolites, chloroethylene oxide and 2-chloroacetaldehyde. *Int J Cancer* 16:639-644
- IARC (1978) IARC working group on the evaluation of the carcinogenic risk of chemicals to humans. IARC, Vol 19, Lyon
- Infante PF, Wagoner JK, McMichael AJ, Waxweiler RJ, Falk H (1976) Genetic risks of vinyl chloride. *Lancet* i:734-735
- Juehe S, Lange CE (1972) Sklerodermieartige Hautveränderungen, Raynaud-Syndrom und Acroosteolysen bei Arbeiten der PVC-herstellenden Industrie. *Dtsch Med Wochenschr* 97:1922-1923
- Kappus H, Bolt HM, Buchter A, Bolt W (1975) Rat liver microsomes catalyze covalent binding of ¹⁴C-vinyl chloride to macromolecules. *Nature* 257:134-135
- Lange CE, Juehe S, Stein G, Veitman G (1975) Further results in polyvinyl chloride production workers. *Ann N Y Acad Sci* 246:18-21
- Lange CE, Juehe S, Veitman G (1974) Über das Auftreten von Angiosarkomen der Leber bei zwei Arbeitern der PVC-herstellenden Industrie. *Dtsch Med Wochenschr* 99:1598-1599
- Lefaux R (1966) *Chimie und Toxikologie der Kunststoffe*. Krausskopf-Verlag Mainz
- Lilis R, Anderson H, Nicholson WJ, Daum S, Fischbein AS, Selikoff IJ (1975) Prevalence of diseases among vinyl chloride and polyvinyl chloride workers. *Ann N Y Acad Sci* 246:22-41
- Lilis R, Anderson H, Miller A, Selikoff IJ (1976) Pulmonary changes among chloride polymerization workers. *Chest* 68 [Suppl 2]:299-303
- Lloyd JW (1974) Angiosarcoma of the liver in vinyl chloride/polyvinyl chloride workers. *J Occup Med* 16:809
- Lloyd JW (1975) Angiosarcoma of the liver in vinyl chloride/polyvinyl chloride workers. *J Occup Med* 17:333-334
- Magnusson J, Ramel C (1978) Mutagenic effect of vinyl chloride on *Drosophila melanogaster* with and without pretreatment with sodium phenobarbiturate. *Mutat Res* 57:307-312
- Malaveille C, Bartsch H, Camus AM, Montesano R (1975) Mutagenicity of vinyl chloride, chloroethylene oxide, chloroacetaldehyde, and chloroethanol. *Biochem Biophys Res Commun* 63:363-370
- Maltoni C, Lefemine G (1975) Carcinogenicity bioassays of vinyl chloride: Current results. *Ann N Y Acad Sci* 246:195-219
- McCann J, Simmon V, Streitwieser D, Ames BN (1975) Mutagenicity of chloroacetaldehyde, a possible product of 1,2-dichloroethane (ethylene dichloride), vinyl chloride, and cyclophosphamide. *Proc Natl Acad Sci USA* 72:3190-3193
- Miller A (1975) Pulmonary function defects in non-smoking vinyl chloride workers. *Environ Health Perspect* 11:247-250
- Miller A, Teirstein AS, Chuang M, Selikoff IJ, Warshaw R (1975) Changes in pulmonary function in workers exposed to vinyl chloride and polyvinyl chloride. *Ann N Y Acad Sci* 246:42-53
- Monson RR, Peters JM, Johnson MN (1974) Proportional mortality among vinyl chloride workers. *Lancet* ii:397-398
- Müller G, Norpoth K, Eckard R (1976) Identification of two urine metabolites of vinyl chloride by GC-MS investigations. *Int Arch Occup Environ Health* 38:69-75
- Müller G, Norpoth K, Wickramasinghe RH (1979) An analytical method using GC-MS for the quantitative determination of urinary thiodiglycolic acid. *Int Arch Occup Environ Health* 44:185-191

- Müller G, Heger M, Norpoth K (1980) Bestimmung der Hydroxyäthyl-Mercaptursäure im Harn Vinylchlorid-Exponierter. Methodische Erfahrungen und analytische Ergebnisse. *Verh Dtsch Ges Arbeitsmed* 20:533-536
- Müller KT, Buchter A, Gross R, Bolt W (1976) Ergebnisse einer Studie an 17 Fällen von Langzeitexposition gegenüber Vinylchlorid. *Med Welt* 27:21-24
- Müller R, Bechtelsheimer H, Gedick P, Marsteller HJ, Leibach WK (1975) Das morphologische Bild der Leberschädigung nach chronischer Vinylchlorid-Exposition. *Leber Magen Darm* 5:204-208
- Nicholson WJ, Hammond EC, Seidman H, Selikoff IJ (1975) Mortality experience of a cohort of vinyl chloride-polyvinyl chloride workers. *Ann N Y Acad Sci* 246:225-230
- Peter S, Ungvary G (1980) Lack of mutagenic effect of vinyl chloride monomer in the mammalian spot test. *Mutat Res* 77:193-196
- Purchase IFH, Richardson CR, Anderson D, Paddle GM, Adams GF (1978) Chromosomal analyses in vinyl chloride-exposed workers. *Mutat Res* 57:325-334
- Rannug U, Johansson A, Ramel C, Wachtmeister CA (1974) The mutagenicity of vinyl chloride after metabolic activation. *Ambio* 194-197
- Rannug U, Göthe R, Wachtmeister CA (1976) The mutagenicity of chloroethylene oxide, chloroacetaldehyde, 2-chloroethanol, and chloroacetic acid, conceivable metabolites of the vinyl chloride. *Chem Biol Interact* 12:251-263
- Reini W, Weber H, Greiser E (1979) Erkrankungen durch Vinylchlorid. Jahresbericht der Gewerbeaufsicht des Landes Nordrhein-Westfalen für das Jahr 1978, S 347-377
- Saric M, Kulcar Z, Zorica M, Gelic I (1976) Malignant tumors of the liver and the lung in an area with a PVC industry. *Environ Health Perspect* 17:189-192
- Short RD, Minor JL, Winston JM, Lee CC (1977) A dominant lethal study in male rats after repeated exposures to vinyl chloride or vinylidene chloride. *J Toxicol Environ Health* 3:965-968
- Spiras R, Kaminski R (1978) Angiosarcoma of the liver in vinyl chloride/polyvinyl chloride workers. 1977 update of the NIOSH register. *J Occup Med* 20:427-429
- Stafford J (1980) Liver angiosarcoma cases (Review data, 8th Jan. 1980). Imperial Chemistries Industries Ltd Plastics Division
- Suzuki Y (1978) Pulmonary tumors induced in mice by vinyl chloride monomer. *Environ Res* 16:285-301
- Szentesi I, Hornyak E, Ungvary G, Czeizel A, Bognar Z, Timar M (1976) High rate of chromosomal aberration in PVC workers. *Mutat Res* 37:313-316
- Tabershaw JR, Gaffey WR (1974) Mortality studies of workers in the manufacture of vinyl chloride and its polymers. *J Occup Med* 18:509-518
- Thomas LB, Popper H (1975) Pathology of angiosarcoma of the liver among vinyl chloride-polyvinyl chloride workers. *Ann N Y Acad Sci* 246:268-277
- Viola PL, Bigotti A, Caputo A (1971) Oncogenic response of rat skin, lungs, and bones to vinyl chloride. *Cancer Res* 31:516-522
- Watanabe PG, McGowan GR, Gehring PJ (1976) Fate of (¹⁴C)-vinyl chloride after single oral administration in rats. *Toxicol Appl Pharmacol* 36:339-352
- Waxweiler RJ, Stringer W, Wagoner JK, Jones J, Falk H, Carter C (1976) Neoplastic risk among workers exposed to vinyl chloride. *Ann N Y Acad Sci* 271:40-48
- Withey JR (1976) Pharmacodynamics and uptake of vinyl chloride monomer administered by various routes to rats. *J Toxicol Environ Health* 1:381-394
- Yllner S (1971) Metabolism of chloroacetate-1-¹⁴C in the mouse. *Acta pharmacol* 30:69-80
- Zajdela F, Croisy A, Barbin A, Malaveille C, Tomatis L, Bartsch H (1980) Carcinogenicity of chloroethylene oxide, an ultimate reactive metabolite of vinyl chloride, and bis-chloromethyl ether after subcutaneous administration, and in initiation-promotion experiments in mice. *Cancer Res* 40:352-356

Received April 15, 1981/Accepted July 17, 1981

