

Environmental Health Perspectives
Vol. 9, pp. 229-237, 1974

NOTICE
This material may be
protected by copyright
law Title 17 U.S. Code

SA-442

229

and increased gastrointestinal cancer risk; (c) particles accumulate in tissues for the lifetime of the individual; (d) particles may, in some way, interact with other substances in the environment to produce a synergistic, multiplicative effect; (e) there may exist a zero threshold for particle tolerance in humans, insofar as cancer is concerned.

Gastrointestinal Cancer in the United States

It has been estimated that upwards of 335,000 men, women, and children died of cancer, of all sites, in the United States in 1971 (4). Cancer is the second leading cause of death in this country, second only to heart diseases. The leading cause of neoplastic death in men is lung cancer; in females, it is cancer of the breast. Although the second leading cause of neoplastic death in women was uterine cancer in 1971, colon and rectum cancers were almost equal in number. In both men and women, over 46,000 deaths from cancer of the rectum and colon were recorded in 1971 (Table 1). The rate of rectum-colon cancer is increasing, so that it has been estimated that before the end of the decade this disease may be the second most common cause of cancer deaths in the United States for both men and women.

Table 1. Deaths from Cancer in the United States: 1971.

	No. of deaths
By sex	
Males	183,000
Females	152,000
Total	335,000 (best estimate)
By site	
Males, lung	53,100
Males, colon-rectum	22,400
Females, breast	30,500
Females, uterus (genital)	25,929
Females, colon-rectum	23,600

* 70-80% of malignant tumors of the large bowel occur in the cecum, sigmoid and rectal regions.

Hyperplastic and neoplastic lesions have been found, on scrutiny by light microscopy of random autopsy material, to be far more numerous and widely disseminated in the bowel, colon and rectum than has been previously reported

(5). However, considering the physical extent and mass of this organ, most malignant lesions tend to be confined to the lower bowel (sigmoid) and the rectum. A question is then whether increased gastrointestinal cancer in the United States is in any way related to the observed increased silicate contamination of foods, beverages, water supplies, pharmaceuticals, or any other substance which is directly ingested by humans.

Association between Mineral Particles and Neoplastic Diseases

A number of studies over the last 25 years has established an association between occupational exposure to some specific species of inorganic particles and increased risk of neoplastic disease in workers. Workmen exposed to asbestos dust (at least five separate mineral species and their chemical ranges) (6-12), hematite miners (hematite dust in the presence of free silica and other gangue silicates, including perhaps asbestos) (13), talc miners and millers (9), miners and millers of uranium ores (14), workmen who process chromium ores (15), and heavy metal processors who are exposed to lead, zinc or copper fumes (16) all have substantially increased risk of developing malignant neoplasms. Most of the reported tumor excess tend to be in the thoracic organs. However, in asbestos exposure, there are other tumors as well (Table 2). Exposure to

Table 2. Epidemiological studies demonstrating an association between exposure to mineral particles and increased risk of gastrointestinal cancer.*

Study	No. men	Material used
Keul (8)	42	Amosite; chrysotile; crocidolite
Selikoff et al. (7)	632	Chrysotile and insulation
Kleinfeld et al. (9)	91	Talc; tremolite; anthophyllite
Enterline and Kendrick (6)	1,843	Chrysotile (textiles)
Elmes and Simpson (12)	170	Mixed fiber; insulation
Newhouse et al. (11)	928	Crocidolite
Selikoff et al. (12)	220	Amosite

* Most of the studies deal with one or more forms of asbestos mineral fiber. At least five of the major fiber types are implicated as being biologically active.

asbest
ing ne
tract.
little a
3).

Tab

Fiber

Amosite
Chrysotile

* Two o
Approx
occurs i

* Data

* Data

Fiber /
Questi

Obs
tissue:
cupati
porta;
asbest
itself,
increa
cupat
amou:
magn
This
pulmo
the d:
such:
one n
that
them:
creas
to th
nitro:
in th
Stale
fact:
sugg:
tabli:
smol
fiber
disea
are:

asbestos fiber leads to increased risk of developing neoplastic tumors of the gastrointestinal tract. Difference in fiber type appears to have little effect on relative excess mortality (Table 3).

Table 3. Asbestos fiber type and increased risk of gastrointestinal cancer.

Fiber	No. cases	Expected	Observed
Amosite ^b	230	1.6	5.0
Chrysotile ^c	632	9.4	29.0

^a Two occupational settings utilizing different fiber types. Approximately three times as much gastrointestinal cancer occurs in both groups.

^b Data of Selikoff et al. (12).

^c Data of Selikoff et al. (7).

Fiber Amounts in Tissues and Disease; Important Questions, No Answers

Observations concerning fiber amounts in tissues and the occurrence of disease in occupationally exposed workmen, raises some important questions. One can ask whether asbestos is such a potent carcinogen, in and of itself, that it may be associated with a threefold increase of gastrointestinal cancer in occupationally exposed workmen when the amount of fiber in the affected tissues is many magnitudes less than observed in their lungs. This may suggest that tissues, other than pulmonary, are more sensitive to the insult of the dust. Perhaps the gastrointestinal tract is such a more sensitive tissue, thereby, making it one more "target" organ. It may also suggest that there are factors other than the fibers themselves which may be responsible for increased tissue response. The question arises as to the effects of "organic additives", such as nitroso compounds, and the role they may play in the etiology of gastrointestinal disease. Synergism, for neoplastic development between particle exposure and cigarette smoking was suggested several decades ago (16), but established epidemiologically for asbestos and smoke only recently (25). Multiple interaction of fiber and organic compound increases neoplastic disease risk in a multiplicative manner. What are the effects of such combinations as particle

to particle, particle to gas, and particle to organic phase in asbestos disease etiology? And, most important, does direct ingestion of small amounts of asbestos fiber, in food and water likely less than one observes for the working environment, pose a serious threat of neoplastic disease for individuals not occupationally exposed to the fiber? Does environmental direct ingestion of small amounts of asbestos fiber approach in magnitude the amounts of fiber inadvertently ingested, or migrated, following occupational exposure?

Is asbestos a unique substance with respect to disease? If so, what are the special properties which render it so unique? Is a narrow chemical or physical entity of asbestos within a broader range, the only biologically active member of that series? Although amosite is a narrowly defined mineral entity within the cumingtonite-grunerite series, it has been implied that all other members of this series are biologically inactive. This appears unlikely because the magnesium member of this series, which alters crystal form slightly, is another amphibole mineral called anthophyllite. Anthophyllite has been shown to be biologically active.

Considerations and Future Work

Future work in the area concerning the biological effects of inorganic particles will have to take into consideration a number of parameters. For example, there is a natural background, and a wide variety of inorganic particles which are normally ingested. Foods contain inorganic components, as well as adventitious minerals within the substance itself; some of which have formerly been considered inert. Opal phytoliths have long been known to occur as siliceous parts of plants (22). Their long-term biological activity is known only in animals (23). Potable water may carry large quantities of dissolved and suspended materials which are ingested. Their effects are unknown. The "natural" background of particles in ambient air is only partially known and is continually fluctuating qualitatively and quantitatively. The identity of many of the particulate components still have to be determined. In addition to these natural airborne materials, particle contaminants and additives in food, water, and beverages add to the body's burden.

In addition to characterizing of these particles, one must systematically examine the affected tissues. To this date, this has not been done. We observe in lung tissues of workmen that each portion of the lung contains a slightly different particle size fraction assemblage. This has not been studied for any of the extrapulmonary organs. Judging from the prevalence of neoplastic lesions in bowel, colon, and rectum, one may expect that the accumulation of particles in the digestive tract is site related. The development of proper tissue preparation techniques and instrumental methods of analysis has been achieved to a large degree (24). The expansion of this research into possible etiologic agents responsible, in part, for the development of malignant lesions in the human gastrointestinal tract, is only at the beginning stage.

Asbestos Exposure and Neoplastic Disease

Asbestos fibers may be easily detected in lung tissues of asbestos workmen (17). These men are generally exposed to highly concentrated dust aerosols so that the inhalation and entrapment of fibers in their lung tissues is certain. Many fibers are eliminated from the lungs by the body's normal defense mechanism. However, elimination never equals the quantity of material which is inhaled. The fate of these fibers which are removed from the lung may be twofold: they are eliminated from the body or they are removed from the lung, but migrate to other body sites (18-20). Migration may include hematogenous and/or lymphatic mechanisms

(19). The size distribution of even the larger amphibole fibers is such that this transport may be accomplished without size obstacle. A statistical study of the size distribution of amosite fibers in the lung of a workman exposed to dust occupationally, has demonstrated that 95% of the fibers in his lung was shorter than 5 μ m in length and the width distribution for these fibers was modally centered at 0.15 μ m size range (21). Even in these cases, the vast majority of fibers may easily migrate within the body. Therefore, asbestos fibers found in the extrapulmonary tissues of workmen may have come to rest in these organs through migration and/or direct ingestion.

We have studied extrapulmonary organs of asbestos workmen by both light and electron microscopy in order to determine the presence or absence of fibrous particles. The results of one such light microscopy study is shown in Table 4. On the basis of this very limited study by optical microscopy, we report that large fibers, those detectable by light microscopic techniques, exist in all extrapulmonary organs examined. However, their amounts are substantially less than light visible objects normally encountered in lung tissue from the same individual. Also, uncoated fibers exceed coated fibers (asbestos bodies) by substantial amounts. Using light visible objects as markers, fibers are observed in extrapulmonary organs, but they are magnitudes less in number than in the lung.

We also have examined a number of such extrapulmonary tissues by electron microscopy. Here also, on the sublight microscopic level, we

Table 4. Presence of asbestos bodies and uncoated fibers in lung and extrapulmonary tissues of workmen occupationally exposed to asbestos dust (light microscopy).^a

Case	Lung		Liver		Pancreas		Kidney		Adrenal		Spleen	
	Body	Fiber	Body	Fiber	Body	Fiber	Body	Fiber	Body	Fiber	Body	Fiber
1	>99	>99	1	5	2	16	0	47	— ^b	—	—	—
2	>99	>99	—	—	—	—	—	—	—	—	15	30
3	>99	>99	2	4	7	48	1	>99	—	—	—	—
4	>99	>99	0	1	—	—	0	69	0	5	—	—
5	>99	>99	0	2	2	10	0	60	—	—	—	—

^a Analyses were made on 175 μ m stacks of seven 25 μ m thick tissue sections. These are ashed in a low-temperature ashier, immersed in an appropriate oil, covered and scanned under phase-contrast at 440 $\times$ magnification. Area scanned about 10 $\times$ 10 mm. Counts stopped at 99.

^b Denotes no tissue available.

observe
microsc
less th
vation
organs
the out
blood
demon
the en
perime
liver.
animal
(19). I
means
body, a
come t
magnit
lung ti

Acknow

The
under
NIEHS
from :

1. Ang
Dul
O.R.
mer
Cha
2. Nic
A.S.
177
Wh
stm
A. Silv
Am
A. Har
ong
vold
K. K
L. L
S. S
188
En
En

observed uncoated fibers. Again, as with light microscopy, their concentration is significantly less than encountered in lung tissues. The observation that asbestos fiber may be present in organs which have limited direct contact with the outside environment, by direct ingestion and blood or lymph routes, emphatically demonstrates the ability of asbestos to infect the entire human body. It has been noted experimentally that asbestos fibers are found in liver, spleen, and kidney of experimental animals after asbestos instillation into the lung (19). Inhalation of asbestos fiber invariably means dissemination of fiber throughout the body, although the number of fibers which may come to rest in extrapulmonary organs may be magnitudes less than one observes entrapped in lung tissues.

Acknowledgement

The author wishes to acknowledge support under a Career Scientist Award from the NIEHS, ES44812 and under a Center Grant from the NIEHS, ES00928.

REFERENCES

1. Anonymous. A Preliminary Report on Asbestos in the Duluth, Minnesota Area. Office of Technical Analysis, Office of Enforcement and General Counsel, Environmental Protection Agency, Washington, D. C., 1974. Chaps. 1 and 3.
2. Nicholson, W. J., Maggiore, C. J., and Selikoff, I. J. Asbestos contamination of parenteral drugs. *Science* 177: 171 (1972).
3. Windum, H., Griffin, J., and Goldberg, E. D. Talc in atmospheric dust. *Environ. Sci. Technol.* 1: 923 (1967).
4. Silverberg, E., and Holleb, A. *Cancer Statistics, 1971*. American Cancer Society Inc. New York, N. Y., 1972.
5. Hammond, E. C. Personal communication concerning ongoing research at the East Orange V.A. Hospital, colon and rectum cancer, 1974.
6. Kral, Z. E. Asbestosis and abdominal neoplasms. *Lancet* 1211 (3, Dec. 1960).
7. Selikoff, I. J., Churg, J., and Hammond, E. C. Asbestos exposure and neoplasia. *J. Amer. Med. Assoc.* 188: 22 (1964).
8. Enterline, P. E., and Kendrick, M. A. Asbestos dust exposures at various levels and mortality. *Arch. Environ. Health* 15: 181 (1967).
9. Kleinfeld, M., et al. Mortality among talc miners and millers in New York State. *Arch. Environ. Health* 14: 663 (1967).
10. Elmes, P. C., and Simpson, M. Insulation workers in Belfast. 3. Mortality 1940-66. *Brit. J. Ind. Med.* 28: 226 (1971).
11. Newhouse, M., et al. A study of the mortality of female asbestos workers. *Brit. J. Ind. Med.* 29: 134 (1972).
12. Selikoff, I. J., Hammond, E. C., and Churg, J. Carcinogenicity of amosite asbestos. *Arch. Environ. Health* 25: 183 (1972).
13. Lamy, P., et al. Les opacités massives chez les mineurs de fer. *J. Franc. Med. Chir. Thoraciques* 13: 283 (1959).
14. Holaday, D. A. Uranium mining hazards. In: *Handbook of Experimental Pharmacology*. H. C. Hodge, J. N. Stannard, and J. B. Harsh, Eds., Springer-Verlag, New York, 1973.
15. Mackle, W., and Gregorius, F. Cancer of the respiratory system in the United States. *Publ. Health Repts.* 63: 1114 (1948).
16. Breslow, L., et al. Occupations and cigarette smoking as factors in lung cancer. *Am. J. Publ. Health* 44: 171 (1954).
17. Langer, A. M., and Pooley, F. D. Identification of single asbestos fibers in human tissues. In: *Proceedings of the International Conference on Biological Effects of Asbestos*, Lyon, France, Oct. 2-6, 1972, Bogovski, P., Gilson, J.C., Timbrell, V., J.C. Wagner, Eds., IARC, Lyon, Scientific Pub. No. 8: 119 (1974).
18. Holmes, A., and Morgan, A. Investigations into the use of radioactive asbestos in studying the translocation of fibers injected intrapleurally into rats. Report Atomic Energy Authority of the UK, Harwell, 1967.
19. Kanazawa, K., Birbeck, R. L., and Roe, F. J. C. Migration of asbestos fibers from subcutaneous injection sites in mice. *Brit. J. Cancer* 24: 96 (1970).
20. Westlake, G. E., Spjut, H. J., and Smith, M. N. Penetration of colonic mucosa by asbestos particles. An electron microscopic study in rats fed asbestos dust. *Lab. Invest.* 14: 2029 (1965).
21. Langer, A. M., and Mackler, A. D. Distribution of asbestos fibers in the lung of an asbestos worker. In preparation.
22. Parry, D. W., and Smithson, F. Opaline silica in the inflorescences of some British grasses and cereals. *Ann. Botany* 30: 119 (1966).
23. Baker, G., and Jones, L. H. P. Opal in the animal body. *Nature* 189: 682 (1961).
24. Langer, A. M., et al. Identification of asbestos in human tissues. *J. Occup. Med.* 16: 287 (1973).
25. Selikoff, I. J., Hammond, E. C., and Churg, J. Asbestos exposure, smoking, and neoplasia. *J. Amer. Med. Assoc.* 204: 106 (1968).