

FINAL REPORT
Epidemiology, 220-6W-08
Medical Department
3M Company
St. Paul, MN 55144

Date: January 30, 2013

Title: Biomonitoring Assessment of the Cottage Grove East Cove Remediation Project

Study
Start Date: September 1, 2011

Protocol Number N/A
IRB Approval N/A

Principal Investigator: Geary W. Olsen, D.V.M., Ph.D.¹
Co-investigators: Barbara A. Gibson, M.D., M.P.H.¹
David J. Ehresman, B.S., MT (ASCP)²

Study Director: Carol A. Ley, M.D., M.P.H.

1. Corporate Occupational Medicine, Medical Department, 220-6W-08, St. Paul, MN 55144
2. Toxicology Laboratory, Tox Assmt & Compl Assurance, Medical Department, Mail Stop 236-1B-22, St. Paul, MN 55144

Summary

A total of 21 subjects, who worked on the 3M Cottage Grove East Cove remediation project, had baseline and end-of-project serum clinical chemistry and perfluoroalkyl (PFOA and PFOS) measurements. There were no statistically significant *increases* in paired PFOA and PFOS concentrations when the end-of-project values were compared to baseline assessments.

Conclusion: Based on the biomonitoring data, there was no substantive occupational exposure to PFOA or PFOS while working on the 3M Cottage Grove East Cove remediation project.

Introduction

Several remediation projects have been completed regarding legacy perfluorochemicals. These included the 3M Cottage Grove Building 15/73 demolition and disposal project, the 3M Woodbury landfill remediation project, the Decatur Building 2/49 demolition and disposal project, and the 3M Cottage Grove Building 25 re-roof project. For each of these projects, baseline and end-of-project assessments were required of 3M employees and contractor workers who entered specified work zones where potential exposure to legacy perfluorochemicals [PFOS (perfluorooctanesulfonate) and PFOA (perfluorooctanoate)] was possible but due to the uniqueness of the work the magnitude was unknown. Both baseline and end-of-project assessments required response to a medical questionnaire and blood (serum) measurements of PFOS, PFOA, and several clinical chemistries. The purpose of this report is to provide a descriptive analysis of the PFOA and PFOS serum concentrations (ng/mL) at baseline and end-of-project time periods for the 3M Cottage Grove East Cove remediation project. The project began in mid-August 2011 and concluded in mid-January 2012.

Methods

1. Informed consent

The purpose of the project was explained in an Informed Consent. Subjects read and signed this informed consent at both baseline and end-of-project assessments. Subjects were informed they could not work on this specific project without such compliance.

2. Clinical chemistries

Clinical chemistries included a lipid panel profile, blood glucose, BUN, creatinine, serum electrolytes, and liver enzyme tests (including alkaline phosphatase, AST, ALT, and total bilirubin). Fasting was not a requirement because of the logistics of collecting blood samples during various times of the day. Clinical chemistries were analyzed by Quest Diagnostics. Individual clinical chemistry analyses at baseline and at end-of-project were medically reviewed by Dr. Gibson. Subjects were informed in writing that this examination program was not a full medical ‘check-up.’ Values out of reference range were indicated with a notation for the subjects to follow up with their primary care physician if they had abnormal test results. However, questions could be directed to Dr. Gibson regarding their clinical chemistry test results should the subject so desire.

3. Analytical measurements of PFOA and PFOS

Serum samples were analyzed for PFOA and PFOS by state-of-the-art high performance liquid chromatography mass spectrometry methods by the 3M Medical Department’s Toxicology

Laboratory under the direction of Dave Ehresman. Medical Department personnel collected blood samples that were processed to provide serum samples for analysis. These samples were assigned unique identification numbers and randomized prior to the samples being delivered for analysis. The 3M Medical Department's Toxicology Laboratory was "blinded" to the identity of all samples received for analysis.

Sample extractions were performed using solid phase extraction (SPE) technique. The extraction and sample clean-up was based on a 100 uL sample size and utilized Waters (Milford, MA) Oasis® hydrophilic-lipophilic balance (HLB) 3.0mL cartridges (Ehresman et al. 2007).

The method used two stable labeled internal standards for quantitation. The internal standards used were a dual labeled PFOS where two ^{18}O molecules were included in the sulfonate group (internal standard, >99% purity, synthesized by Research Triangle Institute, Research Triangle Park, NC) and a dual labeled PFOA molecule, where the carboxyl and alpha carbons were labeled with ^{13}C stable isotope (greater than 97%, provided by DuPont, Wilmington, DE). All quantitations were based on matrix matched extracted standard curves.

A 5 uL injection of the sample eluate was introduced into the High Pressure Liquid Chromatograph (HPLC) which was directly interfaced into the triple quadrupole mass spectrometer (Applied Biosystems/MDS-Sciex Instrument Corporation, Forest City, CA). Standard curves covered the range from 1.0 - 150 ng/mL. Standard curves were evaluated using a quadratic regression model where the standards were weighted at $1/x$, and each curve had an "R" value equal to or greater than 0.9998. Matrix spiked controls (QC samples) evaluated during this study all had acceptable results "within" their previously established ranges. Matrix-matched dilutions were used for samples requiring dilution to bring the samples into the linear range of the assay. Extracted serum and aqueous blanks remained below the lower limit of

quantitation established at 1.0 ng/mL (lowest standard fitted on the standard curve used for this project).

4. Communication

After each blood collection, individual letters were sent to the participants describing their results. In the baseline assessments, two letters were sent. One referred to the clinical chemistries and the other letter provided baseline PFOA and PFOS concentrations. At the end-of-project, two letters were again sent to each participant. The first letter provided the individual's clinical chemistry results. The second letter compared baseline to end-of-project serum PFOA and PFOS concentrations.

Results

A total of 21 3M employees and contract workers had paired baseline and end-of-project assessments. Five subjects had baseline assessments but did not participate in end-of-project assessments.

Table 1 and Table 2 (below) provide the measures of central tendency for baseline and end-of-project PFOA and PFOS concentrations for the 21 subjects with paired measurements.

Values are provided in ng/mL (parts-per-billion).

Table 1. PFOA Baseline and End-of-Project Concentrations (ng/mL)

	Min	Q1	Median	Q3	Max	<u>Arithmetic</u>		<u>Geometric</u>	
						Mean	95% CI	Mean	95% CI
Baseline	1.3	2.3	2.8	5.0	10.4	3.8	2.7 – 4.9	3.2	2.5 – 4.2
End-of-Project	1.0	2.0	2.7	4.6	7.5	3.3	2.5 – 4.1	2.9	2.3 – 3.7

Table 2. PFOS Baseline and End-of-Project Concentrations (ng/mL)

	Min	Q1	Median	Q3	Max	<u>Arithmetic</u>		<u>Geometric</u>	
						Mean	95% CI	Mean	95% CI
Baseline	7.6	11.0	13.8	21.3	33.5	16.4	13.1 – 19.7	15.1	12.4 – 18.2
End-of-Project	6.5	11.1	13.8	18.8	34.2	15.5	12.6 – 18.5	14.4	12.1 – 17.2

As the data are paired samples, a matched-paired statistical analysis was performed where the mean value was determined for the average of the sum of the individual differences for the 28 subjects. See Equation 1.

Equation 1.

$$\text{Mean Difference} = \sum_{i=1}^{21} (\text{end-of-project} - \text{baseline}) / 21$$

Because of the skewed PFOA and PFOS distributions, as seen in Tables 1 and 2, the mean differences of the natural logs of the paired values were also determined. See Equation 2.

Equation 2.

$$\text{Mean} = \sum_{i=1}^{21} [(\ln(\text{end-of-project value})) - (\ln(\text{baseline value}))] / 21$$

Results for *Equation 1 and Equation 2*, and associated statistics of the mean differences of the paired comparisons, are provided in Table 3 (below) for both PFOA and PFOS.

Table 3. Matched-pair Analysis Comparing Baseline to End-of-Project PFOA and PFOS concentration (ng/mL)

Mean Paired Difference	PFOA (ng/mL)	t-ratio	p value	PFOS (ng/mL)	t-ratio	p value
<i>Equation 1</i>	-0.5	-2.7	0.007	-0.9	-1.2	0.13
<i>Equation 2</i>	-0.1	-3.2	0.003	-0.04	-1.0	0.17

The only statistically significant difference was a *decrease* in the mean of the differences for PFOA (Equations 1 and 2). In other words, the end-of-project PFOA concentration was statistically significantly less than the baseline PFOS concentration (ng/mL) when analyzed as paired differences.

Discussion

Comparison of baseline and end-of-project serum concentrations of PFOA and PFOS provided each worker an excellent method to assess his/her exposure experience for the 3M Cottage Grove East Cove remediation project. Given the fact that exposure potential during the course of the remediation project was unknown, the use of biomonitoring data allowed for an assessment the actual exposure. Because the biomonitoring data indicated there were no substantive differences in any increase in exposure to PFOA or PFOS between baseline and end-of-project, the clinical chemistry data were therefore not analyzed in aggregate by paired

statistics. Potential exposure to any other materials would not be known since only PFOA and PFOS were analyzed in the workers' serum.

Conclusion Based on the biomonitoring data, there was no substantive occupational exposures to PFOA or PFOS for the 21 total 3M and contract workers involved with the 3M Cottage Grove East Cove remediation project.

References

Ehresman DJ, Froehlich JW, Olsen GW, Chang SC, Butenhoff JL. 2007. Comparison of human whole blood, plasma, and serum matrices for the determination of perfluorooctanesulfonate (PFOS), perfluorooctanoate (PFOA), and other fluorochemicals. Environ Res 103:176-184.