

file HED

From the desk of

W. B. PATAGEORGE

11/7/75

Copies to:

D. R. Bishop
W. C. Hammann
C. Paton
R. A. Pohl
R. G. Potter
D. L. Sliney
J. C. Weber
W. W. Withers/T. H. Bottini
D. Wood
J. J. Zeman

MONS 092911



DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE
PUBLIC HEALTH SERVICE
CENTER FOR DISEASE CONTROL

NATIONAL INSTITUTE FOR OCCUPATIONAL
SAFETY AND HEALTH
3600 FISHERS LANE
ROCKVILLE, MARYLAND 20852

November 4, 1975

NOV 8 1975

Mr. W. B. Papageorge
Speciality and Process Chemicals
Division
Monsanto Industrial Chemicals Company
800 North Lindberg Boulevard
St. Louis, Missouri 63166

Dear Mr. Papageorge:

Thank you for your assistance in providing us with information
on polychlorinated biphenyls.

Enclosed is a copy of the current intelligence bulletin on poly-
chlorinated biphenyls which was released yesterday afternoon.

Sincerely,

Harvey P. Stein

Harvey P. Stein, Ph.D.
Chemist
Priorities Evaluation Branch
Office of Occupational Health
Surveillance and Biometrics

Enclosure

MONS 092912



DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE
PUBLIC HEALTH SERVICE
CENTER FOR DISEASE CONTROL

NATIONAL INSTITUTE FOR OCCUPATIONAL
SAFETY AND HEALTH
5600 FISHERS LANE
ROCKVILLE, MARYLAND 20852

NOV 3 1975

Dear Colleague:

The enclosed background material has been prepared by this office to inform the occupational health community of current knowledge concerning industrial uses and toxicity of polychlorinated biphenyls.

Your comments and suggestions for changes to future reports are solicited.

Sincerely yours,

J. William Lloyd, Sc.D.
Director
Office of Occupational Health
Surveillance and Biometrics

Enclosure

MONS 092913

BACKGROUND INFORMATION ON POLYCHLORINATED BIPHENYLS

Office of Occupational Health Surveillance and Biometrics

National Institute for Occupational Safety and Health

November 3, 1975

MONS 092914

POLYCHLORINATED BIPHENYLS


Introduction

Reports of adverse health effects in humans and the demonstration of carcinogenic effects in certain animal species have led to the reexamination of the distribution of polychlorinated biphenyls (PCBs) in the environment and the potential health effects of human exposure.

Because the industrial environment represents the major source of potentially high exposures to PCBs, the National Institute for Occupational Safety and Health (NIOSH) has gathered pertinent information on the manufacture, uses, and reported deleterious effects of polychlorinated biphenyls and is advising the occupational health community of these hazards.

Background

Polychlorinated biphenyls (PCBs) describe a group of synthetic chlorinated organic compounds having the following structure:



where each of the ten Z's can represent either a hydrogen or a chlorine atom. There are two hundred nine different chlorinated biphenyls and they are collectively referred to as PCBs although many are not actually polychlorinated. Approximately half of these compounds have been synthesized and characterized.

Mixtures of polychlorinated biphenyls are important industrial products. The most common tradenames for these mixtures are Aroclor,^a Inerteen,^b Kanechlor,^c and Pyranol.^d Other known tradenames for PCB-containing products are listed in Table I. PCB-containing dielectrics (electrical insulators) are generally referred to as "askarels" in the electrical industry.

Mixtures of polychlorinated biphenyls are very resistant to degradation, are thermally stable, and resistant to oxidation, acids, bases, and other chemical agents. They are soluble in most of the common organic solvents and lipids, but only slightly soluble in water, glycerol, and glycols. PCBs are good electrical insulators. Although most individual polychlorinated biphenyls are solids at room temperatures, the mixtures vary in consistency from mobile oils to viscous liquids or sticky resins.

PCBs are generally prepared industrially by the chlorination of biphenyl with anhydrous chlorine in the presence of iron filings or a ferric chloride catalyst. Trace quantities of chlorinated naphthalenes and chlorinated dibenzofurans have been reported in some commercial samples of PCBs and it has been suggested that the presence of these impurities may be of toxicological significance.(1,2,3)

^a Aroclor is a registered trademark of the Monsanto Company.

^b Inerteen is a registered trademark of the Westinghouse Electric Corporation for its brand of PCB containing dielectric fluids.

^c Kanechlor is a registered trademark of the Kanegafuchi Chemical Industry Company, Ltd.

^d Pyranol is a registered trademark of the General Electric Company for its brand of PCB containing dielectric fluids.

Commercial PCBs are generally mixtures of many different chlorinated biphenyls, as shown in Table II, manufactured to meet operational specifications (such as dielectric constant, flash point, fire point, density, percent chlorine, and color); these commercial mixtures may vary chemically from batch to batch.

PCBs have found wide use in industry and have been manufactured in the United States since 1929. The major domestic manufacturer, Monsanto Company, produces PCBs at Sauget, Illinois and reports manufacturing 40 million pounds of PCBs in the United States during 1974 (down from 85 million pounds in 1970). (4) Monsanto's domestic production and sales of PCBs by grade and category from 1957 through the first quarter of 1975 are shown in Table III.

Uses

PCBs are employed in capacitors and transformers because they combine attractive dielectric properties with chemical stability and fire resistance. Approximately twice as many pounds of PCBs are used in the manufacture of capacitors as in the manufacture of transformers.

Prior to the environmental concern surrounding the persistence and ubiquitousness of PCBs, (5) they were more widely used in industry as fluids for heat transfer systems, hydraulic systems, gas turbines, and vacuum pumps, as fire retardants, and as plasticizers in adhesives, textiles, surface coatings, sealants, printing, and carbonless copy paper.

Beginning in 1971, Monsanto voluntarily restricted its domestic sales of PCBs to closed system dielectric applications in capacitors and transformers. (6) Other current domestic applications of PCBs include use in investment casting processes, as heat exchange fluids, and as hydraulic

fluids. Imports of PCBs have been estimated to exceed 375,000 pounds for 1974. Reclaimed PCBs also are reported to be available.(7)

More than 95 percent of all power capacitors contain PCBs. Among their applications are use on electric utility lines, in air conditioners, and in the ballast of fluorescent lamp fixtures. PCBs are employed for safety, reliability, and long life, as well as to achieve size compatibility with equipment and installation requirements. However, non-PCB power capacitors are being manufactured (e.g., General Electric's Econol line and Sprague's Eccol line) which may serve as alternatives.(8)

PCBs are employed in transformers at locations where their proximity to people and/or property demand a fire resistant dielectric. Approximately 5 percent of transformers are PCB filled and each of the transformers so filled contains between 40 and 500 gallons of PCBs (about 235 gallons is average). Possible alternatives to PCB filled transformers may include dry transformers (which are larger) as well as transformers filled with silicone fluids or other materials under evaluation.(9,9)

Chlorinated terphenyls, as well as PCBs, are used in some formulations of wax for investment casting processes.(8) The chemical structure of chlorinated terphenyls resembles that of PCBs. Chlorinated terphenyls have been reported to have toxicological effects similar to those of PCBs.(10)

Approximately 330,000 pounds of chlorinated terphenyls were imported into the United States during 1974(11) and 125,532 pounds during 1973.(12)

Toxicity

PCBs are poorly metabolized and tend to accumulate in animal tissues, including humans.(13,14,15,16,17) The accumulation, particularly in

tissues and organs rich in lipids, appears to be higher in the case of penta and more highly chlorinated biphenyls.(18)

Studies have revealed PCBs in human fat tissue and blood plasma. PCBs, in amounts greater than 2 parts per million, were reported in 198 of 637 (31 percent) samples of human fat tissue taken from the general population of 18 states and the District of Columbia.(14) PCB residues ranging up to 29 parts per billion have also been found in 43 percent of 616 plasma samples collected from volunteers in a southeastern U.S. county.(15)

Human:

The known toxic effects of PCBs in humans include an acnelike skin eruption (chloracne), pigmentation of the skin and nails, excessive eye discharge, swelling of eyelids, and distinctive hair follicles.(19)

For a number of years, chloracne of the face and neck has been reported among workers exposed to chlorinated hydrocarbons. Workers exposed to PCBs in the process of insulating cables,(20) in the production of condensers(21) and in the manufacture of chlorobiphenyls(22) have reported these skin lesions along with systemic effects such as digestive disturbances, edema of the face and hands, burning of the eyes, impotence, and hematuria.(16,22)

The toxic effects of PCBs in humans are further illustrated by a 1968 outbreak of poisoning in Japan that involved over 1,000 people who ingested PCB contaminated rice bran oil for a period of several months. The contamination of the oil (estimated 1,500 to 2,000 ppm) occurred when heat transfer pipes immersed in the oil during processing developed pin-sized holes. The clinical aspects of the poisoning included chloracne,

brown pigmentation of the skin and nails, distinctive hair follicles, increased eye discharge, swelling of eyelids, transient visual disturbance, and systemic gastrointestinal symptoms with jaundice(19). In some patients, symptoms persisted 3 years after PCB exposure was discontinued. Infants born to poisoned mothers had decreased birth weights, and showed skin discoloration due to PCB placental passage. Two stillbirths to PCB exposed women were also reported.(23)

Animal:

The toxic effects of PCBs in animals have been studied extensively. Reports of malignant, non-malignant, and reproductive effects are shown in the following three tables:

MALIGNANT PATHOLOGIC EFFECTS INDUCED BY PCBs IN ANIMALS

<u>PCB Mixture</u>	<u>Treatment</u>	<u>Animal</u>	<u>Effects</u>
Kanechlor 500(24)	500 ppm, 250 ppm	Mice	Liver weight increased with percent chlorine and dosage.
Kanechlor 400	and 100 ppm in		
Kanechlor 300	diet for 32 weeks		Kanechlor 400 and 500 produced liver cell hypertrophy.
			500 ppm of Kanechlor 500 produced hyperplastic nodules in 7 of 12 mice and hepatocellular carcinomas in 5 of 12 mice
			No nodules or carcinomas in controls
Aroclor 1254(25)	300 ppm in diet for 6 & 11 months	Mice	Increased liver weight. Adenofibrosis in all mice fed for 11 months. None of 22 mice fed for 11 months had hepatomas. One of 24 mice fed for 6 months had a hepatoma.
			No hepatomas in control mice.
Aroclor 1260(26)	100 ppm in diet for 21 months	Rats	Hepatocellular alteration in 183 of 184 rats fed PCBs and in 28 of 173 control rats. Neoplastic nodules in 26 of 184 PCB fed rats and none in control rats.
			Hepatocellular carcinomas in 26 of 184 rats fed PCBs and in 1 of 173 control rats.

NON-MALIGNANT PATHOLOGIC EFFECTS INDUCED BY PCBS IN ANIMALS

<u>PCB Mixture</u>	<u>Animal</u>	<u>Route of Administration</u>	<u>Effects</u>
<u>ACUTE EFFECTS</u>			
Unspecified Product Containing 42 % Chlorine(27)	Guinea Pigs Rats Rabbits	Subcutaneous injection Skin application Feeding	Fatty degeneration and central atrophy of the liver.
Aroclor 1242(28) Aroclor 1254	Guinea pigs Rats Mice Rabbits Cat	Inhalation	<u>Aroclor 1242</u> produced no ill effects on basis of mortality, growth, pathology, organ enlargement, liver function, or hematological changes. <u>Aroclor 1254</u> produced no harmful effects regarding growth or mortality, but did produce enlarged livers with microscopic evidence of hepatic cellular injury.
Aroclor 1248(29)	Monkeys	Feeding	Weight loss, hair loss, mouth and eyelid edema, acneform lesions, decreased hemoglobin and hematocrit, severe gastric mucosal ulceration and extreme hypertrophy of the liver.
<u>CHRONIC EFFECTS</u>			
Aroclor 1254(30) Aroclor 1260	Rats	Feeding	Hypertrophy of the liver cells, a brown pigment in the Kupffer cells, lipid accumulation in the cytoplasm of hepatocytes and adenofibrosis.

MONS 092921

REPRODUCTIVE EFFECTS OF PCBS IN ANIMALS

<u>PCB Mixture</u>	<u>Animal</u>	<u>Route of Administration</u>	<u>Effects</u>
Aroclor 1242(31) Aroclor 1254 Aroclor 1260	Chickens	Feeding	Aroclor 1242 and 1254 reduced egg production and hatchability and caused thin eggshells. Aroclor 1260 produced no harmful effects.
Aroclor 1254(32)	Mallards Bobwhites	Feeding	No adverse effects.
Aroclor 1254(33)	Pheasant	Feeding	Reduced egg production and hatchability.
Aroclor 1254(34)	Mink	Feeding	Severely affected reproduction.
Aroclor 1248(29)	Female monkeys	Feeding	Reduced ability to become pregnant. Pregnancies produced small infants with PCBs in tissues.

MONS 092922

Occupational Exposure

It is estimated that 12,000 people are occupationally exposed to polychlorinated biphenyls. The majority of these exposures are in capacitor production and in investment casting processes.

Permissible Occupational-Exposure

The current Occupational Safety and Health Administration, Department of Labor standards for chlorinated biphenyls are 1 mg/cubic meter for 42 percent chlorine mixtures and .5 mg/cubic meter for 54 percent chlorine mixtures. These are based on the Threshold Limit Values (TLV) established by the American Conference of Governmental Industrial Hygienists.(35)

BIBLIOGRAPHY

1. Hutzinger, O., Safe, S., and Zitko, V.: The Chemistry of PCB's, CRC Press, Cleveland, Ohio, p. 22, 1974
2. Bowes, G.W., Mulvihill, M.J., Simoneit, B.R.T., Burlingame, A.L., and Risebrough, R.W.: Identification of Chlorinated Dibenzofurans in American Polychlorinated Biphenyls, Nature, 256:305, 1975
3. Curley, A., Burse, V.W., Jennings, R.W., Villanueva, E.C., and Kimbrough, R.D.: Evidence of Tetrachlorodibenzofuran (TCDF) in Aroclor 1254 and the Urine of Rats Following Dietary Exposure to Aroclor 1254, Bul Environ Contam and Toxicol, 14(2):153-58, 1975
4. Communication from Monsanto Industrial Chemicals Company, St. Louis, Missouri, September, 1975
5. Interagency Task Force on PCBs: PCBs and the Environment, COM-72-10419, Washington, DC, March 20, 1972
6. Polychlorinated Biphenyls: Monsanto Company's Program of Restricted Sales and Controlled Manufacturing Practices, Monsanto Industrial Chemicals Company, St. Louis, Missouri, October, 1974
7. Glenn Schweitzer, Office of Toxic Substances, U.S. Environmental Protection Agency, testimony at an August, 1975, Wisconsin hearing, quoted in Toxic Materials News, p. 139, September 15, 1975
8. Personal communications with representatives of the industry, October 1975
9. Dow Corning Dielectric Liquids for Power Transformers, Dow Corning Corporation, Midland, Michigan, received September, 1975 (booklet not dated)
10. Allen, J.R., Abrahamson, L.J., and Norback, D.H.: Biological Effects of Polychlorinated Biphenyls and Triphenyls on the Subhuman Primate, Environmental Research, 6:344-54, 1973
11. Telephone conversation with a representative of the U.S. International Trade Commission, Washington, DC, September, 1975
12. U.S. Tariff Commission: Imports of Benzoid Chemicals and Products, Washington, DC., p. 24, 1973
13. Biros, F.J., Walker, A.C., and Medbery, A.: Polychlorinated Biphenyls in Human Adipose Tissue, Bull Environ Contam and Toxicol, 5:317-23, 1970
14. Yobs, A.R.: Levels of Polychlorinated Biphenyls in Adipose Tissue of the General Population of the Nation, Environmental Health Perspective, 1:79-80, 1972

15. Finklea, J., Priester, L.E., Creason, J.P., Hauser, T., Hinners, T., and Hammer, D.I.: Polychlorinated Biphenyl Residues in Human Plasma Expose a Major Urban Pollution Problem, *AJPH*, 62(5):645-51, 1972
16. Kimbrough, R.D.: The Toxicity of Polychlorinated Polycyclic Compounds and Related Chemicals, *Critical Reviews Toxicology*, CRC Press, Inc., 2(4):445-98, 1974
17. International Agency for Research on Cancer, IARC Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Man, 7:261-89, 1974
18. Hutzinger, O., et al., op. cit., p. 2
19. Kuratsune, M., Yoshimura, T., Matsuzaka, J., and Yamasuchi, A.: Yusho, a Poisoning Caused by Rice Oil Contaminated with Polychlorinated Biphenyls, *HSMHA Health Reports*, 36:1083-91, 1971
20. Fulton, W.B. and Matthews, J.L.: A Preliminary Report of the Dermatological and Systemic Effects of Exposure to Hexachloro-Naphthalene and Chloro-Diphenyl, Bureau of Industrial Standards, Harrisburg, Pennsylvania, 1936
21. Mayers, M.R. and Silverberg, M.G.: Skin Conditions Resulting from Exposure to Certain Chlorinated Hydrocarbons, *J Indust Hyg Toxicol*, 20:244-58, 1938
22. Schwartz, L.: Dermatitis from Synthetic Resins and Waxes, *Am J Pub Health*, 26:586-92, 1936
23. Miller, R.W.: Cola-colored Babies, Chlorobiphenyl Poisoning in Japan, *Teratology*, 4:211-12, 1971
24. Ito, N., Nagasaki, H., Arai, M., Masayuki, A., Makiura, S., Suginara, S., and Hirao, K.: Histopathologic Studies on Liver Tumorigenesis Induced in Mice by Technical Polychlorinated Biphenyls and Its Promoting Effect on Liver Tumors Induced by Benzene Hexachloride, *JNCI*, 51(5):1637-46, 1973
25. Kimbrough, R.D. and Linder, R.E.: Induction of Adenofibrosis and Hepatomas of the Liver in BALB/cJ Mice by Polychlorinated Biphenyls (Aroclor 1254), *JNCI*, 53(2):547-52, 1974
26. Kimbrough, R.D., Squire, R.A., Linder, R.E., Strandberg, J.D., Montali, R.J., and Burse, V.W.: Induction of Liver Tumors in Rats by Polychlorinated Biphenyl Aroclor 1260, *JNCI*, (in press), December, 1975
27. Miller, J.W.: Pathologic Changes in Animals Exposed to a Commercial Chlorinated Diphenyl, *Pub Health Reports*, pp. 1085-93, August 18, 1944

28. Treon, J.F., Cleveland, F.P., Cappell, J.W., and Atchley, R.W.: The Toxicity of the Vapors of Aroclor 1242 and Aroclor 1254, Am Ind Hyg Quart, 17:204-13, 1956
29. Allen, J.R.: Response of Primates to PCB Exposures, Federation Proceedings, 34:1675-79, 1975
30. Kimbrough, R.D., Lipder, R.E., and Gaines, T.B.: Morphological Changes in Liver of Rats Fed Polychlorinated Biphenyls, Arch Environ Health, 25:354-64, 1972
31. Keplinger, M.L., Fancher, O.E., and Calandra, J.C.: Toxicologic Studies with Polychlorinated Biphenyls, Toxicol Appl Pharmacol, 19:402, 1971
32. Heath, R.G., Spann, J.W., Kreitzer, J.R., and Vance, C.: Effects of Polychlorinated Biphenyls on Birds, Proc 15th Intern Ornithol Congr, Leiden, Holland, p. 476, 1972
33. Dahlgren, R.B., Linder, R.L., and Carlson, C.W.: Polychlorinated Biphenyls: Their Effects on Pinned Pheasants, Environ Health Perspect, 1:89, 1972
34. Ringer, R.K., Aulerich, R.J., and Zabik, M.: Effect of Dietary Polychlorinated Biphenyls on Growth and Reproduction of Mink, ACS Div Water Air and Waste Chemistry, Preprints of Papers 12:149, 1972
35. Federal Register, Vol. 37, No. 202, p. 22140, Wednesday, October 18, 1972

MUNS 092926

TABLE I
 TRADENAMES FOR KNOWN PCB CONTAINING PRODUCTS

<u>Tradename</u>	<u>Tradename Owner</u>
Aroclor	Monsanto Company St. Louis, MO
Chlorextol	Allis-Chalmers Milwaukee, WI
Clophen	Farbenfabriken Bayer GmbH Germany
Dykanol	Federal Pacific Electric Co. Newark, NJ
Fenclor	Caffaro S.P.A. Italy
Inerteen	Westinghouse Electric Corp. Pittsburgh, PA
Kanechlor	Kanegafuchi Chemical Industry Co., Ltd. Japan
Noflamol	Wagner Electric Corporation Newark, NJ
Phenoclor	Prodelec France
Pyralene	Prodelec France
Pyranol	General Electric Co. Schenectady, NY
Santotherm	Mitsubishi-Monsanto Japan
Therminol*	Monsanto Co. St. Louis, MO

*Therminol products now formulated in the U.S. do not contain PCBs.

MONS 092927

TABLE II
DESCRIPTION OF PCB MIXTURES

	Aroclor 1221 ^a	Aroclor 1016 ^a	Aroclor 1242 ^a	Aroclor 1254 ^a	Aroclor 1260 ^b	Kaeclor 300 ^c	Kaneclor 400 ^c	Kaneclor 500 ^c
Approximate Chlorine Content	21%	42%	42%	54%	60%	42%	48%	53%
Components								
Biphenyl	11	<0.1	<0.1	<0.1				
Monochlorobiphenyls	51	1	1	<0.1				
Dichlorobiphenyls	32	20	16	0.5		17	3	
Trichlorobiphenyls	4	57	49	1		60	33	5
Tetrachlorobiphenyls	2	21	25	21		23	44	27
Pentachlorobiphenyls	<0.5	1	8	48	12	0.6	16	55
Hexachlorobiphenyls	none detected	<0.1	1	23	38		5	13
Heptachlorobiphenyls	none detected	none detected	<0.1	6	41			
Octachlorobiphenyls	none detected	none detected	none detected	none detected	8			
Nonachlorobiphenyls					1			

Aroclor is a registered trademark of the Monsanto Company.

Kaneclor is a registered trademark of the Kanegafuchi Chemical Industry Company, Ltd.

It must be emphasized that these are approximate compositions of the PCB mixtures and that a particular product may vary in chemical composition from batch to batch.

^a Weight-weight percent. None detected = less than 0.01%. Source of component compositions: Monsanto Company, quoted in Hutzinger, O., et al., op. cit., p. 23

^b Percent. Source of component compositions: Thruston, A., PCB Newsletter, No. 3, July 1971, quoted in Hutzinger, O., et al., op. cit., p. 23

^c Percent. Source of component compositions: Ito, N., et al., op. cit., p. 1637

TABLE III

PCB MANUFACTURE AND SALES
MONSANTO INDUSTRIAL CHEMICALS COMPANY
1965 THROUGH 1975
(Thousands of Pounds)

	1965	1966	1967	1968	1969	1970	1971	1972	1973	1974	(1st quarter) 1975
U.S. PRODUCTION	60480	65849	75309	82854	76389	85054	34994	38600	42178	40466	8532
DOMESTIC SALES	51796	59078	62466	65116	67194	73061	34301	26408	37742	34406	7986
U.S. EXPORT SALES	4234	6852	8124	11231	10624	13651		6388	8346	5395	1538

DOMESTIC SALES BY CATEGORY

Heat Transfer	1237	1766	2262	2529	3050	3958	3060	752			
Hydraulics/ Lubricants	4616	4258	4643	5765	8039	7403	1552	0			
Misc. Industrial	1841	1779	1426	1283	1079	1627	1155	0			
Transformer	8657	8910	11071	11585	12105	13828	11134	} 25656	37742	34406	7986
Capacitor	23749	28884	29703	29550	25022	26708	14141				
Plasticizer											
Applications	11696	13481	13361	14404	16460	19537	3259	0			
Petroleum											
Additives	--	--	--	--	1439	--	--	0			

DOMESTIC SALES BY PCB GRADE

Aroclor 1221	369	528	442	136	507	1476	2215	171	35	57	10*
Aroclor 1232	7	16	25	90	273	260	171	0	0	0	0
Aroclor 1242	31533	39557	43055	44853	45491	48588	21981	728	6200	6207	2201**
Aroclor 1248	5565	5015	4704	4894	5650	4073	213	807	0	0	0
Aroclor 1254	7737	7035	6696	8891	9822	12421	4661	3495	7976	6185	2115**
Aroclor 1260	5831	5875	6417	5252	4439	4890	1725	305	0	0	0
Aroclor 1262	558	768	840	720	712	1023	1	0	0	0	0
Aroclor 1268	196	284	287	280	300	330	0	0	0	0	0
Aroclor 1016	0	0	0	0	0	0	3334	20902	23531	21955	3660*

* Used primarily in capacitors

** Used primarily in transformers

MONS 092929

TABLE III (continued)

PCB MANUFACTURE AND SALES
MONSANTO INDUSTRIAL CHEMICALS COMPANY
1957 THROUGH 1964
(Thousands of Pounds)

	1957	1958	1959	1960	1961	1962	1963	1964
U.S. PRODUCTION	(1)	(1)	(1)	37919	36515	38353	44734	50833
DOMESTIC SALES	32299	26061	31310	35214	37538	38043	38132	44869
U.S. EXPORT SALES	(2)	(2)	(2)	(2)	(2)	(2)	3647	4096

DOMESTIC SALES BY CATEGORY

Heat Transfer	--	--	--	--	--	157	582	929
Hydraulics/Lubricants	1612	1549	2685	2523	4110	3915	3945	4374
Miscellaneous Industrial	704	755	1569	1559	2114	1681	1528	1692
Transformer	12955	5719	5984	7921	6281	7984	7290	7997
Capacitor	17028	14099	16499	16967	15935	15382	15606	19540
Plasticizer Applications	(1)	3939	4573	6244	9098	8924	9181	10337
Petroleum Additives	--	--	--	--	--	--	--	--

DOMESTIC SALES BY PCB GRADE

Aroclor 1221	23	16	254	103	94	140	361	596
Aroclor 1232	196	113	240	155	241	224	13	13
Aroclor 1242	18222	10444	13598	18196	19827	20654	18510	23571
Aroclor 1248	1779	2559	3384	2827	4023	3463	5013	5238
Aroclor 1254	4461	6691	6754	6088	6294	6325	5911	6280
Aroclor 1260	7587	5982	6619	7330	6540	6595	7626	8535
Aroclor 1262	31	184	359	326	361	432	414	446
Aroclor 1268	--	72	102	189	158	210	284	190
Aroclor 1016	--	--	--	--	--	--	--	--

(1) Production figures and Plasticizer Applications figures unavailable during year indicated.

(2) U.S. Export Sales figures unavailable during year indicated.

SOURCE: Monsanto Industrial Chemicals Company, St. Louis, Missouri, September, 1975.

MUNS 092930