

REPORT

EVALUATION OF THE MUTAGENIC ACTIVITY OF

T-5869

IN AN IN VITRO MAMMALIAN CELL GENE MUTATION TEST

WITH L5178Y MOUSE LYMPHOMA CELLS

(WITH INDEPENDENT REPEAT)

NOTOX Project 115954
NOTOX Substance 38196

- page 1 of 22 -

RECEIVED

NOV - 9 1994

3M

TOXICOLOGY

STATEMENT OF GLP COMPLIANCE

NOTOX B.V., 's-Hertogenbosch, The Netherlands

The study described in this report was conducted in compliance with the most recent edition of:

The OECD Principles of Good Laboratory Practice

which are essentially in conformity with:

The United States Food and Drug Administration. Title 21 Code of Federal Regulations Part 58.


The United States Environmental Protection Agency (FIFRA). Title 40 Code of Federal Regulations Part 160.

The United States Environmental Protection Agency (TSCA). Title 40 Code of Federal Regulations Part 792.

With the exception that the stability of the test substance in the vehicle was unknown.

Study Director

Ing. E.J. van de Waart


.....

Date: 19/04/1994

QUALITY ASSURANCE STATEMENT

NOTOX B.V., 's-Hertogenbosch, The Netherlands.


Study procedures were subject to periodic inspections and general non study specific processes were also inspected at periodic intervals.

This report was audited by the NOTOX Quality Assurance Unit and the methods and results accurately reflect the raw data.

DATES OF QAU INSPECTIONS/ AUDITS	REPORTING DATES
January 24, 1994 February 10, 1994 March 28, 1994	January 24, 1994 February 14, 1994 March 28, 1994

Quality Assurance Manager

C.J. Mitchell B.Sc.




Date: 20-4-94

REPORT APPROVAL

STUDY DIRECTOR:

Ing. E.J. van de Waart


.....

Date 19/04/1994

MANAGEMENT:

Dr. I.C. Enninga
Technical Director


.....

Date 20/04/1994

PREFACE

Sponsor	3M Belgium - Chemical EBC Canadastraat 11 B-2070 ZWIJNDRECHT Belgium
Study Monitor	Mr. R.H. Cox
Testing Facility	NOTOX B.V. Hambakenwetering 3 5231 DD 's-Hertogenbosch The Netherlands
Study Director	Ing. E.J. van de Waart
Technical Coordinator	C.M. Verspeek
Study Plan	Start : February 01, 1994 Completed : February 22, 1994

TEST SUBSTANCE

Identification	T-5869
Description	Cream solid
Batch	2408
Purity	95%
Specific Gravity	1.5
Instructions for test substance storage	At room temperature in the dark
Stability under storage conditions	Stable
Expiry date	January 01, 1996
Stable for at least 4 hours in vehicle	Dimethylsulphoxide: not indicated

VEHICLE

The test substance was suspended or dissolved in dimethylsulphoxide of spectroscopic quality (Merck). Test substance concentrations were prepared directly prior to use. The final concentration of the solvent in the culture medium amounted to 0.8% (v/v).

GUIDELINES

The study procedures described in this report were based on the following guidelines:

- Organisation for Economic Co-operation and Development (OECD), OECD Guidelines for Testing of Chemicals, Guideline no. 476: "Genetic Toxicology: In Vitro Mammalian Cell Gene Mutation Tests", (adopted April 4, 1984).
- European Economic Community (EEC), Directive 87/302/EEC. Annex V of the EEC Directive 67/548/EEC, Part B: Methods for the Determination of Toxicity; "Other Effects-Mutagenicity: In Vitro Mammalian Cell Gene Mutation Test". EEC Publication no. L133 (adopted May 30, 1988).

ARCHIVING

NOTOX B.V. will archive the following data for at least 10 years: protocol, report, test article reference sample, all specimens and raw data.

OBJECTIVE

Purpose of the study

The objective of this study was to evaluate the test substance for its ability to induce forward mutations at the thymidine kinase (TK) locus in L5178Y mouse lymphoma cells. The assay was conducted in the absence and presence of a metabolic system (S9-mix). The TK mutational system detected base pair mutations, frame shift mutations and small deletions.

Justification for selection of the test system

L5178Y mouse lymphoma cells are used because they are sensitive indicators of mutagenic activity of a broad range of chemical classes. The TK mutational system is able to detect base pair alterations, frame shift mutations and small deletions.

Cells deficient in thymidine kinase (TK), due to the forward mutation (TK⁺/- to TK⁻/-) are resistant to the cytotoxic effects of the pyrimidine analogue trifluorothymidine (TFT). TK deficient cells can not incorporate the analogue into its phosphorylated derivative (nucleotide); the nucleotides needed for cellular metabolism are obtained solely from de novo synthesis. In the presence of TK, TFT is converted into nucleotides, which are lethal to the cells. Thus, cells which will survive in culture medium containing TFT are mutated, either spontaneously or by the action of the test substance, giving rise to a TK deficient phenotype.

A test article which induces a positive response in this assay is presumed to be a potential mammalian cell mutagen.

MATERIALS AND METHODS

TEST SYSTEM

Test System	L5178Y mouse lymphoma cells
Rationale	Recognized by the international guidelines as the recommended test system (e.g. EPA, OECD, EEC).
Source	Dr. A.G.A.C. Knaap, Department of Radiation Genetics and Chemical Mutagenesis of the State University of Leiden, The Netherlands (1981). This mouse lymphoma cell line was originally derived from the Fischer L5178Y line, isolated by Clive (1975)* *) Clive, D. and Spector, J.F.S., 1975, Laboratory procedure for assessing specific locus mutations at the TK locus in cultured L5178Y mouse lymphoma cells, Mutation Res., <u>31</u> , 17-29.

Stock cultures of these cells were stored in liquid nitrogen (-196°C). The cultures were checked for mycoplasma contamination.

CELL CULTURE

Cell culture conditions	L5178Y mouse lymphoma cells were cultured in F10 complete culture medium. Cultures were incubated in a humid atmosphere containing 5% CO ₂ in air at 37°C. Cell density was preferably kept below 7×10^5 cells/ml. Cells were exposed to the test substance in F10 complete culture medium without serum, buffered with 20 mM HEPES.
F10 complete culture medium	F10 complete culture medium consisted of Ham's F10 medium without thymidine and hypoxanthine (Gibco), supplemented with 10% horse serum, L-glutamine (2 mM) and penicillin/streptomycin (50 U/ml and 50 µg/ml respectively).
Selective medium	Selective medium consisted of F10 complete culture medium which contained, in addition 0.33% Agar Noble (Difco) and 5 µg/ml TFT (Sigma).
Cloning medium	Cloning medium consisted of F10 complete culture medium, supplemented with 10% horse serum, which was transformed into a gel by addition of 0.33% Agar Noble.

Environmental conditions

All incubations were carried out in a humid atmosphere (80-95%) containing 5% CO₂ in air in the dark at 37°C. The temperature and CO₂-percentage were monitored during the experiment.

REFERENCE SUBSTANCES

Negative control:

The vehicle of the test article.

Positive controls:Without metabolic activation (-S9-mix):

Ethylmethanesulphonate (EMS; CAS no. 62-50-0; purity 98%; Janssen Chimica) (2 mM) was used. EMS causes direct alkylation of DNA.

With metabolic activation (+S9-mix):

Dimethylnitrosamine (DMN; CAS-no. 62-75-9, purity 99%, Janssen Chimica) (0.5 mM) was used. DMN had to be activated by microsomal enzymes present in the S9-mix, resulting in a methyl diazonium ion which could react with cellular DNA.

Solvents for Reference Substances

Hank's balanced salt solution (HBSS) without calcium and magnesium.

Solutions of reference substances were prepared immediately before use.

METABOLIC ACTIVATION SYSTEM

Preparation of S9-homogenate*

Rat liver microsomal enzymes were routinely prepared from adult male Wistar or Sprague Dawley rats, which were obtained from BRL, Switzerland.

The animals were housed at NOTOX in a special room under standard laboratory conditions, as described in the SOP's. The rats were injected intraperitoneally with a solution (20% w/v) of Aroclor 1254 (500 mg/kg body weight) in corn oil. Five days later, they were killed by decapitation; (they were denied access to food for at least 12 hours preceding sacrifice). The livers of the rats were removed aseptically, and washed in cold (0°C) sterile 0.1 M sodium phosphate buffer (pH 7.4) containing 0.1 mM Na₂-EDTA. Subsequently the livers were minced in a blender and homogenized in 3 volumes of phosphate buffer with a Potter homogenizer. The homogenate was centrifuged for 15 min at 9000 g. The supernatant (S9) was transferred into sterile ampules, which were stored in liquid nitrogen (-196°C).

- *) Ames, B.N., Mc Cann, J. and Yamasaki, E., 1975, Methods for detecting carcinogens and mutagens with the Salmonella/mammalian microsome mutagenicity test. Mutation Res., 31, 347-364.

Preparation of S9-mix

S9-mix was prepared immediately before use and kept on ice during the test. S9-mix contained per ml: 1.02 mg $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$; 2.46 mg KCl; 1.7 mg glucose-6-phosphate; 3.4 mg NADP; 4 μmol HEPES and 0.5 ml S9. The above solutions were mixed and filter (0.22 μm)-sterilized (apart from the S9-fraction, which was added after filter-sterilization of the S9-mix components). Metabolic activation was achieved by adding 0.2 ml liver S9-mix to each ml of cell suspension.

EXPERIMENTAL PROCEDURE

Selection of Dose Levels/Cytotoxicity Test

Prior to the actual mutagenicity test cytotoxicity data were obtained by treating 6×10^6 cells, suspended in 6 ml of F-10 medium buffered with 20 mM HEPES, in the absence of serum in a sterile 30 ml centrifuge tube with a range of test substance concentrations in approximately half log steps, both in the absence and presence of S9-mix. The centrifuge tubes were rotated for 3 h on a roller mixer at 37°C.

After the exposure, the cells were separated from the treatment solutions by 3 centrifugation steps (115 g, 8 min), each followed by removal of the supernatant and resuspension of the cells, twice in Hank's balanced salt solution and finally in F-10 medium. The cells in the final suspension were counted in an "Improved Neubauer" haemocytometer. Relative cytotoxicity, expressed as the reduction after approximately 24 h and 48 h growth compared to nontreated control cells, was used to determine a suitable concentration range (4 doses) of the test substance to be applied to cultures prepared for mutagenicity testing. Because the test substance was not toxic and difficult to dissolve in aqueous solutions the highest concentration was determined by the solubility in the culture medium.

Cleansing

Prior to mutagenicity (and cytotoxicity) testing, the cells were grown for 1 day in culture medium containing 10^{-4} M hypoxanthine, 2×10^{-7} M aminopterin and 1.6×10^{-5} M thymidine (HAT-medium) to reduce the amount of spontaneous mutants, followed by a recovery period of 2 days on medium containing hypoxanthine and thymidine only. After this period cells were returned to normal medium at least for 1 day before starting the experiment.

Mutagenicity test*

The test substance was tested both with and without S9-mix in two independent experiments. 6×10^6 cells ($10^6/\text{ml}$), or 12×10^6 cells ($10^6/\text{ml}$) for test substance concentrations expected to be strongly toxic, for each selected dose cells were exposed for 3 h to the test substance in HEPES-buffered Ham's F-10 medium without serum. For this purpose the cell suspensions were placed in 30 ml centrifuge tubes on a roller mixer at 37°C . Solvent and positive controls were included. After exposure, cells were washed twice with HBSS, counted and seeded in F-10 culture medium for expression of the mutant phenotype. The cultures were subcultured at least every other day in order to maintain log phase growth. The expression period for TFT-resistant mutants was 3 days. Immediately after exposure to the test substance 3×200 cells of each dose were plated into P90 petri dishes containing 15 ml cloning medium to determine cell survival (cloning efficiency). The cell survival was counted after 10-14 days with the Artek colony counter or the naked eye.

- *) Clive, D., Johnson, K.O., Spector, J.F.S., Batson, A.G. and Brown, M.M.M., 1979, Validation and characterization of the L5178Y/TK Mouse lymphoma mutagen assay system, *Mutation Res.*, 59, 61-108.
- *) Amacher, D.E., Paillet, S.C., Turner, G.N., Ray, V.A. and Salsburg, D.S., 1980, Point mutations at the thymidine kinase locus in L5178Y mouse lymphoma cells. II. Test validation and interpretation, *Mutation Res.*, 72, 447-474.
- *) Jotz, M. and Mitchell, A.D., 1981, Effects of 20 coded chemicals on the forward mutation frequency at the thymidine kinase locus in L5178Y mouse lymphoma cells. In: *Evaluation of short-term tests of carcinogens*. F.J. de Serres and J. Ashby (Eds.), Elsevier-North Holland.
- *) Van der Hoeven, J.C.M., Bruggeman, I.M. and Debets, F.M.H., 1984, Genotoxicity of quercetin in cultured mammalian cells, *Mutation Res.*, 136, 9-21.
- *) Clive, D., Caspary, W., Kirby, P.E., Krehl, R., Moore, M., Mayo, J. and Oberly, T.J., 1987, Guide for performing the mouse lymphoma assay for mammalian cell mutagenicity, *Mutation Res.*, 189, 143-156.

Mutant selection

After the expression period a total number of 1.5×10^6 cells were plated in ten P90 petri dishes, each containing 15 ml selective medium (TFT-selection). TK-deficient mutants formed microcolonies in 10-14 days, which were counted with the naked eye. Cloning efficiencies at the time of mutant selection were determined as described above. The mutant frequency was expressed as the number of mutants per 10^5 surviving cells.

ACCEPTABILITY OF ASSAY

A mutation assay was considered acceptable if it met the following criteria:

- a) The absolute cloning efficiency of the solvent controls was $\geq 50\%$.
- b) At least three of the four doses of the test substance had an acceptable number of surviving cells (10^6) analysed for expression of the TK mutation.
- c) The spontaneous mutant frequency in the untreated or solvent control was < 5 per 10^5 clonable cells.
- d) The positive controls (ethylmethanesulfonate and dimethylnitrosamine) induced significant (at least 2-fold) increases in the mutant frequencies.

DATA EVALUATION AND STATISTICAL PROCEDURES

No formal hypothesis testing was done.

A test substance was considered positive (mutagenic) in the mutation assay if:

- a) It induced at least a 2-fold increase in the mutant frequency compared to the solvent control in a dose-dependent manner; and
- b) The results were reproducible in an independently repeated test.

A test substance was considered negative (not mutagenic) in the mutation assay if:

- a) None of the tested concentrations showed a mutant frequency at least twice that of the solvent control.
- b) The results were confirmed in an independently repeated test.

The preceding criteria were not absolute and other modifying factors might enter into the final evaluation decision.

RESULTS

CYTOTOXICITY TEST/DOSAGE SELECTION

Preliminary solubility tests indicated that T-5869 precipitated in the exposition medium at concentrations of 100 µg/ml and upwards. Table 1 shows the results of the preliminary cytotoxicity test with T-5869 in the presence and absence of a metabolic activation system (S9-mix). In the absence of S9-mix T-5869 inhibited the growth of the lymphoma cells in suspension at a test substance concentration of 33 µg/ml by 22% and at 100 µg/ml by 72%. In the presence of S9-mix the test substance did not inhibit the growth of the lymphoma cells in suspension.

MUTAGENICITY TEST

Based on the results of the cytotoxicity test (Table 1), the following dose range was selected for mutagenicity testing:

Without S9-mix: 3 to 100 µg/ml culture medium

With S9-mix : 3 to 100 µg/ml culture medium

EMS (2 mM) was used as a positive control in the test without metabolic activation, whereas DMN, (0.5 mM) served as a positive control for the assay with metabolic activation. Tables 2 and 3 show the percentages of cell survival and the mutant frequencies for various concentrations of T-5869. Individual colony counts of cloning and selective plates, and cell counts during subculturing are listed in Tables 4-9 of the appendix.

Both in the presence and absence of S9-mix the test substance induced no significant increase in the mutant frequency in both independent experiments.

The spontaneous mutant frequencies in the solvent-treated control cultures were within our historical control data range (1.9 ± 1.2 in the absence of S9-mix and 1.4 ± 0.7 in the presence of S9-mix; indicated are means $\pm$ S.D. for n=44 and 45 respectively).

Mutant frequencies induced by positive control chemicals were increased by 7- to 9-fold for EMS and by 21- to 23-fold for DMN. It was therefore concluded that the test conditions were optimal and that the metabolic activation system (S9-mix) functioned properly.

CONCLUSION

In conclusion, T-5869 was found to be not mutagenic in the TK mutation test system under the experimental conditions described in this report.

TABLE 1 PRELIMINARY CYTOTOXICITY DETERMINATION IN L5178Y MOUSE LYMPHOMA CELLS

Dose ($\mu\text{g/ml}$)	Cells/ml ($\times 10^5$)			Suspension growth	
	After 0 h	After 24 h	After 48 h	Total ¹	% of control
Without metabolic activation (-S9-mix)					
Solvent control	6.1	4.9	6.1	71.2	100
1	6.4	5.1	6.5	82.9	116
3	7.4	4.5	5.9	76.7	108
10	5.7	5.3	5.6	66.1	93
33	4.9	4.7	6.2	55.8	78
100 ²	5.0	2.5	4.1	20.0	28
With metabolic activation (+S9 mix)					
Solvent control	7.9	5.7	6.8	119.6	100
1	6.6	6.3	5.7	92.6	77
3	7.0	6.4	6.0	105.0	88
10	6.4	7.7	5.8	111.7	93
33	6.3	6.9	6.2	105.3	88
100 ²	6.6	6.8	6.4	112.2	94

Solvent control = DMSO

Subculture = 1.6×10^5 cells/ml
¹ Total growth = Cell count after 0 h $\times$ $\frac{(24 \text{ h cells/ml})}{\text{cells subcultured directly after treatment}}$ $\times$ $\frac{(48 \text{ h cells/ml})}{\text{cells subcultured after 24 h}}$
² Test substance precipitated slightly in the exposition medium

TABLE 2 CYTOTOXIC AND MUTAGENIC RESPONSE OF T-5869 IN THE MOUSE LYMPHOMA L5178Y TEST SYSTEM

EXPERIMENT 1

DOSE ($\mu\text{g}/\text{ml}$)	C.E. AT DAY 0 (% OF CONTROL)	C.E. AT DAY 3 (ABSOLUTE %)	MEAN NO. OF MUTANTS PER PLATE	MUTATION FREQUENCY $\times 10^5$
Without metabolic activation (-S9-mix)				
Solvent control	100	84	2.0	1.6
3	98	77	1.0	0.9
10	101	80	1.2	1.0
33	82	89	1.8	1.3
100 ¹	89	90	1.5	1.1
2 mM EMS	104	79	14.1	11.9
With metabolic activation (+S9-mix)				
Solvent control	100	88	1.0	0.8
3	86	82	1.0	0.8
10	76	84	0.8	0.6
33	96	87	1.1	0.8
100 ¹	80	88	1.9	1.4
0.5 mM DMN	44	49	13.6	18.5

C.E. = Cloning Efficiency

Solvent control = DMSO

EMS = Ethylmethanesulphonate

DMN = Dimethylnitrosamine

¹ Test substance precipitated slightly in the exposition medium

TABLE 3 CYTOTOXIC AND MUTAGENIC RESPONSE OF T-5869 IN THE MOUSE LYMPHOMA L5178Y TEST SYSTEM

EXPERIMENT 2

DOSE ($\mu\text{g/ml}$)	C.E. AT DAY 0 (% OF CONTROL)	C.E. AT DAY 3 (ABSOLUTE %)	MEAN NO. OF MUTANTS PER PLATE	MUTATION FREQUENCY $\times 10^5$
Without metabolic activation (-S9-mix)				
Solvent control	100	86	1.2	0.9
3	111	88	1.1	0.8
10	102	79	0.6	0.5
33	58	88	2.4	1.8
100 ¹	44	95	2.9	2.0
2 mM EMS	83	81	9.3	7.7
With metabolic activation (+S9-mix)				
Solvent control	100	87	0.7	0.5
3	95	81	0.6	0.5
10	75	88	0.9	0.7
33	91	81	0.6	0.5
100 ¹	82	75	0.7	0.6
0.5 mM DMN	33	49	7.9	10.7

C.E. = Cloning Efficiency

Solvent control = DMSO

EMS = Ethylmethanesulphonate

DMN = Dimethylnitrosamine

¹ Test substance precipitated slightly in the exposition medium

APPENDIX

Individual Colony Counts and Cell Counts during Expression Period

FORMULAS AND CALCULATIONS FOR L5178Y MOUSE LYMPHOMA TEST SYSTEM.

$$\text{Mutant frequency per } 10^5 \text{ survivors} = \frac{100}{\text{cloning efficiency}} \times \frac{\text{Total number of mutant colonies on selective plates}}{\text{number of seeded cells}}$$

$$\text{Cloning efficiency} = \frac{\text{Average No. of colonies on cloning plates} \times 100\%}{200}$$

L5178Y MOUSE LYMPHOMA TEST SYSTEM

- Cell counts during expression period
- Cloning efficiency immediately after exposure
- Mutation experiments, individual colony counts

Tables 4-6 Experiment 1

Tables 7-9 Experiment 2

Abbreviations used: DMN, dimethylnitrosamine
EMS, ethylmethanesulphonate

Solvent control: Dimethylsulphoxide

TABLE 4 L5178Y MOUSE LYMPHOMA TEST SYSTEM - CELL COUNTS AND SUBCULTURE DATA

Experiment 1

dose ($\mu\text{g/ml}$)	DAY 0				DAY 2				DAY 3	
	Total amount of	Total amount of	Subculture		Subculture		Subculture		Subculture	
	cells before (treatment $\times 10^6$)	cells after (treatment $\times 10^6$)	$\times 10^6$ ($\%1$)	Total amount ³⁾	Cell count ($\text{c/ml} \times 10^5$)	$\times 10^6$ ($\%2$)	Total amount ⁴⁾	Cell count ($\text{c/ml} \times 10^5$)	$\times 10^6$ ($\%2$)	
Without Metabolic Activation (-S9-mix)										
DMSO	6	5.2	87	3.0	6.6	100	4.0	5.8	100	
3	6	3.9	65	3.0	6.5	98	4.0	6.5	112	
10	6	5.5	92	3.0	5.9	89	4.0	6.0	103	
33	6	4.5	75	3.0	5.2	79	4.0	5.9	102	
100	12	6.6	55	6.0	5.2	79	4.0	6.0	103	
EMS	6	4.1	68	3.0	6.3	95	4.0	6.2	107	
With Metabolic Activation (+S9-mix)										
DMSO	6	4.8	80	3.0	6.6	100	4.0	5.9	100	
3	6	4.7	78	3.0	7.1	108	4.0	5.9	100	
10	6	4.7	78	3.0	5.7	86	4.0	6.4	108	
33	6	3.6	60	3.0	6.9	105	4.0	5.5	93	
100	6	4.5	75	3.0	6.1	92	4.0	5.4	92	
DMN	6	5.4	90	3.0	2.4	36	4.0	4.8	81	

(1) $\frac{\text{cells after treatment}}{\text{cells before treatment}} \times 100\%$

(2) $\frac{\text{cell count}}{\text{cell count of control}} \times 100\%$

(3) cell density 0.4×10^5 c/ml

(4) cell density 1.6×10^5 c/ml

TABLE 5 L5178Y MOUSE LYMPHOMA TEST SYSTEM - CLONING EFFICIENCY DAY 0

EXPERIMENT 1

Dose (μ g/ml)	No. of colonies/ cloning plate			Mean No. of colonies/plate	Cloning efficiency	
	1	2	3		absolute	relative (% of control)
Without S9-mix						
Solvent control	154 ;	151 ;	169	158	79	100
3	161 ;	149 ;	INF	155	78	98
10	143 ;	174 ;	159	159	80	101
33	122 ;	135 ;	INF	129	65	82
100	167 ;	129 ;	126	141	71	89
EMS	163 ;	165 ;	INF	164	82	104

With S9-mix						
Solvent control	187 ;	188 ;	169	181	91	100
3	145 ;	167 ;	INF	156	78	86
10	125 ;	140 ;	145	137	69	76
33	163 ;	167 ;	188	173	87	96
100	148 ;	147 ;	138	144	72	80
DMN	80 ;	76 ;	84	80	40	44

INF = Plate infected with fungi

TABLE 6 L5178Y MOUSE LYMPHOMA TEST SYSTEM - SELECTION DATA AND CLONING EFFICIENCY

EXPERIMENT 1

Dose µg/ml	Number of colonies/Selection plate										Total No.	No. of colonies /cloning plate			Mean	MF
	1	2	3	4	5	6	7	8	9	10		1	2	3		
Without Metabolic Activation (-S9-mix)																
Solvent control	0	2	4	2	2	2	2	3	2	1	20	161	173	171	168	1.6
3	0	3	0	0	1	1	1	1	2	1	10	162	153	148	154	0.9
10	2	0	1	2	0	1	3	1	0	2	12	161	159	156	159	1.0
33	2	1	0	4	2	1	0	4	1	3	18	177	177	181	178	1.3
100	1	0	0	2	2	1	1	3	3	2	15	174	188	177	180	1.1
EMS	22	14	21	16	13	14	9	14	8	10	141	156	161	157	158	11.9
With Metabolic Activation (+S9-mix)																
Solvent control	1	0	0	1	3	0	2	1	1	1	10	177	181	170	176	0.8
3	0	1	0	0	2	0	2	1	2	2	10	162	157	170	163	0.8
10	0	1	0	2	0	1	0	2	1	1	8	163	171	INF	167	0.6
33	3	0	1	2	2	0	0	0	0	3	11	182	167	169	173	0.8
100	1	1	2	3	2	0	4	2	2	2	19	171	178	179	176	1.4
DMN	6	13	15	15	16	14	18	13	12	14	136	94	94	105	98	18.5

MF = Mutant frequency per 10^5 survivors
 INF = Plate infected with fungi

TABLE 7 L5178Y MOUSE LYMPHOMA TEST SYSTEM - CELL COUNTS AND SUBCULTURE DATA

Experiment 2

dose ($\mu\text{g/ml}$)	DAY 0				DAY 2				DAY 3	
	Total amount of	Total amount of	Subculture		Subculture		Subculture		Subculture	
	cells before treatment $\times 10^6$	cells after treatment $\times 10^6$	% ¹⁾	$\times 10^6$ Total amount ³⁾	Cell count $\times 10^5$	% ²⁾	$\times 10^6$ Total amount ⁴⁾	Cell count $\times 10^5$	% ²⁾	
Without Metabolic Activation (-S9-mix)										
DMSO	6	4.3	72	3.0	6.3	100	4.0	5.9	100	
3	6	3.9	65	3.0	6.4	102	4.0	5.3	90	
10	6	3.9	65	3.0	5.2	83	4.0	5.7	97	
33	6	3.1	52	2.9	3.4	54	4.0	5.5	93	
100	12	3.6	30	3.5	2.0	32	4.0	4.8	81	
EMS	6	3.8	63	3.0	5.9	94	4.0	4.7	80	
With Metabolic Activation (+S9-mix)										
DMSO	6	3.8	63	3.0	8.4	100	4.0	4.3	100	
3	6	3.9	65	3.0	8.7	104	4.0	5.2	121	
10	6	4.2	70	3.0	7.5	89	4.0	5.0	116	
33	6	3.8	63	3.0	6.1	73	4.0	5.6	130	
100	6	3.7	62	3.0	7.4	88	4.0	5.1	119	
DMN	6	4.4	73	3.0	3.2	38	4.0	3.5	81	

(1) $\frac{\text{cells after treatment}}{\text{cells before treatment}} \times 100\%$

(2) $\frac{\text{cell count}}{\text{cell count of control}} \times 100\%$

(3) cell density 0.4×10^5 c/ml

(4) cell density 1.6×10^5 c/ml

TABLE 8 L5178Y MOUSE LYMPHOMA TEST SYSTEM - CLONING EFFICIENCY DAY 0

EXPERIMENT 2

Dose (μ g/ml)	No. of colonies/ cloning plate			Mean No. of colonies/plate	Cloning efficiency	
	1	2	3		absolute	relative (% of control)
Without S9-mix						
Solvent control	130 ;	140 ;	129	133	67	100
3	135 ;	154 ;	154	148	74	111
10	140 ;	121 ;	145	135	68	102
33	70 ;	80 ;	82	77	39	58
100	51 ;	60 ;	62	58	29	44
EMS	112 ;	101 ;	116	110	55	83

With S9-mix						
Solvent control	204 ;	180 ;	198	194	97	100
3	192 ;	170 ;	191	184	92	95
10	152 ;	138 ;	149	146	73	75
33	177 ;	179 ;	174	177	89	91
100	159 ;	162 ;	155	159	80	82
DMN	69 ;	67 ;	56	64	32	33

TABLE 9 L5178Y MOUSE LYMPHOMA TEST SYSTEM - SELECTION DATA AND CLONING EFFICIENCY

EXPERIMENT 2

Dose µg/ml	Number of colonies/Selection plate										Total No.	No. of colonies /cloning plate			Mean	MF

	1	2	3	4	5	6	7	8	9	10		1	2	3		
Without Metabolic Activation (-S9-mix)																
Solvent control	0	0	1	4	1	1	0	1	2	2	12	161	179	173	171	0.9
3	0	0	0	2	0	2	4	2	1	0	11	183	170	172	175	0.8
10	0	0	0	0	0	0	2	0	2	2	6	159	149	165	158	0.5
33	0	4	4	4	3	1	1	4	2	1	24	170	185	171	175	1.8
100	1	4	2	4	6	2	3	3	2	2	29	192	199	178	190	2.0
EMS	8	5	5	9	8	6	10	13	18	11	93	157	156	173	162	7.7
With Metabolic Activation (+S9-mix)																
Solvent control	3	0	0	0	2	0	1	0	0	1	7	175	181	162	173	0.5
3	0	0	3	1	0	0	0	2	0	0	6	161	159	167	162	0.5
10	2	0	0	2	0	2	2	1	0	0	9	180	171	175	175	0.7
33	0	0	0	0	1	0	2	0	1	2	6	164	163	160	162	0.5
100	2	0	1	0	1	1	0	1	1	0	7	152	136	160	149	0.6
DMN	6	8	8	9	6	6	8	9	11	8	79	96	91	106	98	10.7

MF = Mutant frequency per 10⁵ survivors