

55

87

TRANSACTIONS
of the
McINTYRE-SARANAC CONFERENCE
on
OCCUPATIONAL CHEST DISEASE

Edited by G. W. H. Schepers, M.D., D.Sc.

Tenney Hall, Saranac Lake, N. Y.
Feb. 7, 8, and 9, 1955

AMERICAN MEDICAL ASSOCIATION PRESS
Chicago 10, Illinois
1955

REFERENCE

Swartz, H. H. Jr.: *Silicosis: Impregnation of* 12:545, 1939.

W. R. La and Wentworth, *et al.* Pneumoconiosis, J. Exp. Med. 1, 1953.

Irwin, D. A. and King, *et al.* Silicosis: Quarries, Sericite and L. M. A. J. 31:237, 1934.

W. R. La and Wentworth, *et al.* Epidemiology of L. M. A. J. 31:237, 1934.

The Dust Diseases in Great Britain

A. I. G. McLAUGHLIN, M.D., F.R.C.P., London

The story of dust diseases in Great Britain goes back a long way. E. L. Collis, in his classical Milroy Lectures of 1915, quoted Herbert Spencer as saying that the starting point of human progress was the "localization of industries." There was such a localization in prehistoric factories for the making of flint implements at Grime's Graves, near Brandon in Suffolk, where the flint knappers still use tools like the deer-horn picks of their prehistoric ancestors. In 1914, Collis showed that these workers have a high mortality from silicosis. He said that it is probable that "the starting point of human progress" was associated with at least one form of pneumoconiosis. It has been long known that the dust of flint, which is nearly pure SiO_2 , is dangerous to health. Over 200 years ago, Thomas Benson, of Newcastle under Lyme, was granted a patent for grinding flints by a wet method. At that time it was said that a man who ground the flints dry could not live longer than two years.

Our greatest localization of industries took place during the Industrial Revolution, and this was really the starting point of our intensive knowledge about the effects of dust on the lung.

The first notable contribution was that of Pearson (1813), who, after many autopsies, decided that the black pigment in the bronchial nodes and the lungs was due to the inhalation of small particles resulting from the burning of coal, wood, and other inflammable materials. The German pathologists, on the other hand, thought that the pigment came from inside the body. Meiklejohn (1951) has described the hitherto little-known and re-

markable contributions of the Scottish physicians of the 1800's on the connection between disease of the lungs and coal dust; he also tells how the incidence of anthracosis diminished with improved ventilation of the mines. The Scottish physicians decided that simple coal dust was comparatively harmless and that stone dust was the really noxious factor.

Charles Turner Thackrah, a physician in Leeds, played a major part in the study of occupational diseases (including the dust diseases) in Great Britain. In his book "The Effects of Arts, Trades, and Professions," the second edition of which was published in 1832, he crystallized the idea that the inhalation of large quantities of dusts of any kind can damage the lungs but that some dusts are more harmful than others. His conclusions were based on first-hand observations of workers in hospitals, in their homes, and in the factories.

Two other great Englishmen in the history of pneumoconiosis about the middle of the 19th century were T. B. Peacock and E. M. Greenhow. Peacock first established miners' disease as an entity and distinguished it clinically from pulmonary tuberculosis before little was known about bacteriology and nothing about x-rays. Greenhow carried out the first large field investigations into the dusty industries of England and Wales, including the heavy-metal industries, the potteries, coal, copper, and lead mining, and even agriculture. In the *Transactions of the Pathological Society of London* (1860-1866) there are to be found excellent descriptions by both of these physicians of the disease, which was later to be called silicosis by Visconti, in 1870. They even found the dust of free silica in the lungs and examined it under polarized light.

For a long time after this excellent work nothing much was done about the dust diseases, but about the beginning of the 20th

Recorded for publication April 7, 1955.
H. M. Medical Inspector of Factories.

century a new interest began to be taken in the problem not only in England but also in other parts of the world. About the same time the tempo of life in general began to increase, and there was an urge for increased speed of production—an urge which has gradually gathered momentum. With the replacement of hand labor by the machine, dusty processes have become more dusty and methods of dust control have lagged behind output. The result has been a remarkable increase of the incidence of the dust diseases.

A noteworthy contribution in 1892 was Artidge's book on the occupational diseases, which was only the second to be published in England.

In 1904, J. S. Haldane and his colleagues ascribed the high mortality among tin miners to the inhalation of rock dust. A little earlier, dust phthisis among slate workers was shown to be related to the dust of slate, particularly when it had a high quartz content.

E. L. Collis (1915) pinpointed the dust of free silica as the main cause of most of the dust diseases. He also drew attention to the role that dust inhalation plays in determining the mortality from lung diseases experienced by the general population. Long ago he showed the influence of air pollution on the mortality from lung diseases in general. Though Collis did this work 40 years ago, he is always up-to-date. He had, and still has, an uncanny knack of seeing to the heart of any problem. Just now in England there is controversy about chronic bronchitis and emphysema and whether they can be held to be caused by the inhalation of dust. It is true that the incidence of chronic bronchitis and emphysema is high among the general population, and the possible causes are legion. It is difficult to decide (in Artidge's words) how much of the malady is "town-made" or "trade-made." But as Collis showed years ago, an agent which irritates the lung parenchyma can also irritate the bronchial mucous membrane. At present there is one industrial pulmonary disease of which the diagnosis is made largely on the presence of chronic bronchitis and emphysema, and that is byssinosis.

84

In the last 50 years in Great Britain (as elsewhere), there has been intensive research into all aspects of the dust diseases, such as causation, pathology, incidence in various industries, and prevention. For the first 25 years or so, much of the impetus came from H. M. Medical Inspectors of Factories, and prominent in the study of the chest diseases were Collis, Middleton, and Merewether. It might be said that they played almost a lone hand, with meager facilities for investigation. However, they were helped greatly by hospital physicians, general practitioners, radiologists, and pathologists, notably Hall, Robertshaw, Kettle, Gloyne, and Cooke. During this period much attention was paid to silicosis, and it, with tuberculosis, was established as being the cause of disability and death in most of the industries where there is exposure to the dust of free silica. *Pari passu*, regulations laying down dust-control measures were issued by the Factory Department for most of the major industries. Notable contributions were Collis's work, already mentioned, on the role of free silica and Middleton's field investigation (with E. L. Macklin) into the grinding industry in 1923. In the late '20's, a series of brilliant observations by Cooke, Gloyne, Burton Wood, Stuart McDonald, and M. J. Stewart on the clinical, radiological, and pathological features of a new disease, which Cooke named asbestosis, led to field investigations by Merewether and Price (1930). These firmly established that the dust of asbestos was dangerous to health, and it was the first time that the dust of combined silica, as opposed to free silica, was found to damage the lungs. The condition, as Legge points out, was first seen in 1906 by Murray at Charing Cross Hospital, but little or no notice of it was taken at that time.

An important landmark was the passing of the Workmen's Compensation (Silicosis) Act of 1918, which came into force in 1919. For the first time compensation for disablement or death was given in respect of an industrial pulmonary disease. The Act provided initial and periodical medical examination of all workers in the refractories industry

and suspension from work on the basis of diagnosis of simple silicosis in association with tuberculosis. The scheme was responsible for the examination of tuberculosis Officers and the administering sanatorium. The National Health Insurance Act of 1925 was revised in 1925 to provide for examination of workers in the silicosis and asbestosis industries. This was the beginning of the National Health Insurance (Silicosis and Asbestosis) Scheme, its first (and only) C. L. Sutherland.

A close association between the Medical Inspectorate and the industries working under the scheme, the Medical Inspectorate, and field investigations which led to the various compensation schemes for members of the Board of Trade, Manchester, Stoke-on-Trent, and other industries, such as the pottery industry (Bryson, 1929), glass (Bryson, 1929), and paper (Bryson, 1949). But perhaps the most important result of the formation of the scheme was the collection of case histories together with occupational and mortuary findings.

At first the Board of Trade was responsible for the scheme, but gradually it came under examination of the various industries first compensation scheme in 1927, and sandstone was the first industry to be included. The scheme was amended in 1927, and sandstone was the first industry to be included. The scheme was amended in 1927, and sandstone was the first industry to be included. The scheme was amended in 1927, and sandstone was the first industry to be included.

Great Britain (as intensive research diseases, such as dence in various in- For the first 25 impetu- came from re of Factories, and if the chest diseases and Merewether. It played almost a lone ies for investigation. ed greatly by hospi- ractitioners, radiolo- otably Hall, Robert- d Cooke. During this was paid to silicosis, s, was established as ity and death in most there is exposure to *ari fossa*, regulations of measures were is- partment for most of Notable contributions idy mentioned, on the Middleton's field in- .. Macklin) into the '23. In the late '20's, ervations by Cooke, Stuart McDonald, and clinical, radiological, res of a new disease, pestosis, led to field in- ither and Price (1930). ed that the dust of ar- to health, and it was list of combined silica, t, was found to damage m, as Legge points out, by Murray at Claring title or no notice of it

mark was the passing of pensation (Silicosis) me into force in 1919. pensation for disable- ven in respect of an in- lisease. The Act pro- dical medical examina- the refractories industry

and suspension from the industry on a diag- nosis of simple silicosis as well as silicosis with tuberculosis. The medical officers re- sponsible for the examinations were the Tu- berculosis Officers of the Local Authority ad- ministering sanatorium benefit under the National Health Insurance Act. The scheme was revised in 1925, and again in 1931, to provide for examinations by a medical board. This was the beginning of the whole-time Silicosis and Asbestosis Medical Board under its first (and only) Chief Medical Officer, C. L. Sutherland.

A close association began between the Medical Inspectorate of Factories and the Silicosis and Asbestosis Medical Board, both groups working under the Home Office. In general, the Medical Inspectorate made the field investigations which established the risk of silicosis (or asbestosis) in an industry or process and which led to the establishment of the various compensation schemes. But the members of the Board (stationed at Sheffield, Manchester, Stoke on Trent, and Cardiff) also carried out field investigations into vari- ous industries, such as sandstone (Sutherland and Bryson, 1929), granite (Sutherland and Bryson, 1929), and potteries (Meiklejohn, 1949). But perhaps the most significant re- sult of the formation of the Board was that there was begun a systematic and valuable collection of case histories with x-ray films, together with occupational details and post- mortem findings.

At first the Board was limited to a few in- dustries, but gradually its scope has increased and more and more dusty industries have come under examination. Workers in the re- fractories industry first came under a com- pensation scheme in 1925, metal grinders in 1927, and sandstone workers in 1929. In 1928, the Various Industries Scheme included the processes of mining, quarrying, drilling, and blasting in silica rock, the crushing and grind- ing of siliceous materials, and also isolated processes in steel foundries (not iron found- ries) and metal works, in potteries, and in tin mines. In 1934 the Various Industries Scheme was amended to include coal miners

with silicosis; in 1935 hematite ore miners were included, and in 1939 slate miners. Workers with asbestosis were first compen- sated in 1931; compensation for byssinosis was introduced in 1941 and was dealt with by a specially constituted board. Pneu- moconiosis, as opposed to classical silicosis, in coal miners was brought under the compensa- tion schemes in 1943 and in coal trimmers in 1946. The Various Industries Scheme was again amended in 1946 to include molders of iron castings who used siliceous parting pow- ders and also blasters of any type of metal castings to free them from adherent sand, even if the blasting abrasive was non-siliceous. The National Insurance (Industrial Injuries) Act was passed in 1946 and altered the whole basis of compensation in that it became the re- sponsibility of the Government but with con- tributions from employers and workers. Ber- yllium "poisoning" became compensable in 1949. In 1954 all foundry workers (iron, steel, and nonferrous) became entitled to com- pensation under the Industrial Injuries Act. A fuller account of the development of com- pensation in Great Britain has recently been written by Meiklejohn (1954). The Silicosis and Asbestosis Medical Board was trans- ferred from the Home Office in 1946 to the Ministry of National Insurance (later com- bined with the Ministry of Pensions). The Board was divided into a series of Pneu- moconiosis Panels, but they work in much same way as the Board did, except that there is now no Chief Medical Officer.

The various compensation schemes men- tioned above reflect the evolution of the study of the industrial lung diseases in Great Brit- ain. There are comparable legal enactments and regulations designed to control the risks in each industry, but they are too numerous to mention in detail.

To "flash back" a little, in 1936 Middleton in his Milroy Lectures, reviewed the position of the dust diseases in Great Britain. He covered all the dusty trades, including coal mining, but at that time only cases of silicosis in coal miners were recognized as being eligi- ble for compensation. Most of the miners,

such as handlayers, drifters, or rippers, had been exposed to fine dust, but Middleton also showed that there was a high incidence of a lung disease among coal miners, particularly in South Wales, which was not classical silicosis. Much of his evidence was drawn from the findings of the Silicosis and Asbestosis Medical Board. Following his paper, there began a series of intensive investigations by the Medical Research Council into the health risks of coal mining, which covered all aspects, such as clinical, radiographic, and environmental (Medical Research Council Special Reports I [1913], II [1943], and III [1945]). The important fact emerging from this work was that coal dust by itself could cause pneumoconiosis, and this was followed, as previously mentioned, by the extension of compensation to coal miners and coal trimmers. In 1915, the Pneumoconiosis Research Unit of the Medical Research Council was set up in Cardiff, and an impressive body of work has emerged on various aspects of coal miners' pneumoconiosis. This work is well known, and I propose, therefore, to limit my remaining observations mainly to the dust diseases resulting from work in places which come under the Factories Acts, with only incidental references to coal mining.

SILICOSIS

Silicosis has been found to be a cause of disability in the following broad groups of industries as well as in coal mining and other forms of mining, such as tin, hematite, copper, barites, and fire clay:

1. Refractories industry. This group includes the making of silica bricks, furnace dismantling and rebuilding, and retort setting and repairing.

2. Pottery industry. Workers in this group have been engaged in the manufacture of both earthenware and china with their various sub divisions and as flint millers and polishers. The substitution of ground flint by alumina for the glazing of bone ware has resulted in a diminution of deaths from silicosis among workers in that process.

3. Stone quarrying, crushing, and dressing. This group includes workers with sandstone, millstone, gneiss, slate, granite, and other igneous rocks. The risk varies with the amount of free silica in the rock and, of course, with the protective measures adopted. Recently some limestone quarrymen have contracted silicosis, but the limestone contained a fairly high proportion of free silica.* In this group could be included rock tunnelers, in whom the onset of silicosis may be very rapid. As regards slate workers, a billiard-table maker has recently received compensation for silicosis.




Fig. 1.—Chart showing diminishing numbers of deaths among metal grinders during the period 1930-1951, inclusive.

4. Metal grinding. The silicosis risk in the grinding of metals has greatly diminished, owing to the replacement of the sandstone grinding wheels with artificial ones composed of Carborundum, alumina, or emery. Apart from the lower "toxicity" of the dusts from these substances, the wheels are much harder than sandstone, and less dust is created. But the grinders of castings (iron, steel, and non-ferrous) are still exposed to a silicosis risk on account of the presence of burned-on sand on the castings. The diminishing number of deaths among grinders is shown in the chart (Fig. 1).

* Doig, A. T.: Unpublished data.

DUST DISEASES

5. Sandblasting. (1936) showed, had position has altered, began to be replaced by abrasives, such as steel shot, sand as an abrasive in the blasting of Castings Regulations. The painting of ships' hulls does not (because legal), but it is thought of silicosis in the job used in the open air construction, e. g., in metal surfaces for the


6. Manufacture of abrasives. In 1930 the abrasives were carried or subacute silicosis, not silicosis. There was a factory in London 1928, when there were workers. This experience control in the process, and 1952 there were none among this type of work to state the incidence of because there is a top (mostly young women) flour is still being used. Efforts are being made to substitute which will manufacturers and the new cases of silicosis in

7. Foundry industry. Foundry industry has great detail during the and the results were put form by McLaughlin in Lung Diseases of Iron Workers"). Pathological as well as clinical and some 3000 workers, showing varying risk of silicosis, pneumoconiosis (to which low) in the various workers.

Steel fettlers (dressers) had a severe risk.

...ing, and dressing.
...s with sand, long,
...e, granite, and other
...isk varies with the
...in the rock and, of
...ve measures adopted,
...ne quarrymen have
...the limestone con-
...portion of free silica,*
...included rock tunnel-
...t of silicosis may be
...slate workers, a bil-
...cently received com-

DEATHS IN METAL GRINDERS
1940-1951



* diminishing numbers of
grinders during the period

The silicosis risk in
has greatly diminished.
nent of the sandstone
ificial ones composed
nina, or emery. Apart
ity" of the dusts from
heels are much harder
ss dust is created. But
s (iron, steel, and non-
sed to a silicosis risk on
e of burned-on sand on
diminishing number of
s is shown in the chart

lished data.

5. Sandblasting. This job, as Merewether (1950) showed, had a high silicosis risk. The position has altered materially after sand began to be replaced by other nonsiliceous abrasives, such as steel shot, and since the use of sand as an abrasive was prohibited in 1949 by the Blasting of Castings (and other Articles) Regulations. The process of wet sandblasting of ships' hulls does not come under the Regulations (because legally a ship is not an article), but it is thought that there is also a risk of silicosis in the job. Sandblasting is still used in the open air on works of engineering construction, e. g., in order to prepare large metal surfaces for the application of paint.

6. Manufacture of abrasive soap. Middleton showed in 1936 that the manufacture of abrasive soaps carried with it a risk of acute or subacute silicosis, accompanied by tuberculosis. There was a disastrous experience at one factory in London between 1921 and 1928, when there were 13 deaths among 81 workers. This experience led to rigorous dust control in the processes, but between 1941 and 1952 there were nine deaths from silicosis among this type of worker. It is not possible to state the incidence of disease and disability, because there is a rapid turnover of labor (mostly young women) in the job. Silica flour is still being used as an abrasive, though efforts are being made to find a less harmful substitute which will be acceptable to the manufacturers and the users. There were no new cases of silicosis in this trade in 1953.

7. Foundry industry. The iron and steel foundry industry has been investigated in great detail during the past 10 years or so, and the results were published in 1950 in book form by McLaughlin and others ("Industrial Lung Diseases of Iron and Steel Foundry Workers"). Pathological studies of 64 cases, as well as clinical and x-ray examinations of some 3000 workers, showed that there was a varying risk of silicosis and mixed dust pneumoconiosis (to which reference is made below) in the various categories of foundry workers.

Steel fettlers (dressers or castings cleaners) had a severe risk, largely owing to the

fact that the dust from the pneumatic hammer is not controlled. The main pathological lesion was classical silicosis, with or without tuberculosis. Among iron fettlers, the risk was not so great as among steel fettlers, and the main pathological lesion found was mixed dust pneumoconiosis, though classical silicosis did occur. Steel molders were less affected by disease and disability than iron molders, though statistically the incidence of x-ray abnormalities was greater among steel workers. More deaths occurred among iron molders, and this fact appeared to be related to the use of siliceous parting powders. The use of siliceous parting powders has now been prohibited by the Foundries (Parting Materials) Regulations, 1950. Silicosis still occurs among shot blasters of both iron and steel castings, but the risk is controlled by ventilation and personal protective devices, and the numbers of cases and deaths, in contrast with the experience among steel fettlers, are not increasing. Other categories of iron and steel foundry workers are subject to much less risk than the above groups of workers. New and stringent regulations (Iron and Steel Foundry Regulations) were issued in 1955. They require strict dust control in the dusty processes.

The question of pneumoconiosis among nonferrous foundry workers is now being examined in detail. Isolated cases of silicosis and mixed dust pneumoconiosis have occurred among both casters and dressers of nonferrous metals, but an extensive survey has not yet been carried out. Harding and McLaughlin† have described the pathological, clinical, radiological, and environmental details of six fatal cases.

Table 1 shows the number of new cases of pneumoconiosis (i. e., silicosis and mixed dust pneumoconiosis) in foundry workers diagnosed by the Pneumoconiosis Panels in 1953.

These figures can be regarded as an understatement of the real position. More than half

† Harding, H. E., and McLaughlin, A. I. G.: To be published.

In Table 2 details are given of fatal cases of silicosis and asbestosis investigated fully by the Medical Inspectorate of Factories between 1930 and 1953. The average ages at death and the length of employment in the dusty industries are also given.

It is seen that as regards silicosis the manufacture of scouring powders and sandblasting were the most dangerous industries, with the lowest average age at death and the shortest average period of employment. Another interesting point is that tuberculosis occurred in about half of the cases of silicosis, whereas it was found in only about two-fifths of the cases of asbestosis. The average age at death and the length of exposure in the cases of asbestosis were much lower than in the silicotic group as a whole but more on the level of sandblasters and makers of scouring powders.

The connection between asbestosis and cancer of the lung is becoming clearer, and in one series of 100 autopsies on asbestosis cases there were 25 cases of cancer of the lung. Wyers (1949) has pointed out that the x-ray

appearances of asbestosis. Some change in that is ground-glass shadows disappear. He thinks it probably of the lessened exposure period, owing to the more exhaust ventilation to the use of personal protection of asbestosis, however, as workers who do asbestosis boilers. In this process, asbestosis is being sprayed, it adequate protective measure because the workers are

PRESENT NO.

How many cases of pneumoconiosis, asbestrosis, coal miner's

Industry
Potteries
Sandstone
Grinding of limest. etc.
Refractories
Miscellaneous
Coal mining
Other mining
Asbestos
Union (thymovais)
Total
Nonoccupational

Industry	Number of Deaths
Potteries	1
Sandstone	1
Grinding of lucites, etc.	1
Refractories	1
Miscellaneous	1
Coal mining	1
Other mining	1
Asbestos	1
Cotton (textile mills)	1
Total	10
Nonoccupational	1

	Deaths, No.	Average Age at Death, Yr.	Duration of Employment		
			Longest, Yr.	Shortest, Yr.	Average, Yr.
Silicosis					
Pottery					
Silicosis	730	61.7	63.0	2.8	36.1
Silicosis with tuberculosis	301	56.2	67.0	3.0	31.7
Sandstone					
Silicosis	207	60.9	62.0	0.0	32.0
Silicosis with tuberculosis	220	37.3	58.0	3.0	37.4
Grinding of Metals					
Silicosis	183	50.5	61.0	11.0	35.9
Silicosis with tuberculosis	211	64.0	50.0	2.8	33.0
Sandblasting					
Silicosis	73	50.4	42.0	1.7	13.4
Silicosis with tuberculosis	10	40.2	40.0	2.0	12.1
Manufacture of scouring powders					
Silicosis	10	40.8	37.0	2.3	8.2
Silicosis with tuberculosis	6	40.8	11.2	2.0	7.0
Miscellaneous					
Silicosis	283	33.1	37.0	1.3	21.2
Silicosis with tuberculosis	213	37.6	50.0	0.7	21.3
Total					
Silicosis	1,313	50.4	62.0	1.3	31.1
Silicosis with tuberculosis	1,201	34.4	57.0	0.7	31.2
Asbestosis					
Asbestosis	230	49.5	49.0	0.5	10.9
Asbestosis with tuberculosis	37	40.2	34.0	0.3	11.4

etc.) occur in Great Britain. It is mentioned that at the present time it is not possible to give accurate statistics of the populations at risk in the various occupational classes. Both the Registrar General and the Ministry of Labour classify the populations in the different occupational groups too broad to give accurate statistics. It may be possible in some cases to give more accurate information by classifying the populations according to the occupations and trades unions. It may be possible in some cases to give more accurate information by classifying the populations according to the occupations and trades unions. It may be possible in some cases to give more accurate information by classifying the populations according to the occupations and trades unions.

The deaths from all forms of cancer between 1940 and 1959 in England and Wales are given in Table 1.

DEATHS FROM

given of fatal cases of asbestosis fully by factories between age ages at death and in the dusty indus-

silicosis the man- mers and sandblasting industries, with the and the shortest yment. Another in- therentiosis occurred of silicosis, whereas at two-fifths of the average age at death re in the cases of as- r than in the silicotic ore on the level of of scouring powders. een asbestosis and ming clearer, and in es on asbestosis cases cancer of the lung. ed out that the x-ray

interval, 1950-1953

ion of Employment		
	Shortest, yr.	Average, yr.
	2.5	33.1
	3.0	34.7
	9.0	39.0
	9.9	37.4
	11.0	35.9
	11.3	32.0
	1.7	33.4
	2.0	33.1
	2.3	6.3
	2.0	7.0
	1.5	21.3
	0.7	21.8
	1.5	31.1
	0.7	31.2
	0.5	10.6
	0.8	11.1

DUST DISEASES IN GREAT BRITAIN

appearances of asbestosis are undergoing some change in that nodular as opposed to ground-glass shadows are beginning to appear. He thinks it probable that this is because of the lessened exposure to dust over a longer period, owing to the rigorous application of exhaust ventilation to the dusty processes and the use of personal protective devices. Cases of asbestosis, however, are now appearing in workers who do asbestos lagging of pipes and boilers. In this process, particularly if asbestos is being sprayed, it is difficult to apply adequate protective measures, more especially because the workers are usually peripatetic.

PRESENT POSITION

How many cases of pneumoconiosis (silicosis, asbestosis, coal miners' pneumoconiosis,

In all industries there were 8781 deaths from occupational fibrosis of the lung in the 12-year period, and it will be seen that the total yearly figures are going up. Over the same period there were 6807 deaths from non-occupational fibrosis of the lungs. About two-thirds of the total number of deaths in the occupational group occurred in coal miners, who form the largest group (about 700,000) exposed to dust inhalation. The figures rose steeply from 232 in 1940 to 937 in 1951 (in 1952 the figure dropped slightly, to 912). It may well be that part of the increase is due to more accurate diagnosis, or at least to greater interest in the pneumoconiosis problem, among coal miners. In Figure 2 is shown a comparison, based on the crude figures of Table 3, between the coal miners and factory

TABLE 3.—Deaths from All Types of Pneumoconiosis in England and Wales, 1940-1951

Industry	1940	1941	1942	1943	1944	1945	1946	1947	1948	1949	1950	1951	Total
Factories	53	45	47	41	32	41	49	51	45	63	72	62	608
Sandstone	109	82	54	77	57	61	55	65	68	51	29	71	779
Grinding of metals, etc.	41	26	26	21	24	22	22	27	26	37	23	21	265
Refractories	36	32	7	7	6	9	6	10	13	13	6	8	117
Miscellaneous	7	5	7	5	13	13	14	16	27	25	29	129	217
Coal mining	232	316	290	276	311	282	421	577	680	757	846	917	5,808
Other miners	61	40	46	43	30	49	51	50	49	61	42	31	579
Asbestos	11	37	21	8	10	11	16	18	15	17	12	18	161
Cotton (byssinosis)	..	..	6	7	1	10	3	4	8	7	11	5	61
Total	539	423	431	453	459	604	632	818	850	1,008	1,113	1,201	8,781
Nonoccupational	630	501	445	408	357	229	306	616	651	551	679	722	6,607


etc.) occur in Great Britain? It should be mentioned that at the present time it is impossible to give accurate figures of the populations at risk in the various trades and processes. Both the Registrar-General and the Ministry of Labour classify occupations in groups too broad to give specific figures of the populations in the dusty industries, and those in the possession of employers' associations and trades unions are not complete. It may be possible in several years' time to give more accurate information. All that can be done now is to state the numbers of deaths reported to the Registrar-General and to give the figures for one year of the new cases diagnosed by the Pneumoconiosis Panels.

The deaths from all forms of pneumoconiosis between 1940 and 1951, inclusive, in England and Wales are given in Table 3.

workers. In the factory processes, the yearly number of deaths was going down until 1943, but there has been a slight rise in the later years.

In both groups, it should be emphasized that the deaths in each year are related to conditions which obtained in mines and factories some years previously, possibly 10 to 20 years or even longer.

New Cases in Factory Occupations.—The following Table 4 gives the numbers of new cases of pneumoconiosis diagnosed by the Pneumoconiosis Panels in 1953. These figures have been kindly supplied by the Ministry of Pensions and National Insurance, but it should be noted that they differ slightly as regards the numbers in each occupation from the figures originally supplied. The Ministry usually classifies the cases as falling into the

DEATHS FROM ALL TYPES OF PNEUMOCONIOSIS
IN FACTORY OCCUPATIONS

FACTORY OCCUPATIONS



Fig. 2.—Chart showing trends of deaths from fibrosis of the lungs among coal miners and factory workers during the period 1940-1951, inclusive.

category of the last jobs done by the workers, whereas Dr. G. O. Williams, of the Factory Department, has reclassified some of the cases on the basis of all occupations done by each worker in order to determine the most likely cause of the pneumoconiosis.

There were 896 cases of pneumoconiosis diagnosed, and this figure includes 54 cases of byssinosis. The cases with an uncomplicated occupational history numbered 665, and in the

remainder there was exposure for varying periods to other dusts, mainly coal dust. Additional exposure to coal dust occurred mainly in those industries situated in coal-mining areas, such as refractories, stone (sandstone and granite), pottery, slate, and foundry industries. The pottery industry contributed the greatest number of cases (353, mainly silicosis), and the foundry industry came next with 190 cases. There are roughly 25,000 pottery and about 220,000 foundry workers (iron, steel, and nonferrous); so it is probable that the pottery industry has a higher pneumoconiosis risk than foundries. Such general statements are not of much value, because the risk varies from process to process within the same industry. The substitution of ground flint by calcined alumina has reduced the silicosis risk among makers of bisquit ware in potteries, whereas the fettling of steel castings is still a dangerous job.

New Cases in Mining.—Table 3 gives the numbers of new cases of pneumoconiosis occurring in all mining operations in 1953.

These figures show clearly the tremendous problem which faces the coal-mining industry. There is no space to give in detail the varying degrees of severity of pneumoconiosis found in the 4048 cases, but the figures are given in the "Digest of Pneumoconiosis Statistics

TABLE 4.—New Cases of Pneumoconiosis—1953
Factory Occupations

Industry	No Other Dust Exposure	Other Dust Exposure		Total
		Coal	Other	
(1) Refractories	13	10	..	23
(2) Stone workers (sandstone, granite).....	25	10	1	36
(3) Pottery	254	66	13	333
(4) Slate workers (quarrying and splitting)....	62	29	..	91
(5) Foundries	151	27	9	187
(6) Metal grinding (other than foundries).....	5	2	..	7
(7) Subblasting	5	4	..	9
(8) Shot blasting	2	..	..	2
(9) Furnace dismantling, etc.....	17	7	..	24
(10) Boiler scaling	4	1	..	5
(11) Abrasive wheel manufacture.....	3	..	..	3
(12) Asbestos	11	8	3	22
(13) Abrasive soap powders.....	..	..	..	..
(14) Graphite and carbon electrodes.....	..	..	..	..
(15) Tin smelting.....	1	..	..	1
(16) Cotton (byssinosis).....	10	14	..	24
(17) Coal trimming.....	10	..	..	10
(18) Miscellaneous	21	..	..	21
Total.....	615	205	25	845

POST 1953-4

TABLE 3.—New

(1) Coal mining	4048
(2) Pottery	353
(3) Foundry	190
(4) Pottery	100
(5) Pottery	100
(6) Pottery	100
(7) Pottery	100
(8) Pottery	100
(9) Pottery	100
(10) Pottery	100
(11) Pottery	100
(12) Pottery	100
(13) Pottery	100
(14) Pottery	100
(15) Pottery	100
(16) Pottery	100
(17) Pottery	100
(18) Pottery	100
(19) Pottery	100
(20) Pottery	100
(21) Pottery	100
(22) Pottery	100
(23) Pottery	100
(24) Pottery	100
(25) Pottery	100
(26) Pottery	100
(27) Pottery	100
(28) Pottery	100
(29) Pottery	100
(30) Pottery	100
(31) Pottery	100
(32) Pottery	100
(33) Pottery	100
(34) Pottery	100
(35) Pottery	100
(36) Pottery	100
(37) Pottery	100
(38) Pottery	100
(39) Pottery	100
(40) Pottery	100
(41) Pottery	100
(42) Pottery	100
(43) Pottery	100
(44) Pottery	100
(45) Pottery	100
(46) Pottery	100
(47) Pottery	100
(48) Pottery	100
(49) Pottery	100
(50) Pottery	100
(51) Pottery	100
(52) Pottery	100
(53) Pottery	100
(54) Pottery	100
(55) Pottery	100
(56) Pottery	100
(57) Pottery	100
(58) Pottery	100
(59) Pottery	100
(60) Pottery	100
(61) Pottery	100
(62) Pottery	100
(63) Pottery	100
(64) Pottery	100
(65) Pottery	100
(66) Pottery	100
(67) Pottery	100
(68) Pottery	100
(69) Pottery	100
(70) Pottery	100
(71) Pottery	100
(72) Pottery	100
(73) Pottery	100
(74) Pottery	100
(75) Pottery	100
(76) Pottery	100
(77) Pottery	100
(78) Pottery	100
(79) Pottery	100
(80) Pottery	100
(81) Pottery	100
(82) Pottery	100
(83) Pottery	100
(84) Pottery	100
(85) Pottery	100
(86) Pottery	100
(87) Pottery	100
(88) Pottery	100
(89) Pottery	100
(90) Pottery	100
(91) Pottery	100
(92) Pottery	100
(93) Pottery	100
(94) Pottery	100
(95) Pottery	100
(96) Pottery	100
(97) Pottery	100
(98) Pottery	100
(99) Pottery	100
(100) Pottery	100

No cases in other
industries.
Excluded fire clay
dust.

for 1953," issued by
Power (1954). In
more than 50% of
disability, and only
disabled.

The number of cases
diminishing, but the
mining are usually
severe silicosis.

DUSTS OTHER THAN COAL

About the middle
part of the century
dusts, Middleton, in
to dusts, such as talc,
gulf, talc, china clay,
the same year Doig
published a paper on the
the lungs of electricians,
has arisen a world-
and radiopaque dusts
contributors have been
dergrass, Norwalk, in
most interesting outcome
been the alteration of
interpretation of x-ray
because the x-ray features,
and stannosis. In
national histories and
can be mistaken for
lungs. In fact, an ab-
often found in a work-
disability. The study
by the excellent work
(1934) and later Cray

ure for varying percentages of dust. Additionally, occurred mainly in coal-mining, stone (sandstone), and foundry industry contributed 4 (353, mainly silicosis) industry came next with roughly 25,000 pottery foundry workers (5); so it is probable that as a higher pneumoconiosis. Such general value, because the process within the titation of ground as reduced the silicosis of biscuit ware in ing of steel castings

—Table 5 gives the pneumoconiosis conditions in 1953.

the tremendous al-mining industry.

detail the varying pneumoconiosis found figures are given pneumoconiosis Statistics

TABLE 5.—New Cases of Pneumoconiosis—1953
Mining*

Industry	Cases, No.
(1) Coal mining.....	1,045
(2) Tin mining.....	3
(3) Barite mining.....	1
(4) Fire clay mining.....	15
(5) Other clay mining.....	2
(6) Hematite mining.....	4
(7) Lead mining.....	2
(8) Tunneling.....	2
Total.....	1,073

*No cases in chert, oil shale, or stratified ironstone mining.
†Fourteen fire clay miners were also exposed to coal dust.

for 1953," issued by the Ministry of Fuel and Power (1954). It can be said, however, that more than 50% of the cases had only slight disability, and only about 3% were totally disabled.

The number of cases in hematite mining is diminishing, but the small numbers found in tunneling are usually instances of rapid and severe silicosis.

DUSTS OTHER THAN SILICA, ASBESTOS, AND COAL

About the middle 1930's the Factory Department began to turn its attention to other dusts. Middleton, in his 1936 review, referred to dusts, such as tripoli, sillimanite, kieselguhr, talc, china clay, and fullers' earth. In the same year Doig and McLaughlin published a paper on the x-ray appearances of the lungs of electric-arc welders. From this has arisen a world-wide study of the inert and radiopaque dusts, and notable American contributors have been Sander, Enzer, Pendergrass, Vorwald, and Hamlin. Perhaps the most interesting outcome of this study has been the alteration of the approach to the interpretation of x-ray films of the chest, because the x-ray features of siderosis, hantosis, and stannosis, in the absence of occupational histories and clinical examinations, can be mistaken for fibrotic changes in the lungs. In fact, an alarming x-ray picture is often found in a worker who has little or no disability. The study of siderosis was helped by the excellent work of Stewart and Faulds (1934) and later Crow (1937) on the hemati-

te miners, in which group the lesion was found to be siderostegosis, often accompanied by tuberculosis. Crow's later work (1947) in eliminating tuberculosis from the hematite mines and in improved dust control methods is of major importance.

Other occupations and dusts which have been studied include those of boiler sealers, graphite workers, grain dockers, and workers exposed to beryllium and its oxides, to the dust of leather mixed with other dusts, and to bagasse. Attention has also been given to exposures to the dusts of aluminum, manganese, and vanadium and to asthma occurring in workers exposed to the double salts of platinum and also to various wood dusts, such as western red cedar.

Pathology.—A great deal of work has been done on the pathology (gross, histological, and experimental) of coal miners' lungs by Gough, Harding, King, Heppleston, Wright, Gloyne, Nagelschmidt, and others. I might, however, state some conclusions about pathology which Harding and I (with our late colleague, S. Roodhouse Gloyne) have arrived at after studying the lungs of workers in many diverse industries.

Until about 10 years ago, the pathology of the dust diseases was dominated by the classical silicotic nodule (Fig. 3). In the same way, before a diagnosis could be made, the patient had to have x-ray nodulation, or the classical "snowstorm" effect. Classical silicosis usually occurs after the worker has been exposed to dust containing a high proportion of free silica. It is becoming increasingly clear, in our opinion, that even small proportions of free silica in a dust can cause a fibrosis which is composed of nodules not of the classical type (Fig. 4). The arrangement of the fibers is linear and radial, and the outline of the whole nodule is stellate. It looks like a black star. Harding, Gloyne, and I have applied the term mixed dust pneumoconiosis, or mixed dust fibrosis, to this nodule. It has been found in foundry workers, especially in cleaners of iron castings, in persons exposed to the dust of graphite containing about 10% free silica, and in boiler sealers who are ex-



Fig. 3.—Classical silicotic nodule from lung of a coal miner (56 years old); hematoxylin and eosin; $\times 24$.

posed to a mixed dust with a low proportion of silica. The coal nodule of coal miners' pneumoconiosis has much the same appearance as our mixed dust pneumoconiosis nodule. Coal contains varying proportions of free silica. In any case, the common denominator in those cases and occupations in which this type of nodule is found appears to be the presence of a small proportion of free silica in the dust. Where there is a high proportion, as stated above, the classical silicotic nodule is found. But in the same case, there may be both the classical nodule and the

Fig. 4.—Mixed dust pneumoconiosis nodule from a steel fitter's lung; hematoxylin and eosin; $\times 24$.



mixed dust pneumoconiosis nodule. Indeed, there are also nodules which show a transition stage between the mixed dust nodule and the classical one (Fig. 5). It should be mentioned that the x-ray appearances of the lungs of a worker with mixed dust pneumoconiosis differ little, if at all, from those of one with silicosis, and the disease is just as disabling and as fatal.

Boiler Scalers' Pneumoconiosis.—Experience of pulmonary diseases in ships' boiler scalers illustrates well the etiology of mixed dust pneumoconiosis. These men are exposed to a mixed dust which varies according to the type of fuel used to heat the boilers and also according to the source of the water used



Fig. 5.—Transition nodule, from mixed dust pneumoconiosis nodule to silicotic nodule in an iron dresser's lung; hematoxylin and eosin; $\times 24$.

in them. It contains carbonates, silicates, iron, and carbon. There is usually under 5% of free silica, but there is often a high proportion of iron and its oxides. Flue dust contains more iron than does the scale on the water tubes, sometimes as much as 48%.

In the first case of boiler scalers' pneumoconiosis the worker had been scaling boilers for over 40 years (Harding, Tod, and McLaughlin, 1944). In 1938 an x-ray film of his chest showed nodulation in the upper and outer lung fields, and in the lower zones reticulation (or micronodulation). At that time we made a tentative diagnosis of silicosis in the upper zones, with siderosis in the lower zones. Seven years later the patient


died of cancer of the bronchus. At the examination of the lungs (the cancer) the previous nodules in the upper and the lower zones (i.e., reticulation) there was carbon and iron dust fibrosis. In all other cases were nodules of mixed dust. Up to the present time on nine boiler scaler type of fibrosis. Only silicotic nodules in addition also died of cancer of the bronchus; well have been partly of substances present in the

The change from ships' boilers has been a problem. The dust causes symptoms of bronchitis in a high proportion of cases, but that this is due to the dust in the oil soot. Williams, an interesting series of point. My colleagues examining (with clinical environmental details (some 300 of them) in Hampton, where the boiler large passenger liners a litigation has not been compensated we have not damage as was shown in the port of Hull, fishing trawlers are not we have found three cases of pneumoconiosis among cases of cancer of the bronchus; proportion of cases with pneumoconiosis and died

Talc Pneumoconiosis.—Work on talc pneumoconiosis in the United States in 1933; Dreessen and Biddell, 1940; Siegal, 1943, etc.). Merrett in England, studied the features of rubber was dust. But it was not until

node. Indeed, it shows a transitional node and it should be remembered that the lungs of a patient with pneumoconiosis are those of one with just as disabling

coniosis.—Experiences in ships' boiler etiology of mixed cases where men are exposed to soot and to the boilers and to the water used



from mixed dust pneumoconiotic nodule in an iron and eosin: $\times 24$.

carbonates, silicates, is usually under 5% and often a high proportion of silicates. Fine dust does the scale on the inside of the boiler as much as 48%.

Boiler scalers' pneumoconiosis.—Experience with boiler scalding, Tod, and in 1938 an x-ray film was taken in the upper and lower zones (reticulation). At that time the diagnosis of silicosis with siderosis in the lungs later the patient

died of cancer of the lung, and histological examination of the lungs showed (apart from the cancer) the presence of classical silicotic nodules in the upper and outer zones, and in the lower zones (where there was x-ray reticulation) there were mainly deposits of carbon and iron dust in the lungs, with no fibrosis. In all other parts of the lung there were nodules of mixed dust pneumoconiosis. Up to the present time we have had autopsies on nine boiler scalers, and all showed this type of fibrosis. Only one scaler had classical silicotic nodules in addition. But five of them also died of cancer of the lung, and this may well have been partly caused by carcinogenic substances present in the soot.

The change from coal to oil as a fuel for ships' boilers has brought with it another problem. The dust from oil-fired boilers causes symptoms of bronchospasm or asthma in a high proportion of cases, and it is likely that this is due to the presence of vanadium in the oil soot. Williams (1952) has published an interesting series of cases illustrating this point. My colleagues and I are at present examining (with clinical, radiographic, and environmental details) the boiler scalers (some 300 of them) in the port of Southampton, where the boilers of ocean-going and large passenger liners are cleaned. The investigation has not been completed, but up to the present we have not found as much lung damage as was shown in a previous inquiry in the port of Hull, where the boilers of fishing trawlers are mainly scaled. So far we have found three cases of compensable pneumoconiosis among 50 workers and one case of cancer of the lung, together with a proportion of cases with lower degrees of pneumoconiosis and disability.

Talc Pneumoconiosis.—A great deal of work on talc pneumoconiosis has been done in the United States and Canada (Dreessen, 1933; Dreessen and Dalla Valle, 1935; Riddell, 1940; Siegal, Smith, and Greenburg, 1943, etc.). Merewether (1933-1934), in England, studied the x-ray and clinical features of rubber workers exposed to talc dust. But it was not until 1949 that we were

able to study the histological appearances of a proved case of talc pneumoconiosis (McLaughlin, Rogers, and Draham, 1949). A man, 51 years of age, had worked for 37 years in a rubber-tire factory, where he had been exposed to a fairly high concentration of talc dust. Moderately advanced pneumoconiosis of both lungs was found at autopsy (in addition to incompetence of the aortic valve). Throughout the lung substance were scattered small gray nodules, more numerous in the lower lobes. In some areas the nodules had coalesced to form small masses. The fibers appeared to be arranged concentrically around small vessels, giving the impression of whorling, but not like the appearance of silicosis. There were, in addition, many "curious" or talc bodies, resembling, but easily distinguishable from, asbestos bodies. Much dust, which proved to be talc, appeared in the sections, and it was thought at the time that the particles were all fibers. More recent work on the lung by Nagelschmidt † has shown that the bulk of the dust is in the form of plates with only a few fibers.

At least three other autopsies on cases of talc pneumoconiosis (all from rubber works) have been made, and they show similar features to the first published case. Small surveys have been made on workers exposed to talc dust in various industries, but larger ones have been planned to take place shortly. It is clear that talc dust, though fibrogenic, is not so active as asbestos in damaging the lungs.

Leather Dressers' Pneumoconiosis.—Co-operation between the directors of chest clinics, the mass radiography units, and the Medical Inspectorate of Factories has been instrumental in bringing to light hitherto unsuspected causes of pneumoconiosis. Since the mass radiography campaign was begun in 1948 for the early detection of cases of pulmonary tuberculosis, some 13,000,000 persons have undergone examination. Many factory populations have been surveyed, and

† Nagelschmidt, G.: Personal communication to the author.

included among them have been factories where there is a dust risk. The workers at one leather factory were examined by Dr. Hugh Ramsay, of the Wandsworth Radiography Unit. He drew our attention to the fact that in one department a high proportion of the workers showed abnormal x-ray changes, whereas in other departments, where there was no dust, no abnormalities were seen. In the department with the high proportion of x-ray abnormalities, skins loaded with china clay and calcium carbonate are dressed on rapidly revolving felt wheels covered with a layer of fine Carborundum powder. The job is very dusty, though it is done under exhaust ventilation. The dust is composed of much fine leather dust, with smaller proportions of china clay (kaolin), calcium carbonate, and Carborundum. It was found that the dust contained about 5% of free silica. The x-ray films showed all stages of abnormality, varying from early reticulation (micronodulation) through nodulation to massive shadows. In a few cases there was clinical evidence of disability. One man with an x-ray film showing massive shadows had died from "asthma and pneumonia" a year before the investigation began, and there was no autopsy. In spite of the fact that up to the present time no pathological evidence has been available, it is likely that the condition will fall into the group of the mixed dust pneumoconioses. The dust of china clay, which for years has been thought to be comparatively harmless, is becoming more and more suspect. Even apart from china clay, the presence of a small proportion of free silica is prima facie evidence that the dressing of such skins is a hazardous occupation.

Beryllium Pneumonitis and Granulomatosis.—It is remarkable that British experience of beryllium pneumonitis and granulomatosis is not as extensive as that in the United States, though many workers have been exposed to the dust of fluorescent lamp powders containing beryllium oxide and also the dust and fumes from the alloys. Only one death has occurred, and the Factory Department has collected clinical, x-ray, and environ-

mental details of five nonfatal cases, one of whom is now very ill. The other four cases are either recovering, or their condition is stationary. It is also remarkable that five out of six affected workers were chemists engaged in the development of the beryllium lamp powders. At the factory where the fatal case of beryllium granulomatosis occurred, 150 of the other workers were examined clinically and radiographically and no more cases were found, though there had been considerable exposure to the dust of the lamp powders in the early days. It is undeniable that beryllium oxide is toxic, but examination of the histology of the lungs of the one fatal case has led me to hold the unorthodox view that beryllium was not the sole cause of the condition. There were two types of lesion: one of a granulomatous type, in which there were giant and epithelioid cells, and the other in which were many fibrotic nodules indistinguishable from silicosis. Under polarized light many doubly refractile particles were seen in the nodules, and an eminent crystallographer stated that these were cristobalite, one of the most active forms of free silica. This is not surprising, because one of the ingredients of the lamp powder is silica gel (25%), which in the preparation of the powder is heated up to about 1100 C. The changed methods of preparing the lamp powder have eliminated both the beryllium oxide and the free silica, so that the risks both of beryllium granulomatosis and of silicosis have also been eliminated.

"Pneumoconiosis" from Vegetable Dusts.—Apart from cotton, not a great deal of work has been done on the vegetable dusts. The illnesses noted among cotton workers include mill fever, a transitory illness which affects nearly all new workers in cotton, flax, and hemp mills and also in malt houses; "Monday fever" (or feeling); weavers' cough, and byssinosis.

Byssinosis develops after about 20 years as a natural progression from "Monday fever" and in its final stage has the characteristics of chronic bronchitis and emphysema. As Schilling (1954) says, though it was de-

scribed by Greenhalgh, its etiology is still of but characteristic is and breathlessness usually extends to other disease progresses. "symptoms are very remain worse on Monday. It causes no specific lung fields." In in the cotton card and most from the disease, ers who clean the card ally affected. As men- ers with byssinosis are special scheme.

Bagassosis.—Only try handles bagasse (the sugar), and bag, mainly in those work- bagasse in a dry state, acute bronchiolitis, severe dyspnea, and x- showing generalized: ten cases of this ac- the notice of the Fac- of these have been in and Hamilton-Paterson Taylor (1942), and (1946). The last ca Since we got the firm under water, there h cases.

Farmers' Lung.—F lung was first described and after that Fawcitt a great deal of work on features similar to he occurs mainly among v handling moldy hay de is regarded as being Fawcitt was firmly of cases of the condition time to time by physici districts.

Pneumoconiosis in C ner, Hermon, and Day, the clinical features are pearances in a group

about 20 years as
"Monday fever"
he characteristics
emphysema. As
sugh it was de-

Pneumocoinosis in Grain Dockers.—Danner, Hermon, and Vagball described in 1946 the clinical features and abnormal x-ray appearances in a group of 55 grain dockers.

Vanadium Pneumonitis.—Wyers (1946) recorded his observations on workers exposed to vanadium pentoxide dust, and his results

are similar to those described by Sjöberg (1949) in Sweden. The effects are a combination of systemic poisoning and irritation of the pulmonary tissue, leading in some cases to bronchospasm and pneumonia. Doig § and Williams (1952) have described the marked bronchospasm occurring in workers exposed to the soot of oil fuel, which is thought to be caused by the high percentage of vanadium in the soot.

SUMMARY

The position as regards the dust diseases in Great Britain may be summarized as follows:

1. During the past 50 years, the risk of silicosis in most of the major industries has been firmly established. Until the late 1920's, free silica was the only dust which was thought to damage the lungs. Asbestosis, caused by the dust of a combined silica mineral, came into prominence about that time. In the middle 1930's, attention was given to the effects of other dusts on the lungs, such as iron and its oxides, and of radiopaque dusts, china clay, sillimanite, kieselguhr, talc, and mixed dusts containing free silica.

During the middle 1930's also, the effects of coal dusts on the lungs were given increasing attention, and extensive surveys were carried out, culminating in the formation of the Pneumoconiosis Research Unit of the Medical Research Council in 1945. The coal miners numerically constitute the greatest problem of dust disease in Great Britain.

2. Despite the combined efforts of inspectors of factories and mines, physicians, chemists, engineers, and research workers, the deaths from pneumoconiosis have continued to rise yearly. The reasons for this increase may be found in more accurate diagnosis of the condition and in the publicity given to occupational diseases of the lungs, leading to more frequent mention of pneumoconiosis by physicians on death certificates. But a more likely cause is the urge for increased speed of production, and the introduction of machines

which make a dusty process more dusty and cause a lag in dust-control methods.

3. Compensation for industrial pulmonary disease began soon after 1918, and gradually most of the processes and industries which damage the lungs are being brought under the provisions of the various Acts of Parliament. The Industrial Injuries Act of 1946 removed the responsibility for compensation from employers and insurance companies to the Government. Since then cases taken under Common Law against the employers have increased rapidly.

4. Numerous legal provisions, contained both in Acts of Parliament and in Regulations, have been brought into force. Routine inspection by inspectors of factories and mines has been instrumental in limiting the numbers of cases of disability and death brought about by inhalation of the dangerous dusts.

5. It is expected that more accurate information about the incidence of the dust diseases will be obtained in the near future, when statistical studies of the information obtained by the Pneumoconiosis Panels have been carried out, and when the report of the Registrar-General about the occupational incidence of disease at the time of the 1951 Census becomes available. At the present time no reliable information exists about the populations at risk in each industry and process.

6. Surveys of the population by mass radiography to detect early tuberculosis are bringing to light information about the effects of dust in some hitherto unsuspected occupations.

7. Many problems about dusts and the lungs are still unsolved, and some would say that we have only just begun to attack the fringes of the problem. Nevertheless, a great deal has already been accomplished.

Assistance in compiling the information contained in this paper was rendered by Dr. P. K. Walker and the members of the Pneumoconiosis Panels of the Ministry of Pensions and National Insurance: Drs. E. R. A. Merewether, A. T. Doig, and G. O. Williams, of the Factory Department,

DUST DISEASES 15

Ministry of Labour and
E. Hardon, lately of the
G. Nagelschmidt, of the
Laboratories, Sheffield.

BIBLIOGRAPHY

- Arlidge, J. T.: Hygiene in Occupations, London.
Cawpwell, J. M.: Acute Soot with Hay, Brit. M. J. 2:478, 1942.
Castleden, L. I. M., and J. Lagasosis: An Indust. M. J. 2:478, 1942.
Collis, E. L.: Industrial Special Reference to Health 23:232, 1914-15.
Cooke, W. E.: Pulmonary 2:1024, 1927.
Craw, J.: Blood Examination of Haematite Iron 19:8, 1937.
Control and Elimination of Coast Haematite Iron Indust. Med. 4:30, 1945.
Davies, T. A. L.: Mangan. Indust. Med. 3:111, 1945.
— and Harding, H. E.: Further Clinical and tions, Brit. J. Indust. 1.
Doig, A. T., and McLaughlin: appearances of the Lungs, Lancet 1:771, 1945.
Clearing of X-Ray Silicosis, Lancet 1:789, 1945.
Dreesen, W. C.: Effects on the Lungs, J. Indust. — and Dalla Valle, J. M. Dust in Two Georgia Pub. Health Rep. 59:1.
Dunbar, L.: Observations in Graphite Workers, 1945.
Observations on the Pneumoconiosis, Brit. — Hermon, R., and Pneumoconiosis in Dockers Seeds, Brit. J. Radiol. — and Pagnall, D. J. Graphite Workers, 1949.
Fawcett, R.: Fungoid Con. J. Radiol. 9:172 and Occupational Diseases of Cultural Workers, Brit. Gillison, J. A., and Taylor, Notes of Four Cases.

§ Doig, A. T.: Unpublished data.

M. McLAUGHLIN

more dusty and methods.

rial pulmonary IS, and gradually industries which brought under Acts of Parliament for compensation see companies to hen cases taken at the employers

isions, contained and in Regula- force. Routine of factories and d in limiting the ility and death of the dangerous

re accurate infor- of the dust dis- the near future, the information -sis Panels have e report of the e occupational ime of the 1951 At the present m exists about ch industry and

on by mass radi- tuberculosis are n about the ci- tio unsuspected

dusts and the some would say n to attack the theless, a great lished.

information con- ed by Dr. P. K. e Pneumoconiosis ons and National ther, A. T. Doig, istry Department,

DUST DISEASES IN GREAT BRITAIN

Ministry of Labour and National Service; Dr. H. E. Harding, lately of Sheffield University, and Dr. G. Nagel-Smith, of the Safety in Mines Research Laboratories, Sheffield.

BIBLIOGRAPHY

- Attridge, J. T.: Hygiene, Diseases, and Mortality of Occupations, London, Percival & Co., 1892.
- Campbell, J. M.: Acute Symptoms Following Work with Hay, *Brit. M. J.* 2:1143, 1932.
- Castleden, L. I. M., and Hamilton-Paterson, J. L.: Bagassosis: An Industrial Lung Disease, *Brit. M. J.* 2:478, 1942.
- Collis, E. L.: Industrial Pneumoconiosis, with Special Reference to Dust-Phthisis, *Pub. Health* 25:252, 1914-1915.
- Cooke, W. E.: Pulmonary Asbestosis, *Brit. M. J.* 2:1024, 1927.
- Craw, J.: Blood Examinations in Pulmonary Fibrosis of Haematite Iron Ore Miners, *Tubercle* 19:8, 1937.
- Control and Elimination of Silicosis in the West Coast Haematite Iron Ore Industry, *Brit. J. Indust. Med.* 4:30, 1947.
- Davies, T. A. L.: Manganese Pneumonitis, *Brit. J. Indust. Med.* 3:111, 1946.
- and Harding, H. E.: Manganese Pneumonitis: Further Clinical and Experimental Observations, *Brit. J. Indust. Med.* 6:82, 1949.
- Doig, A. T., and McLaughlin, A. I. G.: X-Ray Appearances of the Lungs of Electric Arc Welders, *Lancet* 1:771, 1936.
- Clearing of X-Ray Shadows in Welders' Siderosis, *Lancet* 1:789, 1948.
- Dreesen, W. C.: Effects of Certain Silicate Dusts on the Lungs, *J. Indust. Hyg.* 15:66, 1933.
- and Dalla Valle, J. M.: Effects of Exposure to Dust in Two Georgia Tale Mills and Mines, *Pub. Health Rep.* 50:131, 1935.
- Dunbar, L.: Observations on Pulmonary Disease in Graphite Workers, *Brit. J. Radiol.* 18:33, 1945.
- Observations on the Development of Graphite Pneumoconiosis, *Brit. J. Radiol.* 21:182, 1948.
- Hermon, R., and Bagnall, D. J. T.: Pneumoconiosis in Dockers Dealing with Grain and Seeds, *Brit. J. Radiol.* 19:506, 1946.
- and Bagnall, D. J. T.: Pneumoconiosis in Graphite Workers, *Brit. J. Radiol.* 22:573, 1949.
- Fawcett, R.: Fungoid Conditions of the Lungs, *Brit. J. Radiol.* 9:172 and 334, 1936.
- Occupational Diseases of the Lungs in Agricultural Workers, *Brit. J. Radiol.* 11:378, 1935.
- Gillison, J. A., and Taylor, F.: Bagassosis: Further Notes of Four Cases, *Brit. M. J.* 2:577, 1942.
- Gloyne, S. R.: Presence of the Asbestos Factor in the Lungs of Asbestos Workers, *Tubercle* 10:491, 1929.
- Marshall, G., and Hoyle, C.: Pneumoconiosis Due to Graphite Dust, *Thorax* 4:31, 1949.
- Greenhow, E. H.: Report of Medical Officer of the Local Government Board, Appendixes IV and VI, 1866-1891.
- Haldane, J. S.; Martin, J. S., and Thomas, R. A.: Report to Secretary of State for the Home Department on the Health of Cornish Miners, Command Report 2691, H. M. Stationery Office, London, 1904.
- Harding, H. E.; Tod, D. L. M., and McLaughlin, A. I. G.: Disease of Lungs in Boiler Scalers, With a Case Report and Review of Literature, *Brit. J. Indust. Med.* 1:247, 1944.
- Pneumoconiosis in a Boiler Scaler, *Brit. J. Indust. Med.* 4:100, 1947.
- and Oliver, G. E.: Changes in Lungs Produced by Natural Graphite, *Brit. J. Indust. Med.* 6:91, 1949.
- Hart, P. D'A., and Aslett, E. A.: Chronic Pulmonary Disease in South Wales Coalminers, Special Report 243, Medical Research Council, London, 1942.
- Hunter, D., and Perry, K. M. A.: Bronchiolitis Resulting from the Handling of Lagasse, *Brit. J. Indust. Med.* 3:64, 1946.
- McDonald, S.: Histology of Pulmonary Asbestosis, *Brit. M. J.* 2:1025, 1927.
- McLaughlin, A. I. G.; Groux, J. L. A.; Barrie, H. J., and Harding, H. E.: Iron Oxide Dust and the Lungs of Silver Finishers, *Lancet* 1:337, 1945.
- Rogers, E., and Dunham, K. C.: Tale Pneumoconiosis, *Brit. J. Indust. Med.* 6:184, 1949.
- Cheeseman, E. A.; Garrad, J.; Gloyne, S. R.; Goodall, K. L.; Harding, H. E.; Jukes, M. H.; Lawrie, W. B.; Perry, K. M. A.; Sutcliffe, C. L., and Woods, H.: Industrial Lung Diseases of Iron and Steel Foundry Workers, H. M. Stationery Office, London, 1950.
- Macklin, E. L., and Middleton, E. L.: Report on Grinding of Metals and Cleaning of Castings, Home Office, H. M. Stationery Office, London, 1925.
- Medical Research Council: Chronic Pulmonary Diseases in South Wales Coal Miners, Parts I-III, Special Report 243, 1942; Special Report 244, 1943, and Special Report 250, 1945.
- Meiklejohn, A.: Silicosis in the Potteries: Some Observations Based on 750 Necropsies, *Brit. J. Indust. Med.* 6:220, 1949.
- History of Lung Diseases of Coal Miners in Great Britain, 1800-1875, *Brit. J. Indust. Med.* 8:127, 1951.

- Department of Compensation for Occupational Diseases of the Lungs in Great Britain. *Erit. J. Indust. Med.* 11:158, 1954.
- Moreweather, E. M. A.: Annual Report, Chief Inspector of Factories, London, p. 63, 1933, and p. 37, 1934.
- Risk of Silicosis in Sandblasters, *Tubercle* 17: 385, 1936.
- and Price, C. W.: Report on Effects of Asbestos Dust on the Lungs and Dust Suppression in the Asbestos Industry. H. M. Stationery Office, London, 1930.
- Middleton, E. L.: Industrial Pulmonary Disease Due to the Inhalation of Dust, with Special Reference to Silicosis. *Lancet* 2:1 and 59, 1930.
- Peacock, T. B.: Diseases of the Organs of Respiration. *Tr. Path. Soc. London* 12:36, 1861.
- Pearson, G.: *Philos. Tr.* London, 103:159, 1813.
- Pneumoconiosis Statistics for 1953. Digest of, Ministry of Fuel and Power, H. M. Stationery Office, London, 1954.
- Riddell, A. R.: Silicosis: Studies and Reports of I. L. O. (Series F, 17), Geneva, p. 39, 1940.
- Schilling, R. S. F.: Byssinosis in the Lancashire Cotton Industry, *Tr. A. Indust. M. Off.* 4:61, 1954.
- Siegal, W.; Smith, A. R., and Greenburg, L.: Dust Hazard in Tremolite Talc Mining, Including Radiological Findings in Talc Workers, *Am. J. Roentgenol.* 49:11, 1945.
- Sjöberg, S. G.: Vanadium Pentoxide (V_2O_5) Intoxication, *Nord. med.* 41:200, 1949.
- Stewart, M. J., and Haddow, A. C.: *J. Path. & Bact.* 32:172, 1929.
- and Fields, J. S.: Pulmonary Fibrosis of Magnetite Miners, *J. Path. & Bact.* 39:233, 1934.
- Sutherland, C. L., and Bryson, S.: Report on Occurrence of Silicosis Among Sandstone Workers, Mines Department, H. M. Stationery Office, London, 1929.
- Report on the Occurrence of Silicosis Among Granite Workers, H. M. Stationery Office, London, 1930.
- Thackeray, C. T.: Effects of Arts, Trades, and Professions, and of Civic States and Habits of Living, on Health and Longevity, Ed. 2, London, Longman, 1832.
- Williams, N.: Vanadium Poisoning from Cleaning Oil-Fired Boilers, *Brit. J. Indust. Med.* 9:50, 1952.
- Wood, W. E., and Glynn, S. R.: Pulmonary Asbestosis: A Review of One Hundred Cases, *Lancet* 2:1383, 1934.
- Wyers, H.: Some Toxic Effects of Vanadium Pentoxide, *Brit. J. Indust. Med.* 3:177, 1945.
- Asbestosis, *Postgrad. M. J.* 25:631, 1949.

Clearance from the L

ROY E. ALBERT, M.D.,
and
LAWRENCE C. ARNETT,

One of the important hazards associated with the industry is that of air-borne dust and radon gas. This is not a new problem; uranium mines have had this experience with it in the European uranium was an extremely high cancer.¹

The control of radon sense an economic sense greater the expenditure to prevent the less the degree of exposure. The problem is to find reasonable levels which would insure safety of being overly conservative expensive. Two approaches have been made. The concentration of radon (3% the Schneeberg and Jáchymov have been used as the basis maximum permissible curies/liter.² In practice curies/liter has been the extent. The other approach of establishing safe work is based on a maximum dosage rate to the lungs per week.³ An estimate of concentration of the particulate which if inhaled over a working week would do

Recorded for publication