

difficulty with some of the terms. chapter it would have been well to the difference in general between and chronic effects, as with many ces there is little similarity and the effect is usually the more serious. subject of workmen's compensation place in a book on industrial health, velopment of occupational disease ion should have been mentioned. o general types of occupational dists (all-inclusive and specific) with advantages or disadvantages are a paragraph or two.

chapter on "First-Aid Treatment of es" treats only the traumatic in- Burns are taken care of in another r, but under first-aid a few words have been helpful describing what to acute cases of poisoning, cyanide, spills, etc., where the patient is still ing but where prompt action is neces-

oes combine in one compact volume a readable style much information ustrial safety and health that other- would require reference to many s. For the beginning students of inal engineering, industrial safety and i, or personnel relations, it should a helpful book. The operating er and manager will want more com- information on the particular expo- in his plant and the industrial dian, if interested in prevention, will more on the engineering methods of ol.—S. W. Gurney.

TRIAL CANCER. By E. W. Baader, "NEUE ERGEBNISSE AUF DEM GEBIETE KREBSKRANKHEITEN". Adam and er, Leipzig, 1937 (pp. 104-128). aluable study, including the history dustrial cancer. Less well known of industrial cancer are included: the and nasal cancers seen in England in i workers, the lung cancers of genera- orkers in Japan, and the liver cancer iers in the Curische Hafl, caused by ech, *Opisthorchis felineus*.—L. Teleky.

THE JOURNAL OF INDUSTRIAL HYGIENE AND TOXICOLOGY

VOLUME 21

FEBRUARY, 1939

NUMBER 2

THE SYSTEMIC EFFECTS RESULTING FROM EXPOSURE TO CERTAIN CHLORINATED HYDROCARBONS*

LEONARD GREENBURG, MAY R. MAYERS AND ADELAIDE ROSS SMITH
Division of Industrial Hygiene, New York State Department of Labor, New York City

CHLORINATED naphthalenes and diphenyls, because of their electrical, heat and moisture-resisting properties, and because they are non-inflammable, are used extensively for insulating wire and in the manufacture of electrical condensers. The chlorinated naphthalenes are naphthalenes in which one or more of the hydrogen atoms has been replaced by chlorine. There is, thus, a series of these substances beginning with monochloronaphthalene and going on to the octochlor derivative. In industry they usually occur in mixtures in which more than one chlorinated

product is present. On the whole, the higher the chlorination, the more toxic this material becomes. In the manufacture of chlorinated diphenyls, C_6H_6 is converted into $C_{12}H_{10}$ which, in turn, is chlorinated to $C_{12}Cl_{10}$; the substitution products range from the monochlor to the decachlor diphenyl.

A rather characteristic acneform skin eruption resulting from exposure to these substances has been recognized for a great many years—indeed, ever since they began to be manufactured about 25 years ago. These skin eruptions came into some prominence in Germany during the war and have been attracting sporadic interest in this country ever since. An investigation of this condition, as it appeared among a group of young workers engaged in the manufacture of electrical condensers, was reported in a recent issue of this *Journal* (1).

Experience has shown that the medical practitioner is still somewhat unfamiliar with the skin eruptions, even

* Received for publication September 13, 1938.
The chlorinated naphthalenes are sometimes referred to as "Halowax" by purchasers and users. While the Halowax Corporation manufactures these substances, it also manufactures many others. Since the term "Halowax" is merely a trade name for substances manufactured by this Company, it should not be used indiscriminately. In all instances, substances should be designated by their chemical names only.



though they are by no means uncommon in the industries where the chlorinated naphthalenes and diphenyls are used. The reason for this is that any one physician is likely to see such cases only rarely, unless he happens to be practicing in the immediate vicinity of one of these factories where large numbers of workers are employed. That the relatively rare systemic effects resulting from such exposure almost invariably go unrecognized is not surprising under the circumstances. It is of the greatest importance however, that physicians become acquainted as promptly as possible with the clinical and pathological pictures presented by these patients, particularly as regards systemic effects, since failure to detect early clinical manifestations of toxicity, and to remove susceptible individuals promptly from further exposure, may, on occasion, result fatally.

REVIEW OF LITERATURE ON SYSTEMIC EFFECTS*

Experimental studies.—The first mention in the literature of systemic effects from chlorinated naphthalenes is that of Lehmann in 1919 (2). He found that animals which were fed or which inhaled these substances lost appetite and at death showed "peculiar" lesions in the liver.

In 1936, Flinn and Jarvik (3) experimented on rabbits with three different compounds; (A) a mixture of tri- and tetrachloronaphthalene, (B) a mixture of tetra- and penta-chloronaphthalene and (C) a mixture of penta- and hexa-chloronaphthalene. They also used sublimates of (B)

and (C) given off at 192°C and 172°C respectively. Large doses, approximately 15 mg. per kg, were injected subcutaneously each day.

Animals receiving compound (A) and the sublimate of compound (B) showed no lesions attributable to these substances when killed at the end of 2 months. The 30 animals, however, which had received the higher chlorinated compounds (B) and (C) and the sublimate of (C), all died in from 12 to 26 days. Autopsies uniformly showed extensive damage.

The next important study to be published was that of Drinker, Warren and Bennett (4). These investigators administered chlorinated hydrocarbons by inhalation, subcutaneously and by mouth, to white rats. Mixtures of (a) tri- and tetrachloronaphthalene; (b) penta- and hexachloronaphthalene alone, and (c) with 10% refined chlorinated diphenyl; and (d) chlorinated diphenyl were used.

Briefly, these experiments showed, that the tri- and tetrachloronaphthalenes produced relatively unimportant pathological changes in the liver until extremely high concentrations were used. Animals exposed for 6 weeks to the higher chlorinations, however, in relatively low concentrations, regularly showed minor degrees of liver damage even though, as a group, they gave no clinical evidences of such toxicity while alive. Exposed to still higher concentrations, the rats lost weight and appetite, and began dying after 8 days' exposure—many with severe jaundice. Examination of the liver of these animals at autopsy revealed marked central fatty degeneration with necrosis of liver cells.

It is of considerable interest that

* A review of the literature on skin manifestations appeared in the article by Mayers and Silverberg mentioned above (1).

192°C and 172°C doses, approximately, were injected day.

compound (A) of compound (B) attributable to these led at the end of animals, however, the higher chlorinated (B) and (C) and (D), all died in from autopsies uniformly image.

stant study to be of Drinker, Warren

These investigations of chlorinated hydrocarbons, subcutaneously in white rats. Mixture of tetrachloronaphthalene and hexachloronaphthalene (c) with 10% refined nyl; and (d) chlorinated are used.

experiments showed, tetrachloronaphthalene relatively unimportant changes in the liver until concentrations were exposed for 6 weeks to injections, however, in concentrations, regularly degrees of liver damage a group, they gave no signs of such toxicity exposed to still higher the rats lost weight and began dying after 8—many with severe degeneration of the liver at autopsy revealed fatty degeneration of liver cells.

considerable interest that

the exposed rats, which gave no clinical evidences of disease, promptly died from acute yellow atrophy of the liver when given a very small dose of carbon tetrachloride—a dose well tolerated by control animals.

Two other points of interest were brought out by the study: (1) in the rats, which on autopsy showed even marked pathological lesions of the liver, no abnormalities were found in the other organs—a finding not uncommon in experiments on animals, but rather rare in humans who die from acute yellow atrophy of the liver; (2) even the less important pathological changes induced in the liver by these chlorinated hydrocarbons in the lower concentrations were found to be very persistent, being present even 2 months after cessation of exposure.

Clinical Reports.—Clinical reports of cases of systemic poisoning from the chlorinated naphthalenes are as yet rare in spite of the length of time that these substances have been in use. No doubt the infrequency of such reports has been, in part, due first, to the fact that cases of systemic poisoning are unusual occurrences—the element of individual susceptibility appearing to play an even more important rôle than usual in this situation—and second, to failure on the part of physicians to recognize cases of poisoning. Until recently there has been general lack of knowledge of the toxicological properties of these chlorinated hydrocarbons, and there is still relatively little information available regarding the clinical picture of industrial poisoning from them.

The danger that lies in such a lack of knowledge is exemplified in a situation that recently came to the atten-

tion of the Labor Department in New York State, in which a physician had been treating a severe case of jaundice in a young woman exposed to chlorinated naphthalenes. After a long and serious illness, when the girl was on her way to recovery, the physician expressed the opinion that within a few weeks she would be able to return to her former work. Whatever the cause of the jaundice in this case, there is good reason to believe, on the basis of the experimental work previously cited, that further exposure to such substances would have entailed a grave risk to the patient's life.

In 1934, Courtois-Suffit (5) reported on the work of Touraine who, with his associates, examined 60 workers exposed to trichloronaphthalene and found mild digestive disturbances and dizziness in 13, but nothing of a more serious nature.

In 1935, Schwartz suggested the possibility of systemic disease from exposure to these substances in a talk before the American Public Health Association (6).

In 1936, three fatal cases of jaundice in chlorinated naphthalene workers were recognized in this country. These were reported by Flinn (3) and Drinker (4) who summarized the cases briefly.

All three of the men were young and in none could any predisposing cause other than their industrial exposure be found to account for their illness. Two of the men who had worked side by side died within 2 months of each other. Both had been exposed to mixtures of penta- and hexachloronaphthalene, and one had been exposed to a mixture of tetra- and penta-chloronaphthalene with 10% chlorinated diphenyl. In both, the

diagnosis of acute yellow atrophy of the liver was made on autopsy. In the third case no autopsy was reported, but death occurred after an acute illness characterized by jaundice. In one case dermatitis characteristic of the effect of chlorinated naphthalenes had preceded the jaundice.

In addition to these fatal cases, Drinker (4) also mentions four cases of non-fatal jaundice among individuals with similar exposure. No details are given.

CASE HISTORIES

Because of the obvious need for more clinical data in regard to the effects of chlorinated naphthalene exposure, we are reporting 3 cases in persons who, after exposure in the course of their work to these known hepatotoxic substances, died of acute yellow atrophy of the liver, and in whose cases no other etiological factors could be discovered even after very careful investigation. The first case was seen in consultation at the Lincoln Hospital in New York City by Adelaide Ross Smith. The second and third cases were seen in consultation at the New Haven Hospital by Dr. Leonard Greenburg, (now Executive Director of the Division of Industrial Hygiene of the New York State Department of Labor) when he was Commissioner of Health of New Haven, Conn. The plant conditions under which the patients had worked were carefully investigated.

Case 1. H. F.*

A 17 year old girl was admitted to the Lincoln Hospital, New York City, on the

* Grateful acknowledgement is made of the courtesy extended by the staff of Lincoln Hospital and the Medical Examiner's Office in granting permission to use the records in this case.

service of Drs. Kenneth Taylor, Edwin Hauser and Scott Johnson on April 26, 1932 in a semi-comatose condition. She was intensely jaundiced on admission.

Her past medical history, obtained from members of the family, was entirely negative with the exception of a tonsillectomy in 1930. Until the onset of the present illness she had been unusually healthy and free from symptoms of any kind.

The occupational history was as follows: After being graduated from grammar school she attended high school for 1 year and then obtained a job, her only one, with a concern manufacturing electrical condensers for use in radios. She worked at this one place for 7 months and stopped working there 7 days before her admission to the hospital.

Her work consisted of soldering and labelling condensers. It is possible, from the information given, that she may also have assisted in the sealing operations, but this could not be definitely ascertained. In any event, in the soldering of the condensers she was exposed to the fumes of tri- and tetrachloronaphthalene with which the condensers were originally impregnated. At the same time she was exposed to fumes of the higher chlorinated naphthalenes from the sealing operations conducted in close proximity to the tables at which the soldering was done.

The present illness began about 5 months before admission to the hospital—or approximately 2 months after starting work in the plant—at which time she noticed several pigmented areas on her face. These continued to increase in severity and extent, and caused her to visit the outpatient department of the Lincoln Hospital. Here, sugar was found in her urine and she was referred to the skin clinic where a diagnosis of acute catarrhal jaundice was made. Her rectal temperature, at that time, was 100°. A diffuse papulo-pustular eruption was also present on her face. On being questioned about it, the patient stated that it had been present to her knowledge for about 2 months.† From then on she suffered from

† Investigation revealed the fact that some of her co-workers were also suffering from a similar acneiform eruption, and one girl discontinued her work because of it. None of these other girls suffered any systemic disease.

increasi
petite w
sively n
tion inc.
Abou
mother :
About 3
of sever
abated.
a puffine
feet and
She co
1 week b
of admis
moveme:
tary boi
with any
seemed t
change i
the hosp:
This was
curred.
state of :
Summ:
patient ;
colored &
white. &
developed
puffiness
men. Sh
quently
contracti
ciated w:
head. T)
114, respi:
The fir
tient is s
More care
neck reve
tion begin
ing down
Besides t
were three
rest, appe
coal—one
one on tl
these blac
of aggre
did not sh
tion. It r
teristic of
The skin
of the bod
ish sublin

Taylor, Edwin
April 26, 1932
She was
obtained from
entirely nega-
amputation in
present illness
healthy and free
of any other
disease was as follows:
from grammar school
1 year and then
with a concern
condensers for use
at this one place for
7 days
the hospital.
of soldering and
it is possible, from
that she may also
operations, but
ascertained.
of the con-
to the fumes of tri-
with which the
impregnated.
exposed to fumes
caphthalenes from
conducted in close
at which the solder-
can about 5 months
to the hospital—or ap-
after starting work in
she noticed several
face. These con-
severity and extent,
the outpatient de-
Hospital. Here
urine and she was
where a diagnosis
was made. Her
at that time, was 100°.
On being questioned
it had been
for about 2
on she suffered from
the fact that
were also suffering
eruption, and one
her work because of it.
other girls suffered any

increasing weakness and nausea. Her appetite was poor; her skin became progressively more jaundiced, and the pigmentation increased.

About 6 weeks before admission her mother noticed that her eyes were yellow. About 3 weeks later she began to complain of severe headaches which continued unabated. This was followed in a few days by a puffiness and swelling of the face, hands, feet and abdomen.

She continued at her regular work until 1 week before admission. On the morning of admission she began to have convulsive movements of the abdomen with involuntary bobbing of her head, unassociated with any pain. At the same time there seemed to develop memory defects and a change in mental status. On the way to the hospital in the ambulance she vomited. This was the first time vomiting had occurred. She arrived in the hospital in a state of prostration.

Summary of physical examination.—The patient gave the impression of being a colored girl although she was, in reality, white. She looked well nourished and well developed for her age. There was general puffiness of the face, hands, feet and abdomen. She was hiccupping, and was frequently disturbed by a series of tetanic contractions of the abdominal muscles associated with involuntary bobbing of the head. The temperature was 100°, pulse 114, respiration 26.

The first admission note stated, "patient is so jaundiced her face is black." More careful examination of the face and neck revealed a negroid type of pigmentation beginning at the hair line and extending down to, and including, the neck. Besides this general pigmentation there were three areas somewhat darker than the rest, appearing very much like dabs of charcoal—one on each cheek below the eyes and one on the chin. On close examination these black spots appeared to be the result of aggregations of comedones. The scalp did not share in the general dark pigmentation. It was of a yellowish color characteristic of jaundice.

The skin of the thorax and the lower part of the body was dark brown with a yellowish subtinge. Over the skin of the abdo-

men there was a girdle-like area of deeper pigmentation beginning at the level of the umbilicus and extending to the pelvis. There was also a line of deeper pigmentation in the garter region on the posterior surface of the left thigh about 1-inch above the popliteal fold.

Examination of the eyes revealed deep jaundice of the sclerae and the conjunctivae. The pupils were equal and regular, and reacted to light and accommodation. There was no nystagmus or strabismus. The blephoral conjunctivae were redder than normal. The discs were somewhat pale, though well defined. There was no exophthalmus. The normal ratio in width of artery to vein was preserved. The vessels were not contracted, and did not present the hair-line appearance believed to be associated with arsenic or quinine poisoning.

Other positive physical findings were: slight injection of the throat; slight edema of the feet and hands; an apical systolic murmur; abdomen distended and tympanic; liver on percussion small, with tympany extending to the very costal edge.

Laboratory findings: These were as follows:

Blood count on admission—White cells, 9,100; Red cells, 4,200,000; Polys., 80%; Lymphs, 20%; Hemoglobin, 67%; Bleeding time, 4.5 minutes; Clotting time, 2.5 minutes; Platelets, 220,000.

Blood chemistry: Non-protein nitrogen, 25, later 45, mg./100 cc.; Protein, 6.9 mg./100 cc.; Urea N. 20.83 mg./100 cc., Creatinin, 2 mg./100 cc.; Glucose, 88 mg./100 cc.; Cholesterol, 145 mg./100 cc.; Calcium, 8 mg./100 cc.; CO combining power, 52; Albumin, 2.5 mg./100 cc.; Icteric index, 230; Van den Berg, immediate direct reaction; Wasserman, negative.

Urine—Sp. gr. 1.008; Alkaline. Albumin negative. Sugar, a trace. Acetone, 1+. Bile pigment present. Urobilinogen a trace. Few red and white cells.

Stools—Urobilin present.

Spinal fluid—Pressure normal. Cells: 10, Lymphocytes. Clear. Glucose present. Globulin absent.

X-ray examination—No evidences of lesions of the heart and lungs; bones of skull, ribs, shoulder joints, clavicles, tibia and fibula. The gastric contents, and later,

at autopsy, the organs were analyzed for arsenic, phosphate, lead, tin and antimony. A trace of phosphate was reported, probably from inorganic phosphates ingested before admission.

Diagnosis—The patient presented a picture which was distinctly puzzling to all physicians who saw her and no conclusive diagnosis was reached at the time. The various diagnoses suggested included: obstruction of the common bile duct, toxic jaundice possibly due to some dye, obstruction of the inferior Vena Cava, blood dyscrasia, Addison's diabetes, hemochromatosis, industrial poisoning of unknown origin possibly due to arsenic, causing acute glomerular nephritis and acute hepatitis; and finally acute yellow atrophy of the liver and pancreas possibly due to an unknown industrial poison.

Course—The clinical course was steadily down-hill. At 11 p.m. on April 27th the patient became unconscious and went into a coma. There was Cheyne-Stokes breathing. A peculiar odor was present which was variously identified by hospital physicians as ethylene or garlic. The pupils were immobile to light. Deep reflexes were greatly depressed. There was no Babinski. Abdominal reflexes disappeared. The skin became dusky throughout. The liver dullness seemed to diminish. On April 28th the temperature rose to 106°; the pulse to 140, and the patient died.

Anatomical diagnosis—An autopsy was performed at the Medical Examiner's office on April 29, by Dr. Chas. H. Hochman, Assistant Medical Examiner. The report in full was as follows:

Body is that of a white adult female, well developed and nourished. Cyanosis of lips, ears and fingernails. Marked pigmentation about eyes, lower lips, neck and abdomen. General icteric tinge to entire body. No evidence of violence. Some edema about ankles. Hair is black. Eyebrows black. Brown iris. Conjunctivae and sclerae icteric. Rigor mortis present. Postmortem lividity of dependent parts. Scalp incised and reflected, calvarium removed. Brain found congested. Meninges bile tinged. On section, no evidence of intracranial injury or hemorrhage. Dura stripped and reveals no evidence of fracture.

Body opened in usual midline incision. Sternum removed. Some remnants of thymus still present. Lungs are free. Bronchi contain a bloody mucous. Mucosa injected, has yellowish tinge. Hemorrhagic infarct about size of hen's egg in left lower lobe. This is airless and dark red in color, firm. There are also similar smaller areas in other lobes. Heart is small and shows some sub-epicardial hemorrhages. Some hypertrophy of left ventricle. Valves thinned. Arteries natural. All are bile stained. No significant findings in myocardium or coronary arteries.

Esophagus is natural. Stomach is distended, contains some dark material. The rugae are somewhat hypertrophied. The duodenum contains green chyme. The ampulla is patent. The jejunum, ileum and large gut are natural. Many petechial hemorrhagic areas noted in the mesentery with some enlargement of mesenteric lymph nodes.

Liver is small, weight 720 gm. Capsule wrinkled. Right lobe on section shows areas of yellow surrounded by red areas.

Left lobe much firmer although liver cuts firmly throughout. All ducts patent. Gall bladder small. Wall thickened throughout.

Pancreas is natural in size, somewhat congested. Spleen is natural in size. On section, firm, dark red in color. Follicles are somewhat indistinct. Kidneys show swollen cortex. Evidence of parenchymatous degeneration of epithelial cells. Capsule strips easily. No granulation. Degenerative changes in cells of glomeruli only. No glomerulitis. Uterus is small. Endometrium bile-stained. Ovaries are natural in size; right contains corpus luteum cyst. Bladder wall is somewhat thickened. Mucosa is injected.

Microscopic examination—Liver—From size of a through and through transverse section one suspects that the entire liver was about $\frac{1}{2}$ normal size. Liver segment is very firm and composed of two distinct and peculiar types of tissue: (a) That constituting major portion of parenchyma is red in color; firm and elastic in consistency; (b) Other areas are light yellow in color and are scattered indiscriminately throughout reddish areas. Yellow areas vary from $\frac{1}{2}$ to 2 in. in diameter. External surface

Fel

of
smlow
red
str.
dis:
the
seer
blo-
and
nun
seer
yell
genT
of t
port
duc
The
rege
seer
has

ing

proc

S

cori

Som

deej

E

myo

K

erat

glon

glon

A

dege

P.

of cc

S:

dent

volv

liver

and

simi

A

mitt

2, 19

tory

E

ual midline incision. Some remnants of thyngs are free. Brony mucous. Mucosa vish tinge. Hemor-size of hen's egg in is airless and dark here are also similar lobes. Heart is small, ib-epicardial hemor-ophy of left ventri-l. Arteries natural. No significant findings rony arteries. ral. Stomach is dis-e dark material. The hypertrophied. The green chyme. The he jejunum, ileum and ral. Many petechial sted in the mesentery at of mesenteric lymph

ight 720 gm. Capsule be on section shows unded by red areas. ner although liver cuts All ducts patent. Gall thickened throughout. ral in size, somewhat a natural in size. On ed in color. Follicles stinct. Kidneys show dence of parenchyma-of epithelial cells. ily. No granulation. s in cells of glomeruli itis. Uterus is small. stained. Ovaries are at contains corpus lu-r wall is somewhat is injected. mination—Liver—From nd through transverse a that the entire liver l size. Liver segment posed of two distinct f tissue: (a) That con-tion of parenchyma is d elastic in consistency; ight yellow in color and riminately throughout llow areas vary from ster. External surface

of liver, judging from this portion, was smooth.

A typical picture of clear subacute yellow atrophy is revealed. Slides show that red area is composed of so-called "red atrophy" in which all the liver cells have disappeared, their place being occupied by their normal supporting stroma, which seems to have been spared, great numbers of blood cells, early fibroblastic proliferation and older scar tissue. In these red areas, numerous bile ducts are seen. These also seem to have been spared destruction. The yellow areas are composed of actively regenerating islands of liver cells.

The toxin evidently has destroyed most of the liver cells, leaving intact their supporting stroma, blood vessels and bile ducts. The latter seem to be regenerated. Those liver cells spared destruction have regenerated and have formed yellow islands seen in the gross picture. *There evidently has been one or more attacks of hepatitis judging from the different ages of the pathological process in various parts of liver.**

Skin—Shows an increased density of corium which stains deeply with eosin. Some of the epidermal cells are slightly deeper in corium than normal.

Heart—Shows mild parenchymatous myocardial degeneration.

Kidneys—Show severe epithelial degeneration involving epithelium of tubules and glomeruli. No inflammatory reaction of glomeruli.

Adrenals—Show severe parenchymatous degeneration of cells.

Pancreas—Shows intense degeneration of cells of acini and of islets.

Summary—An unknown toxin has evidently caused a severe diffuse cytotoxicity involving most of organs, predominately the liver. *The latter has evidently undergone and recovered from previous attacks of a similar nature.**

Case 2—F. D.

A young man, 24 years of age, was admitted to the New Haven Hospital on May 2, 1934, complaining of jaundice. The history given was as follows:

He had worked in a wire factory coating

* Authors' italics.

wire with wax† from July to the middle of December, 1933. At that time he felt run down and became jaundiced. He was seen in the dispensary in January and found to have an enlarged palpable liver. He then spent several weeks in another hospital with slight improvement in the jaundice.

In March, 1934, he returned to the job in the wire factory. Following this, the jaundice became quite intense with increase in general malaise, anorexia, attacks of dizziness and loss of weight. Three days before admission vomiting had occurred. He had taken no drugs and there had been no exposure to carbon tetrachloride or chloroform.

Summary of physical examination—The temperature was 99°, pulse 74, respiration 20, blood pressure 130/74. The patient was a thin well-developed man who appeared moderately ill. There was bright yellow jaundice of the entire body. The pupils were widely dilated, the nasal septum deviated, causing obstruction on the right, the mucous membranes injected and the tongue moderately coated. The tonsils were enlarged and cryptic. Lungs were clear, heart normal, radial vessels soft, abdomen tympanitic, soft non-tender. The liver edge was felt at the costal margin. The organ was soft and non-tender. The upper edge of the liver dullness was in the 6th interspace.

Laboratory findings—These were as follows:

Urine—was dark yellow and was negative except for the presence of bile and urobilin.

Blood count—red cells, 4,380,000; hemoglobin, 85%; white cells, 6,800; Polys., 65%; Lymphs, 29%; Large Monos., 4%; Eos., 2%. Subsequent examinations showed a slight rise in white cells to 10,000.

Stool—showed bile, considerable undigested food, small amount of fat.

Kahn test—negative.

Liver function tests—Icteric index—30 to 50. Bromsulfonphthalein showed marked

† The waxes employed in this process are the higher chlorinated naphthalenes. These are used in a molten state in a bath through which the wire to be coated is passed. The process is partially but not entirely enclosed. Exhaust ventilation is in use.

impairment: 100% retention in 5 min., 85% retention in 30 min.

Blood chemistry: Non-protein nitrogen, 28 mg./100 cc.; Urea N, 11 mg./100 cc.; Serum total proteins, 6.15 mg./100 cc.; Serum total albumin, 2.66 mg./100 cc.; Serum total globulin, 3.49 mg./100 cc. A/G ratio, 0.76; Blood sugar, 68.0 mg./100 cc.; Blood calcium, 10.32 mg./100 cc.; Blood phosphorus, 4.12 mg./100 cc.; Serum fatty acids, 11.3 mg./100 cc.; Serum lipoid phosphorus, 7.5 mg./100 cc.; Serum total cholesterol, 10.5 mg./100 cc.; Serum free cholesterol, 63.0 mg./100.

Sugar tolerance test was within normal limits.

X-ray of abdomen showed hepatic flexure of colon unusually high.

Course. The patient was put on a high carbohydrate diet and his general condition improved but the jaundice persisted. He was discharged on May 16, 1934 with the diagnosis of toxic hepatitis.

He was admitted again on June 1, 1934 complaining of abdominal swelling of 3 days' duration, weakness, anorexia and edema of the legs which had developed soon after his discharge. The physical examination showed deep jaundice, systolic murmur, abdomen distended and tympanitic with dullness in the flanks, moderate pitting edema of the legs. Examinations of the urine, blood and stool showed no important variations from the original findings with the exception of a drop in white cell count to 6,400 with 74% polys., and a decrease in serum albumin to 1.88%. Serum CO₂ content was 54.54. Serum chloride was 98.0 mg./100 cc.

The patient's course after the second admission was rapidly downhill. The distention could not be controlled and he soon passed into complete coma. A convulsion occurred on June 8th and on June 10th he died.

Anatomical diagnosis.* Extensive necrosis, fibrosis and regeneration of liver, acute entero-colitis with edema, fibrosis of pancreas, acute pancreatitis, jaundice, ascites, edema of lower extremities, focal

* Autopsies on this and the following case were performed on the pathological service of the New Haven Hospital by Dr. H. M. Zimmerman. Grateful acknowledgement is made of his courtesy in permitting use of the records.

pneumonia (bilateral), subpleural hemorrhages, cloudy swelling of kidneys.

Case 3—C. C.

A young man 22 years of age was admitted to the New Haven Hospital on February 28, 1935 with the complaints of jaundice, abdominal pain, nausea and vomiting of bloody material. The history given was as follows:

He had worked in the same wire coating plant as the previous patient (F. D.). The present illness had begun with jaundice 2 months previously with no other symptoms. This continued for about 1½ months. Two weeks before admission he became more jaundiced and concomitantly developed upper abdominal pain, malaise, nausea and finally vomiting—the vomitus becoming bloody in character. He was treated by his family doctor with no relief. His condition became worse and he finally became delirious and incoherent. Hospitalization was advised.

Summary of physical examination—The patient was comatose, irrational and vomiting bloody material. Positive findings of significance were as follows: Temperature not elevated. Blood pressure 102/68. Generalized jaundice, petechiae over the extensor surfaces of the arms; tenderness over the upper quadrants of the abdomen. No liver dullness percussable.

Course—His condition became rapidly worse, coma setting in soon after admission. The vomiting continued. The patient died about 24 hours after admission.

Anatomical diagnosis—Extensive necrosis and regeneration of liver; jaundice, acute lymphadenitis of portal nodes; ascites; perienteritis of jejunum; cloudy swelling of heart and kidneys; acute pulmonary congestion; healing exanthematous rash of fore-arms.

DISCUSSION

These three cases show the occurrence of similar pathological changes in the liver in three young adults known to have been working with chlorinated naphthalenes and diphenyls—all were exposed directly or indirectly to the higher chlorinated hydrocarbons. In

Vol. 21, no. 2

suppurative hemor-
rhage of kidneys.

Case 1. The patient was admitted
on February 28,
with jaundice, ab-
dominal pain, and vomiting of
bilious material. The story given was as

follows: The patient was coating
the same wire coating
machine as patient (F. D.). The
patient had jaundice 2
months before admission.
The patient had been
about 11 months.
The patient had become
increasingly de-
pressed, with pain, malaise,
vomiting, and the vomitus
was bilious. He was
unable to eat with no relief.
The patient was finally
admitted to the hospital. Hos-

pital examination—The
patient had jaundice and vomit-
ing. The patient had
positive findings of
jaundice. Temperature
was 102/68.
The patient had
petechiae over the
arms; tenderness
of the abdomen.

The patient became rapidly
worse after admission.
The patient died
on admission.

Extensive necro-
sis of liver; jaundice,
portal nodes; as-
sessment of jejunum; cloudy
kidneys; acute pul-
monary edema; exanthematous

1938-1939

How the occur-
rence of pathological changes in
the young adults known
to be exposed with chlorinated
hydrocarbons—all were
of the type of indirectly to the
chlorinated hydrocarbons. In

all three cases, a most careful investi-
gation failed to reveal any other pre-
disposing cause for the condition. In
the first case in particular, that of a
healthy young girl on her first job,
the absence of any conditions predis-
posing to liver damage either preced-
ing or following her exposure to the
substances in question is especially
clear.

It is of interest to note that two of
the cases apparently had suffered from
at least one previous attack of hepati-
tis followed by a certain degree of
improvement before the onset of the
fatal attack.

In view of the fact that Drinker,
Warren and Bennett found no lesions
in organs other than the liver in their
experimental animals, it is noteworthy
that such lesions were conspicuous in
all three of the cases here reported.
Such lesions, are, indeed, usual find-
ings in cases of acute yellow atrophy
of the liver in humans regardless of
etiology. On the basis of present
knowledge, it would be impossible to
say whether they are the result of
primary intoxication or are merely
secondary to liver damage. In the
light of the animal experiments, how-
ever, one would incline to the latter
view and speculate as to the part
played by length of exposure, in the
final pathological picture.

It has been mentioned that in
Drinker's (4) experiments, rats with
no evidence of clinical disease while
being exposed to the chlorinated naph-
thalenes, promptly died from acute
yellow atrophy of the liver when given
doses of carbon tetrachloride so small
as to be harmless to control animals.
This prompts speculation as to
whether death from acute yellow
atrophy of the liver in workers simi-

larly exposed occurs only in those
having some pre-existing substratum
of liver damage—such as might follow
an attack of catarrhal jaundice, for
example—and that this might ac-
count, in part at least, for the fact
that only a very occasional worker out
of a large group will suffer from sys-
temic effects of exposure to these sub-
stances. In the three cases reported
however, no history was given sug-
gesting that any hepatic disorder prior
to exposure had occurred.

The presence of a papulo-pustular
eruption in two cases is of interest.
In one it antedated the systemic symp-
toms. This type of eruption is char-
acteristic of the dermatitis caused by
chlorinated naphthalenes, and until
recently was the only disturbance at-
tributed to them.

The presence of "aggregations of
comedones" such as were found in
Case I is also characteristic of the
skin eruptions produced by these sub-
stances. The question as to whether
or not the skin eruption in such a case
is in any way connected with the onset
of systemic effects cannot be answered,
since thus far no correlation has been
established between skin lesions and
systemic disease (1).

It is interesting, therefore, that in
Case I the skin eruption apparently
antedated all evidences of systemic
disease. Had the girl been promptly
removed from further exposure, when
it was first observed, it is possible that
her life would have been saved; or had
the physicians first called upon to
treat the jaundice (of whatever origin)
in her case, or in the second case here
reported, recognized the danger of
continuing exposure to the chlorinated
naphthalenes, these deaths might pos-
sibly have been averted.

RECOMMENDATIONS FOR MEDICAL CONTROL

1. The authors wish to stress the need for the conscientious reporting by physicians of all illnesses occurring among workers exposed to the chlorinated naphthalenes and diphenyls, particularly cases which have been carefully worked up, so that the clinical disease entities resulting from such exposure can become further clarified and thus more readily recognized in the future. Fewer errors in the diagnosis and management of these cases would occur, and workers' lives could undoubtedly be saved in this way.

2. Persons suffering from the typical acneform eruptions should be removed from further exposure.

3. Persons who have, at any time in the past, had any liver disease—even a mild catarrhal jaundice—should not work with these substances; nor should workers with a history of typhoid fever, malaria, gall-stones or other diseases known to affect the liver adversely.

4. Persons receiving arsphenamine treatment for syphilis; or those who are taking drugs believed to be injurious to the liver in susceptible

persons, should not be further exposed in their work to potential liver poisons.

5. Persons working with the chlorinated naphthalenes and diphenyls, if requiring a general anesthetic for an operation, should not be given chloroform or avertin, and vice versa, individuals who have recently received such anesthetics should not immediately thereafter go back to their former work or to work with other substances believed to be potentially toxic to the liver.

6. Pregnant women should not be exposed because the liver, in pregnancy, appears to be peculiarly susceptible to injury.

7. Experience seems to indicate that by proper attention to ventilation and medical supervision of workers the chlorinated naphthalenes and diphenyls can be used in industry with safety.

SUMMARY

The systemic effects resulting from exposure to certain chlorinated naphthalenes are discussed, and the literature of the subject briefly summarized. Three liver deaths in workers handling these substances are presented in some detail, with autopsy findings. Recommendations for prevention are given.

BIBLIOGRAPHY

1. MATERS, M. R., AND SILVERBERG, M. G.: Skin conditions resulting from exposure to certain chlorinated hydrocarbons. *This J.*, 20, 244 (1938).
2. LEHMANN, K. B.: *Kurzes Lehrbuch der Arbeits- und Gewerbehygiene*. S. Hirzel, Leipzig, 1919. p. 251.
3. FLINN, F. B., AND JARVIE, D. E.: Action of certain chlorinated naphthalenes on the liver. *Proc. Soc. Exper. Biol. and Med.*, 56, 118 (1936).
4. DRINKER, C. K., WARREN, M. F., AND BENNETT, G. A.: The problem of possible systemic effects from certain chlorinated hydrocarbons. *This J.*, 19, 283 (1937).
5. COURTOIS-SUFFIT: Étude sur l'intoxication professionnelle par le trichloronaphthalène. *Ann. de med. legale*, 14, 422 (1934). Abstr. of paper by TOURAINE, A., AND MÉNÉTRÉL, B.: Dermatoses professionnelles par le naphthalène et ses dérivés. *Prat. med. franc.*, 15, 335 (1934).
6. SCHWARTZ, L.: Dermatitis from synthetic resins and waxes. *Am. J. Pub. Health*, 26, 586 (1936).