

UNITED STATES DEPARTMENT OF THE INTERIOR
HAROLD L. ICKES, Secretary
BUREAU OF MINES
JOHN W. FINCH, Director

Bulletin 403

PLAINTIFF'S EXHIBIT

CAP-1475

ASBESTOS

BY
OLIVER BOWLES



UNITED STATES
GOVERNMENT PRINTING OFFICE
WASHINGTON : 1937

For sale by the Superintendent of Documents, Washington, D. C. - - - Price 15 cents

ASARCO ALV 0006244

CONTENTS

	Page		Page
Introduction.....	1	Foreign deposits.....	30
Definition.....	1	Canada.....	30
Varieties.....	1	Rhodesia.....	34
Physical properties.....	2	Union of South Africa.....	36
Outstanding characteristics.....	2	Cape of Good Hope.....	37
Fibrous character.....	2	Transvaal.....	37
Fineness of fiber.....	3	Chrysotile.....	37
Color and luster.....	3	Amosite.....	39
Heat resistance.....	3	Crocidolite.....	39
Heat conductivity.....	3	Anthophyllite.....	39
Acid resistance.....	4	Natal.....	39
Electrical resistance.....	4	Other African occurrences.....	40
Specific gravity.....	4	Soviet Russia.....	40
Chemical composition.....	4	Italy.....	42
Synthetic asbestos.....	5	Cyprus.....	43
History.....	5	Other foreign deposits.....	44
Early records.....	5	Argentina.....	44
History of commercial pro- duction.....	7	Australia.....	45
History of asbestos product manufacture.....	8	Western Australia.....	45
Origin.....	10	South Australia.....	45
Mode of occurrence.....	11	Queensland.....	46
Uses.....	11	New South Wales.....	46
General distribution.....	14	Tasmania.....	46
Production and consumption.....	15	Austria.....	46
History of production in the United States.....	15	Brazil.....	46
World production.....	16	Bulgaria.....	47
World consumption.....	17	China.....	47
Domestic deposits.....	19	Czechoslovakia.....	47
Chrysotile.....	19	Finland.....	47
Alaska.....	19	France.....	48
Arizona.....	20	Germany.....	48
California.....	21	India.....	48
Montana.....	23	Ireland.....	49
Oregon.....	23	Mexico.....	49
Vermont.....	24	Newfoundland.....	49
Wyoming.....	25	New Zealand.....	49
Other occurrences.....	25	Philippine Islands.....	49
Amphibole.....	26	Portugal.....	49
Alaska.....	26	Spain.....	49
California.....	26	Switzerland.....	49
Georgia.....	26	Turkey.....	50
Idaho.....	27	Venezuela.....	50
Maryland.....	28	Other countries.....	50
Montana.....	28	Political and commercial control.....	50
North Carolina.....	28	Political control.....	50
Oregon.....	29	Commercial control.....	51
Texas.....	29	International trade.....	51
Virginia.....	29	Exports from principal pro- ducing countries.....	51
Washington.....	29	Imports into the United States.....	54
Other occurrences.....	30	Exports from the United States.....	58
		Prospecting and exploration.....	59

	Page		Page
Mining methods.....	59	Milling methods—Continued.	
Arizona.....	59	Antophyllite milling.....	77
Vermont.....	60	Marketing.....	78
Georgia and Maryland.....	60	Prices.....	79
Canada.....	60	Manufacture of asbestos prod-	
Rhodesia.....	62	ucts.....	82
Union of South Africa.....	62	A widespread and diversified	
Russia.....	63	industry.....	82
Cyprus.....	63	Asbestos-products plants.....	82
China.....	64	United States.....	82
Milling methods.....	64	Canada.....	85
Classification of fiber.....	64	Great Britain.....	83
General features of milling		Germany.....	85
methods.....	65	Austria.....	85
Canada.....	66	France.....	86
Preparation of crudes.....	66	Belgium.....	86
Milling practice.....	66	Italy.....	86
Preliminary crushing.....	66	Czechoslovakia.....	86
Drying.....	66	Netherlands.....	86
Storage.....	67	Finland.....	86
Fiberizing.....	67	Scandinavia.....	86
Screening.....	68	Australia.....	87
Air suction.....	68	Russia.....	87
Wet process.....	68	Other countries.....	87
Grading.....	69	Fabrication of asbestos prod-	
Classification.....	69	ucts.....	87
Arizona.....	71	Fabrics.....	87
Vermont.....	72	Shingles and lumber.....	89
Union of South Africa.....	72	Paper and millboard.....	89
Chrysotile.....	72	Compounded packings.....	90
Crocidolite.....	73	Magnesia insulation.....	90
Amosite.....	74	Asbestos cement.....	90
Rhodesia.....	74	Cold-molded articles.....	90
Russia.....	75	Noncorrosive filters.....	91
Cyprus.....	77	Selected bibliography.....	91
Australia.....	77		

ILLUSTRATIONS

Fig.			
1. Typical cross-fiber veins of Canadian asbestos.....	10		
2. Asbestos consumption compared with automobile production and building construction, 1926-36.....	14		
3. Asbestos production in the three leading foreign centers and in the United States, 1916-35.....	17		
4. Crude chrysotile asbestos from Arizona.....	20		
5. Veins of asbestos on drift face 500 feet below surface, Thetford Mines, Quebec.....	30		
6. Asbestos veins in the New Amianthus mine, Barberton district, Transvaal, South Africa.....	38		
7. Steel chutes and chain-chute gates, Asbestos Corporation, Ltd., mine, Thetford Mines, Quebec.....	60		
8. Mill of Vermont Asbestos Corporation near Eden, Vt.....	72		
9. Average value per short ton of Canadian crudes and spinning fibers, 1921-35.....	80		
10. Average value per short ton of Canadian single, millboard, and paper stocks, 1921-35.....	81		

By C

IN

The mineral asbestos, being allied to cotton and wool by resistance; therefore, it supplies no substitutes are yet available. The features of the asbestos industry, mining, milling, utilization, in the United States leads all countries, but domestic source of raw material required. As great an extent on foreign deals at length with the asbestos industry.

Several reports now available in the countries. Among them are Ross; Asbestos in the Union series of articles on the Russian. These have been drawn upon hereby made for the helpful the magazine Asbestos also having all phases of the industry in the bibliography at the end and have been supplemented with the industry for a concise, world-wide history of the asbestos industry.

"Asbestos" is not the name rather, a commercial term applied to minerals. These minerals differ are diverse in strength, flexibility,

The following varieties of Chrysotile is a hydrous mineral represented by the chemical composition as massive serpentine. Chrysotile is characterized by

¹ Work on manuscript completed Jan. 1936.
² Assistant chief engineer, Nonmetallic

CONTENTS

Page		Page
59	Milling methods—Continued.	77
59	Anthophyllite milling.....	78
60	Marketing.....	79
6	ices.....	
60	Manufacture of asbestos prod-	82
62	ucts.....	
63	A widespread and diversified	82
63	industry.....	82
64	Asbestos-products plants.....	82
64	United States.....	85
64	Canada.....	85
64	Great Britain.....	85
65	Germany.....	85
66	Austria.....	86
66	France.....	86
66	Belgium.....	86
66	Italy.....	86
66	Czechoslovakia.....	86
67	Netherlands.....	86
67	Finland.....	86
68	Scandinavia.....	87
68	Australia.....	87
68	Russia.....	87
69	Other countries.....	87
69	Fabrication of asbestos prod-	87
71	ucts.....	87
72	Fabrics.....	89
72	Shingles and lumber.....	89
72	Paper and millboard.....	90
73	Compounded packings.....	90
74	Magnesia insulation.....	90
74	Asbestos cement.....	90
75	Cold-molded articles.....	91
77	Noncorrosive filters.....	91
77	Selected bibliography.....	91

STRATIONS

adial. Asbestos.....	10
ed with automobile production and	14
the leading foreign centers and in the	17
Arizona.....	20
10 feet below surface, Thetford Mines,	30
amianthus mine, Barberton district,	33
es, Asbestos Corporation, Ltd., mine,	60
ration near Eden, Vt.....	72
Canadian crudes and spinning fibers,	80
Canadian single, millboard, and paper	81

ASBESTOS¹

By OLIVER BOWLES²

INTRODUCTION

The mineral asbestos, because of its unique fibrous character, is allied to cotton and wool but has the advantage of heat and fire resistance; therefore, it supplies many industrial needs for which no substitutes are yet available. This report covers the essential features of the asbestos industry, including occurrence, production, mining, milling, utilization, international trade, and marketing. The United States leads all countries in the manufacture of asbestos products, but domestic sources furnish only a small fraction of the raw material required. As the United States is dependent to so great an extent on foreign supplies of raw materials, the bulletin deals at length with the asbestos industry in foreign lands.

Several reports now available deal with the industry in individual countries. Among them are Chrysotile Asbestos in Canada, by J. G. Ross; Asbestos in the Union of South Africa, by A. L. Hall; and a series of articles on the Russian industry, by Walter A. Ru Keyser. These have been drawn upon freely, and full acknowledgment is hereby made for the helpful information thus secured. The file of the magazine Asbestos also has furnished many valuable data touching all phases of the industry. Many other books and articles listed in the bibliography at the end of this bulletin have been utilized and have been supplemented by the author's intimate acquaintance with the industry for about 20 years. The intent is to present a concise, world-wide, historical, technical, and economic treatment of the asbestos industry.

DEFINITION

"Asbestos" is not the name of a distinct mineral species but is, rather, a commercial term applied to fibrous varieties of several minerals. These minerals differ widely in composition, and the fibers are diverse in strength, flexibility, and consequent usefulness.

VARIETIES

The following varieties of asbestos are generally recognized:

Chrysotile is a hydrous magnesium silicate, a fibrous serpentine, represented by the chemical formula $H_3Mg_3Si_2O_{10}$. It has the same composition as massive serpentine, with which it is always associated. Chrysotile is characterized by a strength and flexibility that permit

¹ Work on manuscript completed Jan. 26, 1937.

² Assistant chief engineer, Nonmetal Economics Division, U. S. Bureau of Mines.

the use of the longer fibers for spinning. The bulk of the world production of both long and short fibers is of this variety.

Other varieties of asbestos belong to the amphibole group of minerals. *Anthophyllite*, having the composition $(\text{Fe}, \text{Mg})\text{SiO}_3$, is the type most abundant in the United States. Usually, it is brittle and of low tensile strength and is more resistant to the action of acids and heat than chrysotile. It is used for nonspinning purposes.

Amosite, a variety of iron-rich anthophyllite occurring in Africa, has unusually long fibers. The best-grade material is used for spinning.

Crocidolite, also called "blue asbestos", having the composition $\text{NaFe}(\text{SiO}_3)_2\text{FeSiO}_3$, occurs in commercial quantities chiefly in Africa. It is used for both spinning and nonspinning purposes. Crocidolite occurring in quartz is sold as the semiprecious "tiger-eye."

Fibrous *tremolite*, $\text{CaMg}_3(\text{SiO}_3)_4$, *actinolite*, $\text{Ca}(\text{MgFe})_2(\text{SiO}_3)_4$, and *hornblende*, a complex silicate similar in composition to actinolite, generally are regarded as the minerals to which the term "asbestos" was applied originally. Except for Italian tremolite, they have little commercial value. *Mountain leather* and *mountain cork* are varieties of amphibole consisting of flexible sheets of interlaced fibers. *Mountain wood* is a compact fibrous mass of amphibole resembling dry wood. The last three varieties are mineral curiosities and have no commercial value.

PHYSICAL PROPERTIES

OUTSTANDING CHARACTERISTICS

The most characteristic properties of asbestos—those that give it commercial value—are fibrous structure and fire resistance. The longest and strongest fibers have qualities possessed by no other mineral, in that they may be spun into yarn or thread, woven into cloth, or felted into sheets and packings in almost the same way as animal and vegetable fibers, but unlike cotton or wool, they consist of incombustible matter.

FIBROUS CHARACTER

Although all asbestos is fibrous, the quality of the fiber in the various types and in different deposits is quite diverse. Chrysotile, amosite, and crocidolite normally possess high tensile strength, flexibility, and elasticity, but these qualities vary greatly in material from different localities. Chrysotile occasionally is harsh and somewhat brittle. Crocidolite is said to be stronger than chrysotile. The fibers of amosite are exceptionally long. Anthophyllite is generally brittle and of low tensile strength, although the fiber from some deposits has considerable strength and toughness. This type usually is more flexible than fibrous tremolite or actinolite. Tremolite is generally too weak and brittle for textile use; Italian tremolite, however, is said to be of spinning grade.

Length of fiber is important in asbestos strong and flexible enough for spinning and weaving. The fiber is graded on the basis of length, and materials of identical quality may range in price from \$20 to \$500 a ton, according to length.

Fineness of fiber is accomplished by fiber is due to a highly cleavage planes being size of the ultimate subdivision, therefore definitely, the fineness of the tools used and t

Varieties of asbestos markedly in the ease can be separated readily with difficulty and the mill has measured ant ester,³ and the smallest asbestos is measured in able and by the amount fineness desired.

Chrysotile occurs in When fiberized into a tically white, unless gray or yellowish gray fiberized. Crocidolite when the fibers are gray, greenish gray, unless staining impurities is silky or pearly.

Cirkel states that of 2,000° to 3,000° no visible effect on when chrysotile is heat lization is driven off lite, and amosite con higher temperatures though having a low magnetic mass.

Asbestos does not insulation is due to structure, which are nonconductors of heat

¹ Merrill, G. P., Proceedings, 282-283.

² Cirkel, Fritz, Chrysotile Canada Dept. Mines, Mines 1

³ Cirkel, Fritz, Work cited.

⁴ Dana, C. S., System of Mineralogy

s for spinning. The bulk of the world's short fibers is of this variety.

s belong to the amphibole group of minerals. Their composition $(\text{Fe}, \text{Mg})\text{SiO}_3$ is the same as that of the amphibole group of minerals in the United States. Usually, it is brittle and is more resistant to the action of acids than is used for nonspinning purposes.

It is used for nonspinning purposes.

a-rich anthophyllite occurring in Africa.

The best-grade material is used for

"blue asbestos", having the composition $(\text{Fe}, \text{Mg})\text{SiO}_3$ in commercial quantities chiefly in the spinning and nonspinning purposes. It is sold as the semiprecious "tiger-

$(\text{SiO}_3)_n$, actinolite, $\text{Ca}(\text{MgFe})_2(\text{SiO}_3)_n$, silicate similar in composition to actinolite as the minerals to which the term is usually applied. Except for Italian tremolite, they are. Mountain leather and mountain cork consisting of flexible sheets of interlaced a compact fibrous mass of amphibole last three varieties are mineral curiosities.

ICAL PROPERTIES

ING CHARACTERISTICS

properties of asbestos—those that give it its structure and fire resistance. The have qualities possessed by no other mineral into yarn or thread, woven into cloth, stockings in almost the same way as animal milk. Cotton or wool, they consist of

IOUS CHARACTER

fibrous, the quality of the fiber in the mineral deposits is quite diverse. Chrysotile, usually possess high tensile strength, flexible qualities vary greatly in material from chrysotile occasionally is harsh and somewhat to be stronger than chrysotile. The fibers are long. Anthophyllite is generally brittle, although the fiber from some deposits is tough. This type usually is more tough or actinolite. Tremolite is generally textile use; Italian tremolite, however, is

not in asbestos strong and flexible enough.

The fiber is graded on the basis of optical quality may range in price from 10¢ per length.

FINENESS OF FIBER

Fineness of fiber is measured by the degree of separation accomplished by fiberizing processes. The fibrous structure of asbestos is due to a highly developed prismatic cleavage, the spacings between cleavage planes being infinitesimal or, at least, measurable by the size of the ultimate particle of fiber, which is incalculably small. Subdivision, therefore, theoretically may be continued almost indefinitely, the fineness that can be attained depending on the delicacy of the tools used and the skill of manipulation.

Varieties of asbestos and occurrences of the same variety vary markedly in the ease with which they can be fiberized. Some types can be separated readily into slender fibers, while others are divided with difficulty and therefore appear as coarse-fibered material. Merrill has measured anthophyllite fibers down to 0.002 mm in diameter,³ and the smallest measured diameter recorded by Cirkel is 0.00075 mm.⁴ Apart from strength and flexibility, the quality of asbestos is measured to some extent by the fineness of fiber attainable and by the amount of processing required to reduce it to the fineness desired.

COLOR AND LUSTER

Chrysotile occurs in various shades of green or yellowish green. When fiberized into a fluffy mass, all types of chrysotile are practically white, unless stained by impurities. Anosite ranges from gray or yellowish gray to white and, if relatively pure, is white when fiberized. Crocidolite is lavender blue, and this color is maintained when the fibers are separated. Anthophyllite and tremolite are gray, greenish gray, or white, and the separated fibers are white unless staining impurities are present. The luster of asbestos usually is silky or pearly.

HEAT RESISTANCE

Cirkel states that Canadian chrysotile withstands temperatures of 2,000° to 3,000° F. and that a temperature of 5,000° F. produces no visible effect on some varieties.⁵ It has been found, however, that when chrysotile is heated to a high temperature its water of crystallization is driven off and it becomes brittle. Anthophyllite, tremolite, and amosite contain low percentages of water and withstand higher temperatures than other varieties. However, crocidolite, although having a low water content, is easily fused into a black magnetic mass.

HEAT CONDUCTIVITY

Asbestos does not have low heat conductivity. Its value for heat insulation is due to its nonflammability and also to its fibrous structure, which adapt it for manufacture into coverings that are nonconductors of heat because of their porous nature.

¹ Merrill, G. P., Proceedings of the United States National Museum: Vol. 18, 1895, pp. 232-283.

² Cirkel, Fritz, Chrysotile Asbestos—Its Occurrence, Exploitation, Milling, and Uses: Canada Dept. Mines, Mines Branch 69, 2d ed., 1910, p. 35.

³ Cirkel, Fritz, Work cited, p. 30.

⁴ Dana, E. S., System of Mineralogy: Sixth ed., New York, 1911, p. 671.

ACID RESISTANCE

Tremolite and anthophyllite are highly resistant to acids. It is claimed that crocidolite resists chemicals and sea water remarkably well. Chrysotile is affected more readily by acids and other chemicals than are the amphibole varieties.

ELECTRICAL RESISTANCE

Varieties of chrysotile lowest in iron content are the most suitable for electrical insulation. Arizona chrysotile is superior for this use. Apparently the iron content of amphibole does not affect its use for this purpose because crocidolite, which upon analysis shows a content of about 35 percent FeO, has high electrical resistance. This seeming contradiction may be explained by the fact that iron is present in crocidolite as a silicate, whereas in chrysotile it commonly appears as iron oxide impurities.

SPECIFIC GRAVITY

The specific gravity of chrysotile, as given by Dana,⁶ is 2.22, but that of the commercial fiber is always higher because of the presence of impurities. Canadian chrysotile ranges from 2.54 to 2.59, and the Arizona fiber, being lower in iron, has a specific gravity of 2.47. Anthophyllite ranges from 3.1 to 3.2 and crocidolite is still heavier, ranging from 3.2 to 3.3.

CHEMICAL COMPOSITION

The chemical formulas for the various kinds of asbestos are given in the section on varieties. Theoretical composition, as calculated from chemical formulas, rarely corresponds with actual composition because more or less impurity is always present. The following table shows analyses of representative commercial fibers.

Analyses of representative asbestos samples, percent

	SiO ₂	Al ₂ O ₃	Fe ₂ O ₃	FeO	MgO	CaO	Na ₂ O	H ₂ O	Total
Chrysotile:									
Canada ¹	40.36	0.21	1.35	0.66	43.86	-----	-----	13.45	99.89
Arizona ²	41.56	1.27	-----	.64	42.05	-----	-----	14.31	99.83
Barberton ³	40.05	1.90	1.60	.40	38.35	0.15	*0.40	16.60	99.70
Shabani ⁴	40.96	1.70	-----	2.44	38.73	-----	*.10	16.07	100.00
Russia ⁵	39.28	1.75	.40	5.37	40.05	1.74	-----	11.52	100.11
Anthophyllite: Georgia ⁶	56.40	1.15	-----	11.40	28.65	.50	-----	1.63	99.76
Crocidolite: Cape Colony ⁷	51.10	-----	-----	35.80	2.30	-----	6.90	3.90	100.00
Amosite: Penge ⁸	49.72	5.72	-----	37.00	3.77	1.65	-----	2.29	100.15
Tremolite: Natal ⁹	58.80	5.32	-----	-----	22.75	10.65	-----	.50	98.02

¹ Ross, J. G., Chrysotile Asbestos in Canada: Canada Dept. Mines, Mines Branch 707, 1931, p. 20.

² Diller, J. S., Mineral Resources of the United States, 1919. Pt. 2, p. 302.

³ Hall, A. L., Asbestos in the Union of South Africa: Union of South Africa Geol. Survey Mem. 12, 1916, p. 31.

⁴ Includes K₂O.

⁵ Ru Keyser, Walter A., Chrysotile Asbestos in the Bajenova District, U. S. S. R.: Eng. and Min. Jour., vol. 134, no. 8, August 1933, p. 338.

⁶ Hopkins, O. B., Asbestos, Talc, and Soapstone Deposits of Georgia: Geol. Survey of Georgia Bull. 29, 1914, p. 79.

⁷ Hall, A. L., Work cited, p. 19.

⁸ Hall, A. L., Work cited, p. 22.

⁹ Hall, A. L., Work cited, p. 23.

¹⁰ Includes Fe₂O₃.

Varieties of amphibole asbestos combined chemically as an essence may be present in any variety in the form of an oxide. The iron impurity in Arizona, Canadian, African, or Russian

SYNTH

Various attempts have been made to produce natural asbestos. Products such as slag wool, and glass wool have been made but lack its flexibility. Wool made in 1936 and later was with that of asbestos and are German inventor claimed in 1911 that of the same composition as chrysotile and at a cost lower

EAS

The peculiar property of its structure attracted attention. Small amounts were used by the ancients to "serve the ashes of the dead" and "mortal linen" was so difficult to weave. A specimen of cremation cloth in a Roman sarcophagus at the Vatican. According to tradition, the ancients wove asbestos into mats. In lamp wicks and to protect the

The word "asbestos" is a Greek word now known by that name in both ancient and modern Greek. The word "asbestos" means "inextinguishable" conveying quite the opposite of the characteristic feature of asbestos. reference to the "perpetual" Virgins. "Lithos amiantos" for asbestos, meant a rock doubtless referred to the clearest into the fire.

Another common name applied to asbestos because a well-known source. Karystos, southern Euboea, Strabo referred to stone from

¹ Some of the historical data on the history of the asbestos is given in the August, October, and November 1935 magazine Asbestos.

ACID RESISTANCE

Amphibole varieties are highly resistant to acids. It is resistant to chemicals and sea water remarkably more readily by acids and other chemical varieties.

ELECTRICAL RESISTANCE

The lowest in iron content are the most suitable. Arizona chrysotile is superior for iron content of amphibole does not affect because crocidolite, which upon analysis 35 percent FeO, has high electrical resistance. This may be explained by the fact crocidolite as a silicate, whereas in chrysotile on oxide impurities.

SPECIFIC GRAVITY

Chrysotile, as given by Dana,⁷ is 2.22, but is always higher because of the presence of iron. Canadian chrysotile ranges from 2.54 to 2.59. Crocidolite, which is lower in iron, has a specific gravity of 3.1 to 3.2 and crocidolite is still to 3.3.

CHEMICAL COMPOSITION

For the various kinds of asbestos are given. Theoretical composition, as calculated, usually corresponds with actual composition. The following are representative commercial fibers.

Analysis of asbestos samples, percent

Al ₂ O ₃	Fe ₂ O ₃	FeO	MgO	CaO	Na ₂ O	H ₂ O	Total
0.21	1.35	0.66	43.86	-----	-----	13.45	99.59
1.27	-----	.64	42.05	-----	-----	14.31	99.83
1.90	1.40	.40	38.35	0.15	0.40	16.60	99.70
1.70	-----	2.44	38.73	-----	4.10	16.07	100.00
1.75	.40	5.37	40.05	1.71	-----	11.52	100.11
1.15	-----	11.40	28.68	.50	-----	1.63	99.76
-----	-----	35.50	2.30	-----	6.90	3.50	100.00
5.72	-----	37.00	3.77	1.65	-----	2.29	100.15
5.32	-----	-----	22.75	10.65	-----	.50	98.02

Canada: Canada Dept. Mines, Mines Branch 707, 1931, p. 20.

United States: 1919. Pt. 2, p. 302.

South Africa: Union of South Africa Geol. Survey Mem. 12, 1918.

Asbestos in the Bajenova District, U. S. S. R.: Eng. and Min. Jour.,

Asbestos Deposits of Georgia: Geol. Survey of Georgia Bull. 29,

Varieties of amphibole asbestos, except tremolite, contain iron combined chemically as an essential constituent of the mineral. Iron may be present in any variety of asbestos as an impurity, chiefly in the form of an oxide that may be detrimental for some uses. The iron impurity in Arizona fiber is considerably lower than that of Canadian, African, or Russian chrysotile.

SYNTHETIC ASBESTOS

Various attempts have been made to manufacture a product similar to natural asbestos. Products known as mineral wool, rock wool, slag wool, and glass wool have some of the properties of chrysotile asbestos but lack its flexibility and strength. Special forms of glass wool made in 1936 and later are said to have flexibility comparable with that of asbestos and are capable of being woven and spun. A German inventor claimed in 1932 that he had made an artificial product of the same composition and physical properties as Canadian chrysotile and at a cost lower than asbestos mining costs in Canada.

HISTORY

EARLY RECORDS

The peculiar property of incombustibility combined with a fibrous structure attracted attention to asbestos more than 2,000 years ago. Small amounts were used by the Romans for winding sheets to preserve the ashes of the dead when bodies were cremated, but the "immortal linen" was so difficult to weave that only the most distinguished patricians and kings were honored with asbestos shrouds. A specimen of cremation cloth containing ashes was found in 1702 in a Roman sarcophagus and was deposited in the library of the Vatican. According to tradition, the ancient Chinese and Egyptians wove asbestos into mats. In ancient temples, asbestos was used for lamp wicks and to protect altar fires.

The word "asbestos", evidently first applied by Pliny to the fibrous mineral now known by that name, was employed erroneously, for in both ancient and modern Greek it refers to quicklime. The Greek word "asbestos" means "inextinguishable" or "unquenchable"—words conveying quite the opposite meaning from "incombustible", a characteristic feature of asbestos. Possibly Pliny had in mind Plutarch's reference to the "perpetual" asbestos lamp wicks used by the Vestal Virgins. "Lithos amiantos", the original term used by the Greeks for asbestos, meant a rock unstained, untainted, or undetiled, and doubtless referred to the cleansing of asbestos cloth by throwing it into the fire.

Another common name applied to asbestos was "Karystios lithos" because a well-known source of flexible mineral fiber was near Karystos, southern Euboea, Greece. Early in the first century, Strabo referred to stone from this locality that was carded and

⁷ Some of the historical data on this and following pages was obtained from a series of articles on the history of the asbestos-manufacturing industry that appeared in the August, October, and November 1935 and January, May, and August 1936 issues of the magazine Asbestos.

woven into handkerchiefs. Solinus and Plutarch also mention a fibrous mineral from this place. These ancient writers evidently knew little of the deposits, but as two exposures of serpentine occur east of Karystos the fiber probably was chrysotile.*

Pausanias, who lived in the second century A. D., refers to incombustible "Karpasian flax." Some writers think this term indicates that the source of the fiber was Karpasos, northeastern Cyprus, but no asbestos-bearing rock is known in that vicinity. Evans points out that the word "Karpasos" means "cotton" and that probably it was applied to the mineral because of its adaptability to textile use.⁹ Cyprus, however, was a well-known source of supply of asbestos in ancient times. Although it is difficult to determine from early references the exact location of the deposit, probably it was southeast of Mount Troodos in a village known as Amianto, the identity of which is lost. During recent years chrysotile asbestos has been produced in considerable quantities near a town now known as Amiandos. This is probably the site of ancient Amianto, for it is situated near Mount Troodos, 18 miles from the seaport Limasol. William Lithgow, a Scotsman, writing about the minerals of Cyprus early in the seventeenth century, referred to "the admirable stone Amiante, whereof they make Linnen cloth that will not burn, being cast into the fire, but serveth to make it neat and white."

The asbestos used by the Romans doubtless was a long-fibered variety identified as tremolite and occurring in northern Italy. The use of Italian fiber presumably led to the claim widely made in literature that the original "amiantos" consisted of fibrous amphibole (tremolite, actinolite, or hornblende). A recent letter to the author from the chief mineral inspector in Rome, however, casts some doubt as to whether all Italian fiber used by the ancients was tremolite. A significant excerpt from this communication reads as follows:

For a long time it was believed that the Italian "amianti" were all the tremolite type because the first examinations of the material were based on samples of tremolite. It was only in recent years that, after further studies, the producers recognized that their products were for the most part chrysotile with a long, flexible fiber.

A tremolite asbestos glove examined by the writer in the Museum of Natural History, London, England, in 1935 did not appear to consist of strong fiber. A piece of so-called "asbestos" cloth of unknown origin in this museum is not a fabric; it is a felted sheet that may be a natural mountain leather.

Very little mention of asbestos was made throughout the Middle Ages. According to report, Charlemagne, who reigned from 768 to 814, had an asbestos tablecloth, which he would throw into the fire for cleansing. To mystify his guests was apparently the only reason for its use.

Marco Polo refers to "amiantos" cloth shown to him during his travels through Siberia in 1250. He was told by the superstitious people of that region that it was made of the skin of salamanders, but he was too scientifically minded to accept such statements. After much questioning and search, he learned how the fibers were obtained

and prepared. His description consisted of pounding it so that the fiber was cut into threads. The fact that the Chinese were producing asbestos at Minuisinck in the thirteenth century is further evidence that it may have applied to chrysotile.

Further evidence that it may have applied to chrysotile is the uses of the material for lamp wicks, practice, asbestos was woven into and similar fabrics. It is brittle and cannot be used aside from the tremolite variety. South Africa—varieties of Chrysotile, on the other hand, adapted for textiles. It seems reasonable to conclude that it was used in the manufacture of it.

The next important meeting of the Royal Society was the meeting of the Royal Society in 1710 and 1720, and the first asbestos was discovered and handbags were made, through lack of

Asbestos was discovered in 1710 and 1720, and the first asbestos was discovered and handbags were made, through lack of

HISTORY

The utilization of asbestos began in Italy. Abo studies and experiments in the manufacture of asbestos. Several companies in Italy. Serious problems were encountered and heat-insulating began devising suitable machinery. Exhibits at the Unive publicity to the product.

In 1860 asbestos was and a specimen of fine asbestos in London in 1562. To work them profitably began with the discovery of the asbestos in the Coleridge Mines and Coleridge produced in that year. Italian asbestos, and easy availability of a rapid development. \$ of 1,400 tons a year was sold for only \$80 a ton, advanced to \$300 a ton, wasteful and costly h.

* Evans, John W., The Identity of the Amiantos or Karystian Stone of the Ancients with Chrysotile: Mineral. Mag. (London), vol. 14, May 1906, pp. 143-48.
⁹ Evans, John W., Work cited, p. 143.

fa. Solinus and Plutarch also mention a dis place. These ancient writers evidently ts, E. as two exposures of serpentine occur r pr only was chrysotile.*

n the second century A. D., refers to incom-
-7 Some writers think this term indicates
er was Karpasos, northeastern Cyprus, but
is known in that vicinity. Evans points
asos" means "cotton" and that probably it
al because of its adaptability to textile use.⁹
well-known source of supply of asbestos in
it is difficult to determine from early refer-
of the deposit, probably it was southeast of
ge known as Amianto, the identity of which
ars chrysotile asbestos has been produced in
ear a town now known as Amiandos. This
ient Amianto, for it is situated near Mount
the seaport Limasol. William Lithgow, a
the minerals of Cyprus early in the seven-
to "the admirable stone Amiante, whereof
that will not burn, being cast into the fire,
at and white."

the Romans doubtless was a long-fibered
olite and occurring in northern Italy. The
ably led to the claim widely made in litera-
amiantos" consisted of fibrous amphibole
hornblende). A recent letter to the author
spectator in Rome, however, casts some doubt
fiber used by the ancients was tremolite. A
this communication reads as follows:

elieved that the Italian "amianti" were all the
rst examinations of the material were based on
only in recent years that, after further studies,
the products were for the most part chrysotile

ove examined by the writer in the Museum
don, England, in 1935 did not appear to
A piece of so-called "asbestos" cloth of un-
eum is not a fabric; it is a felted sheet that
in leather.

asbestos was made throughout the Middle
rt, Charlemagne, who reigned from 768 to
ecloth, which he would throw into the fire
y his guests was apparently the only reason

"amiantos" cloth shown to him during his
n 1250. He was told by the superstitious
t it was made of the skin of salamanders,
y minded to accept such statements. After
ch, he learned how the fibers were obtained

r of the Amiantos or Karystian Stone of the Ancients
London), vol. 14, May 1908, pp. 143-48.
p. 143.

and prepared. His description of the process of preparation, which consisted of pounding in a mortar to eliminate impurities, suggests that the fiber was chrysotile, and this conclusion is substantiated by the fact that the best-known asbestos of Siberia is chrysotile occurring at Minuisinck near the Mongolian border.

Further evidence that the term "amiantos" as used by the ancients may have applied to chrysotile as well as to amphibole is found in the uses of the material. With the exception of twisted fiber used for lamp wicks, practically all the early references are to textile use. Asbestos was woven into handkerchiefs, tablecloths, winding sheets, and similar fabrics. Most varieties of amphibole asbestos are weak and brittle and cannot be spun or woven. The principal exceptions, aside from the tremolite of Italy, are the crocidolite and amosite of South Africa—varieties unknown until comparatively recent years. Chrysotile, on the other hand, is strong, flexible, and silky and well-adapted for textiles. Judging from the probable sources of supply it seems reasonable to conclude that it was employed in ancient times in the manufacture of incombustible fabrics.

The next important reference to asbestos was in 1676 when, at a meeting of the Royal Society, a Chinese merchant exhibited a handkerchief of "salamander's wool" or "linum asbesti." An asbestos napkin belonging to Ferdinand III was exhibited in Vienna in 1679.

Asbestos was discovered in the Ural Mountains of Russia between 1710 and 1720, and the first factory for making asbestos products was operated during the reign of Peter the Great. Textiles, socks, gloves, and handbags were made here for 50 or 60 years, but the enterprise failed, through lack of demand and poor transportation facilities.

HISTORY OF COMMERCIAL PRODUCTION

The utilization of asbestos on a commercial scale had its real beginning in Italy. About 1808, a noblewoman of Valtellina sponsored studies and experiments that brought her many honors and led to the manufacture of asbestos thread, fabrics, and paper of high quality. Several companies were formed between 1860 and 1875. No serious problems were involved in the manufacture of rope packings and heat-insulating board, but much difficulty was encountered in devising suitable machinery for the fabrication of spun products. Exhibits at the Universal Exposition in Paris in 1878 gave wider publicity to the products then manufactured.

In 1860 asbestos was discovered near St. Joseph in Quebec, Canada, and a specimen of fine silky fiber from this region was exhibited in London in 1862. The deposits were few and small, and attempts to work them profitably failed. A new era in the asbestos industry began with the discovery, in 1877, of the famous deposits near Thetford Mines and Coleraine. Mining began in 1878 and 50 tons were produced in that year. The fiber could be worked more easily than Italian asbestos, and its popularity in London, together with the easy availability of a growing market in the United States, led to rapid development. Seven quarries with an aggregate production of 1,400 tons a year were reported in 1885. The highest-grade fiber sold for only \$80 a ton in that year, but by 1900 the price had advanced to \$300 a ton. Thereafter, owing to depressed prices and to wasteful and costly hand methods of mining and preparation, the

recent years the coloring of shingles has attained higher refinement. and asbestos-cement siding shingles have been developed. Flat sheeting for siding and roofs came into use shortly after 1903, and corrugated sheeting was first made in 1905. Production of this great group of building materials has attained large proportions during the past 15 or 20 years.

Asbestos-cement pipes were first manufactured at Genoa, Italy, in 1913 by the Eternit process. Their manufacture began in America in 1929 and the first shipment was made in 1930.

ORIGIN

Asbestos originates for the most part from rocks consisting largely of olivine, such as peridotite or dunite, or from pyroxenite. Most chrysotile deposits result from alteration of olivine to serpentine, with subsequent prismatic crystallization providing the fibrous structure. Contact as well as regional metamorphism evidently plays an important part in the process because in various regions, notably the Canadian and Rhodesian fields, intrusive rocks are associated with the deposits and doubtless definitely influenced their development. Prismatic crystallization occurs in fractures. More detailed discussions of origin will be found in later sections describing asbestos deposits by State and country.

Chrysotile occurs in two forms—cross fiber and slip fiber. The former is the characteristic occurrence in all fissure-filled veins, the fibers crossing the veins approximately at right angles to the walls. In some veins the fibers cross from wall to wall, hence the fiber length is governed by the width of the vein. In others the fibers may be broken into two or more lengths by partings parallel with the vein walls. Slip fiber, on the other hand, parallels the vein wall and usually shows the effects of pressure and movement along a fault plane. Figure 1 shows typical cross-fiber veins of Canadian asbestos.

The chrysotile occurring in sedimentary rocks in Arizona and in the Carolina district of the Transvaal originated in a quite different way. The fiber in these localities is regarded as a product of alteration of magnesian limestone metamorphosed by diabase intrusions. It occurs in cross-fiber veins.

Under some conditions peridotites and pyroxenites are altered to fibrous amphiboles, of which the most common is anthophyllite. In some places it occurs as slip fiber in shear zones, but more commonly the altered rock takes the form of pockets or lenses consisting almost entirely of irregularly arranged bundles of fiber. The rock is called amphibolite, and the asbestos is of the mass-fiber type. In the irregular belt of olivine- and pyroxene-bearing rocks that extends from the Gaspé Peninsula in Canada to southern Georgia the alteration was principally into massive serpentine and chrysotile in the northern part, that is, in the area lying in Quebec, Canada, and in Vermont; but in its southern extension only sporadic occurrences of chrysotile are found because the alteration was more generally to anthophyllite and talc.

The softest and most easily fiberized anthophyllite occurs generally near the surface of the deposits; with increasing depth the fiber becomes harsher, more brittle, and less easily worked into a fluffy mass. The development of the best fiber therefore appears to depend upon

Bulletin 100




Figure 1. Cross-section of asbestos veins.

ASBESTOS

shingles has attained higher refinement, shingles have been developed. Flat sheet-metal use shortly after 1903, and commercial in 1905. Production of this great mass has attained large proportions during

re first manufactured at Genoa, Italy, in 1845. Their manufacture began in America in 1930.

ORIGIN

Most is made from rocks consisting largely of hornblende or dunite, or from pyroxenite. Most is formed from alteration of olivine to serpentine, crystallization providing the fibrous structure. Regional metamorphism evidently plays an important part because in various regions, notably the Adirondacks, intrusive rocks are associated with asbestos and definitely influenced their development. Asbestos occurs in fractures. More detailed discussion is in later sections describing asbestos.

Two forms—cross fiber and slip fiber. The occurrence in all fissure-filled veins, the veins are approximately at right angles to the walls. From wall to wall, hence the fiber length is the width of the vein. In others the fibers may be cut by partings parallel with the vein wall. In other hands, parallels the vein wall and pressure and movement along a fault. The cross-fiber veins of Canadian asbestos. In sedimentary rocks in Arizona and the Transvaal originated in a quite different manner. It is regarded as a product of mesozone metamorphosed by diabase in hornblende veins.

Hornblende and pyroxenites are altered to anthophyllite. In the most common is anthophyllite. In hornblende in shear zones, but more commonly in pockets or lenses consisting almost entirely of bundles of fiber. The rock is called anthophyllite. In the iron-bearing rocks that extends from Canada to southern Georgia the alteration is to serpentine and chrysotile in the north. In Quebec, Canada, and in Vermont only sporadic occurrences of the alteration was more generally to

fibers. Anthophyllite occurs generally in bands; with increasing depth the fiber becomes less easily worked into a fluffy mass. The fiber therefore appears to depend upon

Bulletin 400




FIGURE 1.—Typical cross-fiber veins of Canadian asbestos.

Bureau of Mines

the action of weathering agencies. Consequently, deposits of soft, easily worked anthophyllite tend to take the form of pockets or lenses of limited depth, and if workings are to be confined to fiber of this character the limitation of reserves must be considered with care. If, however, the harsher, more solid asbestos masses can be fiberized successfully into marketable products, the reserves may be much more extensive. Most of the deposits that have been developed show little promise of large reserves.

The crocidolite and amosite of South Africa originated from rocks quite distinct from peridotite or pyroxenite. They were derived from sediments rich in iron and silica, known as banded iron stones. Prismatic crystallization in cross-fiber veins probably was assisted to some extent by contact metamorphism.

MODE OF OCCURRENCE

From the discussion of origin, it is evident that there are three types of asbestos deposits—cross fiber, slip fiber, and mass fiber. Cross-fiber asbestos consists of innumerable strands extending across veins from wall to wall, except where interrupted by longitudinal seams. Most commercial deposits of chrysotile, amosite, and crocidolite are of this type, but anthophyllite rarely, if ever, occurs in cross-fiber veins. In slip-fiber deposits the strands more or less parallel the vein walls. They frequently show slickensided surfaces because they occupy shear zones where the rock has been subjected to movement and pressure. Slip fiber is common in the chrysotile deposits near Eden, Vt., and predominates in the anthophyllite deposits at Pylesville, Md., and near Bedford and Rocky Mount, Va.

In mass-fiber deposits there is no definite orientation of fibers. Almost the entire mass of the rock is composed of bundles of fibers or needles that sometimes show a radiating structure. This type of occurrence is confined almost exclusively to the anthophyllite variety and is well-exemplified by deposits at Sall Mountain, Ga., and Kamiah, Idaho.

USES

The uses of asbestos are so numerous and important that one could write a book on that subject alone. On the basis of use, asbestos falls into two principal classes—spinning and nonspinning fiber. Spinning fiber comprises the longer grades of chrysotile, crocidolite, amosite, and, exceptionally, tremolite. Nonspinning fiber comprises the shorter grades of these varieties and both the long and short grades of anthophyllite and related amphibole varieties. Spinning asbestos is used for weaving into textiles, but under the general classification of textile uses are included prepared yarns, listing, tape, rope, cord, wick, and thread. The largest use of asbestos textiles is for the manufacture of automobile brake-band linings and clutch facings. As much as 70,000,000 linear feet of brake-band lining has been made in a single year in the United States. Asbestos cloth treated with rubber is made into gaskets for use between the abutting or flange ends of pipes, or between adjoining surfaces, such as manhole and handhole plates, to make the joints tight enough to prevent passage of air, steam, or oil. Asbestos cloth is used, also, for the manufacture of fireproof theater curtains and scenery: blank-

ets; mattresses; firemen's suits, gloves, and shoes; aprons; mittens; leggings; helmets; awnings; conveyor belts for carrying hot materials; and woven sheet-packings. Braided and twisted packings are closely related to textiles.

An important use for nonspinning asbestos is for compressed packings consisting of felted masses of fiber combined with rubber or other binders. They are of various forms known as compressed sheet, coil, spiral, and high-pressure packings.

Another large group of uses is for heat insulation; in fact, asbestos is one of the most important fireproofing and heat-insulating materials known. Among the best-known products is 85-percent magnesia pipe covering. Asbestos paper is used for making 3-ply air-cell and similar pipe covering; as sheeting between floors; lining for stoves, heaters, filing cabinets, soldiers' helmets, mufflers for automobiles, drum controllers in sulphite mills, automobile radiator covers; tubes for electric wires; table mats, pads, stove mats, and other household appliances; and in many other ways. Asbestos paper is also used extensively in the manufacture of asbestos felt roofing and built-up roofs and in making asbestos-protected metal roofing. Millboard is used not only as a building material but as a joint packing for steam pipes and as a lining for safes, stoves, electric switch boxes, hoods of automobiles, and ovens, where a material of greater thickness than paper is required. An important use of millboard is for gaskets on automobiles and steam machinery. Asbestos is used for lagging steam boilers to prevent heat radiation. Large quantities of the shorter fibers are used in plastic fireproof cements for boiler, pipe, and furnace covering. An asbestos-cement coating is sometimes applied to the surface of wallboard for both heat insulation and fireproofing. Numerous household appliances, such as pot holders, mats, and table covers, may be classed in the group of heat-insulating materials.

Large quantities of the shorter grades of asbestos are used in the manufacture of building materials. Roofing shingles made of portland cement and asbestos are used extensively. Compressed sheets of asbestos combined with cement are used for corrugated sheeting, wallboard, millboard, lumber, and floor tile. Corrugated asbestos sheets have special merit in constructing chemical plants or other buildings exposed to a corrosive atmosphere. Asbestos-cement wall tiles in a wide range of colors were introduced about 1930. Asbestos-cement products are used increasingly in Europe as roofing, ceilings, partitions, paneling, linings of interior and exterior walls, water pipes, and gutters. It is claimed that the pipes and gutters are waterproof, require no painting, and compare favorably in strength with cast iron. It has been found that wooden flooring may be nailed to a subflooring consisting of concrete to which about 10 percent of asbestos has been added. Asbestos-composition products are becoming popular for floor tile in homes and on ships.

Miscellaneous uses are innumerable and diverse. There is a marked tendency toward a more extensive use of molded brake bands in automobiles. Of the brake bands made in 1929, about 84 percent were woven and 16 percent molded. In 1931, 70 percent were woven and 30 percent molded, and in 1933 these percentages were, respectively, 59 and 41. This trend indicates a relatively wider use of nonspinning asbestos at the expense of spinning grades. However, in

1935 there was a slight reverse and 39 percent molded.

Asbestos-cement pipe for gas mains and sewers is relatively nonconductive of so elastic that traffic vibrates over; they are easily placed; advantageous for carrying so rust. A standard European 20 percent asbestos.

Asbestos is used in fireproofing the surface of metal sheetings, for covering fire hose. A process has been perfected by means of metallurgical process will promote wider

Asbestos paper and mill Short-fiber asbestos mixed asphalt improves the surface

Asbestos is used as a pc suitable for common scour imparting a high luster.

In the chemical laboratories stoppings in combustion tubes and for wicks which when produce colored flames. are combined with the fib cloth of greater strength. a forcement permits the use as iron, Monel, nickel, and choice depending on the operating conditions.

Asbestos is used as a fil articles in which a b i bestos compounds are

The uses of anthophy strength. It is too weak for asbestos-cement shingle strength of the fiber imp of its superior resistance laboratories. The strong tough slip fiber occurring the most suitable for filter: Italian tremolite is also u: may be employed for oth previous paragraph.

The principal uses known cements to cover boilers, battery boxes, and as an acoustical and other wall for mineral wool in hot consistent market for it l never exceeded a few hun

134342*—37—2

suits, gloves, and shoes; aprons; mittens; gaskets; conveyor belts for carrying hot materials; Braided and twisted packings are

Asbestos is for compressed packings of fiber combined with rubber or of various forms known as compressed pressure packings.

Asbestos is for heat insulation; in fact, as important fireproofing and heat-insulating material, the best-known product is 85-percent asbestos paper is used for making 3-ply roofing; as sheeting between floors; lining bins, soldiers' helmets, mufflers for automobiles, sulphite mills, automobile radiator covers; table mats, pads, stove mats, and other in many other ways. Asbestos paper is used in manufacture of asbestos felt roofing and asbestos-protected metal roofing. Mill-building material but as a joint packing for safes, stoves, electric switch boxes, ovens, where a material of greater thickness is required.

An important use of millboard is for steam machinery. Asbestos is used for fireproof heat radiation. Large quantities of plastic fireproof cements for boiler, pipe, and asbestos-cement coating is sometimes applied for both heat insulation and fireproof appliances, such as pot holders, mats, and glassed in the group of heat-insulating

Higher grades of asbestos are used in the materials. Roofing shingles made of portland cement are used extensively. Compressed sheets of asbestos are used for corrugated sheeting, and floor tile. Corrugated asbestos is used in constructing chemical plants or other buildings. Asbestos-cement wall tiles were introduced about 1930. Asbestos is used increasingly in Europe as roofing, ceilings, and of interior and exterior walls, water pipes, and gutters are lined with asbestos, and compare favorably in strength with wooden flooring may be found that concrete to which about 10 percent asbestos-composition products are added in homes and on ships.

Asbestos is used in a number of other ways. There is a very extensive use of molded brake bands made in 1929, about 84 percent were made of asbestos. In 1931, 70 percent were woven and in 1933 these percentages were, respectively, indicates a relatively wider use of non-spun grades. However, in

1935 there was a slight reversal of the trend, 61 percent being woven and 39 percent molded.

Asbestos-cement pipes are being used increasingly for water and gas mains and sewers. They are resistant to acids, noncorrosive, relatively nonconductive of heat and electricity, light in weight, and so elastic that traffic vibrations are unlikely to cause leakage. Moreover, they are easily placed and joined. They are particularly advantageous for carrying solutions that must be kept free from iron rust. A standard European pipe consists of 80 percent cement and 20 percent asbestos.

Asbestos is used in fireproof paints, in arc welding, for protecting the surface of metal sheeting, as a constituent of asphalt roof coatings, for covering fire hose, and in the manufacture of gas burners. A process has been perfected whereby asbestos may be cemented to steel by means of metallic adhesives. Commercialization of this process will promote wider use of asbestos-protected metals.

Asbestos paper and millboard are used for electrical insulation. Short-fiber asbestos mixed in proportions of 4 to 6 percent with asphalt improves the surface of roads.

Asbestos is used as a polishing agent. A rough asbestos cloth is suitable for common scouring, and an acid-refined asbestos cloth for imparting a high luster.

In the chemical laboratory asbestos is used for acid filters, for stoppings in combustion tubes, for fireproof supports and protectors, and for wicks which when saturated with various mineral salts will produce colored flames. A filter cloth in which noncorrosive wires are combined with the fiber has an advantage over the all-asbestos cloth of greater strength, and also of lower cost because the wire reinforcement permits the use of lower grades of asbestos. Metals such as iron, Monel, nickel, and lead-coated steel are used for the wires, the choice depending on the nature of the solution to be filtered, and operating conditions.

Asbestos is used as a filler in insulators and various other molded articles in which a bituminous binder such as gilsonite is used. Asbestos compounds are used to some extent in automobile bodies.

The uses of anthophyllite are limited because of its lack of strength. It is too weak for spinning purposes, and it cannot be used for asbestos-cement shingles or other products to which a high tensile strength of the fiber imparts essential qualities. However, because of its superior resistance to acids, it is well suited for use in chemical laboratories. The stronger grades are used for chemical filters. A tough slip fiber occurring near Pylesville, Harford County, Md., is the most suitable for filter use of any yet found in the United States. Italian tremolite is also used for filters. Anthophyllite and tremolite may be employed for other chemical-laboratory uses mentioned in a previous paragraph.

The principal uses known for anthophyllite are for making plastic cements to cover boilers, pipes, and furnaces, as a filler in rubber and battery boxes, and as an admixture in cement and plastic flooring, acoustical and other wall plasters, and stucco. Its use as a substitute for mineral wool in house insulation is contemplated. No large, consistent market for it has yet been developed, and production has never exceeded a few hundred tons a year.

134342*—37—2

Many industries use asbestos in hundreds of minor ways. The catalogs of some asbestos-product companies contain lists of several hundred articles in which asbestos is a more or less essential ingredient.

From the foregoing enumeration of uses, it is apparent that large quantities of asbestos are employed in making brake bands, clutch facings, gaskets, packings, and insulation for automobiles. Therefore, fluctuations in automobile production are likely to be reflected in the amount of asbestos consumed. A second important field of utilization is for heat insulators, roofing, siding, millboard, wall-board, lumber, paper, and roof coatings, all of which are used extensively in the building trades. Asbestos consumption, therefore, is influenced greatly by the volume of building construction. Figure 2




FIGURE 2.—Asbestos consumption compared with automobile production and building construction, 1926-36. Unlike units are reduced to percentages of the 1923-25 average. Statistics of asbestos are from the Bureau of Mines, automobiles from the U. S. Bureau of the Census, and building contracts from the Federal Reserve Board.

shows the relation that exists between asbestos consumption and these two important industrial activities.

The chart indicates that during the depression years 1930 to 1933 asbestos consumption stood at a relatively higher level than either automobile production or building construction. Doubtless, this is due to the fact that during these years asbestos building materials were used in considerable quantities for the repair and maintenance of existing buildings and that a substantial volume of brake-band linings and clutch facings was used to recondition old cars and trucks.

GENERAL DISTRIBUTION

The United States has never attained importance as a producer of asbestos, the domestic output being only 1½ to 4 percent of the quantity required to supply raw material for its extensive asbestos-product manufacturing industries. Production is centered in four principal countries—Canada, Rhodesia, Union of South Africa, and

the Soviet Union. Since commercial commodity the P nage produced, but r ince is of the shortc of South Africa and Rh as sources of the higher War. Russia ranked next during the revolutionary almost at a standstill. each year since then pro Smaller quantities of Finland, and several oth Chrysotile is the lead but substantial quantitie also are produced in the of anthophyllite and su United States, Italy, Bu

PRODU

HISTORY OF P


Scattered deposits in nished a limited tonnag supplied small quantiti fiber has been produced duced principally in A contain considerable re are not favorable for and costly. Arizona h some interruptions sinc but furnish short fiber intermittently since 19

The following table to the United Sta the United States have since 1880.

Asbestos

Year	Short tons	Value
1880.....	150	\$4,312
1881.....	200	7,000
1882.....	1,200	35,000
1883.....	1,000	30,000
1884.....	1,000	30,000
1885.....	300	9,000
1886.....	200	6,000
1887.....	150	4,500
1888.....	100	3,000
1889.....	30	1,800
1890.....	71	4,560
1891.....	66	3,960
1892.....	104	6,416
1893.....	50	2,500
1894.....	325	4,463
1895.....	795	13,525
1896.....	504	6,100
1897.....	580	6,450
1898.....	605	10,300

bestos in hundreds of minor ways. The product companies contain lists of several ch asbestos is a more or less essential meration of uses, it is apparent that large employed in making brake bands, clutch and insulation for automobiles. Therobile production are likely to be reflected consumed. A second important field of sulators, roofing, siding, millboard, wall-roof coatings, all of which are used exten-des. Asbestos consumption, therefore, is olume of building construction. Figure 2



Comparison with automobile production and building construction is shown in Figure 2. The data are reduced to percentages of the 1923-25 average. The Bureau of Mines, automobiles from the U. S. contracts from the Federal Reserve Board.

between asbestos consumption and these activities.

During the depression years 1930 to 1933 at a relatively higher level than either building construction. Doubtless, this is these years asbestos building materials activities for the repair and maintenance at a substantial volume of brake-band was used to recondition old cars and

L DISTRIBUTION

It attained importance as a producer of asbestos being only 1½ to 4 percent of the raw material for its extensive asbestos industries. Production is centered in four Rhodesia, Union of South Africa, and

the Soviet Union. Since asbestos first became an important commercial commodity the Province of Quebec, Canada, has led in tonnage produced, but a large proportion of the output of this Province is of the shorter, nonspinning grades. Since 1916, the Union of South Africa and Rhodesia have become increasingly important as sources of the higher-grade spinning fibers. Before the World War, Russia ranked next to Canada as a producer of asbestos, but during the revolutionary period the industry in that country was almost at a standstill. In 1926, however, activity was revived, and each year since then production and exports have increased greatly. Smaller quantities of asbestos are produced in Cyprus, Australia, Finland, and several other countries.

Chrysotile is the leading product in all the producing countries, but substantial quantities of blue (crocidolite) and amosite asbestos also are produced in the Union of South Africa. Minor quantities of anthophyllite and similar amphibole varieties are mined in the United States, Italy, Bulgaria, Finland, and several other countries.

PRODUCTION AND CONSUMPTION

HISTORY OF PRODUCTION IN THE UNITED STATES

Scattered deposits in California and other Western States furnished a limited tonnage of asbestos many years ago. Georgia has supplied small quantities of anthophyllite, and the same or similar fiber has been produced at times in other States. Chrysotile is produced principally in Arizona and Vermont. The Arizona deposits contain considerable reserves of high-grade spinning fiber, but they are not favorable for low-cost mining, and transportation is difficult and costly. Arizona has produced a small tonnage annually with some interruptions since 1913. The Vermont deposits are extensive but furnish short fiber almost exclusively. Vermont has produced intermittently since 1908.

The following table, compiled from data reported by producers to the United States Geological Survey until 1924 and thereafter to the United States Bureau of Mines, shows the history of production since 1880.

Asbestos sold in the United States, 1880-1936

Year	Short tons	Value	Year	Short tons	Value	Year	Short tons	Value
1880	150	\$4,312	1899	681	\$11,740	1918	998	\$118,657
1881	200	7,000	1900	1,054	16,310	1919	1,161	248,265
1882	1,200	26,000	1901	747	13,498	1920	1,648	673,231
1883	1,000	30,000	1902	1,005	16,200	1921	531	356,963
1884	1,000	30,000	1903	867	16,760	1922	67	10,120
1885	300	9,000	1904	1,450	25,740	1923	227	9,626
1886	200	6,000	1905	3,109	42,975	1924	300	42,526
1887	150	4,500	1906	1,695	28,365	1925	1,255	51,700
1888	100	3,000	1907	653	11,399	1926	1,338	154,731
1889	30	1,800	1908	936	19,624	1927	2,961	336,582
1890	71	4,560	1909	3,085	62,603	1928	2,239	331,178
1891	66	3,960	1910	3,693	68,357	1929	3,155	351,004
1892	104	6,416	1911	7,604	119,935	1930	4,242	259,284
1893	50	2,500	1912	4,403	87,959	1931	3,228	115,267
1894	325	4,453	1913	1,100	11,000	1932	3,559	105,292
1895	795	13,525	1914	1,247	18,965	1933	4,745	130,677
1896	504	6,100	1915	1,731	76,952	1934	5,087	158,347
1897	350	6,450	1916	1,638	180,994	1935	8,920	292,927
1898	605	10,300	1917	1,938	291,014	1936	11,012	309,994

The following table shows sales for a series of years by varieties of fiber. For the period 1926 to 1928, inclusive, and, again, subsequent to 1931, the number of operators reporting was so small that separation by varieties could not be done without revealing individual figures.

Asbestos sold by producers in the United States, 1921-31, by varieties

Year	Chrysotile		Amphibole		Total	
	Short tons	Value	Short tons	Value	Short tons	Value
1921.....	438	\$313,266	393	\$23,700	831	\$336,966
1922.....	25	3,320	42	6,500	67	10,120
1923.....	69	4,433	155	5,193	227	9,626
1924.....	173	33,941	127	8,885	300	42,826
1925.....	93	40,750	1,165	10,950	1,258	51,700
1926.....	(1)	(1)	(1)	(1)	1,358	134,730
1927.....	(1)	(1)	(1)	(1)	2,951	139,900
1928.....	(1)	(1)	(1)	(1)	2,239	351,100
1929.....	1,963	317,584	1,172	33,420	3,135	351,004
1930.....	3,633	273,292	569	15,992	4,242	289,284
1931.....	2,537	111,706	371	7,259	3,228	118,965

¹ Bureau of Mines not at liberty to publish figures.

WORLD PRODUCTION

The following table, compiled by the Foreign Mineral Service Division, United States Bureau of Mines, shows production of asbestos by countries for 5 years.

World production of asbestos, 1931-35, in metric tons¹

[Compiled by M. T. Latus, Foreign Mineral Service Division]

Country	1931	1932	1933	1934	1935
Argentina.....		17			
Australia.....					
New South Wales.....	8	20	13		37
South Australia.....	6				
Western Australia.....	116	112	270	157	145
Bulgaria.....				3	3
Canada.....	149,047	111,552	143,667	141,502	190,931
China.....	264	250	236	290	(1)
Chosen.....			12	4	6
Cyprus.....	3,628	1,626	4,640	7,712	7,634
Czechoslovakia.....	(1)	(1)	1,200	2,100	2,600
Finland.....	551	756	1,340	1,735	1,742
France.....	500	300	400	400	(1)
Greece.....	10	9	14	30	(1)
India, British.....	6	91		25	61
Italy.....	632	1,264	3,267	2,252	(1)
Japan.....	1,000	1,000	1,000	1,000	1,000
Southern Rhodesia.....	21,810	14,303	27,361	29,224	35,644
Swaziland.....		5			
Turkey.....	4	58	120	4	104
Union of South Africa.....	14,221	10,950	14,411	15,960	20,653
U. S. S. R. (Russia).....	64,674	59,800	71,700	92,200	93,509
United States (sold or used by producers).....	2,925	3,229	4,305	4,615	8,092

¹ In addition to countries listed, Bolivia, Brazil, and Madagascar are reported to produce small quantities of asbestos.

² Rail and river shipments.

³ Exclusive of sand and gravel, production of which is reported as follows: 1931, 6,540 tons; 1932, 3,151 tons; 1933, 5,850 tons; 1934, 4,238 tons; 1935, 2,744 tons.

⁴ Data not available.

⁵ Exports.

⁶ Approximate production.

These figures include both long and short fibers of all varieties of asbestos. The output in Rhodesia and the Union of South Africa

consist
fil.
fil.
the otl
that i
has si
the ea
the sl
ducti-
Th
show
perie
stant

metric tons

F1

a

1:

1:

t

vs sales for a series of years by varieties 1926 to 1928, inclusive, and, again, subse- of factors reporting was so small that id be done without revealing indi-

in the United States, 1921-31, by varieties

Sotile	Amphibole		Total	
	Value	Short tons	Value	Short tons
\$313,268	393	\$23,700	531	\$336,968
3,320	42	6,800	67	10,120
4,433	158	5,193	227	9,625
33,941	127	8,585	300	42,525
40,750	1,165	10,950	1,228	31,700
(0)	(0)	(0)	1,338	134,731
(0)	(0)	(0)	2,981	336,852
(0)	(0)	(0)	2,239	351,173
317,584	1,172	33,420	3,155	351,004
273,292	580	15,992	4,242	259,284
111,708	371	7,259	3,223	115,967

h figures.

ILD PRODUCTION

compiled by the Foreign Mineral Service au of Mines, shows production of asbes-

asbestos, 1931-35, in metric tons¹

tus, Foreign Mineral Service Division]

	1931	1932	1933	1934	1935
	8	20	13	157	25
	116	112	270	157	143
	149,047	111,582	143,657	141,502	190,331
	264	250	236	200	(1)
	3,628	1,226	4,640	7,712	7,634
	(1)	(1)	1,200	2,100	2,000
	381	736	1,540	1,733	1,742
	300	300	400	4,400	(1)
	10	9	14	30	(1)
	6	91	25	25	41
	432	1,254	3,257	2,232	(1)
	7,400	1,000	1,000	1,000	1,000
	21,510	14,303	27,381	29,224	35,644
	5	5	120	4	104
	14,221	10,950	14,411	15,960	20,653
	44,474	39,800	71,700	92,300	95,500
	2,925	3,229	4,305	4,615	8,092

razil, and Madagascar are reported to produce small quantities

of which is reported as follows: 1931, 6,540 tons; 1932, 3,151 tons; 1933, 1,000 tons.

long and short fibers of all varieties of odesia and the Union of South Africa

consists principally of spinning fiber and the better grades of mill fiber, because the local demand is small and the value of the very short fibers too low to justify shipment to foreign markets. Russia, on the other hand, has developed important asbestos-products industries that utilize large quantities of short fiber from the Urals. Canada has small local demand for short fiber, but the industrial centers of the eastern United States provide a ready market, and for this reason the shorter grades constitute a large proportion of the asbestos production.

The output of most countries reached a low level in 1932 but has shown a rapid increase since that date. The Russian industry experienced only a minor decline during the depression, and a substantial increase in output from year to year since 1932 is indicated.




FIGURE 3.—Asbestos production in the three leading foreign centers and in the United States, 1916-35.

Figure 3 shows production in the three leading foreign centers and in the United States for a 20-year period. Canadian production indicated in the chart does not include the lowest grade product designated "sand, gravel, and stone (waste rock only)." For other districts the entire marketed production is included.

The minor importance of the United States among world producers, the leading position of Canada, and the recent rapid gain in prominence of Russia are apparent. On the basis of value of output, Africa and Russia hold a relatively stronger position than is indicated on the chart, because Canada produces a preponderance of the lower-priced short fibers.

WORLD CONSUMPTION

It is more difficult to determine the consumption of asbestos than its production. Production plus imports minus exports of raw asbestos for any year may not give a true picture of consumption, because stocks may be held over for some time and reexported or material consumed may consist of stocks imported in a previous year. The

following table, taken from Asbestos,¹¹ furnishes interesting data on consumption. The figures were compiled by "Tropag" Asbest- und Erzimport G. m. b. H., Hamburg, Germany, from data supplied by the ministerial and consular offices of the countries concerned.

World consumption of asbestos, 1931-33 (long tons)

Consumption	1931	1932	1933	Consumption	1931	1932	1933
United States.....	121,539	86,311	106,755	Poland.....	321	342	529
Canada.....	14,800	15,000	16,500	Norway.....	1,000	794	987
Russia.....	140,000	144,000	145,000	Sweden.....	968	1,193	1,200
Japan.....	7,300	7,650	9,000	Estonia.....	63	21	32
England.....	19,561	17,404	24,386	Finland.....	62	42	68
France.....	10,000	16,000	14,354	Latvia.....	59	35	135
Belgium.....	12,609	5,190	11,239	Lithuania.....	80	11	150
Italy.....	5,134	3,781	5,000	Jugoslavia.....	1,292	474	303
Spain.....	3,277	4,132	6,059	Rumania.....	65	65	145
Portugal.....	58	160	160	Australia.....	1,300	1,400	1,750
Netherlands.....	266	157	522	India.....			
Denmark.....	705	736	1,750	Africa.....	1,200	1,100	1,750
Germany.....	9,808	7,581	12,613	South America and other countries.....			
Austria.....	5,016	3,466	1,267	Total, long tons.....	253,893	205,179	255,095
Hungary.....	2,394	1,597	781	Total, metric tons.....	257,955	208,462	259,177
Czechoslovakia.....	4,678	7,044	2,208				
Switzerland.....	1,145	563	832				

¹ Estimated.

The difference in total consumption for 1933, as given in this table (259,177 metric tons), and total world production in that year, as compiled by the United States Bureau of Mines (276,140 metric tons), may be explained by variations in stocks and materials in transit.

The following table presents a compilation of apparent world consumption for 1934 based on production plus imports minus exports. Because of variations in stocks, the figures are approximate only. For instance, Canadian consumption as shown in this table is a negative quantity because, with a steadily increasing market demand, exports were drawn from stocks as well as from current production. Canadian consumption ranges from 3,000 to 6,000 tons a year. As indicated in the world totals, consumption balances reasonably well with production.

World consumption of asbestos, 1934

[In metric tons]

Continent and country	Production ¹	Imports ²	Exports ³	Apparent consumption
Asia:				
China.....	290	554		574
Chosen.....	4	90		94
Cyprus.....	7,712		7,712	
India, British.....	25	965		991
Japan.....	1,000	20,515		21,515
Manchuria.....		1,291	95	1,195
Netherland India.....		14		14
Turkey.....	4			4
Total, Asia.....	9,035	23,460	7,808	24,687

¹ Compiled by Foreign Mineral Service Division, U. S. Bureau of Mines.

² Figures as given in The Mineral Industry of the British Empire and Foreign Countries, Statistical Summary, published by The Imperial Institute, London.

³ Exports.

⁴ Approximate production.

¹¹ Asbestos, vol. 15, no. 12, June 1934, p. 16.

Cont

Africa:
Algeria.....
Egypt.....
Southern Rhodesia.....
Union of South Africa.....
Total, Africa.....
North America:
Canada.....
United States.....
Total, North America.....
South America:
Brazil.....
Chile.....
Colombia.....
Total, South America.....
Australia.....

Europe:
Austria.....
Belgium.....
Luxembourg.....
Czechoslovakia.....
Denmark.....
Estonia.....
Finland.....
France.....
Germany.....
Greece.....
Hungary.....
Italy.....
Latvia.....
Netherlands.....
Norway.....
Poland.....
Portugal.....
Rumania.....
Spain.....
Sweden.....
Switzerland.....
U. S. S. R.....
United Kingdom.....
Yugoslavia.....

Total, Europe.....

Total by continent:
Asia.....
Africa.....
North America.....
South America.....
Australia.....
Europe.....
Total, world.....

¹ Estimate

Asbestos in far n develop The tween 1

from Asbestos,¹ furnishes interesting data on figures were compiled by "Tropag" Asbest- und L. burg, Germany, from data supplied by offices of the countries concerned.

Consumption of asbestos, 1931-33 (long tons)

1932	1933	Consumption	1931	1932	1933
26,311	106,755	Poland.....	321	342	529
3,000	6,500	Norway.....	1,000	784	867
4,000	48,000	Sweden.....	968	1,193	1,200
7,630	9,000	Estonia.....	63	21	32
17,404	24,384	Finland.....	62	42	68
8,000	14,354	Latvia.....	50	35	135
1,190	11,239	Lithuania.....	80	11	50
3,791	5,000	Jugoslavia.....	1,222	474	303
1,132	6,059	Rumania.....	65	65	165
1,60	1,60	Australia.....	1,300	1,400	1,750
257	822	India.....			
736	1,730	Africa.....			
7,581	12,613	South America and other countries.....	1,100	1,100	1,750
1,466	1,267	Total, long tons.....	253,893	295,179	255,095
1,397	781	Total, metric tons.....	257,955	298,462	259,177
7,044	2,209				
563	832				

consumption for 1933, as given in this table and total world production in that year, as States Bureau of Mines (276,140 metric by variations in stocks and materials in

presents a compilation of apparent world used on production plus imports minus ex- tions in stocks, the figures are approximate adian consumption as shown in this table is use, with a steadily increasing market de- n from stocks as well as from current pro- mption ranges from 3,000 to 6,000 tons a e totals, consumption balances rea- tion.

Consumption of asbestos, 1934

[In metric tons]

	Production	Imports	Exports	Apparent consumption
	290	384		574
	4	90		94
	7,712		7,712	91
	25	966		991
	1,000	20,515		21,515
		1,291	96	1,195
		14		14
	4			4
	9,035	23,460	7,808	24,687

Division, U. S. Bureau of Mines.
istry of the British Empire and Foreign Countries, Statistical
stitute, London.

334, p. 16.

World consumption of asbestos, 1934—Continued

[In metric tons]

Continent and country	Production	Imports	Exports	Apparent consumption
Africa:				
Algeria.....		37		37
Egypt.....		97		97
Southern Rhodesia.....	29,224		29,085	139
Union of South Africa.....	15,960		15,515	145
Total, Africa.....	45,184	134	44,900	418
North America:				
Canada.....	141,502		143,557	-2,055
United States.....	4,615	109,165	1,514	112,266
Total, North America.....	146,117	109,165	145,071	110,211
South America:				
Brazil.....		38		38
Chile.....		37		37
Colombia.....		53		38
Total, South America.....		103		103
Australia.....	157	2,511		2,668
Europe:				
Austria.....		1,173	96	1,092
Belgium-Luxemburg.....		10,797	516	9,981
Bulgaria.....	3			6
Czechoslovakia.....	2,100	4,938	354	6,484
Denmark.....		1,039		1,039
Estonia.....		23		23
Finland.....	1,735	65		1,820
France.....	1,406	10,220		10,626
Germany.....		20,154	190	19,964
Greece.....	30	76		166
Hungary.....		627		627
Italy.....	2,252	8,421	505	10,168
Latvia.....		22		22
Netherlands.....		258		258
Norway.....		51		51
Poland.....		910		910
Portugal.....		345		315
Rumania.....		122		122
Spain.....		4,702		4,702
Sweden.....		1,703		1,703
Switzerland.....		733		733
U. S. S. R. (Russia).....	92,461	116	33,513	55,764
United Kingdom.....		26,972	425	26,547
Yugoslavia.....		580		580
Total, Europe.....	98,987	95,061	36,339	157,659
Total by continents:				
Asia.....	9,035	23,460	7,808	24,687
Africa.....	45,184	134	44,900	418
North America.....	146,117	109,165	145,071	110,211
South America.....		103		103
Australia.....	157	2,511		2,668
Europe.....	98,987	95,061	36,339	157,659
Total, world.....	299,450	230,434	234,168	295,745

¹ Estimate by Imperial Institute, London.

DOMESTIC DEPOSITS

CHRYSTOLE

ALASKA

Asbestos has been found near Shungnak and in the Jade Hills in far northwestern Alaska, and several attempts have been made to develop the deposits.

The Shungnak occurrences are close to the Kobuk River, between 150 and 200 miles from its mouth. According to Smith and

Mertie.¹² the asbestos occurs in small veins in close association with greenstone and serpentine. The veins, which are most abundant on the east side of Dahl Creek, consist mostly of short fiber, although some long fiber has been found. Samples of the latter were submitted to manufacturers some years ago, and they reported that the color was good but the fibers were too weak for high-grade uses. However, samples of strong, flexible chrysotile more than 6 inches long and apparently of very high grade were obtained in 1932 from a deposit in this locality. The asbestos appears to be of the slip-fiber type; therefore, it may be confined to shear zones. Stewart¹³ states that three thin veins, the largest of which is about 3 inches wide, were uncovered at one point, and that some veins have been traced for at least a quarter of a mile. The deposit is undeveloped. Water transportation is available during the summer months.

The Jade Hills lie north of the Kobuk River not far from its mouth. The asbestos deposits in these hills were worked in a small way for a short time many years ago, and in 1925 and 1926 interest in them was revived and a trader at Kotzebue financed some new prospecting in the region. No definite information is available as to the kind or quality of asbestos found, but it probably is similar to that occurring at Shungnak.

ARIZONA

Asbestos was first discovered in Arizona by Charley Newton in 1872. He, with a small force of pioneers, fought a band of Indians, and during the engagement came upon an asbestos outcrop 3 miles northeast of Chrysotile on Ash Creek, Gila County, where the largest asbestos mine in Arizona is now located. No interest was taken in the deposit for many years. About 1900, chrysotile asbestos was discovered in the Grand Canyon and small quantities were mined on the north side opposite Grand View, about 20 miles west of Grand View, and at other points. After these discoveries interest reverted to the Gila County occurrences, and in 1912 and 1913 many claims were staked. Asbestos was found in numerous localities over an area 60 miles long and 25 miles wide, most of which lies west of the San Carlos and Apache Indian Reservations.

The asbestos occurs in the Apache formation, which, in the Salt River region, is represented chiefly by beds of quartzite and limestone. It apparently originated as an alteration product of the Mescal (Cambrian) dolomitic limestone due to metamorphism by diabase intrusions. It is associated with greenish and mottled serpentine. It occurs in cross-fiber veins, which are most abundant throughout the limestone close to the contact surfaces. Places where the limestone is much broken by the diabase intrusions are particularly favorable for the formation of asbestos. The veins reach a maximum thickness of about 6 inches. The fibers may extend across the full width of the vein or they may be broken into two or more lengths. The veins are erratic and are not persistent for long distances. Intervals between them on the vertical face commonly vary

¹² Smith, Philip S., and Mertie, J. B., Jr., *Geology and Mineral Resources of Northwestern Alaska*: U. S. Geol. Survey Bull. 815, 1930, pp. 344-345.

¹³ Stewart, B. D., *Mining Investigations and Mine Inspection in Alaska: Biennium Ending Mar. 31, 1933*, pp. 21, 22.

Bureau of Mines



FIG

ASBESTOS

occurs in small veins in close association with quartz. The veins, which are most abundant on the west side of the hills, consist mostly of short fiber, although some samples of the latter were submitted for analysis, and they reported that the color was gray, and they reported that the color was too weak for high-grade uses. How- ever, flexible chrysotile more than 6 inches long high grade were obtained in 1932 from a vein. The asbestos appears to be of the slip-fiber type, and that some veins have been confined to shear zones. Stewart¹³ reports, the largest of which is about 3 inches in diameter, and that some veins have been traced for a mile. The deposit is undeveloped. It is available during the summer months. It is located on the Kobuk River not far from its mouth in these hills were worked in a small way years ago, and in 1925 and 1926 in- and a trader at Kotzebue financed some work. No definite information is available on asbestos found, but it probably is similar to that found in Alaska.

ARIZONA

Discovered in Arizona by Charley Newton in 1891, one of the pioneers, fought a band of Indians. He came upon an asbestos outcrop 3 miles from the mouth of the Gila River, where the largest deposit is now located. No interest was taken in it at the time. About 1900, chrysotile asbestos was discovered in the same place, and small quantities were mined and sold. After these discoveries interest reverted to the discovery, and in 1912 and 1913 many claims were made, and numerous localities over an area of 10 miles wide, most of which lies west of the Indian Reservations.

The Apache formation, which, in the Salt River Valley, is chiefly by beds of quartzite and limestone, is altered as an alteration product of the limestone due to metamorphism by the associated with greenish and mottled slip-fiber veins, which are most abundant near the contact surfaces. Places where the veins are intersected by the diabase intrusions are particularly rich in asbestos. The veins reach a thickness of 1 to 2 inches. The fibers may extend across the veins, but they may be broken into two or more pieces and are not persistent for long distances. On the vertical face commonly vary

Jr., *Geology and Mineral Resources of Northwest Alaska*, 1930, pp. 344-345.
Bureau of Mines, *Geological Survey of Alaska: Biennial Report*

Bureau of Mines

Bulletin 403




FIGURE 4.—Crude chrysotile asbestos from Arizona. ("Asbestos.")

from 2 to 6 feet. Conditions are most favorable for mining where several veins are close enough to each other to be worked from one tunnel. Usually much barren rock must be mined to recover the fiber-bearing seams. The occurrences are quite different from the Canadian deposits, where fiber veins traverse massive serpentine irregularly.

The highest-grade fiber, shown in figure 4, is soft and silky, has high tensile strength, and is excellent for spinning purposes. A harsher, less desirable fiber, however, is associated with the soft in many places. Arizona chrysotile is very low in iron, therefore it is generally considered more suitable for electrical insulation than chrysotile from other regions. According to Melhase,¹⁴ the asbestos deposits of Gila County may be grouped in five well-defined districts, as follows:

1. Mogollon Rim district at the northern boundary of the county. It contains some interesting occurrences that are too remote to encourage development.

2. Cherry Creek district, comprising about 75 square miles along the canyon of Cherry Creek. The southern end of the district is about 25 miles north of Globe. Considerable mining has been done, notably at the Riga, Triangle, and Aileen properties.

3. Sierra Ancha district, comprising about 60 square miles adjoining the western boundary of the Cherry Creek district. The principal properties in this area are the Clark, Riga, and American ores.

4. Chrysotile district, including about 100 square miles on both sides of Salt River. Chrysotile, where the Johns-Manville mine, the largest asbestos mine in the State, is located, lies 24 miles north-northeast of Globe. Other important properties are the Regal, the Arizona Asbestos Co., and the Bear Canyon Asbestos Co. More than a dozen other properties have been developed to varying degrees.

5. Globe district, covering about 60 square miles, extending from Copper Hill, 4 miles north of Globe, eastward toward the San Carlos River. The Apache mine is the most important in this area.

The location of the deposits, most of which are many miles from a railroad, and the long and costly rail haul to eastern markets have confined mining chiefly to the highest-grade crude fibers. Haulage charges from most of the mines in the Chrysotile district, the most productive area in Arizona, to Globe range from \$22 to \$35 a ton. Very few of the properties have produced during recent years. Production began in 1913 and has never exceeded a few hundred tons a year, attaining a maximum of 1,200 tons in 1920. A new highway, constructed in 1933, from Springerville to Globe lowers the cost of transportation and may encourage enlarged development.

CALIFORNIA

Asbestos occurrences in California are numerous but the history of their development has been disappointing. The first reference to chrysotile in the Mineral Resources chapters of the United States Geological Survey was in 1901. In that year a small production was

¹⁴Melhase, John. Asbestos Deposits in Arizona: Eng. and Min. Jour. Press, vol. 120, no. 21, Nov. 21, 1923, pp. 805-10.

reported in Riverside County and occurrence in Calaveras County was mentioned. A small output was reported from Placer County in 1905; but, although several new deposits were recorded between 1906 and 1911, none of them bore promise of a large supply. Slip fiber was found in Sierra County in 1912. Samples of chrysotile obtained in Shasta and Napa Counties in 1913 were not of promising quality. Up to 1914, asbestos had been reported in 12 counties. Production in Shasta, Placer, Calaveras, and Alameda Counties aggregated 65 tons in 1914. In 1915, a manufacturer of asbestos pipe covering and shingles at Oakland reported purchase of 80 tons of fiber in Placer, Alameda, and Contra Costa Counties. Three carloads of nonspinning fiber were shipped to Oakland from near Sims in Shasta County in 1916. The asbestos is said to occur in cross-fiber veins, in serpentine, which would indicate that it is chrysotile, but shipments from this locality in 1922 were reported as amphibole.

The most productive deposit in California is $2\frac{1}{2}$ miles northwest of Washington, Nevada County, where the fiber occurs in a large area of serpentine. Mining was begun about 1917. A 300-foot tunnel was driven into a serpentine hill on a 6-percent upgrade and a 150-foot raise was put through to the surface. The rock was mined by a glory-hole method, dropped through the raise, and trammed through the tunnel. A mill having a capacity of 100 tons a day was equipped with stamps, screens, fiberizers, and air-suction pipes. Short fiber and a small quantity of spinning fiber were produced, but activity ceased with the sale in 1923 of a small tonnage of fiber milled in 1921. Much of the output was used in the manufacture of flooring in San Francisco. A deposit that bears promise of commercial importance was reported in the Chicago Park district in 1937.

Another property that attained some prominence is about 6 miles east of Copperopolis, Calaveras County, in a serpentine belt having a maximum width of 2,700 feet. Prospecting and developing conducted in 1920 and 1921 revealed a commercially mineralized area about 1,400 feet long and 400 feet wide. In some places the serpentine rock carries 12 to 15 percent asbestos in cross-fiber veins, but the average of workable rock is much lower. A 5-percent recovery is recorded for one period of mill operation. The fiber is of paper-stock grade or shorter. A small crushing and fiberizing mill, capable of handling about 30 tons of ore a day, began operation in 1921. Underground workings consist of several tunnels, the longest of which was driven 278 feet into the hill. Very little fiber was produced, and no activity has been reported since 1923.

A deposit was developed about 1920 and a mill built in 1921 on a property $2\frac{1}{2}$ miles from Hernandez in southern San Benito County. A small tonnage of fiber was produced from 1923 to 1926, and thereafter activity ceased.

A large asbestos deposit was reported in Napa County, and plans were under way for a \$20,000 mill in 1931, but the project did not materialize. Small quantities of asbestos were produced 9 miles southwest of Monticello in 1932, and a mill was built in 1933.

In 1931, asbestos occurring in irregular veins in dark-green serpentine was mined in an open-cut near Carrville, Trinity County. Several tons of chrysotile of good quality and averaging $\frac{3}{4}$ inch in length were obtained.

In 1928, 7 or 8 tons of asbestos were produced near Lake County.

Asbestos of good quality was produced in 1915 and 1919 at a locality northwest of Coalinga.

Small quantities were produced in 1917, and near Vallecito, Alameda County in 1918.

Asbestos deposits have been reported 31.5 miles from Cisco, Placer County; Edgewood, Siskiyou County; and near Vallecito, Alameda County. The maximum production in the State mineralogist in 1923 was 190 tons. The maximum production in the State mineralogist in 1923 was 190 tons. The maximum production in the State mineralogist in 1923 was 190 tons.

Asbestos product

Year	Tons	Value	Year
1887	30	\$1,500	1903
1888	30	1,500	1904
1889	30	1,500	1905
1890	7	4,200	1906
1891	65	3,900	1907
1892	20	1,500	1908
1893	50	2,500	1909
1894	50	2,500	1910
1895	25	1,000	1911
1896			1912
1897			1913
1898	10	200	1914
1899	30	750	1915
1900	50	1,200	1916
1901	110	4,400	1917
1902			1918

Annual details concealed under "unappor"

A deposit of asbestos near (Idaho line, was developed to the west of the chrysotile occurs with section to that found in Arizona associated with diabase into States Bureau of Mines in 1915. The fiber is brittle than fiber from Canada 1923 and additional machinery has been almost negligible.

Chrysotile asbestos, apparently from Malheur County.

The Montebello Co. (formerly the Montebello Co.) operated the mine in 1915 and 1916, but was permitted to buyers during that time.

²⁵ California Mineral Production and Statistics, California Div. of Mines Bull. 111.

ntly and occurrence in Calaveras County output was reported from Placer County areas. Deposits were recorded between 1887 and 1912. Slip County in 1912. Samples of chrysotile in Calaveras County in 1913 were not of promising quality. Asbestos had been reported in 12 counties. In 1915, a manufacturer of asbestos pipe in Oakland reported purchase of 80 tons of asbestos from Contra Costa County. Three cars were shipped to Oakland from near Sims.

The asbestos is said to occur in cross-bedded rock which would indicate that it is chrysotile. In 1922 were reported as amphibole. A deposit in California is 2½ miles northwest of San Francisco, where the fiber occurs in a large quantity. A 300-foot deep shaft was begun about 1917. A 6-percent upgrade and through to the surface. The rock was crushed, dropped through the raise, and a mill having a capacity of 100 tons per day. Small quantities of spinning fiber were produced with the sale in 1923 of a small tonnage of the output was used in the manufacture of paper in San Francisco. A deposit that bears promise was reported in the Chicago Park dis-

covered some prominence is about 6 miles northwest of San Francisco, in a serpentine belt having a width of 100 feet. Prospecting and developing have revealed a commercially mineralized area 400 feet wide. In some places the serpentine contains 15 percent asbestos in cross-fiber veins, but in other places the rock is much lower. A 5-percent concentration of mill operation. The fiber is of good quality. A small crushing and fiberizing plant producing 30 tons of ore a day, began operation in 1923. The mine consists of several tunnels, the longest 278 feet into the hill. Very little fiber has been reported since 1923.

about 1920 and a mill built in 1921 on the property. In 1926, a mill was built in southern San Benito County. Fiber was produced from 1923 to 1926, but no output was reported.

was reported in Napa County, and plans for a 1,000 mill in 1931, but the project did not develop. Asbestos was produced 9 miles northwest of San Francisco, and a mill was built in 1933. The mine is in irregular veins in dark-green serpentine near Carrville, Trinity County. The fiber is of good quality and averaging ¾ inch in

In 1928, 7 or 8 tons of asbestos were obtained 10 miles from Middletown, Lake County.

Asbestos of good quality was reported from a deposit 27 miles northwest of Coalinga, Fresno County, in 1918. A small tonnage produced in 1918 and 1919 was sold in 1925.

Small quantities were produced 2½ miles east of Ione, Amador County, in 1917, and near Valley Springs, Calaveras County, and in Alameda County in 1918.

Asbestos deposits have been reported at Towle, Iowa Hill, and 3½ miles from Cisco, Placer County; at Goodyears Bar, Sierra County; Edgewood, Siskiyou County; and Tolon, Monterey County. The maximum production in any one year was 410 tons reported by the State mineralogist in 1921. The following table¹⁵ shows production in California, by years, since 1887. The figures in this table represent both chrysotile and amphibole. The entire output prior to 1900 was classed as amphibole.

Asbestos production in California, 1887-1935

Year	Tons	Value	Year	Tons	Value	Year	Tons	Value
1887	30	\$1,800	1903	10	\$162	1919	131	\$6,240
1888	30	1,800	1904	112	2,625	1920	410	19,273
1889	30	1,800	1905	70	3,500	1921	50	1,800
1890	71	4,260	1906	70	3,500	1922	20	300
1891	66	3,960	1907	70	6,100	1923	70	4,750
1892	30	1,830	1908	65	4,500	1924	25	1,550
1893	30	2,300	1909	200	20,000	1925		
1894	30	2,350	1910	125	500	1926		
1895	25	1,000	1911	90	2,700	1927		
1896			1912	47	1,175	1928	15	1,160
1897			1913	31	1,530	1929		
1898	10	200	1914	143	2,560	1930	219	6,175
1899	30	750	1915	145	2,350	1931		
1900	30	1,250	1916	136	10,225	1932		
1901	110	4,400	1917	229	9,903	1933	309	3,274
1902			1918			1934		
						1935		

¹ Annual details concealed under "unapportioned."

MONTANA

A deposit of asbestos near Cliff Lake, Madison County, close to the Idaho line, was developed to some extent between 1917 and 1923. The chrysotile occurs with serpentine in limestone, a similar association to that found in Arizona. The asbestos-bearing zone is closely associated with diabase intrusions. Samples sent to the United States Bureau of Mines in 1929 were found to be weaker and more brittle than fiber from Canada or Arizona. A mill was begun in 1923 and additional machinery was delivered in 1926, but production has been almost negligible.

OREGON

Chrysotile asbestos, apparently of good quality, is reported from Malheur County.

The Montbestos Co. (formerly United Asbestos Products Corporation) operated the mine in 1934 but made no sales. Samples submitted to buyers during that year were said to be satisfactory.

¹⁵ California Mineral Production and Directory of Mineral Producers for 1934: State of California Div. of Mines Bull. 111, August 1935, p. 78.

VERMONT

Although asbestos was discovered in Vermont in 1824, no interest was taken in it until M. E. Tucker found a chrysotile vein while felling trees on the eastern side of Belvidere Mountain in 1892 or 1893. Its resemblance to the Canadian fiber was recognized immediately, and it was later found to occur over a considerable area in Lamoille and Orleans Counties.

The Vermont fiber is chrysotile similar in character and occurrence to that found at Thetford Mines, Quebec, the deposit being regarded as a southward extension of the well-known Canadian belt. The asbestos is of two kinds—cross-fiber veins passing irregularly through serpentine and slip fiber lying parallel to slickensided surfaces. Most of the cross-fiber veins are narrow, those over three-fourths inch in width being uncommon. Keith and Bain,¹⁶ after a detailed study of the veins, decided that the asbestos had been introduced as a vein-filling material in torsion cracks and crush fractures. Matched structures on opposite walls of the veins are common. They found little evidence in support of a replacement theory. The slip fiber ranges from a fraction of an inch to 3 or 4 inches long and is somewhat coarser than the cross fiber; however, it is not suitable for spinning. The claim has been made that the proportion of fiber to rock runs 8 to 12 percent, which is somewhat greater than in the Canadian field. The yield of spinning fiber, however, is very small.

Asbestos-bearing rocks occur in four prominent localities in Lamoille and Orleans Counties. The first occurrence, which is undeveloped, is about 2 miles northeast of Lowell. It consists of numerous cross-fiber veins and some slip fiber in serpentine. The second deposit, northwest of Lowell in the vicinity of Westfield, is in the form of a belt several miles long and a mile wide. Considerable slip fiber and cross fiber occur in places, but none has been mined. In the third locality, which is southwest of Lowell near Mount Belvidere, slip fiber predominates. A mill was erected in this district in 1902 by the New England Asbestos Mining & Milling Co., but was operated only 6 months. The fourth serpentine area, extending about 2 miles eastward from the third, occupies a prominent bluff on the shoulder of Belvidere Mountain. Both slip-fiber and cross-fiber veins occur, the latter providing a small amount of crude asbestos. On this property, in 1907, the Lowell Lumber & Asbestos Co. erected a mill with a daily capacity of about 200 tons of rock. Production was first recorded in 1908, and in 1909 the industry was so active that Vermont became the chief asbestos-producing State. In 1910, the output increased 24 percent over that in 1909. In 1911 the total production of the United States, 7,604 tons, was largely from Vermont. During that year considerable no. 2 crude was obtained. In 1912, Vermont, with a production of 4,408 tons, had the distinction of being the only asbestos-producing State in the Union, but for several years thereafter no activity was noted.

The industry was revived in 1920 by the Asbestos Corporation of America but not in the location formerly most productive. Since then operations have been confined to the opposite side of the deposit

¹⁶ Keith, Stanton B., and Bain, George W., Chrysotile Asbestos. I. Chrysotile Veins: Econ. Geol., vol. 27, no. 2, 1922, pp. 169-188.

at the former q
Milling Co. Si
or no spinning
dently extensive
but there was
company was
Asbestos Corpo
the plant in ope
more than doub
kets led to some
has increased s
acquired by Ve
Vermont Asbes
and with plant
anticipated. In
asbestos. The
One important
employed, also,
ering, paint, at
the output in
asbestos-produce

A small pro
in 1892 and fi
area of approx
8 miles south
square miles 20
The mineral o
are more than
appears in plac
a small mill i
ore a day. A
little fiber has
tos Shingle C
Casper Mount
Occurrences
City, Fremont
have not been
land, Platte Co
other places, b
relatively low.

Although at
Carolina, chry
2 inches wide
in 1902. An
has been repor
of Spruce Pin
Jackson Count
County; and n

MONT

red Vermont in 1824, no interest in a chrysotile vein while fell-Belvidere Mountain in 1892 or 1893. A fiber was recognized immediately, over a considerable area in Lamaille

similar in character and occurrence Quebec, the deposit being regarded as a well-known Canadian belt. The veins passing irregularly through parallel to slickensided surfaces. Most are those over three-fourths inch in diameter. Bain, after a detailed study of the deposits had been introduced as a vein and crush fractures. Matched structures are common. They found little support for the slip fiber range theory. The slip fiber ranges from 1/4 to 1/2 inch long and is somewhat irregular, it is not suitable for spinning. A proportion of fiber to rock runs 3 to 1, greater than in the Canadian field. The fiber is very small.

In four prominent localities in Lamaille the first occurrence, which is under Lowell. It consists of numerous veins in serpentine. The second locality, of Westfield, is in the form of a mile wide. Considerable slip fiber has been mined. In the third locality near Mount Belvidere, slip fiber was erected in this district in 1902 by the Lowell & Co., but was operated in a small area, extending about 2 miles from a prominent bluff on the shoulder of the mountain. p-fiber and cross-fiber veins occur, but of crude asbestos. On this property the Lowell & Asbestos Co. erected a mill with the use of rock. Production was first in 1902. The industry was so active that Vermont was producing State. In 1910, the output was 909 tons. In 1911 the total production was largely from Vermont. During 1912, Vermont was obtained. In 1912, Vermont, the distinction of being the only producer, but for several years there-

20 by the Asbestos Corporation of Vermont formerly most productive. Since 1912 to the opposite side of the deposit

- Chrysotile Asbestos. I. Chrysotile Veins:

at the former quarry site of the New England Asbestos Mining & Milling Co. Slip fiber predominates in this region, there being little or no spinning fiber. The reserves of asbestos-bearing rock are evidently extensive. A mill was built and experimental runs were made but there was no commercial production until 1929. In 1928, the company was reorganized under new ownership as the Vermont Asbestos Corporation of America. The new company quickly put the plant in operation, producing 1,049 tons of mill fiber in 1929, and more than doubled that quantity in 1930. Although depressed markets led to some reduction in output thereafter, since 1932 production has increased substantially. In February 1936 the properties were acquired by Vermont Production Co., Inc., subsequently named the Vermont Asbestos Corporation, a subsidiary of The Ruberoid Co., and with plant additions and improvements an increase in output is anticipated. In 1934, the former operators produced 6 grades of asbestos. The longest grade was shingle stock testing 0-2-10-1. One important use is for manufacture of molded brake linings. It is employed, also, in asbestos-cement shingles, asbestos paper, pipe covering, paint, and boiler covering. The present owners use part of the output in their own products and sell the remainder to other asbestos-product manufacturers.

WYOMING

A small production of chrysotile was reported from Wyoming in 1892 and from 1906 to 1912. Asbestos occurs in a serpentine area of approximately 4 1/2 square miles on Casper Mountain about 8 miles south of Casper and in the Smith Creek area of about 7 square miles 20 miles southeast of Casper, both in Natrona County. The mineral occurs principally as cross-fiber veins, few of which are more than an inch across, but a minor quantity of slip fiber appears in places. In 1910, the Wyoming Consolidated Co. operated a small mill in the Smith Creek area, handling about 30 tons of ore a day. A mill was built on Casper Mountain in 1911. Very little fiber has been produced in either locality, but the Patee Asbestos Shingle Co. reported production of 200 tons in 1934 from its Casper Mountain property.

Occurrences of spinning-grade asbestos near Lander and Atlantic City, Fremont County, were reported in 1919, but the properties have not been developed. A small quantity was obtained at Wheatland, Platte County, in 1920. Chrysotile has been found in several other places, but most of the occurrences throughout the State are relatively low-grade, short fibers.

OTHER OCCURRENCES

Although amphibole is the prevailing type of asbestos in North Carolina, chrysotile also occurs in several places. Chrysotile veins 2 inches wide were found near North Wilkesboro, Wilkes County, in 1902. An open-cut 100 feet long was made, but no production has been reported. Other occurrences have been noted 8 miles west of Spruce Pine, in Yancey County; near Glenville and Sapphire, in Jackson County; on the western slopes of Rich Mountain, Watauga County; and near Elk Creek, Ashe County.

A deposit near Round Hill, Loudoun County, Va., was reported in 1894. Small stringers of asbestos also occur in massive serpentine near Great Falls, Fairfax County.

Chrysotile was found in 1902 near Ishpeming, Marquette County, Mich., and some development work was done.

AMPHIBOLE

ALASKA

Asbestos said to be of the amphibole variety occurs on Admiralty Island and at Chitina on the Copper River. A little development work has been done, but the low price of amphibole fiber discourages development so far from markets.

CALIFORNIA

As early as 1882 asbestos deposits were worked in San Bernardino, San Diego, Calaveras, and Placer Counties, and in 1883 and 1884 occurrences in Butte, Fresno, Los Angeles, Tulare, Mariposa, and Inyo Counties were noted and some of them were developed. Production in 1882 was 1,200 tons; in 1883, 1,000 tons; and in 1884, 1,000 tons. In 1885, a small mine was opened at Windsor in Sonoma County, and fiber of good quality was reported from Del Norte and Yolo Counties. Sales of asbestos in the United States of 30 tons in 1889, 71 tons in 1890, and 66 tons in 1891 were credited entirely to California. In 1892 most of the production was from Wyoming, with small quantities from Oregon and California, but again in 1893 California, with an output of 50 tons, was the only producing State. The mines were idle in 1894 but produced 90 tons in 1895. A small and gradually decreasing output was noted from 1897 to 1900. All the production up to the latter year was classed as amphibole asbestos. Evidently it was used chiefly in San Francisco and Oakland for making pipe and boiler coverings, packings, and roofing.

A slip-fiber anthophyllite was mined for several years near Sims, Shasta County, but no activity has been noted since 1923.

A deposit of amphibole asbestos associated with serpentine near Chandlers, Kern County, was developed in 1928. Several short tunnels and open-cuts were made, but evidently the operation has not passed the development stage. Production during recent years has been almost negligible, and there is no prospect at present of the industry attaining importance.

GEORGIA

Occurrences of amphibole asbestos in Georgia are of two kinds: (1) Mass fiber, consisting of aggregations of short fibers representing a complete alteration of the original igneous rock; and (2) long-fibered asbestos, occurring in veins associated with partly altered peridotite, talc, chlorite, and serpentine.

Georgia has been the most consistent producer of amphibole asbestos of any of the States. The largest quantity has been obtained from deposits on or near Sall Mountain in the vicinity of Helen, Santee, Cleveland, and Nacoochee, White County. These deposits

have produced with some in a small way during earlier anthophyllite type. It is quarried as fiber. The weathering has made it so that rock is less desirable and of the best fiber, therefore, available high-grade material for use in a mill at Gaines. It is employed principally in paint manufacture.

Another production center, 1924 to 1927, is Hollywood, which occurs in a deposit as silky than that at Sall Mountain. Hammer-mill methods of fine grades of asbestos were produced for heat insulation and as filler for boxes, and rubber. Little in 1927. A large deposit of the county about 11 miles north by road from Nacoochee, White County.

From 1923 to 1927 a small deposit from a property on Bear Creek, Barrow County. It was used for heat insulation.

Asbestos was mined about Rabun County. The mineral was altered along fracture lines. Many other occurrences of asbestos have been reported.

Fiber 6 to 7 inches long and about 2 miles southeast of McAdams, Long County.

Long slip fiber occurs in White County. Many other undeveloped deposits have been reported by Hopkins.¹⁷

The only asbestos deposit commercially is in Idaho County. The deposit consists of brittle lenticular masses, some of which are altered. It is regarded as an alteration of asbestos. Asbestos Manufacturing Co. produced it for years prior to 1918. In 1918 it was purchased by the Western Mineral Co. It was reported by this company that the deposit, Idaho, acquired the property in 1925. No sales have been reported in a press report in 1930. The plant, purchased in 1930, was reported in 1925; had a capacity of 2,000 tons.

¹⁷ Hopkins, Oliver B., Asbestos, Talc, and other minerals of Georgia. Bull. 29, 1914, pp. 142-189.

Hill, Loudoun County, Va., was reported of asbestos also occur in massive serpentine Co. near Ishpeming, Marquette County, and work was done.

AMPHIBOLE

ALASKA

The amphibole variety occurs on Admiralty the Copper River. A little development at low price of amphibole fiber discourages markets.

CALIFORNIA

deposits were worked in San Bernardino, Placer Counties, and in 1883 and 1884 in Mono, Los Angeles, Tulare, Mariposa, and some of them were developed. Production; in 1883, 1,000 tons; and in 1884, 1,000 tons. A mine was opened at Windsor in Sonoma. High quality was reported from Del Norte. Deposits of asbestos in the United States of 30,000 and 66 tons in 1891 were credited to 1892 most of the production was from mines from Oregon and California, but with an output of 50 tons, was the only ones were idle in 1894 but produced 90 tons. Gradually decreasing output was noted in production up to the latter year was noted. Evidently it was used chiefly in for making pipe and boiler coverings.

was mined for several years near Sims, and has been noted since 1923. Asbestos associated with serpentine near was developed in 1923. Several short made, but evidently the operation has stage. Production during recent years and there is no prospect at present of the.

GEORGIA

Asbestos in Georgia are of two kinds: aggregations of short fibers representing original igneous rock; and (2) long-fiber veins associated with partly altered serpentine.

consistent producer of amphibole asbestos. The largest quantity has been obtained at Mountain in the vicinity of Helen. Cherokee, White County. These deposits

have produced with some interruptions since 1894 and possibly in a small way during earlier years. The asbestos is of the mass-fiber anthophyllite type. It is claimed that 90 to 95 percent of the rock quarried is fiber. The best asbestos is near the surface, where weathering has made it soft and flexible. The solid, unweathered rock is less desirable and requires different treatment. Resources of the best fiber, therefore, are relatively limited, and most of the available high-grade material is exhausted. The fiber is prepared for use in a mill at Gainesville about 27 miles from the deposits. It is employed principally for fireproofing, for insulation, and in paint manufacture.

Another production center of some importance, particularly from 1924 to 1927, is Hollywood, Habersham County. The mass fiber, which occurs in a deposit about 90 feet wide, is harsher and less silky than that at Sall Mountain. After much experimental work, hammer-mill methods of fiberizing were introduced and several grades of asbestos were prepared. The products were sold for heat insulation and as filler for composition flooring, plaster, battery boxes, and rubber. Little or no production has been recorded since 1927. A large deposit of mass-fiber anthophyllite occurs in this county about 11 miles northwest of Clarkesville. It is 8 to 9 miles by road from Nacoochee, White County, the nearest railway station.

From 1923 to 1927 a small output of anthophyllite was reported from a property on Bear Creek 1½ miles northeast of Statham, Barrow County. It was used principally for filtering acids.

Asbestos was mined about 1880, 2 miles northwest of Burton, Rabun County. The mineral is long-fibered and woody. It resulted from alteration along fracture zones of an extensive peridotite intrusion. Many other occurrences are known in the county.

Fiber 6 to 7 inches long occurs in an 18-inch vein in soapstone about 2 miles southeast of Moreland, Coweta County.

Long slip fiber occurs in talcose rock north of Luthersville, Meriwether County.

Many other undeveloped deposits in the State have been described by Hopkins.¹⁷

IDAHO

The only asbestos deposit in Idaho that has been developed commercially is in Idaho County about 14 miles southeast of Kamiah. The deposit consists of brittle mass-fiber anthophyllite occurring in lenticular masses, some of which are 200 feet long and 35 feet thick. It is regarded as an alteration product of dunite. The Kamiah Asbestos Manufacturing Co. produced in a small way for a number of years prior to 1913. In September 1921 the property was purchased by the Western Mineral Co. of Kamiah, but no production was reported by this company. The Panhandle Asbestos Co. of Lewiston, Idaho, acquired the property in 1923 and reported small shipments in 1925. No sales have been recorded since that year. According to a press report in 1930, the Diatom Products Co. of Seattle, Wash., purchased the plant. The mill, that was apparently last operated in 1925; had a capacity of 240 tons of ore a day and was equipped

¹⁷ Hopkins, Oliver B. Asbestos, Talc, and Soapstone Deposits of Georgia: Geol. Survey of Georgia Bull. 29, 1914, pp. 142-180.

with crusher, rolls, screens, and air separators. The fiber was used for pipe and boiler covering, in wall plaster and paint, and as a binder in cements and asphalts. In 1922 equipment was installed for sawing fire blocks and insulating brick, but the project did not materialize.

MARYLAND

Asbestos occurrences were noted in western Maryland as early as 1883, but no production worthy of mention was recorded until 1917. In that year a slip-fiber anthophyllite deposit was developed near Pylesville, Harford County. The asbestos occurs in gneissoid schists that have been softened by weathering. The veins generally are small, and the usable fiber is confined to a zone near the surface. The good asbestos extends to greatest depths along shear zones where movement and pressure have been greatest. Mining has been conducted to a depth of about 50 feet.

The fiber is unusually strong and tough, and is especially adapted to making acid filters; therefore, it is more valuable than anthophyllite from other localities. It is prepared for market by the Powhatan Mining Co. at Woodlawn, a suburb of Baltimore. Before this deposit was discovered, supplies of acid-filter fiber were obtained from Italy.

A little long-fiber anthophyllite, said to be similar to that mined at Pylesville, is found associated with talc in the Harford Talc & Quartz Co. quarry near Dublin, Harford County.

MONTANA

A deposit of anthophyllite asbestos, said to be extensive, in Gallatin County, about 17 miles south of Gallatin Gateway, is worked for a small production of insulating material. The fiber occurs in a dense, hard form but breaks down readily to a soft, fluffy mass. Occasional pockets and lenses contain strong, silky fiber, much of which is an inch or more in length. It is of spinning quality and is the only amphibole of such grade in the United States known to the Bureau. Peter F. Karst operated the property for several years, but in 1936 it was acquired by The Karstolite Co. This firm mines the asbestos and manufactures it into a wall and ceiling insulation sold under the trade name "Karstolite."

A small shipment of anthophyllite was recorded in 1928 from a property on Rainey Creek, 7 miles northeast of Libby, Lincoln County.

NORTH CAROLINA

About 1928 a mill that would handle 35 tons of ore a day was built at Minneapolis, Avery County, for treatment of mass-fiber anthophyllite occurring in that region. The mill equipment was improved considerably in 1935, and steps were taken to develop a market for prepared asbestos to be used in home insulation and in many other ways. A deposit near Plumtree, in this county, was developed some years ago, but very little fiber has been produced.

A mill was built near Otto, Macon County, in 1928 but has not produced. A small output was reported in 1919 from a deposit of mass-fiber anthophyllite at Cane River, Yancey County. Deposits of anthophyllite occur also near Warlick, Burke County; on the

Nantahala River, Macon County, near Todd, Ashe County, Caldwell County.

Sixty-four tons of sold in 1892. In 1917 of Mount Vernon, Gra Baker County, was developed. Work on a 310-foot tunnel was purposes was reported

Amphibole asbestos; cross-fiber veins traverse Gillespie Counties. It is of a considerable supply usual in amphibole-asbestos, supplied by Sellards and

Anthophyllite asbestos Bedford County. The and olivine. Unlike the best asbestos is the occupying shear zones, inches thick supplied for treating the asbestos, able supply is probably 1911. A small quantity, a preparation used by

A deposit of slip-fiber Franklin County. A few tons were obtained from also near Boyce, Clarke County.

The Asbestos-Talc Plant, Skagit County, talc, serpentine, and a used in boiler coverings, also manufactures "Asbestos", "Asbestocite", roofing paints and plaster. Asbestos has been mined in paints. Deposits have County, near Hamilton County; and 8 miles north

¹ Sellards, E. H., and Baker 1935, p. 252.

and air separators. The fiber was used for wall plaster and paint, and as a substitute for asbestos. In 1922 equipment was installed for producing brick, but the project did not

MARYLAND

Asbestos was noted in western Maryland as early as 1880. The first mention was recorded until 1917. Anthophyllite deposit was developed near

The asbestos occurs in gneissoid schists and is weathered. The veins generally are confined to a zone near the surface. The greatest depths along shear zones where asbestos has been greatest. Mining has been conducted to a depth of 100 feet.

The asbestos is strong and tough, and is especially adapted for use in the manufacture of asbestos. Before this deposit was developed, asbestos fiber was obtained from Italy. Anthophyllite, said to be similar to that mined at Libby, was developed in the Harford Talc & Quartz Company.

MONTANA

Asbestos, said to be extensive, in Gallatin National Park, is worked for a variety of material. The fiber occurs in a dense, easily broken to a soft, fluffy mass. Occasional strong, silky fiber, much of which is an excellent quality and is the only American asbestos known to the Bureau of Mines. The property for several years, but in 1936 the Asbestos Talc Co. This firm mines the asbestos and all and ceiling insulation sold under the

Anthophyllite was recorded in 1928 from a deposit 7 miles northeast of Libby, Lincoln

NORTH CAROLINA

A mill to handle 35 tons of ore a day was built for treatment of mass-fiber anthophyllite. The mill equipment was improved and steps were taken to develop a market for asbestos in home insulation and in many other uses. In this county, was developed some asbestos has been produced.

Macon County, in 1923 but has not been reported in 1919 from a deposit of Cane River, Yancey County. Deposits near Warlick, Burke County; on the

Nantahala River, Macon County; near Bakersville, Mitchell County; near Todd, Ashe County; Brevard, Transylvania County; and in Caldwell County.

OREGON

Sixty-four tons of amphibole asbestos produced in Oregon were sold in 1892. In 1917 a small mine was operated a few miles north of Mount Vernon, Grant County. Asbestos occurring near Hereford, Baker County, was described in 1922 as a soft, long-fibered tremolite. Development work on the property was reported in 1925, and in 1926 a 310-foot tunnel was driven. A small production for experimental purposes was reported in 1929.

TEXAS

Amphibole asbestos approaching tremolite in composition occurs in cross-fiber veins traversing serpentine in four localities in Llano and Gillespie Counties. The veins are very irregular, but there is promise of a considerable supply. The occurrence of cross-fiber veins is unusual in amphibole-asbestos deposits. Further information is supplied by Sellards and Baker.¹⁸

VIRGINIA

Anthophyllite asbestos occurs about 12 miles south of Bedford, Bedford County. The prevailing associated minerals are hornblende and olivine. Unlike the mass-fiber deposits of Georgia and Idaho, the best asbestos is the slip-fiber type, occurring in veinlike masses occupying shear zones. Several veins have been noted, and one 18 inches thick supplied considerable fiber from 1904 to 1911. A mill for treating the asbestos was erected in Bedford in 1903. The available supply is probably small, as very little has been produced since 1911. A small quantity has been used in the manufacture of Tenax, a preparation used by dentists.

A deposit of slip-fiber amphibole occurs also at Rockymount, Franklin County. A shaft was sunk to a depth of 40 feet, and 40 tons were obtained from this mine in 1907. Deposits have been noted also near Boyce, Clarke County, and 8 miles from Roanoke, Roanoke County.

WASHINGTON

The Asbestos-Talc Products Co. developed a property near Burlington, Skagit County, and has produced a small tonnage of mixed talc, serpentine, and asbestos since 1930. The milled product is used in boiler coverings, roofing, and acoustic plaster. The company also manufactures "Asbesto-Fill", a heat-insulating building material, and "Asbestocite", a compound of asbestos fiber and talc used in roofing paints and plastics.

Asbestos has been mined near Pateros, Okanogan County, for use in paints. Deposits have been reported near Leavenworth, Chelan County, near Hamilton, Lyman, and at other points in Skagit County; and 8 miles north of Chewelah, Stevens County.

¹⁸ Sellards, E. H., and Baker, Charles Laurence: Economic Geology of Texas, part 3, 1935, p. 252.

OTHER OCCURRENCES

Many other occurrences of amphibole asbestos were known from 1880 to 1905. The following localities, mentioned in early records, evidently had small importance as sources of supply: Staten Island and Long Island, N. Y.; New Brunswick, N. J.; Media and Colerain, Pa.; Lawrence County, S. Dak.; Stevens Point, Wood County, Wis.; Six Mile and Pickens, S. C.; Baraga County, Mich.; and New Hartford, Conn., where a small mill was operated in 1903. From 1901 to 1906 a small output of asbestos was reported near Dalton, Mass. It was first thought to be chrysotile but was identified definitely as anthophyllite in 1906. Blue asbestos (crocidolite) occurs at Beacon Pole Hill near Manville, Providence County, R. I., but not in commercial quantities.

FOREIGN DEPOSITS

CANADA

The asbestos deposits in the eastern townships of the Province of Quebec, Canada, have been famous since 1878. They are situated 100 to 175 miles from Montreal and somewhat nearer Quebec, occupying a stretch of high ground known as the Notre Dame Hills—an extension into Canada of the Green Mountains of Vermont. The hills continue northeastward to the Gaspé Peninsula, and serpentine is exposed in many places throughout their length. An asbestos deposit was developed at Port Daniel on the Gaspé Peninsula in 1919. Commercial asbestos occurs in two places—the Thetford Mines district, described later, and the Eden (Vt.) district, which has been covered in the discussion of domestic deposits.

The Quebec area, about 70 miles long and a maximum of 5 or 6 miles wide, lies between Danville and East Broughton. Within this section are six producing centers—East Broughton, Robertson, Thetford Mines, Black Lake, Coleraine, and Danville. The geology of the district has been covered in detail by Cirkel.¹⁹ The asbestos, a chrysotile of exceptionally high quality, occurs in a very extensive serpentine formation of Cambrian age appearing in places as mountain masses rising 700 to 1,000 feet above the surrounding country. It is said to be the largest area of serpentine near the eastern coast of North America. The serpentine is regarded as an alteration product of the ultrabasic rock dunite, which consists essentially of olivine and chromite. In the process of alteration, serpentinization takes place first along irregular cracks in the olivine, and the wavy contortions of the filaments led to application of the name "serpentine." As alteration of olivine to serpentine involves an increase in volume of more than 30 percent, progressive disruption and fracturing would occur. Cirkel²⁰ attributed the origin of the chrysotile to columnar crystallization of serpentine in cracks and fissures. The process is somewhat obscure, but he presumed that the serpentine, which thus takes the form of bundles of extraordinarily fine filaments, crystallized from aqueous solutions acting upon the vein walls.

¹⁹ Cirkel, Fritz, *Chrysotile Asbestos—Its Occurrence, Exploitation, Milling, and Uses*: Mines Branch, Canada Dept. of Mines, 2d ed., 1910.

²⁰ Cirkel, Fritz, *Work cited*, pp. 92-95.

Bureau of Mines



FIGURE 5.—Veins of asbestos on a

ASBESTOS

OTHER OCCURRENCES

es c amphibole asbestos were known from
win localities, mentioned in early records,
ortance as sources of supply: Staten Island
New Brunswick, N. J.; Media and Colerain,
s. Dak.: Stevens Point, Wood County, Wis.;
C.: Baraga County, Mich.; and New Hart-
ll mill was operated in 1903. From 1901 to
sbestos was reported near Dalton, Mass. It
chrysotile but was identified definitely as
blue asbestos (crocidolite) occurs at Beacon
Providence County, R. I., but not in com-

FOREIGN DEPOSITS

CANADA

n the eastern townships of the Province of
n famous since 1878. They are situated 100
al and somewhat nearer Quebec, occupying
l known as the Notre Dame Hills—an ex-
ie Green Mountains of Vermont. The hills
to the Gaspé Peninsula, and serpentine is
throughout their length. An asbestos de-
ort Daniel on the Gaspé Peninsula in 1919.
irs in two places—the Thetford Mines dis-
l the Eden (Vt.) district, which has been
of domestic deposits.
t 70 miles long and a maximum of 5 or 6
Danville and East Broughton. Within this
centers—East Broughton, Robertson, Thet-
Coleraine, and Danville. The geology of
ered detail by Cirkel.¹⁹ The asbestos, a
ly h₂ quality, occurs in a very extensive
Cambrian age appearing in places as moun-
1,000 feet above the surrounding country.
t area of serpentine near the eastern coast
serpentine is regarded as an alteration
rock dunite, which consists essentially of
the process of alteration, serpentinization
regular cracks in the olivine, and the wavy
nts led to application of the name "ser-
of olivine to serpentine involves an increase
0 percent, progressive disruption and frac-
tel²⁰ attributed the origin of the chrysotile
n of serpentine in cracks and fissures. The
ure, but he presumed that the serpentine,
of bundles of extraordinarily fine filaments,
s solutions acting upon the vein walls.

¹⁹stos—Its Occurrence, Exploitation, Milling, and Uses:
ins, 2d ed., 1910.
22-25.

Bureau of Mines

Bulletin 403



FIGURE 5.—Veins of asbestos on drift face 200 feet below surface, Thetford Mines, Quebec.

Cirkel²¹ outlines the successive steps of formation of the asbestos as follows:

1. Intrusion of olivine (dunite) through the earth crust from below.
2. Gradual alteration of the rock to serpentine through hydration, and perhaps loss of silica, increase in volume.
3. Slow readjustment of the rock masses, resulting in the formation of joints and slickensides.
4. Subsequent formation of fissures as receptacles for asbestos fiber, through shrinkage of the rock and also through injection of granite dikes.
5. Infiltration of serpentinous solution from the sides of the wall through process of segregation, and subsequent slow crystallization of chrysotile.

Cooke²² has made the most comprehensive study of the origin of chrysotile asbestos veins in Quebec. Three hypotheses as to origin have been advanced: (1) Replacement of massive serpentine by asbestos, (2) fissure-filling by circulating solutions, and (3) solutions or vapors penetrating through pores of the wall rock and depositing asbestos in tight fissures, the growth of the fibers pushing the walls apart. Although some objection may be raised to the last-mentioned theory, Cooke concludes that it furnishes the most adequate explanation of the conditions existing throughout the region. The most serious objection to the theory of fissure-filling by circulating solutions is the numerous occurrence of closed veins showing no connection with channels through which solutions could flow.

In some places the fiber is very soft and flexible, while in others it is somewhat harsh and brittle. To determine a reason for this harshness, Cooke²² made very careful analyses of purified fibers of both varieties. He concluded from the results that the harshness is due primarily to the presence of talc. Keep²³ reached a similar conclusion regarding Rhodesian chrysotile.

Almost the entire output of the Black Lake, Thetford, and Danville areas is derived from cross-fiber veins ranging in width from hair lines to 4 or 5 inches and, rarely, even wider. The bulk of the production is obtained from veins $\frac{1}{4}$ to $\frac{1}{2}$ inch across. Width of veins is important because it governs largely the length of fiber. Fortunately the fiber as a rule breaks easily from the wall rock. Figure 5 shows a typical occurrence of asbestos veins in the Quebec district.

Slip fiber originating along slickensided fault planes as a result of slow readjustment of the rock masses at a later time forms the chief product of the East Broughton area. The fibers are matted together more or less in parallel position, and therefore appear to be longer than they really are. Slip fiber, as a rule, is less valuable than cross fiber.

The percentage of fiber in the rock is low. According to Ross,²⁴ 6,208,970 tons of rock were mined in 1929 and total fiber recovery (excluding asbestic) was 311,204 tons. The yield was 5 percent of the total rock mined and 7.7 percent of the rock milled. The value

²¹ Cirkel, Fritz. Work cited, p. 101.

²² Cooke, H. C., Asbestos Deposits of Thetford District, Quebec: Econ. Geol., vol. 31, no. 4, June-July 1936, pp. 355-376.

²³ Cooke, H. C., The Composition of Asbestos and Other Fibres of Thetford District, Quebec: Trans. Roy. Soc. Canada, 3d ser., vol. 29, sec. 4, May 1935, pp. 7-19.

²⁴ Keep, F. E., Geology of the Shabani Mineral Belt, Bellingwe District: Geol. Survey, So. Rhodesia, Bull. 12, 1929.

²⁵ Ross, J. H., Chrysotile Asbestos in Canada: Mines Branch, Canada Dept. Mines Bull. 707, 1931, table 1, p. 80.

of the asbestos recovered was \$2.12 per ton of rock mined and hoisted. Fiber classed as crude constituted only 1½ percent of the asbestos recovered. The output of spinning fiber, including crudes and the highest-grade mill fiber, was 24,415 tons (7.8 percent of the gross yield). Figures for other years vary little from those given above. The average recovery of fiber from milled rock is about 6 percent. Although Canada produces substantial quantities of spinning fiber, considerably more than 90 percent of its production consists of the shorter (consequently, relatively cheap) grades for which, fortunately, an extensive market exists in the United States.

The reserves of asbestos in Quebec are very large. Some of the workings are more than 400 feet deep, and core drilling has revealed an abundant supply of fiber, at depths of more than 1,700 feet, as high in quality as that near the surface.

The following table shows the annual production and value of asbestos in Quebec since the industry was first established.

Annual production (shipments and sales) of asbestos in Quebec, 1878-1935¹

Year	Production	Total value	Average value per ton	Year	Production	Total value	Average value per ton
1878	30	\$19,500	\$65.00	1907	61,985	\$2,455,919	\$39.62
1879	300	24,700	82.33	1908	65,157	2,551,596	39.16
1880	330	35,100	106.36	1909	64,965	2,296,564	35.35
1881	540	52,650	97.50	1910	50,605	2,667,829	52.71
1882	210	25,750	122.62	1911	102,254	3,026,306	29.60
1883	955	65,750	68.84	1912	111,173	3,059,084	27.52
1884	1,141	75,097	65.81	1913	136,609	3,630,504	26.56
1885	2,440	142,441	58.37	1914	107,401	2,895,935	26.96
1886	3,458	206,251	59.65	1915	113,115	3,544,322	31.34
1887	4,619	226,976	49.15	1916	133,339	5,182,905	39.67
1888	4,404	255,007	57.90	1917	137,242	7,195,558	52.45
1889	6,113	426,584	69.77	1918	142,375	9,019,899	63.35
1890	9,860	1,270,240	128.82	1919	135,622	10,932,259	80.47
1891	9,279	999,878	107.75	1920	179,891	14,749,048	81.99
1892	6,052	390,462	64.16	1921	87,475	5,199,789	59.44
1893	8,331	310,156	48.99	1922	160,339	6,053,068	37.75
1894	6,756	420,825	62.44	1923	216,604	7,364,260	33.97
1895	10,592	423,066	39.84	1924	208,762	6,581,659	31.57
1896	13,202	399,528	30.26	1925	273,522	8,976,645	32.82
1897	13,893	496,340	35.76	1926	273,659	10,095,457	36.92
1898	15,571	581,667	37.36	1927	274,798	10,621,571	38.65
1899	21,408	719,416	33.60	1928	273,033	11,233,361	41.16
1900	33,666	1,274,315	37.85	1929	306,055	13,172,581	43.04
1901	30,634	1,161,570	37.92	1930	242,113	6,390,164	26.40
1902	29,261	916,970	31.34	1931	164,297	4,812,856	29.29
1903	25,479	1,156,970	45.43	1932	122,977	3,039,721	24.72
1904	48,960	1,476,450	30.16	1933	155,367	5,211,177	33.56
1905	61,675	2,143,653	34.76	1934	155,950	4,936,326	31.65
1906				1935	210,467	7,054,614	33.52

¹ Figures for 1878-1922 from The Quebec Asbestos Industry, by Norman R. Fisher: Canadian Min. Jour., Aug. 17, 1923, p. 651. Figures for 1923-35 published by the Quebec Dept. of Colonization, Mines, and Fisheries, Bureau of Mines.

Until 1891, virtually the entire Canadian output was crude fiber, but thereafter both crude and milled asbestos were produced. This change in the character of products is reflected in the sharp drop in average value in 1892 compared with the previous year. The exceptionally high average values from 1918 to 1920 may be attributed to the abnormally high prices obtained for crude fiber on account of war conditions. Crude No. 1, for example, sold for more than \$3,000 a ton during part of this period.

The largest asbestos producer in Canada is Asbestos Corporation, Ltd., operating the King and Beaver pits at Thetford Mines, the

Consolidated at Black Lake, the M Black Lake, the Vimy Ridge, Broughton, and several others. The company operates six mills h 500 tons of rock an hour. Bell As Turner & Newall, Ltd., of England, Thetford Mines. Johnson's Co., the pioneer in the district. It oper ford Mines and another similar oper bec Asbestos Corporation, Ltd., contr facturing Co. of Cincinnati, Ohio, of Broughton. The Canadian Johns- large open-pit power-shovel operat from Danville. A mill for treatm products factory are situated near th Mines, Ltd., operates a pit and mill The mills of the district have a cc tons of rock an hour, and the capital estimated at nearly \$22,000,000.

The Canadian deposits occupy an e as they are within easy reach of ex States. The Quebec industry doubtl indefinitely in the short-fiber market on relatively low-priced products di tion. The future of the spinning-fib bright because the price per ton of tl it a world-wide market range, and c and Soviet Russia has become increas the writer² pointed out the danger to States markets of increasing imports, try, but it was not until 5 years late by the industry to meet this competi and exercising economies in mi. Such consolidations and cost reduc or try by economic conditions, because wi can competitors the struggle for dir cutting that was carried to a point v reduced to bankruptcy, and the margin pany was extremely small. Since the lished on a better competitive basis, b output of Russian mines and the conce tion in the hands of one powerful co still a serious problem for producers i substantial quantities of spinning a progress toward placing her products c establishing, in 1931, a uniform classifi.

From 1916 to 1924 some interest in tl industry in Ontario was aroused. Go tained at two points in Doloro Townshi area of Porcupine. A maximum prod \$91,900 was recorded in 1924. The pla

² Bowles, Oliver, Mineral Industry during 1919 Book Co., New York, 1920, pp. 41-42.

was \$2.12 per ton of rock mined and rude constituted only 1½ percent of the spun spinning fiber, including crudes fiber. As 24,415 tons (7.8 percent of the other years vary little from those given) of fiber from milled rock is about 6 produces substantial quantities of spin- e than 90 percent of its production con- quently, relatively cheap) grades for sive market exists in the United States. n Quebec are very large. Some of the feet deep, and core drilling has revealed at depths of more than 1,700 feet, as the surface.

As the annual production and value of industry was first established.

and sales) of asbestos in Quebec, 1878-1935¹

Average value per ton	Year	Production	Total value	Average value per ton
63.00	1907	61,985	\$2,455,919	\$39.62
63.00	1908	65,157	2,531,596	39.16
65.00	1909	64,963	2,296,584	35.35
65.00	1910	80,605	2,667,599	33.10
65.00	1911	102,224	3,026,306	29.60
72.00	1912	111,175	3,059,084	27.52
65.61	1913	136,609	3,830,504	28.04
58.37	1914	107,401	2,895,935	26.96
59.65	1915	113,115	3,544,362	31.34
49.55	1916	133,339	5,182,905	38.37
57.90	1917	137,242	7,198,536	52.46
69.77	1918	142,375	9,019,869	63.35
128.82	1919	135,862	10,932,299	80.47
107.75	1920	179,891	14,749,048	81.99
64.16	1921	57,475	5,199,759	59.44
48.99	1922	160,339	8,053,068	37.75
55.17	1923	216,804	7,364,260	33.97
42	1924	208,762	6,561,659	31.37
38	1925	273,522	8,976,645	32.82
30.26	1926	278,659	10,095,467	36.22
31.23	1927	274,798	10,621,571	38.65
37.36	1928	273,033	11,238,361	41.16
33.60	1929	306,055	13,172,581	43.04
38.70	1930	242,113	8,390,164	34.65
37.92	1931	164,297	4,512,886	29.29
31.94	1932	125,977	3,039,721	24.72
33.43	1933	158,367	5,211,177	32.90
50.16	1934	155,950	4,936,326	31.65
34.76	1935	210,467	7,054,614	33.52

asbestos industry, by Norman R. Fisher: Canadian Min. Jour., published by the Quebec Dept. of Colonization, Mines, and

entire Canadian output was crude fiber. id milled asbestos were produced. This roducts is reflected in the sharp drop in red with the previous year. The excep- s from 1918 to 1920 may be attributed s obtained for crude fiber on account of 1, for example, sold for more than \$3.00 100 lb.

icer in Canada is Asbestos Corporation, ad Beaver pits at Thetford Mines, the

Consolidated at Black Lake, the Maple Leaf between Thetford and Black Lake, the Vimy Ridge at Coleraine, the asbestos mine at East Broughton, and several other smaller mines as the occasion demands. The company operates six mills having an aggregate capacity of 500 tons of rock an hour. Bell Asbestos Mines, Inc., now owned by Turner & Newall, Ltd., of England, operates a large mine and mill at Thetford Mines. Johnson's Co., which began mining in 1878, is the pioneer in the district. It operates a large pit and mill at Thetford Mines and another similar operation at Black Lake. The Quebec Asbestos Corporation, Ltd., controlled by the Philip Carey Manufacturing Co. of Cincinnati, Ohio, operates a mine and mill near East Broughton. The Canadian Johns-Manville Co., Ltd., conducts a large open-pit power-shovel operation at Asbestos, about 5 miles from Danville. A mill for treatment of the ore and an asbestos-products factory are situated near the quarry. The Nicolet Asbestos Mines, Ltd., operates a pit and mill about 7 miles east of Asbestos. The mills of the district have a combined capacity of over 1,000 tons of rock an hour, and the capital employed in mines and mills is estimated at nearly \$22,000,000.

The Canadian deposits occupy an exceptionally favorable position, as they are within easy reach of extensive markets in the United States. The Quebec industry doubtless will maintain its advantage indefinitely in the short-fiber market, because high freight charges on relatively low-priced products discourage transoceanic competition. The future of the spinning-fiber industry in Canada is not so bright because the price per ton of this type is high enough to give it a world-wide market range, and competition with South Africa and Soviet Russia has become increasingly severe. As early as 1919 the writer² pointed out the danger to Canadian supremacy in United States markets of increasing imports of African fiber into this country, but it was not until 5 years later that active steps were taken by the industry to meet this competition by consolidating companies and exercising economies in mining, manufacture, and marketing. Such consolidations and cost reductions were forced upon the industry by economic conditions, because with the loss of markets to African competitors the struggle for diminishing trade led to price-cutting that was carried to a point where several companies were reduced to bankruptcy, and the margin of profit obtained by any company was extremely small. Since then the industry has been established on a better competitive basis, but with the rapidly increasing output of Russian mines and the concentration of Rhodesian production in the hands of one powerful company, foreign competition is still a serious problem for producers in Quebec whose mines supply substantial quantities of spinning asbestos. Canada made great progress toward placing her products on a better marketing basis by establishing, in 1931, a uniform classification of asbestos.

From 1916 to 1924 some interest in the development of an asbestos industry in Ontario was aroused. Good-quality chrysotile was obtained at two points in Doloro Township south of the gold-producing area of Porcupine. A maximum production of 172 tons valued at \$91,900 was recorded in 1924. The plant was dismantled later, and

² Bowles, Oliver, Mineral Industry during 1919 (chapter on Asbestos): McGraw-Hill Book Co., New York, 1920, pp. 41-42.

no subsequent activity has been reported except the sale of 2 tons valued at \$901 in 1926.

In 1935 considerable development work was done by the Rahn Lake Mines Corporation, Ltd., of North Bay, Ontario, on a chrysotile deposit near Matachewan, Ontario, Bannockburn Township, about 30 miles southeast of Timmins in the Porcupine gold area. Erection of a mill is contemplated.

Occurrences of chrysotile asbestos of good quality have been reported near Lytton, Revelstoke, and North Kootenay, British Columbia, but samples from Lytton proved to be weak, brittle amphibole.

Although most Canadian asbestos is exported, the manufacture of asbestos products within the country is assuming increasing importance. Before 1921 only one company manufactured asbestos products in Canada; by 1923 there were three plants in Quebec and one in Ontario; in 1932 there were five plants in Quebec, six in Ontario, one in Nova Scotia, and one in British Columbia; and in 1933, four in Quebec, five in Ontario, and one each in Nova Scotia and British Columbia. Products manufactured in 1932 were valued at \$1,067,801 and in 1933 at \$757,626, compared with a maximum of \$2,301,924 in 1930. Capital employed by the companies operating in 1933 totaled \$1,777,975.

RHODESIA

The growth of the asbestos industry in Rhodesia has been almost phenomenal. From an initial output of only 55 tons in 1908 it reached more than 42,000 tons in 1929, when of all the minerals produced in the country the value of the output of asbestos was second only to that of gold. Annual production since the beginning of activity is shown in the following table:

Annual production of asbestos in Rhodesia, 1908-36

Year	Short tons	Value	Year	Short tons	Value	Year	Short tons	Value
1908.....	55	£532	1918.....	5,574	£153,654	1928.....	39,960	£970,327
1909.....	272	2,722	1919.....	9,796	423,240	1929.....	42,634	1,186,627
1910.....	332	3,320	1920.....	18,623	459,572	1930.....	37,765	1,070,847
1911.....	460	6,397	1921.....	19,528	795,698	1931.....	24,042	386,494
1912.....	290	5,224	1922.....	14,249	577,699	1932.....	15,766	197,092
1913.....	457	8,612	1923.....	20,364	626,896	1933.....	30,182	555,993
1914.....	2,010	32,190	1924.....	26,141	603,423	1934.....	32,214	402,745
1915.....	6,157	99,059	1925.....	34,349	765,926	1935.....	42,598	646,655
1916.....	9,562	189,890	1926.....	33,344	726,835	1936.....	56,346	836,469
1917.....			1927.....	33,176	794,215			

The fiber is of the chrysotile variety occurring in serpentine. It is similar in character and occurrence to that found in Canada and Soviet Russia. There are three producing districts—the Bulawayo, the Victoria (Mashaba), and the Lamagundi, all in southern Rhodesia. Deposits in other localities are known but have not attained commercial importance.

The Bulawayo district is the most important. Its largest mine, the Shabani, opened in 1916, is a prolific producer, in prosperous years providing as much as 1,200 tons of fiber a month. Other important mines in the area are the Birthday, the Nil Desperandum, and the Pangani. The Croft mine at Filibusi has attained some importance.

The asbestos deposits of the Sh of a serpentine mass about 10½ The zone of fiber-bearing se and a maximum width of ai from alteration of dunite, probal intrusion. The asbestos veins a pentine 20 to more than 200 fe talc schist that forms the footw contact ranges from 25° to 45°, in the flat country the dip is only extends down the dip to an unkr working limit. The fiber, which wide, is rarely more than 3 inch tудinal partings occur in most ve row. The fiber seams generally are continuous for only short dis short, irregular, overlapping veir 7,000,000 tons in 1928.

For many years the district wa: transportation; 400 wagons and 7 supplies and products many mil hindrance was overcome when a Somabula was completed.

The earliest Rhodesian produc 46 miles east of Shabani and 26 i cipal mines, the King's and Gat many years. The fiber is similar i area. A small production has be in the Lamagundi district more Here flexible and brittle fibers o separation requires a complex mil

In Rhodesia the percentage of estimated that 25 to 30 percent of as spinning grade, whereas output is of this grade. i... di as these figures indicate, because s 50 and 60 percent of the Canadian in Rhodesia.

Highly efficient mills for treati junction with the larger mines. were consolidated under a single placed in a strong competitive po: prosperous. Profits of the larges 1925 to 1927, 25 percent in 1928, in 1930, 5 percent in 1931, 3% perc Most of the fiber is exported to Eu of the port of Beira, Portuguese E ness of the region from consumin higher-grade mill fibers are sold. stock are sold principally in Europ poses. Very little Rhodesian asbes stock reaches the American market The advantage of unified control rapidly increasing exports from F

n reported except the sale of 2 tons

ment was done by the Rahn Lake
Fort Erie, Ontario, on a chrysotile
quarry, Bannockburn Township, about
in the Porcupine gold area. Erection

asbestos of good quality have been re-
and North Kootenay, British Colum-
proved to be weak, brittle amphibole.
asbestos is exported, the manufacture of
country is assuming increasing impor-
company manufactured asbestos prod-
were three plants in Quebec and one
five plants in Quebec, six in Ontario,
British Columbia; and in 1933, four
one each in Nova Scotia and British
in 1932 were valued at \$1,067,801
ed with a maximum of \$2,301,924 in
companies operating in 1933 totaled

RODESIA

industry in Rhodesia has been almost
output of only 55 tons in 1908 it
1929, when of all the minerals pro-
of the output of asbestos was second
production since the beginning of
table:

Asbestos in Rhodesia, 1908-36

Short tons	Value	Year	Short tons	Value
8.376	154,654	1908	39,990	£970,327
9,798	425,240	1909	42,634	1,186,627
18,823	459,572	1910	37,765	1,070,847
19,523	735,798	1911	24,042	354,494
14,249	277,699	1912	15,766	197,092
20,364	426,398	1913	30,182	555,993
26,141	603,423	1914	32,214	402,745
34,349	763,926	1915	42,598	646,658
33,344	726,535	1916	56,346	836,469
33,176	794,215			

variety occurring in serpentine. It
ence to that found in Canada and
producing districts—the Bulawayo,
the Lamagundi, all in southern
localities are known but have not at-

most important. Its largest mine,
a prolific producer, in prosperous
tons of fiber a month. Other im-
portant mines, the Nil Desperandum,
mine at Filibus has attained some

The asbestos deposits of the Shabani area occur in the central part
of a serpentine mass about 10½ miles long and 1 to 3 miles wide.
The zone of fiber-bearing serpentine has a length of 2½ or 3 miles
and a maximum width of about 200 yards. The serpentine resulted
from alteration of dunite, probably under the influence of a granite
intrusion. The asbestos veins are best developed in a zone of ser-
pentine 20 to more than 200 feet thick close to the contact of a
talc schist that forms the footwall. In the hills, the dip of the
contact ranges from 25° to 45°, but where the formation is found
in the flat country the dip is only 5° to 10°. The fiber-bearing zone
extends down the dip to an unknown distance beyond the practical
working limit. The fiber, which occurs in seams as much as 6 inches
wide, is rarely more than 3 inches long, because one or more longi-
tudinal partings occur in most veins, except those that are quite nar-
row. The fiber seams generally parallel the dip of the contact but
are continuous for only short distances; they consist of a series of
short, irregular, overlapping veins. Ore reserves were estimated at
7,000,000 tons in 1928.

For many years the district was handicapped seriously by difficult
transportation; 400 wagons and 7,000 oxen were used for conveying
supplies and products many miles over poor roads. In 1928 this
hindrance was overcome when a 63-mile railway from Shabani to
Somabula was completed.

The earliest Rhodesian production was in the Mashaba district
46 miles east of Shabani and 26 miles west of Victoria. The prin-
cipal mines, the King's and Gath's, have been large producers for
many years. The fiber is similar in occurrence to that of the Shabani
area. A small production has been recorded from the Ethel mine
in the Lomagundi district more than 200 miles north of Victoria.
Here flexible and brittle fibers occur closely associated, and their
separation requires a complex milling system.

In Rhodesia the percentage of long fiber is relatively high; it is
estimated that 25 to 30 percent of the fiber produced may be classed
as spinning grade, whereas only about 7.8 percent of the Canadian
output is of this grade. The difference, however, is not as great
as these figures indicate, because short fibers that constitute between
50 and 60 percent of the Canadian production are regarded as waste
in Rhodesia.

Highly efficient mills for treating the fiber are operated in con-
junction with the larger mines. In 1929 all the important mines
were consolidated under a single company. The industry is thus
placed in a strong competitive position and before 1930 was highly
prosperous. Profits of the largest company were 20 percent from
1925 to 1927, 25 percent in 1928, 30 percent in 1929, 11¼ percent
in 1930, 5 percent in 1931, 3¾ percent in 1932, and 5 percent in 1933.
Most of the fiber is exported to Europe and America, chiefly by way
of the port of Beira, Portuguese East Africa. Owing to the remote-
ness of the region from consuming centers, only the spinning and
higher-grade mill fibers are sold. Medium grades classed as shingle
stock are sold principally in Europe and Australia for building pur-
poses. Very little Rhodesian asbestos of lower grade than spinning
stock reaches the American market.

The advantage of unified control was demonstrated in 1931, when
rapidly increasing exports from Russia, combined with decreasing

¹Hall, A. L., *Asbestos in the Union of South Africa*: Union of South Africa Geol. Survey Mem. 12, 2d ed., 1930.
²Hall, A. L., *Work cited*, p. 87.

1 inch long; that less than one-half inch long is rejected. Mines have been operated at various times, but production has never been large.

A second deposit of chrysotile in the Carolina district occurs on the farm Kalkloof, 3 miles south of the Komati River and 47 miles by road from Carolina. This occurrence is of the usual type—cross-fiber veins in serpentine derived from ultrabasic igneous rock. The fiber occurs in many closely spaced, parallel thin seams inclined at a steep angle. The asbestos is of good quality, but a large proportion of it is short mill fiber. Evidently reserves are extensive. A mill was in operation in 1929, but production has been small.

Barberton district.—In 1918 Hall²⁹ directed attention to a chrysotile deposit near Barberton in eastern Transvaal, but serious development did not take place until after 1921. The deposit occurs in a belt of serpentine bounded by quartzites lying about 3 miles from Kaapsche Hoop and 14 to 17 miles by road from Godwin River station. The asbestos-bearing portion of the serpentine has been traced for a length of 3 and a width of 2 miles. Further exploration may extend its boundaries considerably. Underground mines have been developed extensively near both the eastern and western ends of a 2-mile zone containing large reserves of fiber.

The New Amianthus Mines, Ltd. (subsidiary of Turner & Newall, Ltd.), operates near the western end. Here the most productive fiber belt is known as the "ribbon rock" or "ribbon line." In a typical section, the upper 7 feet contains narrow parallel fiber seams so numerous and closely spaced that the fiber content may run as high as 40 percent of the entire rock mass. Beneath this band is a 6-foot mass of serpentine, in which the chrysotile veins are fewer and thicker, providing fiber up to 6 inches long. Figure 6 shows the abundance of fiber veins. The veins dip 10° to 20°. The fiber is of high quality and is easily separated from the massive serpentine. Records for one period show that fiber recovered from the upper beds amounted to 30 percent of the rock mined, while from the lower beds it was 10 percent or less. In 1930, the average recovery for the mine was 15 to 17 percent. In that year about 8 percent of the output was of spinning grade, and 92 percent was classed as shingle stock. The mine is a steady producer, furnishing about 1,000 tons of fiber a month in 1930.


The Munnick-Myburgh mine occupies the eastern end of the serpentine area. The fiber seams are more or less parallel and resemble those at the western end already described. Thin seams are close together, and the thicker seams are more widely spaced. A majority of the seams worked range from one-sixteenth to one-half inch in width. Operations began about 1920, and in 1930 the mine was producing about 200 short tons of fiber a month.

For some years the products of the mines were transported 18 miles to Godwin River station in wagons hauled by oxen or mules. The completion in 1929 of a 5.9-mile aerial tramway over a 2,000-foot escarpment to Elandshoek made possible a reduction in haulage cost to about one-eighth of the amount formerly paid. From this point the asbestos is carried by a short-rail haul to Lourenco Marques, Portuguese East Africa, the port of shipment.

The extent of the area, the high percentage of fiber of superior quality in the rock, and the easy access to a seaport make the Bar-

²⁹ Hall, A. L., Asbestos in the Union of South Africa: Union of South Africa Geol. Survey Mem. 12, 1918, p. 59.

Hollis-100-401



Bureau of Mines

ASBESTOS

one-half inch long is rejected. Mines have been discovered, but production has never been large. In the Carolina district occurs on the north bank of the Komati River and 47 miles by road from the Komati River. The occurrence is of the usual type—cross-fibered, parallel thin seams inclined at a steep angle, of good quality, but a large proportion of it is waste. Reserves are extensive. A mill was in operation but has been small.

W. H. Hall directed attention to a chrysotile deposit in eastern Transvaal, but serious development did not begin until after 1921. The deposit occurs in a belt of quartzites lying about 3 miles from Kaapstad, by road from Godwin River station. The deposit of the serpentine has been traced for a distance of 2 miles. Further exploration may be expected. Underground mines have been developed at the eastern and western ends of a 2-mile belt of fiber.

Barberton Mines, Ltd. (subsidiary of Turner & Newall, Ltd.) operates the eastern end. Here the most productive fiber is the "ribbon line" or "ribbon line." In a typical section narrow parallel fiber seams so numerous that the fiber content may run as high as 10 percent. Beneath this band is a 6-foot mass of chrysotile veins are fewer and thicker. The veins are long. Figure 6 shows the abundance of fiber in the veins. The fiber is of high quality and is massive serpentine. Records for one year from the upper beds amounted to 30 percent, while from the lower beds it was 10 percent. The recovery for the mine was 15 to 17 percent. The output was of spinning class, as shingle stock. The mine is a small mine, producing about 1,000 tons of fiber a month in 1930. The mine occupies the eastern end of the serpentine belt. The veins are more or less parallel and resemble each other. Thin seams are close together, while the thicker ones are more widely spaced. A majority of the veins are from one-sixteenth to one-half inch in thickness, and in 1930 the mine was producing about 1,000 tons of fiber a month.

The products of the mines were transported 18 miles in wagons hauled by oxen or mules. The construction of a 5.9-mile aerial tramway over a 2,000-foot drop made possible a reduction in haulage cost to the amount formerly paid. From this point the material is hauled by a short-rail haul to Lourenço Mar, the port of shipment.

The high percentage of fiber of superior quality and the easy access to a seaport make the Barberton mines one of the most important in the Union of South Africa.

Union of South Africa: Union of South Africa Geol.

Barberton Mines



Barberton Mines

FIGURE 6.—Asbestos veins in the New Amantlous Mines, Ltd., Barberton district, Transvaal, South Africa.

berton district one of the most important asbestos centers in the world. The development of mines in this region is reflected in the rapid increase in production of the Transvaal, beginning in 1923, as shown in the table of production in the Union of South Africa by Provinces.

AMOSITE

This unique variety of asbestos occurs most prominently near Penge, about 30 miles north of Lydenburg in northeastern Transvaal. The belt extends from the Steelpoort River northwestward along the basin of the Olifants River, thence westward to Chuniespoort. The asbestos occurs in cross-fiber veins interbedded with siliceous ferruginous slates associated with intrusive diabase sills. The principal mines, the Egnep and Amosa, about 1½ miles apart, are situated near Penge. Fiber is obtained from surface workings and from several levels underground. A number of smaller mines and open-pit workings have been operated in the western part of the belt near the Malips River. The fiber veins show remarkable continuity over long distances, and reserves are extensive.

The most notable feature of the asbestos is its length, which attains a maximum of approximately 11 inches and averages about 6 inches for the commercial product. When the material was first marketed difficulties were encountered, because standard equipment for manufacturing asbestos products was not well adapted for fiber of such great length. With subsequent adjustments, the material was prepared extensively for spinning, magnesia pipe covering, and manufacture of other asbestos products.

The future of amosite is uncertain. The claim is made that the relatively large production recorded for 1928 and 1929 was stimulated chiefly by a shortage of chrysotile. As a similar shortage is unlikely to occur again for many years, future activity in the amosite fields will depend primarily on the serviceability of this type of fiber.

CROCIDOLITE

Crocidolite occurs in the western part of the amosite belt, a region designated as the Pietersburg district. It is associated closely with amosite, both varieties being obtained from some of the workings. The blue asbestos is obtained chiefly from open pits near and east of the Malips River. The maximum length of fiber is about 3 inches. Production never has been large. In 1928, the last year for which it was recorded, the output was 66 tons.

ANTHOPHYLLITE

Anthophyllite occurs in the Zoutpansberg district, 50 miles west of Waterpoort Siding. The fibrous structure has evidently been developed under the influence of weathering, therefore the persistence of usable fiber at depth is doubtful. Asbestos taken from pits not exceeding 30 feet deep was used some years ago for boiler covering, but there has been little recent production.

NATAL

The most important asbestos deposit in Natal is in the Tugela River Valley east of Kranz Kop, Zululand. Chrysotile asbestos occurs here in vertical cross-fiber veins near contacts of aplite dikes

with green serpentine. Most of the fiber is short and inferior. Production since 1910 has ranged from 2 to 62 tons a year.

A deposit of brittle tremolite asbestos has been worked near the Klip River, 45 miles from Dundee, a region where transportation is very difficult. The small amount of material obtained was used for the manufacture of asbestos brick and for insulation.

OTHER AFRICAN OCCURRENCES

Asbestos has been reported at various other points in Africa. The fiber occurs in Bechuanaland, Nyasaland, and in serpentine in the northwest corner of Swaziland. The latter deposit is evidently not far from the productive Barberton district of the Transvaal. Anthophyllite occurs at Morogoro in what was formerly German East Africa, and chrysotile was reported near Macequece, Portuguese East Africa, in 1929.

SOVIET RUSSIA

Before the World War, Russia ranked next to Canada as an asbestos-producing country. During the revolutionary period the industry was almost at a standstill, but since then it has been revived on an extensive scale. The following table shows production since 1913. The figures were obtained from the annual chapters on asbestos in Mineral Resources of the United States, published by the United States Geological Survey until 1924 and thereafter by the United States Bureau of Mines, and from the corresponding chapters in the Minerals Yearbook, published by the Bureau of Mines.

Asbestos production in Russia, 1913-36

Year	Metric tons	Year	Metric tons	Year	Metric tons
1913	17,494	1921	2,604	1929	29,520
1914	15,691	1922	3,215	1930	54,083
1915	8,779	1923	14,780	1931	64,674
1916	8,192	1924	18,456	1932	59,800
1917	(1)	1925	12,330	1933	71,700
1918	(1)	1926	15,334	1934	92,200
1919	(1)	1927	21,156	1935	95,500
1920	1,475	1928	26,492	1936	

¹ Data not available.

² Year ended Sept. 30.

The great increase since 1929 does not affect world markets as much as might be expected, because increasingly large quantities are used for home consumption. According to recent figures,³⁰ the percentage of Russian asbestos exported dropped from 51.1 in 1922-23 to only 20.5 in 1931. This condition reflects the rapidly expanding manufacture of asbestos products in Soviet Russia. However, these figures tell only part of the story, for they do not indicate the percentage of spinning and higher-grade mill fibers exported. The 50 to 80 percent of the production remaining in the country may consist largely of the shorter fibers, which are of minor importance in world trade. The asbestos region providing the major part of Russian production comprises what is known as the Bajenova district in the Urals. Principal

³⁰ Economic Review of the Soviet Union, Oct. 15, 1932, p. 365.

production is at Asbest, about 90 kilometers (Ekaterinburg). Minor production from the Alapaevsk area in the north and the northwest. The largest mine is outside the Bajenova district the only Minusinsk, on the Yenisei River in eastern border. Here the deposits, said to be of as those of Arizona, have been worked in 1,490 tons being reported in 1905. Asbestos in various parts of the Altai Mountains.

A chrysotile asbestos deposit on the Leirick of the Caucasus, about 98 kilometers, has been described.³¹ The mineral occurs in granite. In general from 2 to 7 mm, with a maximum of 25 mm is said to be of good quality, but the deposit has been worked in a small way and development is anticipated.

A small output was reported from 1907 to 1909.

Chrysotile has been reported in other parts of the Ural Mountains in the Permian Siberia; on the upper course of the river near the source of the Kuban River in Turkestan.

At Asbest, in the Bajenova district, now centered, 20 or more open pits are under control of the State since 1918: Uralasbest Trust since 1921. Asbestos was first discovered in 1710, and systematic development began in 1885. The asbestos-bearing zone is 21 kilometers long and 200 to 1,200 meters wide, which are bounded by schist or gneiss on the east. The asbestos is coarse, some of which attain a length of 10 feet. The highest percentage of asbestos is in the cross-fiber veins, which usually is regarded as being higher in quality than the other, but Ru Keyser³² has pointed out that the Russian asbestos is of a lower quality than the Quebec. This probably accounts for the greater use of asbestos upon weathering. Russian asbestos is relatively shallow overburden and are relatively shallow, which may account for the condition of the fiber compared with the Quebec. While the percentage of asbestos in the total commercial fiber is about the same as in the Quebec.

³¹ Tatarinov, P. The Leirick (Beden) Deposit, Caucasus. Neues Jahrb. Mineral. Geol., Referate II, 1933, p. 335.
³² Ru Keyser, W. A., Chrysotile Asbestos in the Ural Mountains, August 1933, p. 335.

st of the fiber is short and inferior. Produced from 2 to 62 tons a year. Solit bestos has been worked near the Dunas, a region where transportation is nount of material obtained was used for brick and for insulation.

AFRICAN OCCURRENCES

at various other points in Africa. The id, Nyasaland, and in serpentine in the ind. The latter deposit is evidently not Barberton district of the Transvaal. orogoro in what was formerly German was reported near Macequece, Portuguese

VIET RUSSIA

Russia ranked next to Canada as an asbestiferous country during the revolutionary period the industry fell into disuse, but since then it has been revived. The following table shows production since 1917. The figures for the United States, published by the Bureau of Mines, are given until 1924 and thereafter by the United States Geological Survey, and from the corresponding chapters published by the Bureau of Mines.

tion in Russia, 1913-36

Year	Metric tons	Year	Metric tons
-----	2,604	1929	29,420
-----	3,213	1930	34,083
-----	4,780	1931	54,674
-----	8,456	1932	39,800
-----	12,330	1933	71,700
-----	18,334	1934	92,290
-----	21,156	1935	95,500
-----	26,492	1936	

* Year ended Sept. 30.

does not affect world markets as much increasingly large quantities are used ding to recent figures,³⁰ the percentage dropped from 51.1 in 1922-23 to only reflects the rapidly expanding manu- Soviet Russia. However, these figures they do not indicate the percentage of fibers exported. The 50 to 80 percent the country may consist largely of the or importance in world trade. the major part of Russian production Bajenova district in the Urals. Prin-

. Oct. 15, 1932, p. 365.

cial production is at Asbest, about 90 kilometers northeast of Sverdlovsk (Ekaterinburg). Minor production in this district is obtained from the Alapaevsk area in the north and from the Neviansk area in the northwest. The largest mine in the latter is the Kramouralsk. Outside the Bajenova district the only notable production is from Minusinsk, on the Yenisei River in eastern Siberia near the Mongolian border. Here the deposits, said to be of the same general character as those of Arizona, have been worked in a small way, a production of 1,490 tons being reported in 1905. Asbestos also occurs in serpentine in various parts of the Altai Mountains southwest of Minusinsk.

A chrysotile asbestos deposit on the Laba River in the Maikop district of the Caucasus, about 98 kilometers from a railway, has recently been described.¹¹ The mineral occurs in veins in serpentine which have been intruded by granite. In general, the fibers range in length from 2 to 7 mm, with a maximum of 25 mm (10 inches). The asbestos is said to be of good quality, but the reserves are not extensive. The deposit has been worked in a small way since 1930 and a larger development is anticipated.

A small output was reported from the Katun River district from 1907 to 1909.

Chrysotile has been reported in other parts of Russia, as follows: In a peridotite-serpentine belt in the Province of Transbaikalia, southern Siberia; on the upper course of the Yenisei River in the far north; near the source of the Kuban River east of the Crimea; and in Turkestan.

Turkestan.

At Asbest, in the Bajenova district, where the asbestos industry is now centered, 20 or more open pits are worked. They have been under control of the State since 1918 and have been operated by the Uralasbest Trust since 1921. Asbestos was first discovered here about 1710, and systematic development under direction of Baron Girard began in 1885. The asbestos-bearing serpentized intrusion is about 21 kilometers long and 200 to 1,200 meters wide. It consists of peridotites, which are bounded by schist or slate on the west and by granite on the east. The asbestos is confined to ellipsoidal masses of serpentine, some of which attain a length of 3,500 feet and a width of 1,000 feet. The highest percentage of asbestos is found in the central parts of these masses. The cross-fiber veins generally run north and south with a vertical dip. Slip fiber appears in places. Russian asbestos usually is regarded as being higher in iron than the Canadian fiber, but Ru Keyser²² has pointed out that Canadian asbestos is high in ferric iron, whereas in the Russian fiber ferrous iron predominates. This probably accounts for the greater discoloration of the Ural asbestos upon weathering. Russian asbestos deposits have a comparatively shallow overburden and are therefore weathered quite extensively, which may account for the somewhat harsher and less silky condition of the fiber compared with the Canadian. Asbestos occurring below the 50-foot level more nearly resembles that obtained in Quebec. While the percentage of spinning fiber in the Russian deposits is a little higher than that in the Canadian, the percentage of total commercial fiber is about the same.

² Tatarinov, P. The Liba (Beden) Deposit of Chrysotile Asbestos in the North Caucasus: Neues Jahrb. Mineral. Geol., Referate II, 1936, pp. 249-250.
³ Ru Kerser, W. A. Chrysotile Asbestos in the Bajenova District, U. S. S. R. Izv. i Min. Jour., August 1933, p. 353.

Extensive and systematic core-drilling indicates a reserve for the entire Bajenova district of more than 3,000,000 metric tons of fiber within 50 feet of the surface. This estimate is based on only a 2-percent recovery; with a probable 4½-percent recovery this quantity would be more than doubled. As the deposits extend far below the 50-foot level it is apparent that the reserves are very great. The reserves of asbestos in the Soviet Union were estimated on January 1, 1934, at 19,400,000 tons.

Much of the upper rock is soft enough to be mined with picks and shovels. Before the revolution, crude methods of mining and fiber recovery were practiced, and only the longest grades of fiber—crudes, spinning, and shingle stock—were recovered. However, with increased mechanization and development of markets for the shorter grades, most of the fiber in the rock mined is now separated and sold. As the flow of water is heavy below the 50-foot level, the workings consist principally of broad, shallow, open pits, although some underground development has taken place. A narrow-gauge railway was built from Asbest to Bajenova (34 km) in 1927, making possible more intensive development.

There are three major groups of mines in this district. The most northerly, known as the "Proletariat", produces shorts almost exclusively. The two major producing groups are the October (central) and the Ilyinski (southern). Several mills with very crude equipment have been operated for many years. Since 1931, newer and more efficient mills have been built and new equipment has been installed in some of the old ones. A large asbestos-products industry has been established in Russia, and crudes and spinning fibers which in former years constituted a large part of the output have been supplemented by mill products similar to those produced in Canada.

ITALY

Italy is called "the cradle of the asbestos industry", because the mining of asbestos and the manufacture of its products began on an industrial scale in that country. Italy also is unique in that much of the fiber produced in early years consisted of tremolite, a variety of amphibole asbestos little used elsewhere. Before discovery of the Canadian deposits, Italy was the chief source of supply of asbestos for both Europe and America. After the Quebec deposits were developed, the Italian industry experienced little growth. Production from 1898 to 1934 is shown in the following table:

Annual production of asbestos in Italy, 1898-1934, in metric tons

Year	Tons	Year	Tons	Year	Tons
1898.....	131	1911.....	170	1924.....	2,160
1899.....	81	1912.....	169	1925.....	2,105
1900.....	126	1913.....	175	1926.....	2,900
1901.....	243	1914.....	171	1927.....	3,840
1902.....	243	1915.....	163	1928.....	4,950
1903.....	202	1916.....	82	1929.....	2,847
1904.....	182	1917.....	85	1930.....	851
1905.....	220	1918.....	60	1931.....	632
1906.....	209	1919.....	98	1932.....	1,284
1907.....	359	1920.....	165	1933.....	3,267
1908.....	359	1921.....	420	1934.....	2,232
1909.....	190	1922.....	540		
1910.....	175	1923.....	1,538		

One important deposit is in the Susa Torino near the French border. Here it is particularly well-adapted to the manufacture of asbestos in the mountains 8,000 to 9,000 feet above sea level. It is along the Aosta Valley near Ivrea in 146 miles from Turin. Tremolite asbestos mountains in this locality since 1865, but it was not until 1905. The fibers are long and high heat, but owing to the difficulty of separating the fibers, it is well-adapted to weaving. A third source of asbestos is near Sondrio in northern Italy. This is a long-fibered variety similar to that of the Balangero mine situated in Torino.

The Balangero mine has been the chief producer of asbestos of this district is chrysotile, the production in 1925 and nearly 5,000 tons. In 1928, modern mechanical production. The fiber is said to be inferior to that of the Balangero for spinning purposes. It commands prices of 100 francs per ton and is used chiefly for the manufacture of asbestos in Sardinia and Corsica.

Italy is both an importer and an exporter of asbestos. All the fiber required for textile use and the shorter grades are imported, while the longer grades and other nonspinning grades are exported in Corsica.

CYPRUS

The chief asbestos deposits of Cyprus are Troodos in the west-central area. The rock is largely of olivine is altered to serpentinite. The fiber chrysotile asbestos occurs in the form of thin layers that have a maximum thickness of about 1/16 inch. Shearing evidently have exerted a great influence on the fibrous structure. The asbestos content is 1 to 2 percent. The deposit is evidently rich in reserves is available.

The fiber is of the chrysotile variety and is used for spinning, but small shipments of asbestos have been made since 1934. Milled fiber is graded into three grades. The standard grade, designated "No. 1", comprises about 90 percent of product.

About 1925 an aerial ropeway 18 miles long was constructed for conveying the asbestos from Amian to the coast. The construction cost was about £22,000. Exporting asbestos by means of the ropeway was given as 6s. 9d. a ton, whereas in 1924 it was 10s. 6d. per ton. The industry attained considerable development but thereafter the demand fell off greatly. Recoveries were conducted at a substantial rate in following years.

core-drilling indicates a reserve for the more than 3,000,000 metric tons of fiber.

The estimate is based on only a 2-per-cent 75-percent recovery this quantity. As the deposits extend far below the that the reserves are very great. The Soviet Union were estimated on January

soft enough to be mined with picks and iron, crude methods of mining and fiber only the longest grades of fiber—crudes, were recovered. However, with in-development of markets for the shorter rock mined is now separated and sold. Below the 50-foot level, the workings shallow, open pits, although some underground place. A narrow-gauge railway was over (34 km) in 1927, making possible

ups of mines in this district. The most "proletariat", produces shorts almost excluding groups are the October (central)

Several mills with very crude equipment many years. Since 1931, newer and more built and new equipment has been used. A large asbestos-products industry, and crudes and spinning fibers which a large part of the output have been similar to those produced in Canada.

ITALY

of the asbestos industry", because the manufacture of its products began on an annual basis. It is also unique in that much of its production consisted of tremolite, a variety of asbestos found elsewhere. Before discovery of the chief source of supply of asbestos in Canada. After the Quebec deposits were discovered, the industry experienced little growth. Production in the following table:

Asbestos in Italy, 1898-1934, in metric tons

Year	Tons	Year	Tons
1898	170	1921	2,160
1899	169	1922	2,105
1900	175	1923	2,900
1901	171	1924	3,540
1902	163	1925	4,950
1903	82	1926	2,847
1904	85	1927	551
1905	60	1928	832
1906	98	1929	1,284
1907	165	1930	3,267
1908	420	1931	2,252
1909	540		
1910	1,535		

One important deposit is in the Susa Valley region of western Torino near the French border. Here tremolite asbestos, which is particularly well-adapted to the manufacture of acid filters, is mined in the mountains 8,000 to 9,000 feet above sea level. A second deposit is along the Aosta Valley near Ivrea in northeastern Torino, about 46 miles from Turin. Tremolite asbestos has been mined in the mountains in this locality since 1865, but there has been little activity since 1905. The fibers are long and highly resistant to acids and heat, but owing to the difficulty of separating them, they are not well-adapted to weaving. A third source of supply of tremolite asbestos is near Sondrio in northern Lombardy. This asbestos also is a long-fibered variety similar to that of the Torino deposits.

The Balangero mine situated in Torino about 20 kilometers north of Turin has been the chief producer during recent years. The asbestos of this district is chrysotile, the output of which was about 2,000 tons in 1925 and nearly 5,000 tons in 1928, the year of greatest production. In 1928, modern mechanical mining equipment was introduced. The fiber is said to be inferior to Canadian chrysotile for spinning purposes. It commands prices ranging from \$75 to \$150 a ton and is used chiefly for the manufacture of artificial stone.

Asbestos was produced in Sardinia in 1926, and deposits are reported in Corsica.

Italy is both an importer and an exporter of asbestos. Practically all the fiber required for textile use and considerable quantities of the shorter grades are imported, while small quantities of filter asbestos and other nonspinning grades are exported.

CYPRUS

The chief asbestos deposits of Cyprus occur at Amiandos, on Mount Troodos in the west-central area. A peridotite plug consisting largely of olivine is altered to serpentine on the periphery. Short-fiber chrysotile asbestos occurs in the serpentine in irregular veins that have a maximum thickness of about one-half inch. Faulting and shearing evidently have exerted a definite influence on development of the fibrous structure. The asbestos content of the rock runs only 1 to 2 percent. The deposit is evidently extensive, but no information on reserves is available.

The fiber is of the chrysotile variety and generally too short for spinning, but small shipments of spinning fiber were reported in 1934. Milled fiber is graded into three classes—standard, shorts, and fines. The standard grade, designated "shingle stock", is said to comprise about 90 percent of production.

About 1925 an aerial ropeway 18 miles long was constructed for conveying the asbestos from Amiandos to the seaport at Limasol. The construction cost was about \$2,320 a mile. The cost of transporting asbestos by means of the ropeway, including interest on investment and complete amortization of the equipment in 10 years, was given as 6s. 9d. a ton, whereas motor-wagon or animal transport formerly employed on the 37-mile winding road cost 20 to 25s. a ton.

The industry attained considerable importance from 1926 to 1929, but thereafter the demand fell off greatly. It is reported that operations were conducted at a substantial loss in 1932, but the industry recovered greatly in following years.

Production has been in the hands of a single company. The Cyprus Asbestos Co., Ltd., operated for several years prior to 1927, but in that year the Cyprus Trading Corporation, Ltd., was organized and absorbed the former company. In 1932, the newer firm was reorganized as Cyprus & General Co., a British organization, and in 1936 the assets of the latter company were acquired by the Tunnel Asbestos Cement, Ltd., a subsidiary of The Tunnel Portland Cement Co. of West Thurrock, England. The asbestos is used by the company for manufacture of asbestos-cement products in England.

As there are no local asbestos-products plants, the entire output is exported, principally to European countries. Exports began in 1906. Figures for tonnage and value of exports over a period of years are shown in the following table:

Asbestos exported from Cyprus, 1922-36

Year	Long tons	Value	Year	Long tons	Value	Year	Long tons	Value
1922.....	2,283	£22,699	1927.....	10,904	£207,562	1932.....	1,600	£27,214
1923.....	2,115	57,115	1928.....	11,579	231,692	1933.....	4,604	144,053
1924.....	4,372	80,070	1929.....	13,796	292,971	1934.....	7,590	73,562
1925.....	3,290	54,639	1930.....	5,469	116,092	1935.....	7,514	50,174
1926.....	6,541	121,257	1931.....	3,571	66,381	1936.....	9,506	80,343

¹ Reported by Cyprus & General Asbestos Co., Ltd.

Of the fiber exported during 1934, 2,889 tons were reported as destined for the United States.

The following table, taken from the Annual Report of the Inspector of Mines and Labour, 1934, indicates the scope of recent operations and the ratio of fiber to rock.

Rock mined, rock treated, and fiber produced in Cyprus, 1931-34, in long tons

	1931	1932	1933	1934
Rock mined.....	113,738	155,765	354,178	711,646
Rock treated.....	25,966	36,819	76,784	153,863
Asbestos fiber produced.....	1,138	1,520	3,494	7,081

OTHER FOREIGN DEPOSITS

Asbestos deposits of minor commercial importance at present are covered in the following pages, by countries, arranged in alphabetical order.

ARGENTINA

Deposits of asbestos are reported at Alta Gracia, Cordoba; Sierra de la Cortadera, Mendoza; Fiambala, Catamarca; and in the Provinces of San Luis and San Juan. The fiber is said to be too short for spinning or weaving and production has been very small. The only available figures show an output of about 42 short tons in 1926 and of less than a ton in 1927.

AUSTRALIA

WESTERN AUSTRALIA

The most important recent development is at the Sherlock property, in the Roebourne from the port of Balla Balla. Here chrome veins that are restricted to a relatively small area, separated by a dolerite dike, have a thickness of 1,000 feet. The percentage of fiber 2 in to be much higher than in the principal world, and is of good quality. Up to 1936 had been mined by contract laborers and no systematic mining had been followed. Creation of a pumping plant were started.

Other producing areas are Lionel in the Soansville and Cooglegong in the Marble Posits are all in the Pilbarra Goldfield area. Tile occurs in veins up to 30 inches wide with of 6 inches. The veins have been proved. Fiber from this area shipped to London at £150 a ton.

Asbestos was discovered in the Marble deposit has been reported 15 miles west. Veins range from threads up to 6 inches.

The following table compiled by Elford Western Australia for a period of years.

Production of asbestos in Western Australia

Year	Quantity, tons	Value	Locality where mined	Year
1921.....	202	£12,221	Lionel.	1927.....
	32	1,360	Cooglegong.	
1922.....	179	7,350	Lionel.	
	2.5	250	Cooglegong.	
1923.....	111	3,885	Lionel.	1930.....
	3	150	Cooglegong.	
1924.....	73	2,206	Lionel.	1931.....
1925.....	50	1,619	Do.	1932.....
1926.....	91	2,436	Do.	
	14	292	Roebourne.	

SOUTH AUSTRALIA

A promising deposit of crocidolite occurs about 248 miles north of Adelaide. The magnesite limestone. The maximum thickness, but most of the asbestos is shown in a drift. Unfortunately, the ore body and nothing is known of its continuation. It occurs as bunches and veins in dolomite, miles from Adelaide. It is said to

² Elford, Harold S., Australian Nonmetallic Minerals, vol. 25, Sept. 5, 1933, p. 396.

134342°-37-4

ASBESTOS

the hands of a single company. The operated for several years prior to 1927, . Tr ing Corporation, Ltd., was organ- r co. ny. In 1932, the newer firm was neral Co., a British organization, and in company were acquired by the Tunnel sidiary of The Tunnel Portland Cement and. The asbestos is used by the com- stos-cement products in England. os-products plants, the entire output is ean countries. Exports began in 1906. e of exports over a period of years are

ed from Cyprus, 1922-36

Long tons	Value	Year	Long tons	Value
10,904	£207,562	1922	1,600	£27,214
11,579	231,692	1923	4,604	144,088
13,796	292,971	1924	7,590	173,562
5,400	116,092	1925	7,514	50,174
3,571	46,381	1926	9,506	50,343

. Ltd.

g 1934, 2,389 tons were reported as from the Annual Report of the In- 1934, indicates the scope of recent to rock.

produced in Cyprus, 1931-34, in long tons

	1931	1932	1933	1934
	113,738	155,765	354,178	711,646
	25,966	36,819	76,754	153,865
	1,138	1,520	3,494	7,081

EIGN DEPOSITS

nmmercial importance at present are : countries, arranged in alphabetical

INTINA

ed at Alta Gracia, Cordoba; Sierra bala, Catamarca; and in the Prov-. The fiber is said to be too short duction has been very small. The put of about 42 short tons in 1926

FOREIGN DEPOSITS

45

AUSTRALIA

WESTERN AUSTRALIA

The most important recent development in Western Australia is the Sherlock property, in the Roebourne district, about 24 miles from the port of Balla Balla. Here chrysotile asbestos occurs in veins that are restricted to a relatively narrow zone. Two main veins, separated by a dolerite dike, have been proved for a length of 1,000 feet. The percentage of fiber 2 inches long and over is said to be much higher than in the principal producing regions of the world, and is of good quality. Up to 1934, about 1,000 tons of fiber had been mined by contract laborers and shipped to Europe, but no systematic mining had been followed. Shaft sinking and the erection of a pumping plant were started in 1934.

Other producing areas are Lionel in the Nullagine district, and Soansville and Cooglegong in the Marble Bar district. These deposits are all in the Pilbarra Goldfield area. At Soansville chrysotile occurs in veins up to 30 inches wide with a maximum fiber length of 6 inches. The veins have been proved over a length of 1,200 feet. Fiber from this area shipped to London commanded a price of £120 to £150 a ton.

Asbestos was discovered in the Marble Bar area in 1909. A large deposit has been reported 15 miles west of Cooglegong. Chrysotile veins range from threads up to 6 inches wide.

The following table compiled by Elford,²² shows production in Western Australia for a period of years.

Production of asbestos in Western Australia, 1921-32

Year	Quantity, tons	Value	Locality where mined	Year	Quantity, tons	Value	Locality where mined
1921	202	£12,221	Lionel.	1927	10.9	£304	Lionel.
	32	1,360	Cooglegong.	1928	11.7	792	Soansville and Lionel.
1922	179	7,350	Lionel.	1929	191	8,568	Sherlock.
	2.5	250	Cooglegong.		63	6,113	Soansville.
1923	111	3,865	Lionel.	1930	65	4,228	Pilbarra.
	3	150	Cooglegong.		17	500	West Pilbarra.
1924	73	2,208	Lionel.	1931	108	1,446	Roebourne.
1925	30	1,619	Do.	1932	110	1,762	Do.
1926	91	2,436	Do.				
	14	292	Roebourne.				

SOUTH AUSTRALIA

A promising deposit of crocidolite occurs 10 miles north of Hawker, about 248 miles north of Adelaide. Apparently it is derived from magnesian limestone. The maximum length of the fiber is about 2 inches, but most of the asbestos is short and matted. Two shafts 40 feet apart have been sunk to a depth of 26 feet and connected with a drift. Unfortunately, the ore body is faulted at the 26-foot level, and nothing is known of its continuation. Bluish gray crocidolite occurs as bunches and veins in dolomitic limestone at Robertstown, 83 miles from Adelaide. It is said to have excellent heat-insulating

²² Elford, Harold S., Australian Nonmetallic Minerals. II. Asbestos: Chem. Eng. and Min. Review, vol. 25, Sept. 5, 1933, p. 396.

qualities. Other occurrences have been noted at Kapunda, Truro, Mount Baker, Gumeracha, and Williamstown.

Chrysotile of good quality occurs at Mimbic near Cowell. Veins attaining a maximum width of 2 inches appear in serpentine traversing marble. According to report, the asbestos-bearing rock would yield 0.5 percent crude and 15 to 20 percent mill fiber. Annual production ranging from 5 to 21 tons has been noted in South Australia for various years.

QUEENSLAND

Asbestos has been noted in many localities in Queensland, mainly within the serpentine belt northeast and north of Rockhampton, extending from Balnagowan near the Fitzroy River to Marlborough. The deposits at Princhester and Marlborough appear to be the most important, though little is yet known of their quality or extent. A small amount of fiber has been mined, and it is reported that it sold for £25 a ton. On Marlborough Creek near its junction with the Fitzroy River well-defined veins having a maximum fiber length of 1½ inches are reported. Queensland asbestos is said to be coarse in texture and of low strength. It is well adapted for asbestos-magnesite flooring, for boiler covering, and as an ingredient of paints.

NEW SOUTH WALES

A chrysotile-asbestos deposit was worked in a small way near Barabara for several years following 1916. Short-fiber chrysotile occurs in a belt of serpentine about ½ mile wide. The area is greatly faulted, which makes mining difficult. A small mill, similar to those in Quebec, was operated for a time, and fiber recovery is said to have been 5.75 percent of the rock treated. Crocidolite occurs near Gundagai and Broken Hill, but the percentage of recoverable fiber is said to be low.

TASMANIA

An occurrence of asbestos at Beaconsfield was discovered more than 30 years ago, a production of 200 tons being recorded in 1899. It is of the chrysotile variety, occurring in irregular and noncontinuous veins in a serpentine belt 2 miles long and 1½ miles wide. The fiber rarely reaches 1 inch in length; commonly, it is ⅜ inch long. The asbestos-bearing area is not very large, and the veins so far uncovered are too scattered and irregular to encourage deep working. A limited production of asbestos of fair quality is possible. The fiber has a higher iron content than Canadian asbestos. A mill for ore treatment was erected in 1917, but very little fiber has been produced.

A promising deposit of asbestos in serpentine is reported on the western shore of Macquarie Harbor, 16 miles south of Strahan. It is undeveloped.

AUSTRIA

Short-fibered amphibole, known as "microasbestos", is mined in Austria near the Hungarian frontier. It is mixed with both asphalt and concrete used for road surfaces.

BRAZIL

An amphibole-asbestos deposit is reported near Rio Branco, Pernambuco, and asbestos of uncertain type occurs in Minas Geraes.

BULGARIA

Amphibole asbestos of the anthophyllite area in the eastern Rhodope Mountains, production of several hundred tons was reported.

CHINA

Asbestos said to be of the chrysotile variety in the Provinces in China, including Chihli, Jehol, Szechuan, Hupeh, and Kwangtung. The are in the Laiyuan district, Chihli (Hopeh "Hopei" in the footnote reference is not an asbestos occurs in numerous generally p limestone. Maximum fiber length is about has evidently resulted from metamorphism limestone under the influence of granite similar in origin to the Arizona deposits described by Hou.³⁴ Reserves in the are tons. The longer grades are of spinning been worked since 1914. Production in The fiber is conveyed on horseback to Yi from there to Tientsin by train. It supplies the latter city. Four such plants were reported are mostly for domestic consumption Japan.

Asbestos mines about 5 miles from Chihli worked in a small way for more than 20 shipped to factories at Osaka and Tokyo.

CZECHOSLOVAK

According to the Bureau of Foreign a sole producer of asbestos in the country Verwertung von Asbest Gesellschaft m. of 2,600 metric tons in 1935, 2,100 reported. No data are available as to product, nor has the Bureau of Mines : bestos occurrence in this country.

FINLAND

Asbestos production is confined to a east-central Finland, where 1,000 to : were mined annually until recent years dently increased. According to a Brit Jopson, production in 1934 was 14,00 9,200 tons in 1933 and 6,000 tons in 19 reported by Borgström.³⁶ Finnish asbe cal composition: SiO₂, 56.57; Al₂O₃, CaO, 0.57; H₂O+, 4.89; H₂O-, 0.14.

³⁴ Hou, T. F., Notes on the Asbestos Deposit (English); Nat. Geol. Survey China Geol. Bull. 2

³⁵ Bureau of Foreign and Domestic Commerce, Apr. 15, 1936, p. 9.

³⁶ Borgström, L. H., Finnish and Foreign Asbe 1935, pp. 105-107.

ASBESTOS

urrences have been noted at Kapunda, Truro, and Williamstown.

Quality occurs at Mimbric near Cowell. Veins of 2 inches appear in serpentine traversing the report, the asbestos-bearing rock would be 15 to 20 percent mill fiber. Annual production of 5 to 21 tons has been noted in South Australia.

QUEENSLAND

Asbestos is found in many localities in Queensland, mainly in the northeast and north of Rockhampton. Asbestos occurs near the Fitzroy River to Marlborough. Asbestos and Marlborough appear to be the most important. It is not known of their quality or extent. Asbestos has been mined, and it is reported that it sold near Marlborough Creek near its junction with the other veins having a maximum fiber length of 1/2 inch. Queensland asbestos is said to be coarse in texture. It is well adapted for asbestos-magnetite, and as an ingredient of paints.

NEW SOUTH WALES

Asbestos was worked in a small way near Barrowfield 1918. Short-fiber chrysotile occurs about 1/2 mile wide. The area is greatly irregular. A small mill, similar to those of a time, and fiber recovery is said to have been treated. Crocidolite occurs near Gunter. The percentage of recoverable fiber is said to be 10 percent.

TASMANIA

Asbestos at Beaconsfield was discovered more than 100 years ago. It is found in irregular and noncontinuous veins, 1/2 mile long and 1 1/2 miles wide. The fiber length; commonly, it is 1/8 inch long. The veins are very large, and the veins so far underground irregular to encourage deep working. Asbestos of fair quality is possible. The fiber is better than Canadian asbestos. A mill for ore has been built, but very little fiber has been produced. Asbestos in serpentine is reported on the west coast, 16 miles south of Strahan. It is said to be 10 percent.

AUSTRIA

Asbestos known as "microasbestos", is mined in the Tyrol. It is mixed with both asphalt and mica.

BRAZIL

Asbestos is reported near Rio Branco. Peridotite type occurs in Minas Geraes.

BULGARIA

Amphibole asbestos of the anthophyllite type occurs over a large area in the eastern Rhodope Mountains of southern Bulgaria. A production of several hundred tons was reported unofficially in 1933.

CHINA

Asbestos said to be of the chrysotile variety occurs in a number of Provinces in China, including Chihli, Jehol, Chahar, Suiyuan, Shensi, Szechuan, Hupeh, and Kwangtung. The most important deposits are in the Laiyuan district, Chihli (Hopei) Province. (The spelling "Hopei" in the footnote reference is not now recognized.) Here the asbestos occurs in numerous generally parallel cross-fiber veins in limestone. Maximum fiber length is about 1 1/2 inches. The asbestos has evidently resulted from metamorphism of a siliceous magnesian limestone under the influence of granite intrusions, therefore it is similar in origin to the Arizona deposits. The deposits have been described by Hou.³⁴ Reserves in the area are estimated at 400,000 tons. The longer grades are of spinning quality. The deposit has been worked since 1914. Production in 1927 was 180 metric tons. The fiber is conveyed on horseback to Yixian (106 kilometers) and from there to Tientsin by train. It supplies manufacturing plants in the latter city. Four such plants were reported in 1927. Their products are mostly for domestic consumption, though some are shipped to Japan.

Asbestos mines about 5 miles from Chinchu, Manchuria, have been worked in a small way for more than 20 years. The output has been shipped to factories at Osaka and Tokyo, Japan.

CZECHOSLOVAKIA

According to the Bureau of Foreign and Domestic Commerce,³⁵ the sole producer of asbestos in the country is "Asbest" Gewinnung und Verwertung von Asbest Gesellschaft m. b. H. at Dobsina. An output of 2,600 metric tons in 1935, 2,100 in 1934, and 1,200 in 1933 has been reported. No data are available as to the nature or quality of the product, nor has the Bureau of Mines any previous record of an asbestos occurrence in this country.

FINLAND

Asbestos production is confined to a locality in Tuusniemi Parish, east-central Finland, where 1,000 to 1,500 tons of amphibole fiber were mined annually until recent years, when production has evidently increased. According to a British report on Finland by Keith Jopson, production in 1934 was 14,000 metric tons, compared with 9,200 tons in 1933 and 6,000 tons in 1932. According to an analysis reported by Borgström,³⁶ Finnish asbestos is of the following chemical composition: SiO₂, 56.57; Al₂O₃, 1.02; FeO, 5.72; MgO, 30.73; CaO, 0.57; H₂O+, 4.89; H₂O-, 0.14. This corresponds closely with

³⁴ Hou, T. F., Notes on the Asbestos Deposit of Laiyuan District, Hopei Province (in English); Nat. Geol. Survey China Geol. Bull. 25, March 1935, pp. 33-43.

³⁵ Bureau of Foreign and Domestic Commerce, Foreign Metals and Minerals: Circ. 2, Apr. 15, 1938, p. 9.

³⁶ Borgström, L. H., Finnish and Foreign Asbestos: Tek. Fören. Finland Förh., vol. 55, 1935, pp. 105-107.

the analysis of Georgia anthophyllite shown on page 4. The product of the mines, together with imported chrysotile fiber, is used in a factory at Helsingfors for the manufacture of asbestos board, Eternit tiles, packings, clothing, and gloves. A small quantity of actinolite has been mined in northern Finland.

FRANCE

Although France uses large quantities of asbestos in its manufacturing plants, a very small production of raw fiber has been recorded. It is interesting to note, therefore, that plans have been made to open an asbestos quarry in the Chateau-Ville-Vieille-et-Chateau-Queyras Commune in the Guil Valley at an altitude of about 2,000 meters.²⁷

A deposit of asbestos in Corsica was reported in 1921.

GERMANY

Discovery of a greenish white chrysotile in Bayerischer Wald, East Bavaria, was reported in 1935. This is said to be the only asbestos occurrence known in the country and is exceedingly important from the German standpoint because of increasing demands for home consumption. Imports of raw fiber to supply German asbestos-products plants increased from 7,582 tons in 1932 to 20,154 in 1934.

INDIA

The Hassan district of Mysore State has been the most productive area in India. Both chrysotile and amphibole fibers are available. A maximum production of over 1,700 long tons was obtained in 1920, but no output has been recorded since 1929. A small production has been noted from Seraikela State, Bihar and Orissa from 1921 to 1932. The material is evidently tremolite. Asbestos occurs in many other parts of India; but no estimates of reserves are available and none of the deposits seems capable of a large production of high-grade fiber. The development of a substantial asbestos-producing industry in India will depend largely upon the establishment of domestic asbestos-products plants that will absorb the short fibers. The following table, compiled from data assembled by Coulson,²⁸ shows production from 1906 to 1934.

Asbestos production in India, 1906-34

Year	Long tons	Value	Year	Long tons	Value	Year	Long tons	Value
1906.....	20.6		1920.....	1,818.0	£7,272	1928.....	158.5	£1,622
1908.....	18.0		1921.....	315.5	884	1929.....	318.4	1,206
1910.....	3.0	£ 6	1922.....	242.0	701	1930.....	33.2	88
1914.....	5.0	23	1923.....	245.0	659	1931.....	6.0	5
1916.....	5.0	22	1924.....	125.3	1,354	1932.....	90.0	677
1917.....	145.0	303	1925.....	16.0	361	1933.....		
1918.....	357.0	965	1926.....	58.4	786	1934.....	24.6	
1919.....	388.0	1,439	1927.....	67.7	1,011			

²⁷ Bureau of Foreign and Domestic Commerce, Foreign Trade Notes: Vol. 4, no. 5, May 27, 1935, p. 4.

²⁸ Coulson, A. L., Asbestos in the Ceded Districts of the Madras Presidency: With Notes on Its Occurrence in Other Parts of India: Mem. Geol. Survey of India, vol. 64, pt. 2, 1934, p. 249.

FOREIGN DEPOSITS

IRELAND

An asbestos deposit at Avoca, County Wick 1931. No data are available on variety, etc.

MEXICO

An unusually good sample of mountain cork matted, amphibole asbestos—was obtained by in 1931 from an undeveloped deposit near San ing to report. surface exposures indicate that tos has also been reported from the State of P

NEWFOUNDLAND

A deposit of chrysotile asbestos discovered thought to represent a continuation of the Qu available as to its quality or extent.

NEW ZEALAND

Chrysotile asbestos in veins as wide as 3 in tine area at least 3 miles long near Moun Takaka district of New Zealand. Both cross available. The cross-fiber veins occur princ inches thick surrounding cores of massive se veins. The fiber yield of the shells is estim: The deposit evidently is extensive and is un

PHILIPPINE ISLANDS

A deposit of amphibole asbestos occurs in Luzon Island. A considerable quantity wa pipe covering and similar uses.

In 1936 the Bureau of Mines received s this deposit, which, upon examination, prove strength and flexibility. Some of it wa spinning fiber, although it is not as s g asbestos. No data are available as to the deposit.

PORTUGAL

Asbestos of uncertain quality, probably extent, is reported from two places in the Alemtejo, southern Portugal. No product

SPAIN

Two occurrences of chrysotile asbestos 1920—one in the Pyrenees and the other tains. A production of 165 metric tons v but no later activity has been noted.

SWITZERLAND

Asbestos occurs in the Cantons of Gr Graubunden, Switzerland, but production about 400 tons was obtained under the 1919. The most important deposits are

lite shown on page 4. The prod-
ported chrysotile fiber, is used in a
ifa. of asbestos board, Eternit
s. A small quantity of actinolite
d.

VCE

antities of asbestos in its manu-
duction of raw fiber has been re-
therefore, that plans have been
in the Chateau-Ville-Vieille-et-
e Guil Valley at an altitude of

was reported in 1921.

ANY

ysotile in Bayerischer Wald. East
ais is said to be the only asbestos
nd is exceedingly important from
increasing demands for home con-
supply German asbestos-products
1932 to 20,154 in 1934.

LA

tate has been the most productive
d amphibole fibers are available.
00 long tons was obtained in 1920,
ice 1929. A small production has
Bihar and Orissa from 1921 to
molite. Asbestos occurs in many
ater e reserves are available and
a o. large production of high-
a substantial asbestos-producing
rgely upon the establishment of
that will absorb the short fibers.
om data assembled by Coulson,³⁸
f.

in India, 1906-34

Long tons	Value	Year	Long tons	Value
318.0	£7,372	1923.....	156.3	£1,622
315.5	864	1929.....	318.4	1,306
242.0	701	1930.....	33.2	38
247.0	639	1931.....	6.0	5
123.3	1,334	1932.....	90.0	677
16.0	361	1933.....		
38.4	759	1934.....	24.6	
67.7	1,011			

re. Foreign Trade Notes: Vol. 4, no. 3, May
Districts of the Madras Presidency: With
ndia: Mem. Geol. Survey of India, vol. 64,

IRELAND

An asbestos deposit at Avoca, County Wicklow, was reported in 1931. No data are available on variety, quality, or extent.

MEXICO

An unusually good sample of mountain cork—a variety of spongy, matted, amphibole asbestos—was obtained by the Bureau of Mines in 1931 from an undeveloped deposit near San Luis Potosi. According to report, surface exposures indicate that it is extensive. Asbestos has also been reported from the State of Puebla.

NEWFOUNDLAND

A deposit of chrysotile asbestos discovered near Port au Port is thought to represent a continuation of the Quebec belt. No data are available as to its quality or extent.

NEW ZEALAND

Chrysotile asbestos in veins as wide as 3 inches occurs in a serpentine area at least 3 miles long near Mount Arthur in the upper Takaka district of New Zealand. Both cross fiber and slip fiber are available. The cross-fiber veins occur principally in shells 6 to 30 inches thick surrounding cores of massive serpentine with few or no veins. The fiber yield of the shells is estimated at 10 to 25 percent. The deposit evidently is extensive and is undeveloped.

PHILIPPINE ISLANDS

A deposit of amphibole asbestos occurs in Ilocos Norte Province, Luzon Island. A considerable quantity was mined about 1920 for pipe covering and similar uses.

In 1936 the Bureau of Mines received samples of asbestos from this deposit, which, upon examination, proved to be chrysotile of fair strength and flexibility. Some of it probably could be classed as spinning fiber, although it is not as strong as Canadian or Arizona asbestos. No data are available as to the extent or location of the deposit.

PORTUGAL

Asbestos of uncertain quality, probably in deposits of limited extent, is reported from two places in the Beja district. Province of Alemtejo, southern Portugal. No production is recorded.

SPAIN

Two occurrences of chrysotile asbestos in Spain were reported in 1920—one in the Pyrenees and the other in the Cantabrian Mountains. A production of 165 metric tons was recorded for that year, but no later activity has been noted.

SWITZERLAND

Asbestos occurs in the Cantons of Grisons, Tessin, Valais, and Graubunden, Switzerland, but production is small; a maximum of about 400 tons was obtained under the war stimulus of 1918 and 1919. The most important deposits are at Poschiavo, Canton of

Grisons, where a small amount of chrysotile of spinning quality is available. Both chrysotile and tremolite are said to occur in Graubunden.

TURKEY

A small quantity of asbestos has been produced near Kutaia, western Turkey. An output of 58 tons was recorded for 1932, with a value of about \$24 a ton. Very promising samples of cross-fiber chrysotile were obtained by the Bureau of Mines in 1934 from a deposit said to occur near Sarikamis, Province of Kars, Turkish Armenia. The samples indicated that the percentage of fiber to rock runs very high and that a fair proportion of crude asbestos is obtainable. No details as to the extent of the deposit or its availability are known.

VENEZUELA

A deposit of chrysotile asbestos apparently of good quality occurs about 8 miles from Tinaquillo, State of Cojedes, about 145 miles from Caracas. The nearest railhead is Valencia, 40 miles distant, and the nearest seaport is Puerto Cabello, 65 miles away.

The occurrence is unusual, as the fiber is associated with clay and shattered rock. A little development work has been done and small shipments have been made, but much systematic exploratory work must be conducted before the commercial value of the deposit can be determined.

OTHER COUNTRIES

Asbestos occurrences have been reported in Bolivia: Gorbea, Chile: Chosen (Korea), Japan; Madagascar: Spitzbergen; and near Alexandretta, Syria.

POLITICAL AND COMMERCIAL CONTROL

Russia only of the four leading producing countries has important asbestos-products manufacturing plants. The other principal producing countries—Canada, Rhodesia, and the Union of South Africa—utilize a very small proportion of the asbestos they produce. On the other hand, the principal asbestos-consuming countries—the United States, England, France, Germany, Belgium, and Japan—produce no asbestos, or only small tonnages. A large proportion of all asbestos mined, therefore, enters international trade, and all the principal nations are interested in foreign sources of supply and their availability in times of emergency.

POLITICAL CONTROL

Three of the four large asbestos-producing countries of the world—Canada, Rhodesia, and the Union of South Africa—are in the British Empire. Moderately important deposits in Cyprus and Australia and small deposits in India, New Zealand, and Newfoundland also are under British political control.

The largest deposits outside the British Empire are those of the Soviet Union, while others of considerable importance are under the dominion of the United States, Italy, and Finland. Relatively small deposits are controlled politically by Argentina, Austria, China, Czechoslovakia, Spain, Switzerland, Turkey, and Venezuela.

COMMERCIAL CONTROL

All asbestos-producing companies are controlled by United States capital.

Commercial control of the Canadian company, Asbestos Corporation, Ltd., is Canadian, and United States capital. Its operator, probably is controlled by the Asbestos Corporation, Ltd., owned by the Bell Asbestos Mines, Inc., a New York firm, and the Bell Asbestos Mines, Inc., capital. The Bell Asbestos Mines, Inc., is now controlled by the Mattison Co., Ambler, Pa., is now controlled by the Bell Asbestos Mines, Ltd., London, England. The ownership therefore, evidently is divided among Canadian capital. The amounts of capital the United States are approximately investment is less.

British capital predominates in the companies of Rhodesia and the Union of South Africa.

The Soviet Union controls, both activities in the asbestos industry of the Soviet Union.

The one company operating in China is capital.

For many years a British company controlled Italy, but since 1929 an Italian company, Balangero Mines, where the chief production is asbestos.

Practically all the deposits in Australia or Australian capital.

INTERNATIONAL TRADE

The United States is the largest consumer and imports over 95 percent of the asbestos. About 1 percent of Canadian production enters the United States in proximity to the extensive United States market. A distinct advantage over all other countries. For many years England, France, Germany, and most of their supplies from Canada. England's needs. Other European countries obtain quantities from Russia, and fluctuate between Canada and Africa.

EXPORTS FROM PRINCIPAL COUNTRIES

A recent report from Irving N. L. in Johannesburg, shows the distribution of destination during 1934 and 1935 exports from the Union of South Africa in 1935 about 52 percent of the total to the United Kingdom, nearly 22 percent to the United States and the remaining 26 percent to other countries. The United States received less than 5 percent, considerably more per ton than any other country.

mt. chrysotile of spinning quality is and tremolite are said to occur in

TURKEY

has been produced near Kutaia, west-58 tons was recorded for 1932, with a very promising samples of cross-fiber the Bureau of Mines in 1934 from a Karikamis, Province of Kars, Turkish stated that the percentage of fiber to it a fair proportion of crude asbestos to the extent of the deposit or its

VENEZUELA

stos apparently of good quality occurs lo, State of Cojedes, about 145 miles railhead is Valencia, 40 miles distant, Puerto Cabello, 65 miles away. is the fiber is associated with clay and opment work has been done and small ut much systematic exploratory work commercial value of the deposit can

OTHER COUNTRIES

en reported in Bolivia: Gorgea, Chile: adagascar: Spitzenbergen; and near

COMMERCIAL CONTROL

each producing countries has manufacturing plants. The other principal countries are Rhodesia, and the Union of small proportion of the asbestos they and the principal asbestos-consuming England, France, Germany, Belgium, os, or only small tonnages. A large l. therefore, enters international trade, are interested in foreign sources of in times of emergency.

LEGAL CONTROL

os-producing countries of the world—on of South Africa—are in the British it deposits in Cyprus and Australia ew Zealand, and Newfoundland also rol. the British Empire are those of the onsiderable importance are under the Italy, and Finland. Relatively small ally by Argentina, Austria, China, land, Turkey, and Venezuela.

COMMERCIAL CONTROL

All asbestos-producing companies in the United States apparently are controlled by United States capital.

Commercial control of the Canadian mines is diverse. The largest company, Asbestos Corporation, Ltd., is controlled by English, Canadian, and United States capital. Johnson's Company, the pioneer operator, probably is controlled by Canadian capital. The Quebec Asbestos Corporation, Ltd., owned by the Philip Carey Manufacturing Co., Cincinnati, Ohio; the Canadian Johns-Manville Co., Ltd., a subsidiary of Johns-Manville, Inc., New York; and the Nicolet Asbestos Mines, Ltd., a New York firm, are dominated by United States capital. The Bell Asbestos Mines, Inc., operated by the Keasbey & Mattison Co., Ambler, Pa., is now controlled by Turner & Newall, Ltd., London, England. The ownership of the Canadian companies, therefore, evidently is divided among English, United States, and Canadian capital. The amounts of capital invested by England and the United States are approximately the same, while the Canadian investment is less.

British capital predominates in all the asbestos-producing companies of Rhodesia and the Union of South Africa.

The Soviet Union controls, both politically and commercially, activities in the asbestos industry of the Urals.

The one company operating in Cyprus is controlled by British capital.

For many years a British company controlled most of the production of Italy, but since 1929 an Italian company has operated the Balangero Mines, where the chief production is now centered.

Practically all the deposits in Australia are controlled by English or Australian capital.

INTERNATIONAL TRADE

The United States is the largest consumer of asbestos in the world and imports over 95 percent of its supplies. Normally 70 to 75 percent of Canadian production enters the United States. Because of its proximity to the extensive United States markets, Canada enjoys a distinct advantage over all other countries in the field of short fibers. For many years England, France, Germany, and Belgium imported most of their supplies from Canada. Africa now furnishes much of England's needs. Other European countries are securing increasing quantities from Russia, and fluctuating supplies are obtained in Canada and Africa.

EXPORTS FROM PRINCIPAL PRODUCING COUNTRIES

A recent report from Irving N. Linnell, American consul general in Johannesburg, shows the distribution of African asbestos by country of destination during 1934 and 1935. The following table shows exports from the Union of South Africa. It may be observed that in 1935 about 52 percent of the total quantity was shipped to the United Kingdom, nearly 22 percent to other European countries, and the remaining 26 percent to countries outside of Europe. The United States received less than 5 percent of the total, but paid considerably more per ton than any other country, due to the fact that

the United States imports from this source only the higher-grade fibers.

Exports of asbestos from the Union of South Africa, 1934 and 1935

Country of destination	1934		1935	
	Short tons	Value	Short tons	Value
United Kingdom.....	8,652	£112,019	12,294	£138,736
Belgium.....	559	7,625	1,124	10,737
France.....	1,028	17,301	1,195	17,335
Germany.....	903	18,860	1,493	26,361
Holland.....	365	7,107	455	7,672
Italy.....	1,139	20,562	666	12,652
Spain.....	624	7,938	250	3,641
Sweden.....	50	736	5	119
Denmark.....			2	39
Portugal.....			10	239
Japan.....	1,327	23,195	2,301	30,497
United States.....	715	13,884	1,114	23,043
Canada.....	9	133	26	369
Australia.....	1,692	24,133	2,206	29,775
India.....	289	1,777	298	1,817
Algeria.....			20	369
Chile.....	73	925	253	3,562
Uruguay.....	24	447	24	1,447
Argentina.....			10	199

- Estimated.

The following table shows similar data for Rhodesia, but the figures are somewhat uncertain because a large percentage of the fiber is shipped through Portuguese East Africa and the ultimate destination is unknown. Evidently 85 percent or more of the 1935 exports reached European ports. The small figures recorded in this table for European countries outside of the United Kingdom indicate that most or all of their requirements of Rhodesian fiber are included under Portuguese East Africa or Optional Ports. Only about 3 percent of the shipments reached the United States. However, that shipped to the United States averaged about \$145 a ton, whereas European shipments averaged about \$90 a ton.

Exports of asbestos from Rhodesia, 1934 and 1935

Country of destination	1934		1935	
	Short tons	Value	Short tons	Value
United Kingdom.....	11,764	£212,014	13,120	£206,995
Australia.....	1,100	20,051	1,450	26,437
India.....	850	11,625	760	11,986
Portuguese East Africa ¹	15,395	291,830	19,028	335,254
Japan.....	1,721	26,515	1,075	23,743
Dutch East Indies.....			50	682
Chile.....	47	423	97	971
United States.....	1,076	24,276	1,418	40,955
Union of South Africa ²	154	1,665	366	3,968
Holland.....	27	384		
Italy.....	2	34		
Germany.....	5	60		
Belgium.....	46	567		
Sweden.....			600	9,552
Optional Ports ³			7,305	132,193

¹ Most of that entering Portuguese East Africa is destined to the Continent of Europe, but ultimate country of destination is unknown at time of shipment.

² Probably for reshipment, but destination unknown.

³ Destined for Continent of Europe, but ultimate country of destination unknown.

As indicated in the following table of 50 percent of the total is consigned to countries, Japan, and, during recent years, substantial buyers. The decline in exports may be due to the pronounced expansion of products manufacturing industries, which titles of the raw asbestos produced. Exports to the United States as shown in this table figures for imports into the United States by the Bureau of Foreign and Domestic Commerce in Russian production has not as might be expected, because Russian asbestos-products-manufacturing industry of its raw material in its own plants.

Exports of asbestos from U. S.

[Compiled by B. B. Waldbauer, Foreign Trade Commission]

Country of destination	Metric tons
Austria.....	
Belgium and Luxemburg.....	3.
Czechoslovakia.....	
Denmark.....	
Finland.....	
France.....	1.
Germany.....	14.
Italy.....	3.
Japan.....	3.
Monrovia.....	
Netherlands.....	
Norway.....	
Poland.....	
South America.....	
Sweden.....	
Switzerland.....	
Turkey.....	
United Kingdom.....	1.
United States.....	2.
Other.....	
Total.....	3.

¹ As given in Foreign Trade (Vneshechnaya Torgovlya) Statistics.

² During these years 1 ruble was worth about 57 cents.

The following table shows exports of asbestos by country of destination, in 1934 and 1935. and 58 percent in value of the total exports to the United States. In 1935, the figures for exports to the United States are relatively small, because the price of asbestos is relatively small, because the price of a heavy transportation expense.

s source only the higher-grade

of S. Africa, 1934 and 1935

1934		1935	
Short tons	Value	Short tons	Value
8,652	£113,019	12,234	£138,736
359	7,625	1,124	10,737
1,028	17,301	1,195	17,335
943	18,360	1,493	24,341
365	7,107	455	7,672
1,139	20,562	666	12,652
624	7,938	250	3,641
50	736	5	119
		2	39
		10	239
1,327	23,195	2,301	30,497
715	13,884	1,114	23,043
9	133	26	369
1,692	24,133	2,206	29,775
299	1,777	298	1,317
		20	289
73	325	253	3,562
24	447	24	147
		10	199

data for Rhodesia, but the large percentage of the fiber is mica and the ultimate destination more of the 1935 exports reached recorded in this table for Euro-Kingdom indicate that most or fiber are included under Portugal. Only about 3 percent of the However, that shipped to the to whereas European ship-

desia; 1934 and 1935

1934		1935	
Short tons	Value	Short tons	Value
11,764	£212,014	13,120	£206,995
1,100	19,051	1,450	26,437
550	11,825	750	11,366
13,395	241,330	19,028	335,254
1,721	26,315	1,075	23,743
		50	682
47	423	97	971
1,076	24,276	1,418	40,955
154	1,965	366	3,968
27	384		
2	34		
3	90		
49	567		
		800	9,552
		7,305	132,193

sed to the Continent of Europe, but ultimate

of destination unknown.

As indicated in the following table of exports from Russia, nearly 50 percent of the total is consigned to Germany. Other European countries, Japan, and, during recent years, the United States are substantial buyers. The decline in exports in 1935 compared with 1934 may be due to the pronounced expansion in the domestic asbestos-products manufacturing industries, which consume increasing quantities of the raw asbestos produced. Exports of asbestos from Russia to the United States as shown in this table do not check closely with figures for imports into the United States from Russia as compiled by the Bureau of Foreign and Domestic Commerce. The great increase in Russian production has not affected world trade as much as might be expected, because Russia is developing an important asbestos-products-manufacturing industry and thus uses a great deal of its raw material in its own plants.

Exports of asbestos from U. S. S. R. (Russia),¹ 1934-35

[Compiled by B. B. Waldbauer, Foreign Mineral Service Division]

Country of destination	1934		1935	
	Metric tons	Rubles ²	Metric tons	Rubles ²
Austria.....	136	19,000	44	6,000
Belgium and Luxemburg.....	3,565	336,000	1,581	139,000
Czechoslovakia.....	410	43,000		
Denmark.....	104	10,000	72	5,000
Finland.....	43	8,000	15	1,000
France.....	1,045	109,000	944	84,000
Germany.....	14,741	1,492,000	12,018	1,255,000
Italy.....	3,454	379,000	1,562	147,000
Japan.....	3,550	310,000	2,417	146,000
Mongolia.....	2	200	4	1,000
Netherlands.....	147	18,000	118	10,000
Norway.....	77	9,000		
Poland.....	74	11,000	15	2,000
South America.....	268	25,000		
Sweden.....	771	80,000	690	69,000
Switzerland.....	49	6,000		
Turkey.....	16	800	14	1,000
United Kingdom.....	1,647	149,000	1,453	118,000
United States.....	3,040	192,000	3,099	151,000
Other.....	566	33,200	463	41,000
Total.....	33,715	3,248,000	25,109	2,277,000

¹ As given in Foreign Trade (Vneshnyaya Torgovlya) Statistical Summary, Moscow.

² During these years 1 ruble was worth about 87 cents.

The following table shows exports of Canadian asbestos, by country of destination, in 1934 and 1935. In 1934, 71 percent in quantity and 58 percent in value of the total exports were destined to the United States. In 1935, the figures were 77 and 66 percent, respectively. Exports from Canada to the United States as given in this table differ only by a small amount from the figures for imports from Canada into the United States as compiled by the Bureau of Foreign and Domestic Commerce. The trans-Atlantic trade in short fibers is relatively small, because the price obtainable is too low to justify a heavy transportation expense.

Exports of asbestos from Canada, 1934-35¹
 (Compiled by B. B. Waldbauer, Foreign Mineral Service Division)

Country of destination	1934		1935		1936	
	Fiber		Sand and waste		Total	
	Short tons	Value	Short tons	Value	Short tons	Value
Australia.....	998	\$49,859	—	—	998	\$49,859
Belgium.....	3,548	191,519	435	38,968	4,003	200,487
France.....	3,949	243,416	540	10,073	4,509	253,491
Germany.....	3,436	441,188	2,497	50,787	5,933	491,975
Italy.....	618	58,090	—	—	618	58,090
Japan.....	18,469	679,723	—	—	18,469	679,723
Netherlands.....	734	35,800	579	11,541	1,313	47,341
Spain.....	162	7,687	—	—	162	7,687
United Kingdom.....	4,618	316,468	2,080	44,620	6,698	361,088
United States.....	44,542	1,996,915	68,171	964,429	112,713	2,961,344
Other.....	153	8,326	656	9,855	809	18,211
Total.....	83,267	4,029,191	74,978	1,100,305	158,245	5,129,496

¹ As given in The Trade of Canada, Ottawa.

IMPORTS INTO THE UNITED STATES

The following table, compiled from Bureau of Mines statistics, shows imports of asbestos into the United States for 6 years. It includes only the crude and mill fibers that constitute, on the basis of value, the principal international trade in asbestos. Figures for earlier years may be found in the Minerals Yearbook and in Mineral Resources of the United States, published by the Bureau of Mines.

Imports of asbestos into the United States, 1930-35 (crude and mill fiber only)

Country of shipment	1930		1931		1932	
	Short tons	Value	Short tons	Value	Short tons	Value
Africa.....	3,210	\$730,274	1,337	\$189,349	1,326	\$117,774
Canada.....	70,957	3,656,195	47,340	2,123,746	31,064	1,285,799
Germany.....	390	83,897	110	10,700	1	30
Italy.....	46	8,752	19	9,855	85	9,627
Soviet Russia.....	4,534	660,559	2,188	104,271	—	—
United Kingdom.....	410	113,664	953	166,465	1	399
Belgium.....	—	—	—	—	—	—

Imports of asbestos into the United States

Country of shipment	1933	
	Short tons	Value
Africa.....	2,325	\$234,557
Canada.....	48,916	2,337,946
Germany.....	1	679
Italy.....	17	8,929
Soviet Russia.....	176	17,339
United Kingdom.....	1	587
Morocco.....	—	—

Countries of shipment as given in the necessarily the countries of origin. Asbestos was chiefly of Russian origin, and fiber Kingdom, Belgium, and Morocco origins. The following table shows imports, by years. For Africa and Russia the figure because the origin of fiber shipped from it not be traced exactly. Also, for both things, relative to short fibers, shipments of usual sources have been disregarded.

Sources of asbestos imported into the United States (fiber only)

Source	1930		
	Short tons	Value	Short
Africa.....	3,020	\$813,958	4
Canada.....	70,957	3,656,195	—
Soviet Russia.....	4,924	744,456	—
Italy.....	48	8,752	—
Total.....	79,549	5,223,361	—

Source	1933		
	Short tons	Value	Short
Africa.....	2,325	\$234,557	—
Canada.....	48,916	2,337,946	—
Soviet Russia.....	182	18,166	—
Italy.....	17	8,929	—
Total.....	51,440	2,469,598	—

The following table shows the percentage value of imports of crudes and mill fiber source during recent years.

from Canada, 1934-35

Mineral Service Division]

19.

Due	Sand and waste		Total	
	Short tons	Value	Short tons	Value
19,359			998	\$19,859
11,319	455	\$8,968	4,003	\$20,487
11,416	540	10,075	4,509	\$23,491
11,188	2,497	50,787	7,833	\$91,973
3,090			618	\$8,090
2,723			18,489	\$79,723
3,800	379	11,541	1,313	\$7,341
7,887			162	\$7,887
4,468	1,080	\$4,620	6,098	\$61,098
4,915	68,171	\$94,429	112,713	\$2,961,344
4,326	556	9,685	809	\$8,211
4,191	74,378	1,100,305	158,245	\$3,129,496

1935

4,632			2,004	\$99,632
1,006	33	\$14,407	3,647	\$85,013
142	220	6,200	4,101	\$60,342
1,062	1,438	28,605	6,351	\$65,867
435			806	\$74,435
397			15,597	\$23,397
723	700	\$4,776	2,371	\$25,501
328			710	\$7,328
760	1,594	\$75,516	8,178	\$66,085
368	32,410	\$440,995	153,569	\$4,520,361
714	570	4,782	577	\$21,496
176	100,925	\$1,365,461	200,211	\$4,565,457

THE UNITED STATES

from Bureau of Mines statistics, the United States for 6 years. It fibers that constitute, on the basis of trade in asbestos. Figures for Minerals Yearbook and in Mineral published by the Bureau of Mines.

1930-35 (crude and mill fiber only)

Value	1931		1932	
	Short tons	Value	Short tons	Value
\$730,274	1,537	\$189,349	1,326	\$117,774
2,654,195	47,340	\$2,123,746	31,064	\$1,285,799
53,697	110	10,700	1	30
4,752	19	9,355	55	9,627
660,559	2,198	104,271		
113,664	953	166,465	1	399

Imports of asbestos into the United States, 1930-35—Continued

Country of shipment	1933		1934		1935	
	Short tons	Value	Short tons	Value	Short tons	Value
Africa.....	2,324	\$234,557	1,794	\$198,479	2,128	\$294,231
Canada.....	48,916	2,337,946	43,053	2,026,161	60,032	3,015,247
Germany.....	5	669				
Italy.....	17	8,929	16	9,775	22	11,464
Soviet Russia.....	176	17,339	657	26,434	4,632	213,698
United Kingdom.....	1	567	22	5,016	312	23,451
Morocco.....					22	2,131

Countries of shipment as given in the preceding table are not necessarily the countries of origin. Asbestos shipped from Germany was chiefly of Russian origin, and fiber shipped from the United Kingdom, Belgium, and Morocco originated chiefly in Africa.

The following table shows imports, by countries of origin, for 6 years. For Africa and Russia the figures are approximate only, because the origin of fiber shipped from nonproducing countries cannot be traced exactly. Also, for both this table and the one following, relative to short fibers, shipments of small quantities from unusual sources have been disregarded.

Sources of asbestos imported into the United States, 1930-35 (crude and mill fiber only)

Source	1930		1931		1932	
	Short tons	Value	Short tons	Value	Short tons	Value
Africa.....	3,620	\$843,958	2,290	\$355,814	1,327	\$118,173
Canada.....	70,957	3,856,195	47,340	2,123,746	31,064	1,285,799
Soviet Russia.....	4,924	744,456	2,298	114,971	1	30
Italy.....	48	8,752	19	9,855	55	9,627
Total.....	79,549	\$5,253,361	51,947	\$2,604,386	32,477	\$1,413,629

Source	1933		1934		1935	
	Short tons	Value	Short tons	Value	Short tons	Value
Africa.....	2,325	\$235,144	1,516	\$203,495	2,352	\$319,313
Canada.....	48,916	2,337,952	43,053	2,026,161	60,032	3,015,247
Soviet Russia.....	182	18,088	657	26,434	4,632	213,698
Italy.....	17	8,929	16	9,775	22	11,464
Total.....	51,440	\$2,600,033	45,542	\$2,265,865	67,038	\$3,590,222

The following table shows the percentages of both quantity and value of imports of crudes and mill fibers obtained from each foreign source during recent years.

Percentage of crude and mill-fiber asbestos obtained from each of the principal foreign sources, 1930-35

Source	1930		1931		1932	
	Quantity	Value	Quantity	Value	Quantity	Value
Africa	4.5	16.1	4.4	13.7	4.1	5.4
Canada	89.2	89.6	91.1	81.6	95.6	91.0
Soviet Russia	6.2	14.1	4.4	4.4		
Italy	.1	.2	.1	.3	.3	.6

Source	1933		1934		1935	
	Quantity	Value	Quantity	Value	Quantity	Value
Africa	4.5	9.0	4.0	9.0	3.5	9.0
Canada	95.0	89.9	94.5	89.4	89.5	84.7
Soviet Russia	.4	.7	1.4	1.2	6.9	6.0
Italy	.1	.4	.1	.4	.1	.3

It may be observed that imports from Africa are much higher in value than in quantity. This is due to the fact that only the higher-priced crude fibers can be imported profitably from such a distant source. On the other hand, the percentage in quantity of Canadian fiber exceeds its percentage in value, because the deposits are near consuming centers, permitting the importation of large quantities of the shorter lower-priced mill fibers.

As many manufacturers of asbestos products are interested chiefly in sources of supply of the higher-grade spinning fibers, an attempt has been made to estimate the tonnages received in recent years from the principal producing centers. The following table shows the approximate tonnages of spinning asbestos received annually from Canada, Africa, and Russia, and the percentage of the total imports each country furnished.

Principal sources of spinning asbestos imported into the United States, 1924-35

Year	Canada ¹		Africa		Russia	
	Short tons	Percent of total imports	Short tons	Percent of total imports	Short tons	Percent of total imports
1924	12,463	80	3,051	20	56	(?)
1925	17,543	79	4,390	20	191	1
1926	17,661	78	4,747	21	349	1
1927	14,953	66	6,096	27	1,782	7
1928	15,165	62	6,778	25	2,339	10
1929	18,363	64	8,778	31	1,436	5
1930	8,473	50	3,620	21	4,924	26
1931	6,770	60	2,290	20	2,298	20
1932	4,380	77	1,327	23	(?)	(?)
1933	(?)		2,325		182	
1934	(?)		1,816		657	
1935	(?)		2,352		4,632	

¹ Figures for Canadian tonnages are approximate. The following text explains methods of determination.

* Figures for Canadian
* Less than 0.5 percent.

³ Less than 1 ton.

* Data not available.

For the purpose of calculation, Canadian Crude No. 1, Crude No. 2, Crude run-of-mill fiber, which, since 1920, has been published annually, but import figures were not until 1927. From 1924 to 1927 it is assumed fiber bore the same relation to total of spinning fiber bore to total production. 3 percent of the total Canadian production, exclusive of "sand, gravel, and stone", was imported is assumed that 85 percent of the spinning was shipped to the United States.

Since 1927 the tonnages of crude and Canada have been recorded, but spinnings separately. To determine this tonnage the various grades of mill fibers (spinnings) are imported in the ratio of their production to the mill stock produced is of spinning grade. The percent of the mill stock imported is spinning fiber thus calculated is then added to the crudes to arrive at the total imports of spinning fiber from the United States.

Approximately all imports from the U.S. are regarded as spinning fiber. Russian imports are separated into crudes and mill fiber, the latter being nonspinning.

By referring to the preceding table, it is seen that the quantity of nonspinning cotton supplied about 80 percent of the total in 1927, however, this percentage dropped to only 50. In 1932 the Dominion again furnished about 80 percent of the total quantity from all sources of imports, but the total quantity from all sources was low.

Africa supplied about 20 percent of imports during the period 1927-29, when the price was higher. Since 1926, imports from Japan and Russia supplied more than one-fourth of the United States. Practical cessation of imports in 1932 may be attributed to the embargo investigation conducted by the United States, although the embargo was lifted. Imports in 1935 were again almost as large as in 1927, in comparison with Canada.

Unfortunately, comparison with Cane years subsequent to 1932, because since the mill fiber are not broken down by class holding its position because African imported one-third of the 1927-29 average.

The following table shows sources of asbestos in the United States of the shorter grades of asbestos cloth. Until recently Canada supplied nearly 100 percent of the total but since 1932 Cyprus, Soviet Russia, and Czechoslovakia have furnished 4 to 5 percent of the total.

tos obtained from each of the principal
ces, 1930-35

Year	1931		1932	
	Quantity	Value	Quantity	Value
1931	4.4	13.7	4.1	5.4
1932	21.1	81.6	95.6	91.0
1933	4.4	4.4	3.3	3.6
1934	1.1	1.3	1.1	1.3
Year	1934		1935	
	Quantity	Value	Quantity	Value
1934	4.0	9.0	3.5	9.0
1935	94.5	89.4	59.5	54.7
1936	1.4	1.2	6.9	6.0
1937	1.1	1.4	1.1	1.3

from Africa are much higher in to the fact that only the higher-ly profitably from such a distant centage in quantity of Canadian e, because the deposits are near uporation of large quantities of

as products are interested chiefly rade spinning fibers, an attempt res received in recent years from The following table shows the asbestos received annually from percentage of the total imports

ported into the United States, 1924-35

Year	Africa		Russia	
	Short tons	Percent of total imports	Short tons	Percent of total imports
1924	3,251	20	56	(?)
1925	4,390	20	191	1
1926	4,747	21	349	1
1927	6,296	27	1,782	1
1928	6,774	28	2,339	10
1929	8,774	31	1,426	5
1930	3,220	21	4,921	29
1931	2,290	20	2,298	20
1932	1,125	13	(?)	(?)
1933	1,255	15	637	15
1934	2,352	20	4,632	20

following text explains methods of determination.

For the purpose of calculation, Canadian spinning fiber includes Crude No. 1, Crude No. 2, Crude run-of-mine, and the highest-grade mill fiber, which, since 1920, has been designated "spinning fiber." Figures for the production of Canadian asbestos by grades are published annually, but import figures were not segregated by grades until 1927. From 1924 to 1927 it is assumed that imports of spinning fiber bore the same relation to total imports that production of spinning fiber bore to total production. In other words, if 85 percent of the total Canadian production, exclusive of material classed as "sand, gravel, and stone", was imported into the United States, it is assumed that 85 percent of the spinning fiber produced in Canada was shipped to the United States.

Since 1927 the tonnages of crude and mill fiber imported from Canada have been recorded, but spinning fiber still is not reported separately. To determine this tonnage, therefore, it is assumed that the various grades of mill fibers (spinning, shingle, and paper stock) are imported in the ratio of their production; that is, if 25 percent of the mill stock produced is of spinning grade, it is assumed that 25 percent of the mill stock imported is spinning fiber. The quantity of spinning fiber thus calculated is then added to imports of Canadian crudes to arrive at the total imports of spinning stock from Canada.

Approximately all imports from the other foreign countries are regarded as spinning fiber. Russian imports in 1931, however, were separated into crudes and mill fiber, the latter being regarded as nonspinning.

By referring to the preceding table, it may be observed that Canada supplied about 80 percent of the total imports from 1924 to 1926. In 1927, however, this percentage dropped to 66, and in 1930 it was only 50. In 1932 the Dominion again furnished over three-fourths of imports, but the total quantity from all sources was exceptionally low.

Africa supplied about 20 percent of the total each year, except during the period 1927-29, when the proportion was considerably higher. Since 1926, imports from Russia have been considerable; in 1930 Russia supplied more than one-fourth of the requirements of the United States. Practical cessation of imports from Russia in 1932 may be attributed to the embargo placed on them during an investigation conducted by the United States Tariff Commission. Although the embargo was lifted, imports in 1933 were small, but by 1935 they were again almost as large as in the peak year 1930.

Unfortunately, comparison with Canada cannot be made for the years subsequent to 1932, because since that year Canadian statistics for mill fiber are not broken down by classes. Apparently Canada is holding its position because African imports in 1935 were less than one-third of the 1927-29 average.

The following table shows sources of imports into the United States of the shorter grades of asbestos classed as "stucco and refuse." Until recently Canada supplied nearly 100 percent of the short fibers, but since 1932 Cyprus, Soviet Russia, Finland, and Italy together have furnished 4 to 5 percent of the total.

Imports into the United States of asbestos classed as "stucco and refuse",
1930-35

Source	1930		1931		1932	
	Short tons	Value	Short tons	Value	Short tons	Value
Canada.....	129.111	\$1,810,588	2	\$2,767	63.606	\$827,744
Soviet Russia.....	2	84	1,619	\$1,105,936	35,423	7,719
Italy.....	17	409	27	315	43	1,105
Africa.....						
Total.....	129.130	1,511,141	94.413	1,144,674	64,278	836,571

Source	1933		1934		1935	
	Short tons	Value	Short tons	Value	Short tons	Value
Canada.....	63,999	\$354,647	70,007	\$1,000,402	94,204	\$1,470,845
Soviet Russia.....	531	40,503	1,935	63,005	181	534
Cyprus.....	2,274	37,393	2,463	43,611	4,628	87,844
Italy.....	339	7,764	246	1,774	523	5,202
Finland.....	37	1,404	35	1,920	11	446
Africa.....	11	339	100	1,417		
Total.....	63,091	942,052	71,792	1,112,129	99,547	1,565,191

EXPORTS FROM THE UNITED STATES

Very little unmanufactured asbestos is exported from the United States. Recent exports, shown in the following table, have consisted principally of shingle stock of foreign origin blended in the United States and shipped mainly to Japan, Mexico, and Europe. The exceptionally high value of exports in 1928 is due to a large shipment of Russian asbestos to Germany. Statistics of the Department of Commerce also show shipments of about 1,000 tons of foreign crude and unmanufactured asbestos annually under "exports of foreign merchandise." These represent transshipments of Canadian material via American ports.

Asbestos (unmanufactured) exported from the United States, 1924-36

Year	Short tons	Value	Year	Short tons	Value	Year	Short tons	Value
1924.....	1,270	\$92,163	1929.....	709	\$105,467	1934.....	1,669	\$94,182
1925.....	1,109	70,846	1930.....	771	95,318	1935.....	550	57,896
1926.....	1,104	85,922	1931.....	1,714	122,391	1936.....	3,744	310,197
1927.....	309	48,744	1932.....	1,767	94,936			
1928.....	850	346,632	1933.....	1,378	88,521			

Although little unmanufactured asbestos is exported, foreign trade in asbestos products is extensive. The following table shows exports by kinds in 1934 and 1935. Brake linings and textile yarns and packings are the largest items.

Manufactured asbestos products exported from kinds

Product	Quantity
Brake linings:	
Molded and remolded.....	linear feet.. 1.
Not molded.....	short tons..
Paper, millboard, and rollboard.....	do.
Pipe covering and cement.....	do.
Textiles, yarn, and packing.....	do.
Magnesia and manufactures.....	do.
Asbestos roofing.....	squares..
Other manufactures.....	short tons..

¹ Quantity not recorded.

PROSPECTING AND EXPLORATION

Asbestos is usually associated with gneiss and is recognized by its green or yellowish gray color. Therefore, the formations in which asbestos is so resistant to weathering than most other minerals are the knolls or ridges above the level of the surface. The weathering of the fiber judged by visual inspection. Such effects usually are few inches only. Exploration at depth by means of test pits which are enlarged into shafts or shot-core drills. Deposits are of test pits which are enlarged into shafts or shot-core drills. Estimation of asbestos-bearing rock is difficult, and is usually made only by making actual mill runs.

MINING METHODS

Asbestos deposits in the various parts of the world and methods of mining depend greatly on the nature of the deposit. In some places the rock is blasted in place and the material is then quarried in much the same way as for the production of crushed stone. In other cases elaborate systems of underground mining are followed. General methods followed in the past are briefly:

ARIZONA

In Arizona the asbestos occurs in veins of diorite intrusions. The veins pinch and swell in size and direction that depends definitely on the mining of many veins in drifts and tunnels, and conditions of two or more veins of workable size are often found in one tunnel. The soft, flexible fiber is much harsher, more brittle asbestos may be associated with it.

asbestos classed as "stucco and refuse",
1930-35

1931			1932		
Value	Short tons	Value	Short tons	Value	Short tons
\$10,589	82,767	\$1,108,936	63,606	5827,744	
84	1,619	35,423	629	7,710	
469	27	315	43	1,109	
\$11,141	84,413	1,144,674	64,278	836,571	

1934			1935		
Value	Short tons	Value	Short tons	Value	Short tons
\$54,647	70,007	\$1,000,402	94,204	\$1,470,845	
40,503	1,938	63,005	181	834	
37,395	2,463	43,611	4,623	37,944	
7,764	246	1,774	523	5,202	
1,404	38	1,920	11	446	
339	190	1,417			
\$42,022	74,732	1,112,129	99,547	1,565,191	

THE UNITED STATES

Asbestos is exported from the United States in the following table, have come of foreign origin blended in the exports to Japan, Mexico, and Europe. Statistics of the Department of Commerce show that about 1,000 tons of asbestos annually under "exemption" represent transshipments of asbestos.

Asbestos from the United States, 1924-36

Year	Value	Year	Short tons	Value
1929	\$105,407	1934	1,669	\$94,182
1930	95,315	1935	550	57,296
1931	122,391	1936	3,744	310,197
1932	94,936			
1933	86,121			

Asbestos is exported, foreign trade. The following table shows exports of asbestos linings and textile yarns and

Manufactured asbestos products exported from the United States 1934-35, by kinds

Product	1934		1935	
	Quantity	Value	Quantity	Value
Brake linings:				
Molded and semimolded.....	(1)	\$607,193	(1)	\$451,338
Not molded.....	1,641,333	255,018	1,426,320	231,381
Paper, millboard, and rollboard.....	602	96,154	767	130,578
Pipe covering and cement.....	1,389	128,190	1,233	143,357
Textiles, yarn, and packing.....	619	594,625	715	662,321
Magnesia and manufactures.....	1,277	241,410	922	146,068
Asbestos roofing.....	26,457	75,254	31,141	106,678
Other manufactures.....	799	146,670	921	190,208

(1) Quantity not recorded.

(2) Revised figure.

PROSPECTING AND EXPLORATION

Asbestos is usually associated with serpentine, a mineral easily recognized by its green or yellowish green color and greasy luster; therefore, the formations in which asbestos occur are traced easily. Asbestos is so resistant to weathering that outcrops frequently occupy knolls or ridges above the level of the surrounding country. The exposure may be widened by stripping or trenching, and the quality of the fiber judged by visual inspection. Surface fiber may be discolored by weathering, but such effects usually are confined to a depth of a few inches only. Exploration at depth may be conducted with diamond- or shot-core drills. Deposits are commonly explored by means of test pits which are enlarged into active quarries or mines if favorable results are obtained. Estimation of the fiber content of an asbestos-bearing rock is difficult, and satisfactory results can be secured only by making actual mill runs.

MINING METHODS

Asbestos deposits in the various localities differ widely in character, and methods of mining depend greatly on conditions of occurrence. In some places the rock is blasted in open quarries and loaded with power shovels in much the same way as limestone or trap rock is quarried for the production of crushed stone. In other deposits elaborate systems of underground mining have been developed. General methods followed in the principal districts are outlined briefly.

ARIZONA

In Arizona the asbestos occurs in veins in the Mescal limestone near diabase intrusions. The veins pinch and swell irregularly and are so erratic in size and direction that development cannot be planned as definitely as in the mining of many other minerals. The fiber is mined in drifts and tunnels, and conditions are most favorable where two or more veins of workable size are close enough to be worked in one tunnel. The soft, flexible fiber is most desired, but veins of harsher, more brittle asbestos may be associated closely with the softer variety.

Modified room-and-pillar mining methods are followed. Much of the waste rock in each room is built up in dry walls about every 10 feet for roof support. Excess waste and fiber-bearing rock are wheeled to chutes. Cobbing and sorting are done in the stopes on a contract basis. Two grades of fiber are usually produced, and contractors are paid 2 cents a pound more for the long than for the short fiber recovered. Mexican labor is found to be most satisfactory.

Several thousand feet of tunnels have been driven on Ash Creek and at other points in the southern part of the State. Production costs at the best mines are said to be about 40 cents per pound of fiber produced. From 25 to 60 pounds of asbestos are recovered for each running foot of a 4½- by 7-foot tunnel.

VERMONT

The chrysotile-asbestos quarries of Vermont are situated at an elevation of 2,500 feet on the south side of Belvidere Mountain approximately 15 miles from Hyde Park, the nearest railway station. As the rock outcrops at this point, very little stripping is necessary. Whatever overburden is encountered is removed in stoneboats drawn by horses because of the steep slope of the mountainside. The quarry face is about 150 feet high, and individual benches are 12 to 15 feet high. Blast holes are sunk with hammer drills. Two electric power shovels with 1¼-cubic yard dippers are used for loading into 3 motor trucks with 10-ton steel rear-dump bodies. The trucks dump the rock into a chute, which feeds a 28- by 36-inch jaw crusher.

GEORGIA AND MARYLAND

Mass-fiber anthophyllite, the variety found in Georgia, usually occurs in pockets or lenses confined, for the most part, to surface zones; therefore it is quarried in shallow open pits. As operations are small, the material is loaded by hand into small cars that convey it to the mill or to the transshipping point. Quarries at Sall Mountain and Hollywood, Ga., are of this type. Slip-fiber anthophyllite, on the other hand, occurs in narrow shear zones, which necessitate excavation in narrow opencuts or underground workings. The Pylesville (Md.) operations are examples.

CANADA

Stripping is a serious problem in the Quebec district, as overburden ranges from 6 to 8 feet to a maximum of 125 feet in depth. For many years, asbestos was obtained from large open-pit quarries served by overhead cableways. About 1927, methods were modified by the introduction of new equipment—cranes or power shovels were introduced for loading, locomotives were used for hauling cars on the quarry floor, and cable hoists removed the cars from the pits up inclined planes or through tunnels. Power-shovel loading, crushing, and subsequent recovery of crude fiber from picking belts are gradually replacing hand cobbing of crudes in the quarry. This method reduces costs but also tends to reduce recovery of crude. The principal reasons for the change were (1) decreasing markets for crudes

Bureau of Mines



FIGURE 7.—Steel chutes and chain-chute gates, Asbestos Co.

ASBESTOS

ining methods are followed. Much of is up in dry walls about every cess waste and fiber-bearing rock are and sorting are done in the stopes on a f fiber are usually produced, and con- id more for the long than for the short r is found to be most satisfactory. nnels have been driven on Ash Creek uthern part of the State. Production id to be about 40 cents per pound of 3 pounds of asbestos are recovered for 7-foot tunnel.

VERMONT

ries of Vermont are situated at an south side of Belvidere Mountain ap- de Park, the nearest railway station. int, very little stripping is necessary. itered is removed in stoneboats drawn lope of the mountainside. The quarry d individual benches are 12 to 15 feet h hammer drills. Two electric power lippers are used for loading into 3 rear-dump bodies. The trucks dump ds a 28- by 36-inch jaw crusher.

AND MARYLAND

a variety found in Georgia, usually fined, for the most part, to surface in shallow open pits. As operations l by and into small cars that convey ppit int. Quarries at Sall Moun- f this type. Slip-fiber anthophyllite, arrow shear zones, which necessitate s or underground workings. The examples.

CANADA

am in the Quebec district, as over- to a maximum of 125 feet in depth. btained from large open-pit quarries About 1927, methods were modified ment—cranes or power shovels were tives were used for hauling cars on s removed the cars from the pits up els. Power-shovel loading, crushing, e fiber from picking belts are gradu- crudes in the quarry. This method educe recovery of crude. The prin- e (1) decreasing markets for crudes

Bureau of Mines

Bulletin 463



FIGURE 7.—Steel chutes and chain-chute gates, Asbestos Corporation, Ltd., mine, Thetford Mines, Quebec.

due to the introduction of molded brake linings; (2) increasing supplies of crudes from Rhodesia, the Union of South Africa, and Russia; (3) a decreasing proportion of crudes in the rock; and (4) lower prices for crudes.

A still later development in the Quebec district is the introduction of shrinkage stoping. The stone is drawn through chutes directly into boxes trammed by hand or electric locomotives through tunnels leading to the open pit, from which the material is hoisted by high-speed cableways.

The most recent innovation is an elaborate block-caving system at the King Mine of the Asbestos Corporation, Ltd., patterned after the system employed in some of the Arizona copper mines. A haulage level is established 500 feet below the surface. The overlying fiber-bearing rock is worked in blocks 160 feet square. Four grizzly drifts 40 feet apart are driven beneath the block 40 feet above the haulage level. The drifts are timbered and are provided at 20-foot intervals with heavy steel-rail grizzlies with 16-inch openings. "Finger raises" are driven upward at each side of the drift, and 20 feet above the drifts the block is undercut completely by driving drifts and blasting out the intervening pillars. The finger raises are funneled to permit access to large pieces of rock that must be broken at the grizzlies. The boundaries of the block are weakened by projecting raises and connecting drifts. The entire block sinks and is broken up by its own weight. Rock is worked through the grizzlies until the block is exhausted. Satisfactory results have been attained, and mining costs are considerably lower than by the shrinkage-stoping method. Details of the process have been published.³⁹ Figure 7 shows loading chutes on the haulage level.

The glory-hole method also is used. Raises are driven to the surface from an established level, and the rock is broken down in a funnel-shaped opening at the top of the raise. Broken rock is drawn off into cars and removed through a tunnel.

Most drilling is done with hammer drills held in the hands, but tripod drills are used for flat breast holes and for steep bench holes. Tripod drill holes range up to 18 feet deep. Heavy equipment is sometimes employed to sink holes to depths of 40 feet or more. Wet drilling is impractical because the fiber forms a spongy mass that tends to clog the drill hole.

If the fiber-bearing rock is quarried in shallow pits, loaded cars may be hauled in trains directly from the face. An elaborate spiral track lay-out is in use at one pit.

Mining costs are variable, depending upon the percentage of fiber in the rock, hardness of rock, quarry conditions, and many other factors. Ross⁴⁰ gives a direct mining and milling cost of \$10 to \$15 per ton of fiber where the rock is loaded with power shovels, and where recovery is 3 to 12 percent of gross quarry output. Where the rock is hand-sorted, shoveled into boxes, and elevated with crane or cableway hoists, the direct operating costs may range from \$20 to \$60 per ton of fiber produced.

³⁹ Ross, J. G., Block Caving at the King Mine of the Asbestos Corporation, Ltd., Thorford Mines, Quebec: Canadian Min. and Met. Bull. 264, April 1934, pp. 154-218.
⁴⁰ Ross, J. G., Chrysotile Asbestos in Canada: Canada Dept. of Mines Bull. 707, 1931, p. 57.

RHODESIA

Two mining methods—quarrying and shrinkage stoping—are used at the Shabani, the largest asbestos mine in Rhodesia.

The quarrying method is peculiar in that the rock is broken in open pits but removed through vertical shafts. Shafts or winzes are sunk in rows 50 to 60 feet apart and connected with crosscuts at a lower level. The rock around them is drilled and blasted, and the fiber is cobbled from the broken fragments. In order that the fiber may be broken as little as possible, light powder charges are fired in shallow holes. The broken rock is thrown into the shafts, which are kept nearly full, drawn into cars on the level below, and taken to the mill for extraction of the short fiber or recovery of the long fiber not removed in cobbing. The cobbled fiber is thrown down a separate shaft. The shafts are also employed for disposal of waste and for drainage.

The shrinkage-stoping method was introduced about 1926. The ground is removed in blocks 80 feet square, leaving pillars 20 to 30 feet wide, which are subsequently reclaimed.

UNION OF SOUTH AFRICA

The largest asbestos mines in the Union of South Africa are those producing chrysotile in the Barberton district of the Transvaal. The asbestos-bearing serpentine occurs in two zones. The upper one, known as the "ribbon reef", contains short-fiber veins, and the lower one provides long fiber. The belt dips 10° to 20°, and underground mining is practiced. The ribbon reef is stoped first, and at a later stage the rock carrying long fiber is removed from the footwall.

Two large mines have been developed in this territory. The New Amianthus mine is worked on several levels. In 1928 it extended down the dip a distance of 1,650 feet from the outcrop. In the same year the Munnik-Myburg mine comprised about 3,300 feet of underground workings.

In the Carolina district of the Transvaal chrysotile occurs in cross-fiber veins occupying a 5-foot zone of altered dolomite dipping 8° to 15°. Mines have been operated since 1905 and have reached a depth of 300 feet.

Amosite of the Lydenburg district of the Transvaal occurs in banded ironstones that have been highly folded and contorted. The fiber-bearing beds usually stand at angles ranging from 15° to vertical. As they are unsuited for open-pit working, underground mining is generally followed. Where the beds are intersected by transverse valleys, tunnels are driven along the strike, and the fiber-bearing rock is stoped out at various levels. Outcrops are numerous over a wide area, and fiber is obtained from many mines worked at relatively shallow depths.

The largest mines are the Egnep and the Amosa, situated near Penge. In this region the fiber-bearing zone dips from 15° to 22°. The Egnep mine is worked on seven levels. A vertical shaft 215 feet deep serves the seventh level and is connected with the sixth by a crosscut. The Amosa mine, as described by Hall in 1930, comprised 11 drifts, starting from the outcrop and ranging from 80 to 1,140 feet in length down the dip. An inclined shaft 320 feet long corresponds to a vertical depth of 290 feet.

Crocidolite (blue asbestos), which is a territory north of the Olifants River, is workings. Crocidolite also occurs in the Kuruman districts of the Cape of irregular and discontinuous, and systemed generally. Much of the fiber is obtained from shallow mines worked by native workers to depths of 250 feet with drift. Hall describes in considerable detail Africa.⁴¹

RUSSIA

The asbestos deposits of the Bajano burden. Hand methods of removal with shovels or hydraulic methods have extent.

More than 20 shallow, open quarries. Up to 1929 drilling was the only mechanical operations being conducted with hand-milling facilities, hand picking began process, and only one-fourth to one-fifth sent to the mill. The average fiber content only about 6 percent, but because of hand picking, that sent to the mill was.

The first steps toward more complete 1929, with the installation of 2- to 4-ton with fixed foot towers and traveling inclined haulageways served by electric in benches 5 to 6 meters high and class waste.

In 1929 two shafts were sunk and at a depth of 50 meters. They were which proved uneconomical. This was largely superseded in 1930 by transport by locomotives. The cars are taken by electric hoists. Mining 75 cents a metric ton of rock. The methods have been supplied by Russia.

CYPRUS

Chrysotile asbestos occurs on the regular veins traversing serpentine. The expansion accompanying metamorphism resulted in excessive fracturing, which workings consist of open-pit terrace 250 feet. Much of the rock is so fine can be broken with picks without the when compressed-air drills were introduced drilled for blasting by hand methods.

⁴¹ Hall, A. L., *Asbestos in the Union of South Africa*, 12, 2d ed., 1930.

⁴² Ru Keyser, Walter A., *Mining Asbestos in South Africa*, September 1933, pp. 375-381.

RHESIA

g a: shrinkage stoping—are used in that the rock is broken in open shafts. Shafts or winzes are sunk connected with crosscuts at a lower level and blasted, and the fiber is recovered. In order that the fiber may be powder charges are fired in shallow into the shafts, which are kept level below, and taken to the mill recovery of the long fiber not recovered is thrown down a separate shaft. Disposal of waste and for drainage was introduced about 1926. The set square, leaving pillars 20 to 30 reclaimed.

SOUTH AFRICA

The Union of South Africa are those in the district of the Transvaal. The is in two zones. The upper one, has short-fiber veins, and the lower dips 10° to 20°, and underground reef is stoped first, and at a later is removed from the footwall.

Developed in this territory. The New several levels. In 1928 it extended set from the outcrop. In the same comprised about 3,300 feet of under-

ran 1 chrysotile occurs in cross- of altered dolomite dipping 8° to ice 1905 and have reached a depth

district of the Transvaal occurs in highly folded and contorted. The angles ranging from 15° to vertical. working, underground mining is beds are intersected by transverse strike, and the fiber-bearing rock outcrops are numerous over a wide many mines worked at relatively

ep and the Amosa, situated near bearing zone dips from 15° to 22°. in levels. A vertical shaft 215 feet is connected with the sixth by a cribed by Hall in 1930, comprised and ranging from 80 to 1,140 feet ed shaft 320 feet long corresponds

Crocidolite (blue asbestos), which is associated with amosite in the territory north of the Olifants River, is mined in small underground workings. Crocidolite also occurs over a wide area in the Prieska and Kuruman districts of the Cape of Good Hope. The veins are irregular and discontinuous, and systematic mining cannot be followed generally. Much of the fiber is obtained from small open pits or from shallow mines worked by natives. A few mines have been worked to depths of 250 feet with drifts over 1,000 feet long.

Hall describes in considerable detail asbestos mining in South Africa.⁴¹

RUSSIA

The asbestos deposits of the Bajanova district have a thin overburden. Hand methods of removal were first employed, but power shovels or hydraulic methods have superseded them to quite an extent.

More than 20 shallow, open quarries were worked for many years. Up to 1929 drilling was the only mechanical process used, all other operations being conducted with hand tools. Owing to inadequate milling facilities, hand picking became an important concentration process, and only one-fourth to one-fifth of the rock quarried was sent to the mill. The average fiber content of the rock mined was only about 6 percent, but because of the concentration attained by hand picking, that sent to the mill was 22 percent.

The first steps toward more complete mechanization were taken in 1929, with the installation of 2- to 4-ton capacity overhead cableways with fixed foot towers and traveling head towers, supplemented by inclined haulageways served by electric hoists. The rock was blasted in benches 5 to 6 meters high and classified into crudes, mill feed, and waste.

In 1929 two shafts were sunk and connected with a haulage level at a depth of 50 meters. They were designed for glory-hole mining, which proved uneconomical. This method was supplemented and largely superseded in 1930 by electric-shovel loading in open pits with transport by locomotives to inclined haulageways, up which cars are taken by electric hoists. Mining costs are estimated at about 75 cents a metric ton of rock. Details of quarrying and mining methods have been supplied by Ru Keyser.⁴²

CYPRUS

Chrysotile asbestos occurs on the slopes of Mount Troodos in irregular veins traversing serpentine formed by alteration of olivine. The expansion accompanying metamorphism of the original olivine resulted in excessive fracturing, which makes excavation easy. The workings consist of open-pit terraces reaching a maximum height of 250 feet. Much of the rock is so fractured and decomposed that it can be broken with picks without the aid of explosives. Before 1927, when compressed-air drills were introduced, the harder rock was drilled for blasting by hand methods. When 40-pound hand hammer

⁴¹ Hall, A. L., *Asbestos in the Union of South Africa: Geol. Survey of South Africa Mem. 12, 2d ed., 1930.*

⁴² Ru Keyser, Walter A., *Mining Asbestos in U. S. S. R.: Eng. and Min. Jour., vol. 124, September 1933, pp. 375-381.*

drills are used, 3-inch holes ranging in depth from 8 to 12 feet are sunk at a rate of about 10 feet an hour. The larger masses of barren rock are removed as waste. Fiber-bearing material is turned over to facilitate drying in the hot sun, and it is then passed over gravity screens. That passing through an 18-mm-mesh screen and retained on a 5-mm screen (about one-fifth of the total material quarried) is sent to the mills for recovery of the fiber. Oversize rock retained on the upper screen, and sand passing the lower screen are trammed to waste heaps. The quarries are worked from March to December. In 1927 approximately 6,000 workers were employed in the quarries and about 1,500,000 tons of rock were broken. Operations were still more extensive in 1928 and 1929, decreased greatly during the depression years, but recovered substantially in 1934 and 1935. Further details of quarry methods are given by Whitworth.⁴³

Production consisted exclusively of short fiber until 1934, when, according to report, a small quantity of spinning fiber was shipped.

CHINA

Primitive methods are followed in the Laiyuan district. The mining tools used are hammers, chisels, picks, and shovels. No machinery or explosives are employed. Small inclined tunnels are sunk along the veins to a maximum depth of about 30 meters. The fiber is cobbled and sorted by hand. Mining costs are said to range from \$96 to \$137 a ton of fiber produced.

MILLING METHODS

CLASSIFICATION OF FIBER

A brief discussion of the method of classifying asbestos fibers is a necessary preliminary to a description of milling methods.

Canadian and Vermont asbestos fibers are divided into three main groups—crudes, mill fibers, and shorts. The term "crude" is applied to fiber of spinning grade, measuring three-eighths inch or longer, that is hand-cobbled and not passed through a mill. Mill fibers are obtained by crushing and beating the fiber-bearing rock until the asbestos is freed, and then removing the fiber from the rock by screening or air separation. Shorts are the lowest grades of mill products.

Russian asbestos also is classed as crudes and mill fibers, but the term "crude" is applied somewhat loosely; it apparently includes some spinning fiber that has been prepared mechanically.

The term "crude" is not generally applied to African products. Spinning fibers comparable in quality with Canadian crudes are prepared by hand-cobbing alone or in conjunction with simple mechanical crushing, disintegrating, and screening processes. A much larger proportion of the African long fibers has been prepared mechanically during recent years. Mill fibers of the shorter grades constitute the largest part of the tonnage.

Virtually all of the asbestos produced in Cyprus is classed as mill fiber.

⁴³ Whitworth, M., *Cyprus and Its Asbestos Industry*: Mining Mag. (London), vol. 39, September 1928, pp. 143-150.

GENERAL FEATURES OF MILLING

As pointed out on preceding pages, the occurrences are (1) deposits derived from tite-pyroxenite rocks and (2) deposits of limestone intruded by diabase. The first type of asbestos in Vermont, Quebec, the Ural Mountains, and the Transvaal. The second type, which represents of the world output, occurs in Arizona; Minn. Carolina district of South Africa. The first type is associated with asbestos of the first type is materials with the second type are serpentine and talc also is serpentine, differing only in structure. It is obvious that the milling of chrysotile forms of ore concentration. As the valuable consist of the same material, neither chemical nor gravity can be utilized as the basis for separation that makes mechanical separation of chrysotile structure, which permits it to be opened and separate filaments, thus making it amenable to gangue by air suction and screening.

The value of chrysotile asbestos depends on the fiber. When fiber three-eighths inch or longer, fiber three-fourths inch long is worth at least \$100 a ton. Thus, it is apparent that breaking fibers into three-fourths of their value. The underlying the milling of asbestos, therefore, is to break the fiber from rock with minimum breakage of spare the fiber unnecessarily rough treatment. Modern mills break the rock effectively. Modern mills separate the fiber after each crushing process as soon as it is formed, and to keep hard. If fiber already freed from rock enters the mill with sand and rock fragments, it will be lost. Asbestos milling consists essentially of a series of recrushing in stages, each step being followed by separation of fiber from rock.

Another peculiarity of asbestos milling is that the known method of testing the quality of the product appears to contain little fiber may yield one that appears promising may show a disastrous result. As both the valuable mineral and the asbestos are useless. A laboratory-scale crushing unit might give results that differ widely from a mill operating on the same rock. Although the recoverable fiber in a rock cannot be determined, the yield can be calculated from the mill output. The value of a fiber-bearing rock can be determined by a test run in a commercial mill.

In the following discussion of milling methods, world, Canadian practice is considered first, followed by the pioneers in the asbestos industry, and finally a basis for practice in other countries.

ing in depth from 8 to 12 feet are our the larger masses of barren material is turned over to and it is then passed over gravity 18-mm-mesh screen and retained of the total material quarried) is fiber. Oversize rock retained on the lower screen are trammed to work from March to December. rs were employed in the quarries re broken. Operations were still decreased greatly during the de- tially in 1934 and 1935. Fur- iven by Whitworth.⁴³ of short fiber until 1934, when, y of spinning fiber was shipped.

the Laiyuan district. The min- picks, and shovels. No machin- small inclined tunnels are sunk h of about 30 meters. The fiber ing costs are said to range from

METHODS

OF FIBER

of classifying asbestos fibers is a on of milling methods. bers are divided into three main ts. term "crude" is applied ng three-eighths inch or longer. through a mill. Mill fibers are he fiber-bearing rock until the ng the fiber from the rock by s are the lowest grades of mill

crudes and mill fibers, but the loosely: it apparently includes repared mechanically. y applied to African products. lity with Canadian crudes are in conjunction with simple med screening processes. A much long fibers has been prepared Mill fibers of the shorter grades nage. duced in Cyprus is classed as

industry: Mining Mag. (London), vol. 39,

GENERAL FEATURES OF MILLING METHODS

As pointed out on preceding pages, the two major types of chrysotile occurrences are (1) deposits derived from alteration of peridotite-pyroxenite rocks and (2) deposits derived from dolomitic limestone intruded by diabase. The first type is the principal source of asbestos in Vermont, Quebec, the Ural Mountains, Rhodesia, and the Transvaal. The second type, which represents a small fraction of the world output, occurs in Arizona; Minusinsk, Siberia; and the Carolina district of South Africa. The principal gangue material associated with asbestos of the first type is serpentine, and the materials with the second type are serpentine and limestone. As chrysotile also is serpentine, differing only in structure from the gangue, it is obvious that the milling of chrysotile must differ from all other forms of ore concentration. As the valuable mineral and the gangue consist of the same material, neither chemical composition nor specific gravity can be utilized as the basis for separation. The property that makes mechanical separation of chrysotile possible is its fibrous structure, which permits it to be opened or divided by impact into separate filaments, thus making it amenable to separation from the gangue by air suction and screening.

The value of chrysotile asbestos depends largely on the length of the fiber. When fiber three-eighths inch long is worth \$100 a ton, fiber three-fourths inch long is worth approximately \$400 a ton. Thus, it is apparent that breaking fibers in half destroys approximately three-fourths of their value. The most important principle underlying the milling of asbestos, therefore, is the separation of fiber from rock with minimum breakage of fibers. The best machines spare the fiber unnecessarily rough treatment and at the same time break the rock effectively. Modern mills are designed to remove the separated fiber after each crushing process, to separate the sand as soon as it is formed, and to keep hard, barren rock out of the mill. If fiber already freed from rock enters the next crushing unit along with sand and rock fragments, it will be broken into shorter grades. Asbestos milling consists essentially of coarse crushing, drying, and recrushing in stages, each step being followed by screening and air separation of fiber from rock.

Another peculiarity of asbestos milling is the absence of any known method of testing the quality of mill feed. A rock that appears to contain little fiber may yield a fair return; conversely, one that appears promising may show a disappointingly low recovery. As both the valuable mineral and the gangue are serpentine, an assay is useless. A laboratory-scale crushing and fiber-separating unit might give results that differ widely from those of a commercial mill operating on the same rock. Although the percentage of recoverable fiber in a rock cannot be determined in advance, the actual yield can be calculated from the mill output. A fair estimate of the value of a fiber-bearing rock can be obtained only by making a test run in a commercial mill.

In the following discussion of milling methods throughout the world, Canadian practice is considered first because Canadians were the pioneers in the asbestos industry, and their methods have established a basis for practice in other countries.

CANADA

PREPARATION OF CRUDES

In Canada the associated rock is broken free of the larger-fiber veins by sledging, and the masses of fiber are sorted by hand. The fiber with some adhering rock is dried on steam coils and then taken to the cobbing sheds, where the fiber veins are flattened with a hand hammer, freeing them from adhering rock. The separated rock, dust, and short fibers are dropped into a receptacle under the bench, and the remaining high-grade fibers are classed as Crude No. 1, material three-fourths inch long and over, and Crude No. 2, material three-eighths to three-fourths inch long. To separate further the short fiber and rock dust, Crude No. 1 is screened on a flat shaking screen with three-eighths-inch holes and No. 2 on one with three-sixteenths-inch holes. After being screened, the fibers are bagged in jute bags of 100 pounds capacity. Five to twenty percent of rock fragments, dust, and short fiber still remain in the crudes as sold, and these impurities are separated at the factory where asbestos products are manufactured. Rejects accumulated in preparing crudes are utilized as mill fibers or sold as screenings. Crudes constitute a small proportion of the recoverable fiber but are the most valuable products.

Ru Keyser⁴⁴ has developed a process of preparing long fiber mechanically, by which rolls are used instead of hand hammers. The fibers are graded on screens that are so operated that the fibers are held in a horizontal plane. The claim is made that fibers thus prepared are superior in grade and purity to hand-cobbed crudes.

MILLING PRACTICE

Mechanical means of preparing fiber began in Quebec in 1888, and by 1896 important tonnages of milled fiber were sold. The principal processes involved in milling are crushing, drying, storing, fiberizing, screening, air separating, and grading.

PRELIMINARY CRUSHING

Rock from the quarries or mines is dumped into a so-called "sluice" that holds several carloads. The sluice may have a railroad-rail-grizzly bottom that provides a bypass for fines. The primary-crusher feed from the sluice is controlled by finger gates—that is, suspended rails that may be moved up or down. A jaw crusher with a 36-inch by 24-inch opening set for a 4- to 6-inch discharge is the most popular primary breaker. Larger or smaller units may be employed as needed. Crushers discharge to picking belts, where liberated crudes are removed by hand.

DRYING

The quarry rock contains considerable moisture, in winter even snow and ice, and it must be dried for proper milling. Standard rotary driers generally are used. The shells are 40 to 60 feet long, 4 to 6 feet in diameter, and inclined at an angle of 7°. The material is lifted by a series of blades and allowed to fall through the cur-

rent of hot air that circulates through range from 30 to 60 tons of rock an hour cents a ton.

Stack driers also are used and are calculated. Rock is elevated to the top of a 50-ft and allowed to fall through a rising current to interrupt the speed of the descent.

STORAGE

A large supply of rock in storage is fed to the mill on belt conveyors carried beneath the storage bins. By drawing rock at the same time material of comparatively uniform quality is obtained. Bins having capacities of 200 tons are in use.

FIBERIZING

Although crushing, drying, and storing form throughout the Canadian district, no flow sheet can be applied generally. No flow sheet can be applied must be modified to suit the rock. A mill would be too severe for fiber enclosed in a casing may be done with jaw or gyratory or hammer mills. A straight impact mill is preferred to a grinding or shearing mill, less desirable than impact mills.

The following flow sheet, given by the best modern practice.

Grizzly in rock sluice.
Jaw crusher.
Drier.
Grizzly, fines to screens fitted
Gyratory crusher.
Screens with suction.
Rolls.
Screens with suction.
Jumbo.
Screens with suction.
Jumbo.
Screens with suction.
Jumbo.
Screens with suction.
Jumbo.
Screens with suction.
Rotary graders and flat cleaners
Floats bin.
Sand to dump.

It will be observed that the principle of each stage of crushing is followed strictly.

According to this flow sheet, crushing is accomplished with gyratory crushers. Jumbo, which is manufactured locally, is 40 feet long and 24 to 30 inches in diameter. The horizontal shaft consists of arms that support the beaters. The beaters are set at intervals

⁴⁴Ru Keyser, Walter A., *Mechanical Cobbing of Chrysotile Asbestos*: Eng. and Min. Jour., vol. 134, no. 6, June 1933, pp. 235-237.

⁴⁵Ross, J. G., *Chrysotile Asbestos in Canada*: Can. Jour. Min., p. 42.

CANADA

TWO CRUDES

Rock is broken free of the larger-fiber lumps of fiber are sorted by hand. The rock is dried on steam coils and then taken to fiber veins are flattened with a hand adhering rock. The separated rock, placed into a receptacle under the bench, fibers are classed as Crude No. 1, and over, and Crude No. 2, material 1/2 inch long. To separate further the No. 1 is screened on a flat shaking holes and No. 2 on one with three-inch screened, the fibers are bagged in it. Five to twenty percent of rock still remain in the crudes as sold, rated at the factory where asbestos

Rejects accumulated in preparing or sold as screenings. Crudes contain recoverable fiber but are the most

process of preparing long fiber measured instead of hand hammers. That it is so operated that the fibers are a claim is made that fibers thus prepared purity to hand-cobbed crudes.

MILLING PRACTICE

Long fiber began in Quebec in 1888, and milled fiber were sold. The principal crushing, drying, storing, fiberizing, and

PRIMARY CRUSHING

Rock is dumped into a so-called "sluice" the sluice may have a railroad-railway bypass for fines. The primary-controlled by finger gates—that is, raised up or down. A jaw crusher with for a 4- to 6-inch discharge is the larger or smaller units may be discharge to picking belts, where hand.

DRYING

considerable moisture, in winter even dried for proper milling. Standard

The shells are 40 to 60 feet long, dried at an angle of 7°. The material allowed to fall through the cur-

Crushing of Chrysotile Asbestos: Eng. and Min.

rent of hot air that circulates through the cylinder. Capacities range from 30 to 60 tons of rock an hour and costs from 1/2 to 18 cents a ton.

Stack driers also are used and are said to be much more economical. Rock is elevated to the top of a 50-foot stack 7 feet in diameter and allowed to fall through a rising current of hot air. Grid bars interrupt the speed of the descent.

STORAGE

A large supply of rock in storage insures regular mill feed. Rock is fed to the mill on belt conveyors carried in concrete tunnels beneath the storage bins. By drawing rock from several gates at the same time material of comparatively uniform size and fiber content is obtained. Bins having capacities of 25,000 to 150,000 tons are in use.

FIBERIZING

Although crushing, drying, and storing methods are fairly uniform throughout the Canadian district, milling methods vary greatly. No flow sheet can be applied generally because milling must be modified to suit the rock. A method suitable for hard rock would be too severe for fiber enclosed in soft rock. Secondary crushing may be done with jaw or gyratory crushers, rolls, Symons cones, or hammer mills. A straight impact that frees and fluffs the fiber is preferred to a grinding or shearing action. Therefore, rolls are less desirable than impact mills.

The following flow sheet, given by Ross,⁴⁵ probably represents the best modern practice.

Grizzly in rock sluice.
Jaw crusher.
Drier.
Grizzly, fines to screens fitted with suction.
Gyratory crusher.
Screens with suction.
Rolls.
Screens with suction.
Jumbo.
Screens with suction.
Jumbo.
Screens with suction.
Jumbo.
Screens with suction.
Jumbo.
Screens with suction.
Rotary graders and flat cleaning screens.
Floats bin.
Sand to dump.

It will be observed that the principle of removing the fiber after each stage of crushing is followed strictly.

According to this flow sheet, crushing beyond the primary stage is accomplished with gyratory crushers and Jumbo machines. The Jumbo, which is manufactured locally, is a horizontal drum 6 to 8 feet long and 24 to 30 inches in diameter. Beaters attached to a horizontal shaft consist of arms that support manganese-steel hammers. The beaters are set at intervals of 6 or 8 inches along the

⁴⁵ Ross, J. G., Chrysotile Asbestos in Canada: Canada Dept. of Mines Bull. 707, 1931, p. 42.

shaft and clear the inside of the shell by about $\frac{1}{4}$ to 1 inch. They are turned at an angle to direct the broken rock toward the discharge end. The shaft rotates at 400 to 800 revolutions per minute, according to the nature of the rock treated. Instead of the Jumbo machine, some mills employ various types of cyclones, which beat the rock to a powder. The Laurie cyclone consists of a cast-iron chamber in which two beaters shaped like ship propellers are driven in opposite directions at a speed of 1,700 to 2,000 revolutions per minute. In the Pharo cyclone the beaters revolve in the same direction and the rate of discharge is controlled by gates. The Torrey cyclone is somewhat like the Jumbo, except that the axis of the beater arms is vertical.

SCREENING

Shaking screens actuated by eccentrics generally are used, although vibrating screens of the Hummer type also are employed. Usual sizes are 3 to 5 feet wide and 8 to 12 feet long. Perforated steel plates are more generally used than wire mesh. Screens with holes up to three-eighths of an inch and larger are used for scalping, and smaller openings down to one thirty-second of an inch for the very fine materials. Wire screens ranging from 3 to 24 meshes to the inch are commonly used for cleaning fiber.

Fines pass through the openings, but the coarser rock fragments and longer fibers remain on the screen. The shaking action brings the fiber to the surface of the bedded-rock fragments, and it is removed by air-suction equipment that will be described later. Thus, each screen furnishes three products—fines that pass through as waste, coarse rock fragments discharged at the end, and fiber collected by suction. Oversize rock fragments pass to the next fiberizing unit.

AIR SUCTION

Fiber segregated on the surface of a layer of rock fragments on the screen is removed through suction hoods that operate on the principle of a vacuum cleaner. The opening of the suction hood is 3 to 4 inches wide and extends across the full width of the screen at its lower end. The cross-section area of the narrow opening is the same as that of the pipe into which it converges, which ranges from 12 to 20 inches in diameter. Large fans produce the necessary vacuum. A 10- to 15-horsepower motor is required to drive a fan giving the necessary lifting power through a pipe 15 inches in diameter. The air current sweeps the fiber from the screens, carries it through pipes, and aspirates it in collecting chambers, where the diminished rate of the air current permits it to fall. The dust-laden air from the fans is blown into dust chambers, and the fine materials that settle there are returned to the mill, where the longer fibers are separated and the extremely short grades are bagged and sold as "floats."

WET PROCESS

Considerable progress has been made in developing a wet process for treating Canadian asbestos. Some claim that wet treatment involves less breakage of fiber and a higher recovery. Experiments were begun in 1921, and a testing plant was built in 1923, but commercial usage has not been attempted.

GRADING

Fiber is graded according to length and 25 inches in diameter. Paddles clamped to a central shaft agitate grades through the screen. One type is a stationary trommel in which the paddles rotate at 5 to 10 revolutions per minute. The tips of the paddles are at an angle, which causes the fiber to travel in a spiral. The machines are so designed that the trommel makes one revolution at 5 revolutions per minute and the paddles at 180 to 250 revolutions per minute. This makes only one finished grade—that which passes through the screen and the oversize is conducted to a second screen. Thus, several trommels may operate on the same grade, never mills the trommel is equipped with screens which make one oversize and two undergrades. Of grades may be made, and special blending. Sand and dust are removed by flat cleaning screens. Finished fiber is conveyed to the store by air suction.

CLASSIFICATION

Methods of grading asbestos fiber by length. To determine with reasonable accuracy the grade and thus be able to modify grading standard testing machine is used. The standard test and thus avoid any dependence upon trade names or descriptions of the standard testing machine "follows."

The machine consists of a nest of four trays, 24½ by 14¾ inches and 3½ inches in superposed one above the other. The trays are numbered 1, 2, 3, and 4. The bottoms of the trays are of the following specifications: Tray 1: 10-mesh wire, 0.105 inch. Tray 2: 20-mesh wire, 0.047 inch. Tray 3: 30-mesh wire, 0.047 inch. Tray 4: 40-mesh wire, 0.047 inch. The trays fall through the three other boxes. The trays rest on a table to which an eccentric gives a movement of 1½-inch travel.

To make a test, 16 ounces of asbestos fiber is covered. The machine is run at 100 revolutions per minute at the shaft of the eccentric. The device this is kept going for exactly 10 minutes. At the end of that time that remains on each tray is weighed. The longest fiber that remains on the first tray is the longest fiber; the longest fiber that remains on the second tray is the second longest fiber; and so on. The shorter fiber, according to 2 and 3 or drops into the pan or low tained on the first screen and the less the grade and, therefore, the greater the grade.

* Ross, J. G., Work cited, p. 49.

shell by about $\frac{1}{2}$ to 1 inch. They the broken rock toward the dis-400 00 revolutions per minute, k-treated. Instead of the Jumbo us types of cyclones, which beat ie cyclone consists of a cast-iron ed like ship propellers are driven of 1,700 to 2,000 revolutions per beaters revolve in the same direc- controlled by gates. The Torrey nbo, except that the axis of the

SING

atrices generally are used, although - type also are employed. Usual o 12 feet long. Perforated steel n wire mesh. Screens with holes larger are used for scalping, and y-second of an inch for the very ing from 3 to 24 meshes to the g fiber.

but the coarser rock fragments reen. The shaking action brings dded-rock fragments, and it is at will be described later. Thus, cts—fines that pass through as arged at the end, and fiber col- gments pass to the next fiberizing

TION

of a layer of rock fragments on tion als that operate on the e op g of the suction hood is s the full width of the screen at ea of the narrow opening is the it converges, which ranges from fans produce the necessary vac- is required to drive a fan giving h a pipe 15 inches in diameter, n the screens, carries it through chambers, where the diminished fall. The dust-laden air from ers, and the fine materials that ll, where the longer fibers are grades are bagged and sold as

ESS

ide in developing a wet process ne claim that wet treatment in- higher recovery. Experiments ant was built in 1923, but com- L

GRADING

Fiber is graded according to length in trommels about 8 feet long and 28 inches in diameter fitted with woven-wire cloth. Paddles clamped to a central shaft agitate the fiber and force the shorter grades through the screen. One type of machine consists of a stationary trommel in which the paddles rotate at 400 to 900 revolutions per minute. The tips of the paddles are twisted at a small angle, which causes the fiber to travel through the machine. Other machines are so designed that the trommel rotates in one direction at 5 revolutions per minute and the paddles rotate in the opposite direction at 180 to 250 revolutions per minute. Some trommels make only one finished grade—that passing through the screen—and the oversize is conducted to a second trommel for further grading. Thus, several trommels may operate in series. At one of the newer mills the trommel is equipped with two sizes of screen cloth, which make one oversize and two undersize grades. Any number of grades may be made, and special grades may be obtained by blending. Sand and dust are removed from the graded fiber on flat cleaning screens. Finished fiber lifted from the cleaning screen by air suction is conveyed to the store room for bagging.

CLASSIFICATION

Methods of grading asbestos fiber by length have been described. To determine with reasonable accuracy the quality of fiber in each grade and thus be able to modify grading methods accordingly, a standard testing machine is used. The purchaser can rely on the standard test and thus avoid any confusion that might ensue from dependance upon trade names or designations. A description of the standard testing machine⁴⁸ follows:

The machine consists of a nest of four wooden boxes measuring $24\frac{1}{2}$ by $14\frac{3}{4}$ inches and $3\frac{1}{2}$ inches in depth. The boxes, which are superposed one above the other, are numbered, from the top down, 1, 2, 3, and 4. The bottoms of boxes 1, 2, and 3 are made of metallic screen of the following specifications: Box 1: $\frac{1}{2}$ -inch opening, diameter of wire, 0.105 inch. Box 2: 4-mesh wire, 0.063 inch. Box 3: 10-mesh wire, 0.047 inch. Box 4 is a receptacle for the fines that fall through the three other boxes. The nest of four boxes or trays rests on a table to which an eccentric with a throw of $\frac{3}{8}$ inch gives a movement of $1\frac{1}{8}$ -inch travel.

To make a test, 16 ounces of asbestos is put on the top tray, which is covered. The machine is run at the rate of 300 revolutions per minute at the shaft of the eccentric, and by means of an automatic device this is kept going for exactly 2 minutes, giving the nest a horizontal shaking movement. At the end of this time the asbestos that remains on each tray is weighed. This gives the grades of the asbestos fiber; the longest fiber naturally stays on the top tray, whereas the shorter fiber, according to its length, remains on screens 2 and 3 or drops into the pan or lowest tray. The more fiber retained on the first screen and the less fiber in the pan, the higher the grade and, therefore, the greater its value. If, for instance, a

⁴⁸ Ross, J. G., Work cited, p. 49.

customer buys spinning fiber of the specification 4-7-1-1, it means that in a sample of 16 ounces, representing the average of the lot shipped, 4 ounces will remain on the top screen, 7 on the second, 4 on the third, and, finally, 1 ounce will go through all the screens into the pan. He will evidently pay more for this material than for paper stock testing 0-0-10-6. This indicates that out of 16 ounces tested, nothing is retained on the first two screens, 10 ounces remain on the third, and 6 ounces go through the latter into the pan. It is evident that the figures of the test represent the proportion, in ounces, of the different lengths of fiber in a pound of asbestos.

Samples for testing usually are taken at the grading machines every half hour. Owing to unavoidable variations in fiber as it comes from the pit, the quality of mill-run fiber usually is maintained a little higher than its designation.

For many years each Canadian company graded its fiber according to its own standards and sold the products under its own trade designations. This practice led to much confusion in marketing because different mills employed similar marks for grades quite unequal in quality and value. It was frequently necessary for buyers to have samples tested before placing orders. In 1931, producers in Quebec agreed upon a uniform classification of fibers whereby they are divided into nine groups, and each group was divided into grades. Crude asbestos was defined as "hand-selected cross-vein material essentially in its native or unfiberized form", and milled asbestos as "all grades produced by mechanical treatment of asbestos ore." Except for the very lowest grades, which are based on the weight per cubic foot, all milled grades are based on the results of tests in the standard testing machine previously described. Ross⁴ gives a complete description of the standard grades of Canadian asbestos as agreed upon in 1931:

Crude asbestos

Class	Standard designation of grade	Description
Group no. 1.....	Crude No. 1.....	Consists basically of crude 3½-inch staple and longer.
Group no. 2.....	Crude No. 2.....	Consists basically of crude 2½-inch staple up to 3½-inch.
	Crude run-of-mine.....	Consists basically of unsorted crudes.
	Crude sundry.....	Consists of crudes other than above specified.

Milled asbestos

Group no. 3.....	Spinning or textile fiber...	Consists of fiber testing 0-8-6-2 and over.
Group no. 4.....	Shingle fiber.....	Consists of fiber testing below 0-8-6-2 to and including 0-1½-9½-5.
Group no. 5.....	Paper fiber.....	Consists of fiber testing below 0-1½-9½-5 to and including 0-0-8-8.
Group no. 6.....	Waste, stucco or plaster...	Consists of material testing below 0-0-8-8 to and including 0-0-6½-9½.
Group no. 7.....	Refuse or shorts.....	Consists of material testing 0-0-5-11 and below, including material testing below 0-0-1-15 and specified as weighing 35 pounds or less per cubic foot, loose measure.
Group no. 8.....	Sand.....	Consists of such asbestos mill products as sand weighing over 35 pounds per cubic foot, loose measure, and under 75 pounds per cubic foot, loose measure, and containing a preponderance of rock.
Group no. 9.....	Gravel and stone.....	Consists of such asbestos mill products weighing 75 pounds and over per cubic foot, loose measure.

⁴Ross, J. G., Work cited, pp. 50-51.

Subdivision of the groups of mill

Group no.	Standard designation of grade	
No. 3. Textile and spinning fibers.....	3D 3F 3K 3M 3R 3T 3Z	8 4 2 2 1 0
No. 4. Shingle fibers.....	4D 4F 4K 4M 4R 4T 4Z	0 0 0 0 0 0
No. 5. Paper fibers.....	5D 5F 5K 5M 5R 5T 5Z	1 1 1 1 1 1
No. 6. Waste, stucco, or plaster.....	6D 6F	1 1
No. 7. Refuse or shorts.....	7D 7F 7K 7M 7R 7T 7Z	1 1 1 1 1 1
No. 8. Sand.....	8-40 8-45 8-55 8-75	1 1 1 1
No. 9. Gravel and stone.....	9	1

The fiber is packed in jute bags of 1 marked with the grade letter or number name.

ARIZONA

The high cost of transportation to charges to markets render the sale of shingle. Spinning fibers are therefore the Arizona deposits. The smaller mines are almost exclusively. The rock is mined by workers on a contract basis. Cobbing is way as in Canada.

Three of the larger mines have mills equipped with crushers, cyclones, and fiber ¾ to 2½ inches long; No. 2, ¾ to 1 inch long; and No. 4, ½ inch and less, according to tests made with a standard

In one of the smaller mills the rock crusher followed by two sets of rolls, rated by an impact screen. The short small fiberizing machine and is cleaned feet long and 4 feet in diameter.

specification 4-7-4-1, it means "average of the lot of 7 on the second, 4 go through all the screens into ore for this material than for indicates that out of 16 ounces of two screens, 10 ounces remain on the latter into the pan. It represents the proportion, in a pound of asbestos, taken at the grading machines, variable variations in fiber as it runs, fiber usually is maintained

company graded its fiber according to products under its own trade name, confusion in marketing marks for grades quite unequal necessary for buyers to have

In 1931, producers in Quebec of fibers whereby they are disposed was divided into grades. selected cross-vein material "es-form", and milled asbestos as "treatment of asbestos ore." which are based on the weight per on the results of tests in the described. Ross⁴⁷ gives a com- of Canadian asbestos as

Subdivision of the groups of milled asbestos

Group no.	Standard designation of grades	Guaranteed minimum shipping test
No. 3. Textile and spinning fibers.....	3D 3F 3K 3M 3R 3T 3Z	8-4-1-1. 7-7-1½-½. 4-7-4-1. 2-9-4-1. 2-8-4-2. 1-9-4-2. 0-8-6-2.
No. 4. Shingle fibers.....	4D 4F 4K 4M 4R 4T 4Z	0-5-10-1. 0-3-12-1. 0-4-9-3. 0-1-3-4. 0-3-9-4. 0-2-10-4. 0-1½-9½-5.
No. 5. Paper fibers.....	5D 5F 5K 5M 5R 5T 5Z	0-1½-10½-5. 0-1½-9½-6. 0-0-12-4. 0-0-11-5. 0-0-10-6. 0-0-9-7. 0-0-8-8.
No. 6. Waste, stucco, or plaster.....	6D 6F	0-0-7-9. 0-0-6½-9½.
No. 7. Refuse or shorts.....	7D 7F 7K 7M 7-20 7-25 7-30 7-35	0-0-5-11. 0-0-4-12. 0-0-3-13. 0-0-2-14. 0-0-1-15. 20 pounds per cubic foot, loose measure. 25 pounds per cubic foot, loose measure. 30 pounds per cubic foot, loose measure. 35 pounds per cubic foot, loose measure.
No. 8. Sand.....	8-40 8-45 8-50 8-55 8-75	40 pounds per cubic foot, loose measure. 45 pounds per cubic foot, loose measure. 50 pounds per cubic foot, loose measure. 55 pounds per cubic foot, loose measure. 75 pounds per cubic foot, loose measure.
No. 9. Gravel and stone.....	9	75 pounds and over per cubic foot, loose measure.

The fiber is packed in jute bags of 100 or 125 pounds capacity marked with the grade letter or number and the manufacturer's name.

ARIZONA

The high cost of transportation to railways and heavy freight charges to markets render the sale of shorter Arizona fibers unprofitable. Spinning fibers are therefore the principal products of the Arizona deposits. The smaller mines produce hand-cobbed crudes almost exclusively. The rock is mined and cobbed by the same workers on a contract basis. Cobbing is conducted in much the same way as in Canada.

Three of the larger mines have milling facilities. The largest mill, equipped with crushers, cyclones, and screens, produces No. 1 mill fiber ¾ to 2½ inches long; No. 2, ¾ to ¾ inch long; No. 3, 1½ to 3½ inch long; and No. 4, 1½ inch and less. The products are graded according to tests made with a standard Canadian testing machine.

In one of the smaller mills the rock first passes through a jaw crusher followed by two sets of rolls. Two grades of fiber are separated by an impact screen. The shorter asbestos is treated with a small fiberizing machine and is cleaned and graded in a trommel 16 feet long and 4 feet in diameter.

708

Description

Quality of crude ¾-inch staple and longer.
Quality of crude ¾-inch staple up to ¾-inch.
Quality of unsorted crudes.
Crudes other than above specified.

709

For testing 0-2-4-2 and over.
For testing below 0-2-4-2 to and including
For testing below 0-1½-9½-5 to and including
For testing below 0-0-8-8 to and including
For testing 0-0-5-11 and below, including
For testing below 0-0-5-15 and specified as weighing
Less per cubic foot, loose measure.
For asbestos mill products as sand weighing
40 pounds per cubic foot, loose measure, and under
45 pounds per cubic foot, loose measure, and under
50 pounds per cubic foot, loose measure, and containing
more than 1% of rock.
For asbestos mill products weighing 75 pounds
per cubic foot, loose measure.

Another small mill produces five grades of spinning fibers and a shingle stock. No grades shorter than shingle stock can be sold profitably.

A considerable quantity of rock containing spinning-fiber veins, which are difficult to cob by hand, are treated in these mills and yield what is known as mechanically cobbled crude fiber, an anomalous designation, according to the Canadian definition of crude, which includes only hand-cobbled fiber. The claim is made, however, that the crude fiber can be graded by machine more accurately than by hand.

VERMONT

The asbestos-bearing rock quarried near Eden, Vt., first is crushed to 5- or 6-inch size in a 28- by 36-inch jaw crusher and then discharged to a 36-inch belt conveyor, which carries the material to a 1,000-ton storage bin. When taken from storage, it is crushed to 3-inch size in a 13- by 30-inch jaw crusher. The product is split to two 4½- by 40-foot rotary driers. A trommel with ¾-inch openings precedes the final crushing unit—a 5½-foot Symons cone. The material that passes through the trommel is combined with the crusher product and conveyed to a second storage bin. The product of the tertiary crusher is treated in a series of Jumbos or fiberizers like those used in Quebec, and the fiber is recovered by screening and air suction. Oversize material passing the hoods is conveyed to a second series of fiberizers. The fiber picked up by air suction in the collectors is further screened, passed through dust removers and graders, dropped into bins according to length, and sacked in 100-pound burlap bags. The principal products manufactured are shingle stock testing 0-2-10-4; millboard, paper, and molded brake-lining fiber testing 0-0-10-6 or 0-0-8-8; and material suitable for roofing, paints, plastics, molded products, and boiler covering testing 0-0-5-11. Equipment was installed in 1936 for salvaging the shorter grades not heretofore recovered, and additional grades now marketed include 0-0-2-14, 0-0-1-15, 0-0-0-16, "no test" and "floats." Special brake-lining fibers are prepared at times. Virtually no crude or spinning fibers are produced. The mill capacity is upward of 100 tons of rock an hour. The demand for the products has more than tripled during the past 4 years. Figure 8 shows the mill as it appeared in 1936.

UNION OF SOUTH AFRICA

CHRYSOTILE

At the New Amianthus mine, Barberton district, Eastern Transvaal, long-fibered chrysotile is hand-sorted and graded by natives as A and B, fiber over 1½ inches long and fiber ¾ to 1½ inches long, respectively; both are designated as crudes. A third grade, E, consisting of hand-selected fiber down to ½ inch long might be classed as crude.

Rock bearing the shorter fibers is crushed, dried, and reduced by rolls. The product of each set of rolls is sieved over 10-mesh screens, the fines going to waste. After passing through four sets of rolls the product is virtually pure fiber except for dust adhering to it. The

Bureau of Mines



FIGURE 8.—Mill of the Vermont Asbestos Co.

STOS

e grades of spinning fibers and a
than shingle stock can be sold

: containing spinning-fiber veins,
re treated in these mills and yield
bbed crude fiber, an anomalous
alian definition of crude, which
he claim is made, however, that
achine more accurately than by

INT

d near Eden, Vt., first is crushed
-inch jaw crusher and then dis-
which carries the material to a
from storage, it is crushed to
crusher. The product is split
friers. A trommel with $\frac{3}{4}$ -
ishing unit—a $5\frac{1}{2}$ -foot Symons
ough the trommel is combined
ed to a second storage bin. The
eated in a series of Jumbos or
, and the fiber is recovered by
material passing the hoods is
ers. The fiber picked up by air
reened, passed through dust re-
bins according to length, and
The principal products manu-
0-2-10-4: millboard, paper,
ng 0-0-10-6 or 0-0-8-8: and
ts, plastics, molded products.
11. Equipment was installed
ades not heretofore recovered,
d ide 0-0-2-14, 0-0-1-15,
s, al brake-lining fibers
crude or spinning fibers are
ed of 100 tons of rock an hour.
e than tripled during the past
appeared in 1936.

AFRICA

E

erton district. Eastern Trans-
orted and graded by natives
nd fiber $\frac{3}{4}$ to $1\frac{1}{8}$ inches long,
udes. A third grade, E, con-
 $\frac{1}{2}$ inch long might be classed

nshed, dried, and reduced by
s sieved over 10-mesh screens,
through four sets of rolls the
for dust adhering to it. The

Bureau of Mines

Bulletin 403




FIGURE 3.—Mill of the Vermont Asbestos Corporation near Eden, Vt.

asbestos is passed through a fiberizer to beat out the dust, then over a dust sieve, and finally over grading sieves. Two principal mill grades are made: F, one-fourth to one-half inch; and G, one-fifth to one-fourth inch. Grade F corresponds to Canadian fiber testing 0-4½-10-1½. The milled fibers are principally shingle and paper stocks. The mill has a capacity of 6,000 tons of rock a month.

At the Munnik-Myburgh mine in the same district four grades of hand-cobbed chrysotile fiber are designated as follows: IXL, over three-fourths inch; XL, one-half to three-fourths inch; X, one-fourth to one-half inch; and XX, one-eighth to one-fourth inch. Hand-cobbed fiber is graded in shaking screens.

Short fiber is treated in a mill with an elaborate flow sheet. Dried and crushed rock is passed through a series of rolls followed by grading screens. All five products obtained are passed through a regrading plant. The grades and their values, as shown on a Canadian standard testing machine, are as follows:

M1-----	2-11-2-1	M4-----	0-0-8½-7½
M2-----	0-3½-6-1½	M5-----	0-0-4½-11½
M3-----	0-3-10-3		

The milled grades constitute about 95½ percent of the total fiber recovery.

CROCIDOLITE

Methods of milling and grading crocidolite vary greatly in different localities. In the Southern Belt of the Cape Provinces the workings are numerous, and many are relatively small. At the smaller workings the fiber is all hand-cobbed. In many places it is hand-sorted into grades according to length, but at the larger workings the hand-cobbed asbestos is graded by machinery. A hand-turned trommel made of perforated sheet iron or covered with wire netting usually is employed. The cobbed fiber from a series of workings may be shipped to a central screening plant where all asbestos one-half inch long or over is recovered, and the minus one-half-inch product is taken to a mill for further crushing and grading. At one mill the rolls travel at slightly different speeds so as to subject the fiber to a tearing action. Double grading trommels sometimes are used. The inner cylinder has one-fourth-inch holes and the outer one-eighth-inch. Material passing the outer cylinder is waste; the oversize from the outer cylinder is designated grade SS and the oversize from the inner cylinder is grade S. The following grades are produced:

- E. Over 2-inch cobbed fiber.
- D. 1¾-inch to 2-inch cobbed fiber.
- C. 1¼-inch to 1¾-inch cobbed fiber.
- B. ½-inch to 1¼-inch cobbed fiber.
- A. ¼-inch to ¾-inch cobbed fiber.
- S. ¼-inch to ½-inch milled fiber.
- SS. Shorts, ¼-inch to ½-inch milled fiber.

In the Northern Belt, material over three-fourths inch long is hand-cobbed. For many years the mill grades were not standardized and the companies had difficulty in marketing their products. Later, a large mill was erected at Kuruman, and the grades manufactured there were established standards that offered such decided

marketing advantages that other producing companies adopted them. In this mill, rock carrying the shorter fibers is passed through crushers, heavy rolls, and disintegrators. The fiber thus separated is classified on grading screens. The following grades are made:

- ES. Over 1¼-inch hand-cobbed fiber.
- No. 1. ¾-inch to 1¼-inch hand-cobbed fiber. Also, 2 grades of plus ¾-inch discolored fiber.
- No. 2. ¾-inch to ¾-inch milled fiber.
- No. 3. ¾-inch, or under, milled fiber.

Of the total fiber recovered, about 13½ percent is hand-cobbed and 86½ percent milled. In both the Northern and Southern Belts about 9 percent of the total fiber is of spinning grade.

Crocidolite of the Pietersburg district, Transvaal, is passed through rolls, fiberizers, and grading trommels. Four grades are produced:

- B. Over 1¼ inch.
- A. ¾ inch to 1¼ inch.
- S. ¾ inch to ¾ inch.
- Paper grade, ½ inch to ¾ inch.

AMOSITE

Because of its unusual length, most of the amosite produced is hand-cobbed. The fiber thus freed from adhering rock is passed through a series of rolls and disintegrators and then graded on shaking screens or trommels, usually into three grades. No. 1 is spinning fiber; no. 2 is used in the manufacture of rope, felts, and similar products; and no. 3 is used for asbestos-cement products such as roofing slates and corrugated sheathing. The fiber is classified both according to length and color. The colors range from whitish gray to yellowish; the gray shades are considered the best. Most of the mills are small.

RHODESIA

The Rhodesian asbestos industry is centered at the large chrysotile mines of Shabani. To avoid damage to the fiber, as much of it as possible is separated from the rock close to the working face; hence, the hand-cobbing and bagging of crude fiber is an important part of quarrying and mining operations. Each cobbing boy recovers about 1,500 pounds of fiber a shift in open quarries and 1,000 pounds a shift in stopes. The cobs are said to average 40 to 50 percent fiber.

The quarry rock first is broken in jaw crushers and rolls, and the products are sorted on picking belts into three groups: (1) Rock containing little or no fiber, (2) rock containing large seams of fiber, and (3) fines and rock carrying small seams of fiber. Product 1 is left on the belt and conveyed to the waste heaps, product 2 is carried on a second belt to the cobbing sheds, and product 3 is conveyed by a third belt to the mill. Hand cobs from both the cobbing sheds and the quarries or stopes are fiberized and graded in one mill, and the fiber-bearing rock is reduced in another.

Some maintain that when fiber-bearing rock is kept wet until it reaches the mill it suffers less damage and is in better condition for milling than when it is handled and rehandled dry. After primary

crushing in the mill, the rock is dried either in rotary driers in which the wood fires are in direct contact or reduced in a series of small grinding pans. Fiber is separated from rock with shakers in the same manner as in Canada. Undersize is carried in the air current is deposited in the pans and passes through the exhaust outlet. The asbestos is taken to a grade further from dust and separated into grades. Grading is done in hexagonal trommels.

Cobbed asbestos is fiberized, cleaned, and the equipment is modified to handle long fiber.

In the Shabani area, the tonnage of fiber is about 3½ percent of the total rock mined. Fiber averages about 4 percent of the total rock, but the rock is leaner than in Quebec, because short-fiber material is discarded as waste in Quebec.

Few data are available on the grading of the proportion of each grade. From the fiber, and most of the remainder is discarded as waste. Paper stock cannot be handled by transportation expense to markets. Grades nos. 1 and 2, are the only grades quoted in reports.

At the King's and Gath's mines asbestos is recovered by hand-cobbing. Marketable fiber obtained is 0.7 to 1.2 percent of the rock.

RUSSIA

In Russia, crude asbestos is sorted and cleaned; but, with increasing demand, a larger proportion of the fiber is recovered from the quarries. For many years the fiber was once recovered from total rock quarried, the elimination of waste raised the grade that mill feed as fiber. During recent years mechanical methods to a large extent have replaced hand-sorting on picking belts.

The preliminary stages of rock crushing in one of the newer mills, the "Gig" 1932. Rock from the quarry is conveyed to a no. 9 Gates gyratory crusher, 6-inch size. It discharges to a picking belt of the primary feed containing too much waste rock. The asbestos is 1½- to 2-inch size in two no. 7 Gates heavy shaking screens. The oversize goes to picking belts, where more barrier

reducing companies adopted them. Short-fiber is passed through rat. The fiber thus separated in the following grades are made:

1. fiber. Also, 2 grades of plus $\frac{3}{4}$ -inch

2. 13½ percent is hand-cobbed and Northern and Southern Belts about mining grade.

3. district, Transvaal, is passed in trommels. Four grades are

TE

Most of the amosite produced is from adhering rock is passed through trommels and then graded on into three grades. No. 1 is for manufacture of rope, felts, and other asbestos-cement products such as roofing. The fiber is classified as follows: The colors range from whitish to black. Most are considered the best. Most

TA

4. cer. 5. ad at the large chrysotile mine to the fiber, as much of it as close to the working face; hence, crude fiber is an important part of the product. Each cobbing boy recovers from open quarries and 1,000 pounds of average 40 to 50 percent fiber. The rock is crushed in jaw crushers and rolls, and the product is divided into three groups: (1) Rock containing large seams of fiber, (2) seams of fiber. Product 1 is waste heaps, product 2 is carried to the mill, and product 3 is conveyed by belt from both the cobbing sheds and graded in one mill, and another.

6. During rock is kept wet until it is dry and is in better condition for rehandled dry. After primary

crushing in the mill, the rock is dried either with steam-heated equipment or in rotary driers in which the products of combustion from wood fires are in direct contact with the rock. The dried rock is reduced in a series of small grinding pans operating like pug mills. Fiber is separated from rock with shaking screens and air suction in the same manner as in Canada. Undesirable brittle fiber is ground in the pans and passes through the screens with the waste rock. Fiber carried in the air current is deposited in a centrifugal settler, and the air passes on through the exhaust fan and dust trap to an outlet. The asbestos is taken to a grading mill, where it is cleaned further from dust and separated into the various market grades. Grading is done in hexagonal trommels.

Cobbed asbestos is fiberized, cleaned, and graded in a mill in which the equipment is modified to handle longer fiber.

In the Shabani area, the tonnage of cobbed fiber ranges from 2½ to 3¼ percent of the total rock mined. Total milled and cobbed fiber averages about 4 percent of the rock quarried. One must not infer that the rock is leaner than in Quebec, where recovery is about 6 percent, because short-fiber material amounting to many thousand tons in Quebec is discarded as waste in Rhodesia.

Few data are available on the grades produced at Shabani or on the proportion of each grade. From 25 to 30 percent is spinning fiber, and most of the remainder is shingle stock. Fiber shorter than paper stock cannot be handled profitably because of the heavy transportation expense to markets. The longest spinning fibers, nos. 1 and 2, are the only grades quoted in the United States market reports.

At the King's and Gaths' mines in the Mashaba district the asbestos is recovered by hand-cobbing only. The tonnage of marketable fiber obtained is 0.7 to 1.2 percent of the rock quarried.

RUSSIA

In Russia, crude asbestos is sorted in the quarry and cobbed as in Canada; but, with increasing mechanization, hand cobbing has declined and a larger proportion of the fiber is prepared mechanically. For many years the fiber was concentrated by hand-sorting in the quarries. Although recoverable fiber averages about 4½ percent of total rock quarried, the elimination of barren rock by hand-sorting so raised the grade that mill feed averaged about 22 percent of the fiber. During recent years mechanical mining equipment has replaced hand methods to a large extent and the fiber is concentrated by hand-sorting on picking belts after the rock is crushed.

The preliminary stages of rock treatment are exemplified best in one of the newer mills, the "Gigant", which began operation in 1932. Rock from the quarry is conveyed to a storage bin and from that to a no. 9 Gates gyratory crusher, which reduces it to about 6-inch size. It discharges to a picking belt, where about 20 percent of the primary feed containing too little fiber to justify milling is thrown out as waste rock. The asbestos-bearing rock is reduced to 1½- to 2-inch size in two no. 7 Gates gyratory crushers discharging to heavy shaking screens. The oversize from the screens passes to picking belts, where more barren rock is eliminated. The good

The most noteworthy feature of the preliminary milling stage is the concentration on picking belts. Because the serpentine tends to break cleanly along the fiber veins, effecting a more or less distinct separation of barren and fiber-bearing rock, conditions particularly favor this method of concentration. Of the original mill feed, consisting of 2,400 tons a day, about 1,440 tons are eliminated as waste, leaving only 960 tons for the later milling processes. Thus, picking belts save operators a great deal of useless milling.

The older mills employed a typically Russian method of milling. The rock was reduced in a series of rolls, and part of the fiber was separated from rock with trommels: the major separation, however, was made on inclined planes provided with suitable gaps. The fiber dropped through the gaps, and the rock fragments jumped over them. Ru Kevser⁴⁸ has described milling processes in some detail.

Little information is available regarding the grades produced and the percentage of each grade. Estimates made in 1931 indicated that fiber one-third inch or more in length constituted about 30 percent of the total output. Since that date, with increasing factory capacity for the manufacture of asbestos products, probably a larger proportion of the shorter grades is produced. Grades entering the German market in 1929 were designated as follows:

- Grade 0. $1\frac{1}{2}$ inch or longer.
Grade 1. $\frac{3}{4}$ inch to $1\frac{1}{2}$ inches.
Grade 2. $\frac{5}{8}$ to $\frac{3}{4}$ inch.
Grade 3. $\frac{3}{8}$ to $\frac{5}{8}$ inch.
Grade 4. $\frac{1}{8}$ to $\frac{3}{8}$ inch.
Grade 5. Less than $\frac{1}{8}$ inch.

⁴⁶ Ru Keyser, Walter A., Asbestos Milling in the Urals: Eng. and Min. Jour., vol. 134, no. 10, October 1933, pp. 415-419.

CYPRUS

Milled asbestos is graded into three fines. The standard grade, designa include about 75 percent of the p is exported. Twelve mills were repoi aggregate capacity is 25,000 tons of fi They are operated during the dry sea

AUSTRAL

ANTHOPHYLL!"

In one mill operated a few years in a gyratory crusher, whence it was carried with air suction for carrying off the fines, followed by reduction in a double roll crusher, the material was crushed in opposite directions. The fiber was carried in a large amount in the suction pipe return, permitting heavy fragments to be returned through the circuit. The fiber was classified by the dried rock in the mill.

In another mill the dried rock is followed by a set of rolls. The material is returned to the fiberizer, and the fiber is conveyed to a bin. The fiber is packed in 100- or 200-pound bales. Other processes are employed in other mills.

at 1-inch size in a jaw
e sl. ng screens in a wet-
rotary driers, which reduce
2 percent. The product is

reliminary milling stage is
cause the serpentine tends
ffecting a more or less dis-
rting rock, conditions par-
tion. Of the original mill
at 1,440 tons are eliminated
er milling processes. Thus,
of useless milling.

ie dried rock is passed over
products are obtained: (1)
size rock conveyed to a set
rolls and are carried to the
a Humboldt disintegrator.
ens, suction pipes, and dis-
recovered. Fiber from the
aking screens for cleaning.
ction fans and classified by
els.

Russian method of milling.
4, and part of the fiber was
major separation, however,
th suitable gaps. The fiber
ck fragments jumped over
g processes in some detail.
2,000,000 tons of rock an-
six grades was nearing com-
sia tal milling capacity
year.

ing the grades produced and
made in 1931 indicated that
nstituted about 30 percent
with increasing factory ca-
products, probably a larger
uced. Grades entering the
as follows:

ie American market include
and shingle stock. Begin-
Crude A.A. was placed on
st grade produced, because

ials: Eng. and Min. Jour., vol. 134,

CYPRUS

In Cyprus, the rock bearing chrysotile asbestos is so decomposed that it breaks down readily into small sizes. Small quantities of hand-cobbed fiber are obtained. Rock reduced in the quarry by hand methods is screened to eliminate barren material over 18 mm (7/10 inch) and fines under 5 mm (1/5 inch); the intermediate product, which averages about one-sixth of the total quarry output, is sent to the mills. The mill feed is crushed and passed over flat shaking screens and the fiber recovered by air suction. It is treated further to remove dust and to classify it according to length. The asbestos recovered constitutes about 3½ percent of the mill feed, and virtually the entire production is short fiber.

Milled asbestos is graded into three classes—standard, shorts, and fines. The standard grade, designated "shingle stock", is said to include about 75 percent of the production. The entire output is exported. Twelve mills were reported in operation in 1935. Their aggregate capacity is 25,000 tons of fiber a year of 6 working months. They are operated during the dry season only.

AUSTRALIA

The Lionel plant in Australia produces four grades, which are numbered in the reverse order from that generally practiced: the smaller numbers are applied to the shorter fibers. No. 1, the lowest grade produced, includes fibers about ¼-inch long; No. 2, about ½-inch; No. 3, about 1 inch; and No. 4, 2 inches and over.

ANTHOPHYLLITE MILLING

Anthophyllite, a variety of amphibole asbestos, has been mined in a small way in the United States for many years. Small mills have been operated in North Carolina, North Carolina, Oregon, Virginia, and Washington. A large percentage of the rock treated is asbestos, fiberization and classification are the principal milling processes. The fiber is much weaker than chrysotile, and therefore severe treatment is to be avoided.

In one mill operated a few years ago the dried rock was broken in a gyratory crusher, whence it went to a hammer mill equipped with air suction for carrying off the loose fiber. This process was followed by reduction in a double-cage mill, the cages running in opposite directions. The fiber was removed by air suction; an enlargement in the suction pipe retarded the velocity of the air current, permitting heavy fragments of unopened fiber to fall. These fragments were returned through a trap to the cage mill in closed circuit. The fiber was classified with vibrating screens.

In another mill the dried rock is first reduced with a jaw crusher, followed by a set of rolls. The crushed asbestos rock is fiberized in a hammer mill equipped with an air separator. Coarse material is returned to the fiberizer, dust is collected in a dust chamber, and the fiber is conveyed to a bin that feeds into a packer. The fiber is packed in 100- or 200-pound sacks. Modifications of these processes are employed in other mills.

134342°—37—8

Anthophyllite mined in Harford County, Md., is peculiarly suited for the manufacture of filters that resist the action of chemicals. It undergoes special treatment, including careful hand selection and washing in acids.

MARKETING

Asbestos is used in so many diverse products that its markets are numerous and widespread. It is used extensively in the United States, England, France, Germany, Belgium, Italy, Netherlands, Spain, Russia, and Japan. The last country has become an important buyer of roofing-grade asbestos. Russia at one time exported almost its entire production, but recently it has become an important manufacturer of asbestos products. Many other countries use smaller quantities. With greater diversity in use, expansion of automobile production, development of new types of building materials, and stronger demands for heat-insulation material, the consumption of asbestos has increased steadily. In recent years, mineral, slag, and glass wools have increased in importance as insulation, and although generally used in fields other than those in which asbestos is employed, they are replacing asbestos to some extent.

Some fiber is handled by agents or jobbers, but most of it is shipped direct from mine to consumer. Usually samples are submitted and fiber length and quality are guaranteed. The larger manufacturers of asbestos products have their own testing machines and can check the producer's classification. The producer should be familiar with manufacturing conditions in order that he may prepare the material to suit each particular use.

Canadian asbestos generally is preferred by users in the United States, although African and Russian fibers may be substituted for many uses. Considerable quantities of the latter fibers are mixed with the Quebec product.

The principal Canadian producers sell direct to consuming industries and also to dealers and agents who distribute it to consumers. Sales frequently are made to consumers only after they have examined and approved samples submitted by dealers or producers. Canadian firms sometimes supply European customers with trial consignments of several tons, particularly if they are introducing new products.

Asbestos is sold in 100-pound or 125-pound bags on a short-ton basis, bags included. Quotations are f. o. b. mines. The weight of a given volume varies with the length of fiber as the longer fibers are bulkier. The volume of a short ton ranges from 60 to 90 cubic feet. A minimum carload of fiber is 20 tons, and of refuse and shorts, 30 tons. Settlements usually are made on the basis of 2 percent discount within 10 days, or 30 days net.

Before the World War, Hamburg was the principal marketing center, but since that time New York has become the most important. The larger producers now have offices in New York, Hamburg, London, and other large cities.

Market requirements are based principally on fiber length, but strength, flexibility, color, chemical composition, and cleanliness may have an important bearing on use. The principal market outlets are indicated in the following brief summary of uses.

The longer and more valuable asbestos upward from about \$70 a ton, are brake linings, textile fabrics, and asbestos grades, ranging from about \$25 to \$70 a ton. Still shorter fibers are used in manufacturing rigid asbestos-cement shingles, lumber and the lowest-grade fibers are employed in cold molded products, and fillers.

European imports from Africa are handled by agents. The importers pay cash or order and sell on credit to manufacturers. They buy directly from producers because they are given 6 months' credit, and importations are on a basis. The Canadian standard test is in effect in Africa.

Several of the largest asbestos organizations; that is, they mine and mill the finished products. At least three large States are of this type, each having mines in the United States. Turner & Newall, the asbestos production of Rhodesia and of South Africa. The company has 20 factories in England. The Russian industry State controls both production of raw and finished products. Because of this control Russian production never enters the open market.

Most of the world supply of raw asbestos is handled, and distribution is conducted in collaboration with buyers, many of whom have expressed concerns. Hence, new producers are not likely to dislocate the market organizations sometimes have difficulty in securing consumers of asbestos ordinarily in advance of actual needs. Central Asbestos Co. makes possible the establishment of Turner & Newall, Ltd., having control of Rhodesia, was able in 1931 to consummate whereby orders were apportioned between the market stabilized on a basis satisfactory to all.

PRICES

Prices of Canadian asbestos have risen in the past 15 years. In 1920, because of a shortage, they rocketed to unprecedented heights, reaching more than \$3,000 a ton. In 1921, the price fell to about one-eighth of the price received in 1920, and by 1925 the highest grade received for 1933 to 1936, the average price received by producers as reported to the Canadian figures, is in enough detail to show values by average price quotations as given in

County, Md., is peculiarly
 hat - ist the action of chemi-
 inc ng careful hand selec-

IG

products that its markets are
 d extensively in the United
 Belgium. Italy, Netherlands.
 ntry has become an important
 a at one time exported almost
 s become an important manu-
 er countries use smaller quan-
 expansion of automobile pro-
 of building materials, and
 material, the consumption of
 cent years, mineral, slag, and
 tance as insulation, and al-
 than those in which asbestos
 to some extent.

jobbers, but most of it is
 . Usually samples are sub-
 are guaranteed. The larger
 e their own testing machines
 tion. The producer should
 tions in order that he may
 ilar use.

red by users in the United
 bers may be substituted for
 the latter fibers are mixed

l di to consuming indus-
 o d. ute it to consumers.
 nly after they have examined
 lers or producers. Canadian
 stomers with trial consign-
 they are introducing new

1-pound bags on a short-ton
 f. o. b. mines. The weight
 of fiber as the longer fibers
 ranges from 60 to 90 cubic
 20 tons, and of refuse and
 made on the basis of 2 per-
 cent.

as the principal marketing
 become the most important.
 New York, Hamburg, Lon-

ipally on fiber length, but
 osition, and cleanliness may
 be principal market outlets
 ary of uses.

The longer and more valuable asbestos crudes and fibers, ranging upward from about \$70 a ton, are used in manufacturing woven brake linings, textile fabrics, and asbestos packing. The next lower grades, ranging from about \$25 to \$70 a ton, are used in manufacturing rigid asbestos-cement shingles, lumber, and corrugated sheathing. Still shorter fibers are used in manufacturing paper and millboard, and the lowest-grade fibers are employed in heat-insulating cements, cold molded products, and fillers.

European imports from Africa are handled principally by dealers or agents. The importers pay cash or open a credit in South Africa and sell on credit to manufacturers. The latter concerns do not buy directly from producers because they are accustomed to buy on 3 to 6 months' credit, and importations are not made easily on such a basis. The Canadian standard testing machine is used to some extent in Africa.

Several of the largest asbestos organizations are of the vertical type; that is, they mine and mill the raw materials and fabricate finished products. At least three large companies in the United States are of this type, each having mines in Canada and factories in the United States. Turner & Newall, Ltd., controls nearly all of the asbestos production of Rhodesia and much of that of the Union of South Africa. The company has 20 large asbestos-products factories in England. The Russian industry is also of this type; the State controls both production of raw materials and manufacture of finished products. Because of this condition, a substantial part of Russian production never enters the open market.

Most of the world supply of raw asbestos is in exceptionally strong hands, and distribution is conducted in close and sympathetic collaboration with buyers, many of whom are also strong, well-organized concerns. Hence, new producers and new sales organizations are not likely to dislocate the market seriously. Furthermore, new organizations sometimes have difficulty in finding a market because consumers of asbestos ordinarily place orders 12 to 18 months in advance of actual needs. Centralized control by large units also makes possible the establishment of marketing agreements. Turner & Newall, Ltd., having control of virtually the entire output of Rhodesia, was able in 1931 to consummate an agreement with Russia whereby orders were apportioned between European consumers and the market stabilized on a basis satisfactory to all concerned.

PRICES

Prices of Canadian asbestos have fluctuated greatly during the past 15 years. In 1920, because of war stimulation, prices skyrocketed to unprecedented heights. Crude No. 1 selling for more than \$3,000 a ton. In 1921, the price dropped to less than half that amount, and by 1925 the highest grades were selling for only about one-eighth of the price received in 1920. The following table shows, except for 1933 to 1936, the average price per short ton, f. o. b. mine, received by producers as reported to the Quebec Bureau of Mines. Since 1932 the Canadian figures, unfortunately, are not presented in enough detail to show values by grades, and for these later years average price quotations as given in Mineral and Metal Markets are

used. The latter figures, however, are generally very much higher than figures for actual sales as given by the Quebec Bureau of Mines.

Average value per short ton of asbestos sold in Canada, 1921-36¹

Year	Crude No. 1	Crude No. 2	Spinning fibers	Shingle stock	Millboard and paper stock
1921	\$1,281.32	\$446.91	\$263.09	\$101.75	\$31.19
1922	648.65	255.32	217.71	81.00	21.65
1923	472.60	237.29	123.37	57.05	22.01
1924	265.97	215.27	110.81	45.12	19.64
1925	364.96	206.12	106.43	50.78	31.03
1926	371.51	229.62	130.22	55.62	33.88
1927	423.65	249.39	129.32	64.80	37.82
1928	534.87	296.65	148.71	73.80	38.73
1929	557.34	331.62	177.30	78.26	38.46
1930	480.44	285.44	141.52	70.64	35.33
1931	431.45	216.35	107.22	55.72	34.66
1932	396.94	192.43	91.36	49.64	30.84
1933	457.50	206.25	100.20	52.71	32.71
1934	450.00	212.50	112.50	55.00	35.00
1935	500.00	212.50	112.50	60.00	35.00
1936	500.00	212.50	115.75	60.00	35.00

¹ Figures for 1921-32 from Quebec Bureau of Mines; for 1933-36, average of quotations, Mineral and Metal Markets.

² Paper and others.

³ Paper only.

Price trends are shown graphically in figures 9 and 10.




FIGURE 9.—Average value per short ton of Canadian crudes and spinning fibers, 1921-35. Data for 1921-32 from value of sales compiled by the Bureau of Mines, Province of Quebec. Data for 1933 to 1935 from price quotations in Mineral and Metal Markets.




FIGURE 10.—Average value per short ton of Canadian shingle stock and millboard and paper stock, 1921-35. Data for 1921-32 from value of sales compiled by the Bureau of Mines, Province of Quebec. Data for 1933 to 1935 from price quotations in Mineral and Metal Markets.

Average quotations for Vermont asbestos, f. o. b. mines, are shown in the following table.

Prices of Vermont asbestos

Year	Shingle stock	Paper stock	Crude stock
1931	\$50.71	\$35.00	\$25.00
1932	42.92	32.92	20.00
1933	42.92	32.71	20.25

The following table shows prices of asbestos per short ton, average quotations in Mineral and Metal Markets. All prices are f. o. b. mines.


generally very much higher
than the Quebec Bureau of Mines.

as sold in Canada, 1921-36¹

Crude No. 2	Spinning fibers	Shingle stock	Millboard and paper stock
\$446.91	\$263.09	\$101.75	\$31.19
265.32	207.71	31.00	21.65
237.20	123.37	37.03	22.01
315.37	110.81	45.12	19.34
206.12	104.43	50.73	31.03
229.62	130.22	58.62	33.88
249.59	129.32	64.80	37.32
296.65	148.71	73.80	38.73
331.82	177.30	75.26	38.56
235.44	141.52	70.64	35.33
216.35	107.22	58.72	34.66
192.43	91.36	49.64	30.84
206.25	100.20	52.71	32.71
212.50	112.50	55.00	35.00
212.50	112.50	60.00	35.00
212.50	118.75	60.00	35.00

-36, average of quotations, Mineral and Metal

figures 9 and 10.



1929 1930 1931 1932 1933 1934 1935
crudes and spinning fibers, 1921-35,
by the Bureau of Mines, Province of
Quebec, as sold in Canada, 1921-36¹




FIGURE 10.—Average value per short ton of Canadian shingle, millboard, and paper stocks, 1921-35. Data for 1921-32 from value of sales compiled by the Bureau of Mines, Province of Quebec. Data for 1933 to 1935 from price quotations in Mineral and Metal Markets. Figures for millboard and paper prior to 1925 represent paper and others.

Average quotations for Vermont asbestos, as given in Mineral and Metal Markets, are shown in the following table. Prices are per short ton, f. o. b. mines.

Prices of Vermont asbestos, 1931-36

Year	Shingle stock	Paper stock	Cement stock	Year	Shingle stock	Paper stock	Cement stock
1931.....	\$50.71	\$35.00	\$20.00	1934.....	\$45.00	\$35.00	\$23.00
1932.....	42.92	32.92	20.00	1935.....	46.23	35.00	23.00
1933.....	42.92	32.71	20.25	1936.....	47.30	35.00	23.00

The following table shows prices of Rhodesian and Russian asbestos per short ton, average quotations for the year, as shown in Mineral and Metal Markets. All prices are c. i. f. New York.

Average price quotations per ton of foreign asbestos

Year	Rhodesian		Russian			
	No. 1	No. 2	Crude No. 1	Crude No. 2	Crude No. 3	Shingle stock
1924	\$361.11	\$227.77				
1925	313.75	193.75				
1926	292.50	222.91				
1927	325.00	250.00				
1928	402.08	322.91				
1929	450.00	350.00				
1930	450.00	350.00				
1931	353.33	300.00	\$210.00	\$160.00	\$50.00	
1932	304.17	204.17	227.50	184.17	113.54	
1933	187.50	131.25	213.54	163.54	125.00	
1934	186.67	136.67	212.80	162.50	125.00	\$35.00
1935	210.00	160.00		182.50	125.75	55.00
1936	210.00	172.50	192.50	167.50	130.00	52.50
1936	210.00	165.00	220.00	186.00	130.00	55.00

Beginning July 1935 a new grade of Russian asbestos designated Crude AA was quoted regularly at \$450 a ton until February 1936, and at \$475 a ton for the remainder of the year.

MANUFACTURE OF ASBESTOS PRODUCTS

A WIDESPREAD AND DIVERSIFIED INDUSTRY

The fire resistance of asbestos combined with its fibrous character adapt it admirably for the manufacture of flexible, heat-insulating products, packings, and gaskets for use in places where fabrics of animal or vegetable origin would be less enduring and would create fire hazards. The addition of asbestos to various fireproof building materials gives them strength and flexibility; and, furthermore, manufacturers of many miscellaneous products find that asbestos, when used as an ingredient, imparts superior qualities. Thus many large industries have grown up in recent years for the manufacture of innumerable products of which asbestos is a major or an essential constituent.

The Bureau of Mines deals primarily with economic, technologic, and safety problems pertaining to raw materials of mineral origin; but, as the manufacture of asbestos products is very important in the United States, while the production of raw asbestos is a small industry, more space than usual is given in this report to manufacturing processes and their resulting products.

ASBESTOS-PRODUCTS PLANTS

UNITED STATES

The United States leads all countries in the manufacture of products of which asbestos is an essential or important constituent. The Bureau of the Census listed 238 establishments in the United States that made asbestos products in 1929, 208 in 1931, 187 in 1933, and 196 in 1935. The industry is centered chiefly in the North Atlantic States, notably in the Philadelphia (Pa.) and eastern New Jersey areas. Important factories are operated, also, in Cincinnati (Ohio), Chicago (Ill.), St. Louis (Mo.), and other midwestern cities; in the

Southern States; and on the Pacific coast; the domestic market: less than 5 percent, 1

The following table shows the quantities of asbestos products manufactured in the United States as given in the Biennial Census of Manufactures. As "other products" are very numerous; companies sell more than 200 different articles, under the last four items in the table covering including wood, steel, copper, cork, and rubbers, which may form a very small proportion

in of foreign asbestos

Russian			
Crude No. 1	Crude No. 2	Crude No. 3	Shingle stock
\$210.00	\$160.00	\$50.00	
227.50	187.17	119.54	
213.54	163.54	125.00	
212.60	162.50	125.00	\$55.00
	162.50	128.75	55.00
192.50	167.50	130.00	52.50
220.00	188.00	130.00	55.00

STOS PRODUCTS

IFIED INDUSTRY

ed with its fibrous character
e of flexible, heat-insulating
: in places where fabrics of
: enduring and would create
to various fireproof building
lity; and, furthermore, man-
acts find that asbestos, when
qualities. Thus many large
s for the manufacture of in-
is : for or an essential.

with economic, technologic, materials of mineral origin; asbestos is a small industry report to manufacturing

PLANTS

12

in the manufacture of prod-
important constituent. The
ments in the United States
8 in 1931, 187 in 1933, and
chiefly in the North Atlantic
-), and eastern New Jersey
also, in Cincinnati (Ohio),
er midwestern cities; in the

Southern States; and on the Pacific coast. Most of the output enters the domestic market; less than 5 percent, in value, is exported.

The following table shows the quantity and value of the principal asbestos products manufactured in the United States in recent years, as given in the Biennial Census of Manufactures. The articles listed as "other products" are very numerous; some manufacturing companies sell more than 200 different articles. Many products included under the last four items in the table consist of various materials, including wood, steel, copper, cork, and rubber, in addition to asbestos, which may form a very small proportion of the total.

Asbestos products manufactured in the United States, 1929-35
[Data from Bureau of Census]

	1920	1931	1933	1935
	Quantity	Value	Quantity	Value
Asbestos textiles and textile products:				
Blank linings.....	110, 246, 179	\$16, 442, 448	56, 079, 620	\$4, 307, 492
Clutch linings.....	38, 690, 113	3, 010, 727	21, 310, 723	1, 790, 701
Do.....	15, 378, 407	6, 173, 569	9, 512, 686	1, 691, 531
Do.....	6, 666, 599	2, 767, 019	1, 113, 970	1, 531, 962
Do.....	2, 226, 852	1, 292, 251	2, 530, 519	1, 098, 751
Tape, beltings, and tubular lagging.....	3, 815, 710	3, 815, 710	1, 735, 925	1, 584, 918
Packing.....	1, 281, 281	6, 741, 122	4, 071, 327	2, 011, 871
Caskets.....	3, 012, 385	1, 088, 005	6, 741, 122	1, 875, 707
Other textiles.....			630, 770	1, 800, 352
Total textiles and products.....		35, 161, 991	18, 510, 219	12, 623, 290
Asbestos building materials:				
Blinches.....	891, 661	5, 277, 308	553, 097	1, 820, 279
Lumber.....	35, 395, 191	3, 091, 420	10, 907, 820	1, 012, 313
Other building materials.....		2, 691, 851	816, 313	620, 510
Total building materials.....		10, 439, 018	5, 617, 532	3, 459, 188
Other asbestos products:				
Blank linings, molded.....	22, 440, 872	2, 531, 616	21, 180, 181	2, 823, 738
Pipe linings (kind unrecycled).....	67, 331, 323	4, 057, 665	30, 083, 820	3, 116, 015
Pipe and boiler covering, air-seal.....	27, 690, 501	2, 516, 429	8, 717, 813	1, 137, 283
Pipe and boiler covering other than air-seal.....	23, 639, 209	3, 015, 253	10, 323, 383	1, 435, 211
Milled blocks, 85 percent magnesia.....	7, 769, 221	2, 315, 375	11, 740, 035	1, 471, 170
Millboard.....	68, 474, 360	917, 810	30, 100, 027	650, 301
Table mats and protectors.....	1, 615, 631	760, 222	33, 856, 201	38, 186, 019
Other products.....		5, 200, 078		700, 082
Gaskets other than asbestos textile.....		15, 741, 911		371, 913
Metallic and semimetallic packings.....		3, 066, 578		4, 590, 772
Brooms and other packing other than asbestos textile.....		7, 992, 768		12, 351, 315
Total other products.....		51, 072, 457		2, 767, 850
Grand total.....		101, 206, 006		2, 406, 069

¹ Asbestos paper and miscellaneous products; includes millboard in 1931 and 1934.
² Includes some gaskets that contain little or no asbestos.
³ Excludes fastener machines.

CANAD.

For many years Canada exported raw asbestos, but lately it has diversified its asbestos products industry. In 1934, 11 plants were manufacturing asbestos products. Four of them, in the centre of Quebec, accounted for 46 percent of the production. The other five were also six plants in Ontario and Quebec. The principal products are brake-band linings, pipe coverings, paper, and roofing.

The following table, compiled from the Dominion Bureau of Statistics, shows the production of paper in the recent years:

Asbestos products plants

Year	No. of
1925	
1926	
1927	
1928	
1929	
1930	
1931	
1932	
1933	
1934	
1935	

GREAT B

The manufacture of asbestos products in Great Britain. In 1929, 57 factories and the estimated value of the raw asbestos used, 44 percent came from the Union of South Africa, small amounts from Cyprus. The range of products is much the same. About one-fourth of the total output

GERM

In 1929, 32 factories in Germany which were devoted chiefly to spinning factories made corrugated sheeting bestos-cement products. The raw Russia, although substantial quantities and Africa. As asbestos has no quantity used annually in Germany except for fluctuation in stocks, it have increased from 7,580 tons in tons in 1934, and about 22,000 tons

ACS

In Austria, the original home of the factories were operated in 1929. The principal product of four factories.

- 1 Asbestos paper and infusible products; includes millboard in 1931 and 1933.
- 2 Includes some gaskets that contain little or no asbestos.
- 3 Excludes heat hose materials.

CANADA

The following table, compiled from data furnished by the Dominion Bureau of Statistics, shows the extent of the industry during recent years:

The following table, compiled from data furnished by the Dominion Bureau of Statistics, shows the extent of the industry during recent years:

Asbestos products plants in Canada, 1925-35

GREAT BRITAIN

GERMANY

AUSTRIA

FRANCE

In 1929, four factories in France produced asbestos-cement roofing tiles and similar products; nine plants were devoted chiefly to the manufacture of spun, woven, and braided products; four factories made high-pressure packings; and two factories made asbestos brake linings.

BELGIUM

Belgium has imported from Canada, Africa, and Russia between 1,200 and 3,000 metric tons of asbestos annually during recent years. Nine factories were in operation in 1929, but only five were reported in 1936. These firms consume over 90 percent of the imports of raw fiber. The principal products made are asbestos-cement shingles, tiles and pipes, and asbestos insulation. The output of millboard and woven products is small. No brake linings are made in the country. A large percentage of the imports of fiber are of shingle grade.

ITALY

In Italy, three large factories and numerous small ones manufactured principally asbestos-cement tiles and sheets in 1929. Thread, cards, cartons, textiles, and filter paper also are made in considerable quantities. Two plants manufacture "Eternit" products. The annual value of Italian asbestos products was estimated in 1932 at 25 million lire (about \$2,061,775 on the basis of the 1935 rate of exchange).

CZECHOSLOVAKIA

In 1929, three factories in Czechoslovakia were devoted to asbestos spinning, weaving, and braiding; one made asbestos board; four, high-pressure packings; and several, asbestos-cement tile. Eight companies manufactured asbestos-cement roofing tiles in 1936. In that year the three largest companies concluded a cartel agreement whereby prices for domestic sales were increased substantially. The claim is made that the new price of 15 crowns (about 62 cents) a square meter gives the companies a small margin of profit.

NETHERLANDS

A plant for making "Eternit" cement-asbestos gas and water conduits was built at Goor, Netherlands, in 1936.

FINLAND

One factory established at Helsingfors, Finland, in 1923 manufactures "Eternit" products, packings, insulating materials, and fabrics. Both domestic and foreign asbestos are used.

SCANDINAVIA

Norway imports 100 to 200 metric tons a year of asbestos from Finland. It is used chiefly for heat insulation. Two factories manufacture asbestos packings, but their requirements of raw fiber are small. Similarly, few asbestos products are manufactured in

Sweden. The country imports about annually, chiefly from Finland and Russia, heat insulation. A small asbestos-pipe mark. In 1934, 1,040 metric tons of imported.

AUSTRALIA

There are quite a number of factories manufacturing of asbestos products. Sydney, Brisbane, Melbourne, and Perth are asbestos-cement sheet siding a pipe. Sheet-metal roofing is used as a substantial substitution of asbestos-cement.

Asbestos production in Australia is small. Asbestos required is imported. The United States and Canada are the chief sources of supply. Figures show annual imports of about \$60,000.

RUSSIA

According to recent data, about 10 percent of asbestos is used in domestic factories of asbestos products. Russia, therefore, is a factor in this industry. One very important asbestos-cement product, fabrics. As a result of research conducted at the Institute, several new products and processes have been developed. The Yaroslavl Rubber Factory produces molded asbestos brake

OTHER COUNTRIES

There are several asbestos factories in Switzerland. Several establishments in the United States manufacture asbestos textiles, insulating materials, and asbestos-cement roofing is becoming an important industry in that country, because recently there has been a demand for shingle-stock fiber.

FABRICATION OF ASBESTOS

The manufacture of asbestos involves many complex processes. Only a few of the operations are presented herein because of the large amount of raw materials.

FIBER

Crude fiber delivered to the manufacturer is first chaser milled, fiberized, and then subjected to air separation. It is then subjected to successive stages of manufacture in which the fibers are prepared by producers need not be detailed.

The processes involved in the manufacture of asbestos are in general those employed in spinning.

rodi asbestos-cement roofing
its devoted chiefly to the
aided products; four factories
o factories made asbestos brake

la, Africa, and Russia between
s annually during recent years.
29, but only five were reported
percent of the imports of raw
are asbestos-cement shingles,
on. The output of millboard
rake linings are made in the
imports of fiber are of shingle

numerous small ones manu-
les and sheets in 1929. Thread,
er also are made in considera-
ure "Eternit" products. The
cts was estimated in 1932 at 25
e basis of the 1935 rate of

AKIA

vakia were devoted to asbestos
e made asbestos board; four,
asbestos-cement tile. Eight
ent roofing tiles in 1936. In
con ed a cartel agreement
increased substantially. The
15 crowns (about 62 cents) a
nall margin of profit.

DS

it-asbestos gas and water con-
in 1936.

fors. Finland, in 1923 manu-
rs, insulating materials, and
bestos are used.

IA

tons a year of asbestos from
sulation. Two factories man-
requirements of raw fiber are
ducts are manufactured in

Sweden. The country imports about 1,000 metric tons of fiber an-
nually, chiefly from Finland and Russia. It is used principally for
heat insulation. A small asbestos-products industry exists in Den-
mark. In 1934, 1,040 metric tons of unmanufactured asbestos were
imported.

AUSTRALIA

There are quite a number of factories in Australia devoted to the
manufacture of asbestos products. They are located chiefly in
Sydney, Brisbane, Melbourne, and Perth, and their principal products
are asbestos-cement sheet siding and roofing and asbestos-cement
pipe. Sheet-metal roofing is used widely in Australia, and a sub-
stantial substitution of asbestos-cement sheets is anticipated.

Asbestos production in Australia is small, hence most of the raw
asbestos required is imported. The Union of South Africa and
Canada are the chief sources of supply. The most recent available
figures show annual imports of about 3,500 tons valued at about
£60,000.

RUSSIA

According to recent data, about two-thirds of the Russian output
of asbestos is used in domestic factories devoted to the manufacture
of asbestos products. Russia, therefore, is becoming an important
factor in this industry. One very large factory at Leningrad makes
asbestos-cement products, fabrics, paper, and insulating materials.
As a result of research conducted at Leningrad by the Asbestos Insti-
tute, several new products and processes of manufacture have been
developed. The Yaroslavl Rubber Combine has started mass pro-
duction of molded asbestos brake linings.

OTHER COUNTRIES

There are several asbestos factories in Poland, Hungary, Spain,
and Switzerland. Several establishments in Japan make asbestos
textiles, insulating materials, and asbestos-cement products. Evi-
dently roofing is becoming an important branch of the industry in
that country, because recently there has been an unusually large
demand for shingle-stock fiber.

FABRICATION OF ASBESTOS PRODUCTS

The manufacture of asbestos products involves many diversified
and complex processes. Only a brief summary of the principal op-
erations is presented herein because this report pertains primarily to
raw materials.

FABRICS

Crude fiber delivered to the manufacturing plant usually is crushed
in a chaser mill, fiberized, and freed from rock impurities by screen-
ing and air separation. It is then ready to enter the various suc-
cessive stages of manufacture into fabrics. Milled spinning fibers
prepared by producers need not undergo this preliminary treatment.

The processes involved in the manufacture of fabrics follow in
general those employed in spinning and weaving cotton, wool, or

silk. Asbestos fibers differ from those of organic origin mainly in the nature of the surface. Wool fibers are covered with scaly bands known as imbrications, and cotton fibers are rough, twisted, and irregular. On the other hand, asbestos fibers have no nodules, twists, or irregularities on the surface that will enable one fiber to cling to another. They are more nearly akin to silk but are smoother and more rodlike. This smooth, slippery condition creates such difficulties in spinning that the manufacture of a 100-percent asbestos yarn is slow and costly; therefore a percentage of some other fiber, usually cotton, is added to act as a vehicle to carry the asbestos through the manufacturing processes. The proportion of cotton added varies with the character of the asbestos and with the nature of the finished product. It rarely exceeds 20 percent, and with long-fibered Canadian asbestos may be as low as 2 percent.

Asbestos used in the manufacture of fabrics may consist of a mixture of varieties from different localities. The proportions depend upon the cost and quality of raw materials and the requirements of the finished product. Most manufacturers in America prefer Canadian fiber as a base, although many products now are made of African fiber alone. The blended fibers, together with the necessary addition of cotton, are mixed thoroughly with revolving beaters. Some manufacturers, however, introduce the cotton at a later stage.

Carding is the next step in the process of manufacture. Carding rolls are covered with leather and fitted with sharp steel bristles. They comb the fibers parallel and remove short fibers, bits of rock, and dust. After passing a succession of carding rolls the fiber emerges as a loose blanket, which may be turned 90° and passed through another carding machine. The blanket then is separated into rovings, which are gathered in a roll on a Jack spool and spun into yarn, as in ordinary textile mills.

Yarns are made in various sizes: a "5-cut" yarn measures about 500 yards to the pound and a "30-cut" yarn about 3,000 yards. When twisted exceptionally hard, it is known as asbestos thread and used in sewing gas mantles, asbestos theater curtains, and asbestos clothing. The spindles of single-ply yarn are transferred to twisting machines and twisted into two- or three-ply yarn, which is wound on spools. Asbestos cord and rope are made by twisting together a greater number of strands.

Where yarn is to be used for brake bands or packings, it usually is reinforced with fine copper, brass, or lead wire. Thus, for brake bands three strands of single-ply yarn and two strands of brass wire of gage nos. 0.006, 0.007, or 0.008 may be twisted together. For packings, a single lead wire or one to three strands of brass wire are twisted with two or three strands of asbestos yarn. Products prepared in this way are known as "metallic yarns."

Yarns are woven into fabrics by well-known processes employed in cotton- or woolen-textile mills. Asbestos cloth is used for theater curtains, fireproof clothing, and many other textile products. Single-ply asbestos yarn also is braided into tape. For electrical insulation it should contain not more than 7 percent carbon and not more than 14 percent cotton. Metallic yarn containing about 16 percent cotton is woven into strips for brake-band linings. Standard widths range from 1 to 6 inches and standard thicknesses from one-eighth to

three-eighths inch. They are processed with various ingredients. The manufacture of asbestos is the most important branch of the industry, as indicated in the table on page 84. Strips of asbestos in 1929 totaled more than 20,000 miles.

Asbestos packings are made in various forms, twisted or braided into valve-stem packings, or compressed into rings, or asbestos cement in other desired forms. They may be mixed with rubber compounds, oil, or flake graphite for some packings.

SHINGLES AND LUMBER

Roofing shingles are made of a mixture of portland cement and 15 percent asbestos. They are manufactured by the so-called "dry" process. The mixture is mixed dry in a pug mill and then pressed into sheets with paddles. The mixture is spread on a belt and sprayed with water at 180° F. to the required thickness, and a rotary cutter cuts the shingles. The shingles are piled in stacks and squeezed in a hydraulic press at 1,000 pounds per square inch, after which they are cut and nailed.

Shingles are also made by the wet process, as in the following paragraph:

Asbestos lumber and shingles may be made by the "Eternit" process first used in Austria. The asbestos and coloring matter are mixed with water and pressed thoroughly with a beater, and the sheets are built up in successive layers. Portland cement thus mixed with asbestos does not lose its capability of setting after the water is removed. For shingles, the sheets are made. Lumber consists of thicker sheets made by crimping that sheet. The sheets are used also for making cement-asbestos tube-shaped articles in more complex equipment.

Asbestos lumber is also made by the dry process, but more time is needed for drying than is required for the thin sheets used for

PAPER AND MATTING

Asbestos of paper-stock grade is mixed with water to make a thin slurry, which is spread on drums covered with slats. Starch, derived partly from the overflow stage, is added to the slurry. The machine is similar to that used in the paper industry or wood pulp. All particles of stone are retained in a sand-catching and knotting machine between rollers to remove

of organic origin mainly in are mixed with scaly bands as rough, twisted, and fibers have no nodules, twists, will enable one fiber to cling to silk but are smoother and condition creates such difficulty of a 100-percent asbestos percentage of some other fiber, vehicle to carry the asbestos.

The proportion of cotton asbestos and with the nature is 20 percent, and with long-w as 2 percent. fabrics may consist of a mixture. The proportions depend on the requirements of materials and the requirements of products now are made of together with the necessary machinery with revolving beaters. the cotton at a later stage. of manufacture. Carding with sharp steel bristles, ve short fibers, bits of rock, of carding rolls the fiber be turned 90° and passed blanket then is separated on a Jack spool and spun

5-cut" yarn measures about in about 3,000 yards. When as a 20s thread and used curtains, and asbestos cloth-are transferred to twisting ply yarn, which is wound made by twisting together

ends or packings, it usually lead wire. Thus, for brake d two strands of brass wire twisted together. For pack- strands of brass wire are asbestos yarn. Products pre- yarns.

known processes employed as cloth is used for theater or textile products. Single. For electrical insulation carbon and not more than ng about 16 percent cotton s. Standard widths range nesses from one-eighth to

three-eighths inch. They are processed with rubber and other ingredients. The manufacture of brake-band linings for automobiles is the most important branch of the asbestos-products industry. As indicated in the table on page 84, strips woven for this purpose in 1929 totaled more than 20,000 miles.

Asbestos packings are made in various ways. The yarn may be twisted or braided into valve-stem packings, the braided forms may be compressed into rings, or asbestos cloth may be cut into gaskets or other desired forms. They may be coated or impregnated with rubber compounds, oil, or flake graphite. Metallic yarn is used in some packings.

SHINGLES AND LUMBER

Roofing shingles are made of a mixture of approximately 75 percent portland cement and 15 percent shingle-grade asbestos. When manufactured by the so-called "dry process," the cement, asbestos, and coloring agent are mixed dry in a cylindrical mixer provided with paddles. The mixture is spread evenly on an 18-inch conveyor belt and sprayed with water at 180° F. Rollers compress it to the required thickness, and a rotary cutter separates it into individual shingles. The shingles are piled in stacks separated by steel pallets and squeezed in a hydraulic press at a pressure of 20,000 pounds per square inch, after which they are cured, trimmed, and punched for nailing.

Shingles are also made by the wet process described in the following paragraph:

Asbestos lumber and shingles may be made by the laminated or "Eternit" process first used in Austria. Cement, shingle-grade fiber, and coloring matter are mixed with a large quantity of water, agitated thoroughly with a beater, and pumped to a paper machine that builds up the sheets in successive laminations to the desired thickness. Portland cement thus mixed with a large quantity of water does not lose its capability of setting at a later stage when most of the water is removed. For shingle manufacture, thin sheets are made. Lumber consists of thicker and larger sheets. Corrugated sheets are made by crimping flat sheets. The laminated process is used also for making cement-asbestos pipes, but the building up of tube-shaped articles is more complicated and requires special equipment.

Asbestos lumber is also made by the dry process described previously, but more time is needed for compression and curing than is required for the thin sheets used for roofing.

PAPER AND MILLBOARD

Asbestos of paper-stock grade is mixed with a large amount of water to make a thin slurry, which is agitated thoroughly by 5-foot drums covered with slats. Starch, flour, or size and sodium silicate, derived partly from the overflow squeezed out of the paper at a later stage, are added to the slurry. This is then conveyed to a paper machine similar to that used in the manufacture of paper from rags or wood pulp. All particles of stone or other impurities are eliminated in a sand-catching and knot-removing machine. The sheets of paper pass between rollers to remove most of the water and are

dried in hot cylinders and wound in rolls. If a two-ply paper is desired, one side of a sheet is coated lightly with sodium silicate, and the two sheets are run together over several hot rolls. Crimped paper is made by passing it over corrugated rolls. In the manufacture of air-cell pipe covering, the tips of the corrugations are coated with sodium silicate, and a flat sheet is added. When this process is repeated, a two-ply, three-ply, or thicker air-cell covering may be made.

Millboard is generally classed with paper because it is manufactured by the same general process. It is simply a thick paper: it bears the same relation to asbestos paper that cardboard bears to wrapping paper. The board usually is built up on rectangular screens rather than on drums.

The manufacture of millboard gaskets is important. The high speeds, temperatures, and pressures attained in modern automobiles and other machinery demand very exacting qualities in the millboard used. It must have uniform density and high strength, and variations in thickness in any part of the sheet must not exceed two one-thousandths of an inch.

COMPOUNDED PACKINGS

Certain types of asbestos packing are made by mixing clean, well-opened fiber with fillers, such as clay, barite, magnesia, iron oxide, graphite, or cellulose, and binding materials, such as gums, resins, lac, or rubber dissolved in benzine. High-quality sheets contain about two parts asbestos to one part filler. The mixture is molded into sheets of the desired thickness. Several sheets may be glued together and reinforced with copper or lead foil.

MAGNESIA INSULATION

For the so-called "85-percent-magnesia" pipe covering a good grade of asbestos—chrysotile, blue asbestos, or amosite—ranging from 0-6-6-4 to 0-5-8-3 is used. Approximately 15 percent of fiber is mixed with 85 percent of basic magnesium carbonate derived from dolomite, magnesite, or bitterns at salt works. These ingredients are mixed thoroughly with water and filter-pressed into cakes, which are dried for 3 days in steam-heated kilns. The blocks are then trimmed, cut in lengths, and bored to form hollow cylinders; these are sawed lengthwise, and the half-cylinder sections are rolled in pairs in glued canvas jackets. For ordinary use the walls have a maximum thickness of about $2\frac{1}{2}$ inches.

ASBESTOS CEMENT

A covering used widely for boiler insulation consists of short-fiber asbestos and a cementing material such as plastic clay. The ingredients are mixed with water to form a paste, which is applied with a trowel.

COLD-MOLDED ARTICLES

Increasing quantities of short-fiber asbestos are used in the manufacture of electrical fittings and household appliances. Mixtures of asbestos, gilsonite, cement, and oil are ground together and com-

SELECTED BIBLIO

pressed in molds: these are baked in oil. Gilsonite imparts a brown color: omitted.

NONCORROSIVE F

Amphibole asbestos, which in general is washed, thoroughly prepared for use in Gooch crucibles or for employ strong acids and alkalis.

SELECTED BIBLIO

- ALLEN, M. A., AND BUTLER, G. M. Asbestos. 1 ASBESTOS. A magazine devoted to the asbestos. July 1919, 16th Floor, Enquirer Bldg., P
BECKER & HAAG. Asbestos. Book published 88 pp.
BERLINBAUT, L. Asbestos Mining Reviving Press, vol. 121, 1926, pp. 164-167.
BOWLES, OLIVER. Asbestos. Chapter in I Edited by J. E. Spurr, 1920, pp. 388-401.
BUREAU OF FOREIGN AND DOMESTIC COMMERCE Trade Inf. Bull. 442, 1926, 22 pp.
— Asbestos; World Production and 10 pp.
CIRKEL, FRITZ. Chrysotile-Asbestos. Its Occ Uses. 2d ed. Canada Dept. Mines. Min
COOKE, H. C. Asbestos Deposits of Their vol. 31, no. 4, 1936, pp. 355-376.
— The Composition of Asbestos and Quebec. Trans. Roy. Soc. Canada, third
COULSON, A. L. Asbestos in the Corded D With Notes on Its Occurrence in Other India, vol. 64, pt. 2, 1934, 266 pp.
DILLER, J. S. The Types and Modes of Occur. Jour. Canadian Min. Ind.
DRESSER, JOHN A. Asbestos in South Africa, vol. 50, 1915.
— The Asbestos Deposits of the Eastern Geol., vol. 4, no. 2, 1909, pp. 132-140.
FISHER, NORMAN R. The Quebec Asbestos vol. 44, 1923, pp. 649-655.
GAMBLE, WILLIAM B. Asbestos. A list of York Public Library, 1929, 72 pp.
HALL, A. L. Asbestos in the Union of South Geol. Survey Mem. 12, 2d ed., 1930, 32-
— On the Asbestos Occurrences Near District. Trans. Geol. Soc. South Africa
HOPEKINS, OLIVER B. Asbestos Deposits of Eng., vol. 50, 1915, pp. 964-973.
— Asbestos, Talc, and Soapstone Deposits Georgia Bull. 29, 1914, 319 pp.
IMPERIAL MINERAL RESOURCES BUREAU. London, 1921, 34 pp.
— The Mineral Industry of the British Asbestos, Statistics (1919-21). London, 1925, 23 pp.
IMPERIAL INSTITUTE. The Mineral Industry of Foreign Countries. Statistical Summary,

ills. If a two-ply paper is de-
tly with sodium silicate, and
se 1 hot rolls. Crimped
rugated rolls. In the manu-
tips of the corrugations are
t sheet is added. When this
y, or thicker air-cell covering

paper because it is manufac-
is simply a thick paper; it
per that cardboard bears to
is built up on rectangular

ets is important. The high
ained in modern automobiles
ing qualities in the millboard
nd high strength, and varia-
set must not exceed two one-

KINGS

made by mixing clean, well-
barite, magnesia, iron oxide,
erials, such as gums, resins,
High-quality sheets contain
ler. The mixture is molded
veral sheets may be glued
lead foil.

TION

"pine covering a good grade
orosite—ranging from
ate 1, 15 percent of fiber is
um carbonate derived from
orks. These ingredients are
ressed into cakes, which are
The blocks are then trimmed,
7 cylinders; these are sawed
are rolled in pairs in glued
alls have a maximum thick-

IT

lation consists of short-fiber
; plastic clay. The ingredi-
te, which is applied with a

LES

estos are used in the manu-
ld appliances. Mixtures of
ground together and com-

SELECTED BIBLIOGRAPHY

pressed in molds; these are baked in ovens, polished, and lacquered.
Gilonite imparts a brown color; if gray is desired the gilonite is
omitted.

NONCORROSIVE FILTERS

Amphibole asbestos, which in general is more resistant to chemicals
than chrysotile, is washed, thoroughly fiberized, and otherwise pre-
pared for use in Gooch crucibles or for other filtering processes that
employ strong acids and alkalis.

SELECTED BIBLIOGRAPHY

- ALLEN, M. A., AND BUTLER, G. M. Asbestos. Univ. Arizona Bull. 113, 1921, 31 pp.
- ASBESTOS. A magazine devoted to the asbestos trade, published monthly since
July 1919, 16th Floor, Enquirer Bldg., Philadelphia, Pa.
- BECKER & HAAG. Asbestos. Book published by Becker & Haag, Berlin, 1923,
88 pp.
- BERLINBAUT, L. Asbestos Mining Reviving in Russia. Eng. and Min. Jour.
Press, vol. 121, 1926, pp. 164-167.
- BOWLES, OLIVER. Asbestos. Chapter in Political and Commercial Geology.
Edited by J. E. Spurr, 1920, pp. 388-401.
- BUREAU OF FOREIGN AND DOMESTIC COMMERCE. Asbestos, Source and Trade.
Trade Inf. Bull. 442, 1926, 22 pp.
- Asbestos; World Production and Trade. Trade Inf. Bull. 20, 1922,
10 pp.
- CIRREL, FRITZ. Chrysotile-Asbestos. Its Occurrence, Exploitation, Mining, and
Uses. 2d ed. Canada Dept. Mines, Mines Branch, 1910, 316 pp.
- COOKE, H. C. Asbestos Deposits of Thetford District, Quebec. Econ. Geol.,
vol. 31, no. 4, 1936, pp. 355-376.
- The Composition of Asbestos and Other Fibers of Thetford District,
Quebec. Trans. Roy. Soc. Canada, third ser., vol. 29, sec. 4, 1935, pp. 7-19.
- COLLSON, A. L. Asbestos in the Ceded Districts of the Madras Presidency:
With Notes on Its Occurrence in Other Parts of India. Mem., Geol. Sur-
vey India, vol. 64, pt. 2, 1934, 266 pp.
- DILLER, J. S. The Types and Modes of Occurrence of Asbestos in the United
States. Jour. Canadian Min. Inst., vol. 14, 1911.
- DRESSER, JOHN A. Asbestos in Southern Quebec. Trans. Am. Inst. Min. Eng.,
vol. 50, 1915.
- The Asbestos Deposits of the Eastern Townships of Quebec. Econ.
Geol., vol. 4, no. 2, 1909, pp. 132-140.
- FISHER, NORMAN R. The Quebec Asbestos Industry. Canadian Min. Jour.,
vol. 44, 1923, pp. 649-653.
- GAMBLE, WILLIAM B. Asbestos. A list of references to material in the New
York Public Library, 1929, 72 pp.
- HALL, A. L. Asbestos in the Union of South Africa. Union of South Africa
Geol. Survey Mem. 12, 2d ed., 1930, 324 pp.
- On the Asbestos Occurrences Near Kaapsche Hoop in the Barbeton
District. Trans. Geol. Soc. South Africa, vol. 24, 1921, pp. 163-181.
- HOPKINS, OLIVER B. Asbestos Deposits of Georgia. Trans. Am. Inst. Min.
Eng., vol. 50, 1915, pp. 964-973.
- Asbestos, Talc, and Soapstone Deposits of Georgia. Geol. Survey of
Georgia Bull. 29, 1914, 319 pp.
- IMPERIAL MINERAL RESOURCES BUREAU. Asbestos, War Period (1913-19).
London, 1921, 34 pp.
- The Mineral Industry of the British Empire and Foreign Countries.
Asbestos, Statistics (1919-21). London, 1924, 16 pp.; Statistics (1920-22).
1925, 23 pp.
- IMPERIAL INSTITUTE. The Mineral Industry of the British Empire and For-
eign Countries. Statistical Summary, 1933-35. London, 438 pp.

- JONES, ROBERT H. Asbestos. Its Properties, Occurrence, and Uses. London. 1890, 236 pp.
- KEEP, F. E. Geology of the Chromite and Asbestos Deposits of the Umvukwe Range, Lomagundi and Mazoe Districts. Southern Rhodesia Geol. Survey Bull. 14, 1930, 105 pp.
- The Chrysotile Asbestos Deposits of Shabani, S. Rhodesia. South African Min. and Eng. Jour., Apr. 12, 1930, pp. 168-170.
- The Shabani Asbestos Deposits. South African Min. and Eng. Jour., Mar. 3, 1928, pp. 5-6.
- KUPFERBURGER, W. Mining Asbestos in the Pietersburg District. South Africa. Eng. and Min. Jour., vol. 130, 1930, pp. 571-574.
- MARCUSE, B. Asbestos (chapter in Marketing of Metals and Minerals, edited by J. E. Spurr and F. E. Wormser). 1925, pp. 235-242.
- Pierre à Cotton. Canadian Geog. Jour., vol. 1, no. 6, 1930, pp. 497-522.
- MINERAL INDUSTRY (The). Edited by G. A. Rousch. A series of volumes published annually from 1892, statistics, technology, and trade. Chapter on asbestos each year.
- MINERAL PRODUCTION OF CANADA. Prepared by Dominion Bureau of Statistics. Published annually.
- MINERAL RESOURCES OF THE UNITED STATES. Pt. II, NONMETALS. Statistical and economic compilation, published each year from 1924 to 1931 by the U. S. Bureau of Mines; before 1924 by the U. S. Geol. Survey; since 1931 published as Minerals Yearbook; chapter on asbestos each year.
- MINERALS YEARBOOK. Statistical and economic compilation, published each year since 1931 by the U. S. Bureau of Mines; before 1932 published as Mineral Resources of the United States; chapter on asbestos each year.
- QUEBEC BUREAU OF MINES, ANNUAL REPORTS. Mining operations and statistics.
- ROSS, J. G. Chrysotile Asbestos in Canada. Canadian Dept. Mines, Mines Branch, no. 707, 1931, 146 pp.
- Itt KEYSER, WALTER A. Asbestos Mining in Quebec. Eng. and Min. Jour., vol. 133, 1932, pp. 17-22.
- Asbestos Mining and Milling in Quebec. Eng. and Min. Jour. Press. vol. 113, 1922, pp. 617-625, 670-677.
- Chrysotile Asbestos in the Bajenova District, U. S. S. R. Eng. and Min. Jour., vol. 134, 1933, pp. 335-339.
- Mining Asbestos in U. S. S. R. Eng. and Min. Jour., vol. 134, 1933, pp. 375-381.
- Asbestos Milling in the Urals. Eng. and Min. Jour., vol. 134, 1933, pp. 415-419.
- SPENCE, HUGH S. Asbestos Industry. Trans. Canadian Inst. Min. and Met., vol. 28, 1925, pp. 94-98.
- TABER, STEPHEN. The Genesis of Asbestos and Asbestiform Minerals. Am. Inst. Min. Eng. Bull. 119, November 1916, pp. 1973-1998. (Includes discussion.)
- U. S. TARIFF COMMISSION. Asbestos. Tariff Inf. Surveys, rev. ed., 1921, 56 pp.
- Report to President. Russian Asbestos, Rept. 67, 2d ser., 1933, 11 pp.

After this publication has served your purpose, no further need for it, please return it to the use of this mailing label to do so with and no postage stamps will be required.

UNITED STATES
DEPARTMENT OF THE INTERIOR
BUREAU OF MINES

OFFICIAL BUSINESS
RETURN PENALTY LABEL
This label can be used only for returning official publications. The address must not be changed.

BUREAU OF MINES

WA

Occurrence, and Uses. London.

" Deposits of the Umrukwe
Southern Rhodesia Geol. Survey

" Shabani. S. Rhodesia. South
pp. 168-170.

" African Min. and Eng. Jour.,

" in the Pietersburg District.
30, 1930, pp. 571-574.

" of Metals and Minerals, edited
pp. 235-242.

" vol. 1. no. 6, 1930, pp. 497-522.

" usch. A series of volumes pub-
nology, and trade. Chapter on

Dominion Bureau of Statistics.

PT. II. NONMETALS. Statistical
year from 1924 to 1931 by the
the U. S. Geol. Survey; since
pter on asbestos each year.

" ic compilation. published each
lines: before 1932 published as
chapter on asbestos each year.

" Mining operations and statistics.

Canadian Dept. Mines, Mines

Quebec. Eng. and Min. Jour.,

" c. Eng. and Min. Jour. Press,

District, U. S. S. R. Eng. and

and Min. Jour., vol. 134, 1933,

" a. Min. Jour., vol. 134, 1933,

Canadian Inst. Min. and Met.,

" nd Asbestiform Minerals. Am.
pp. 1973-1998. (Includes dis-

" f. Surveys, rev. ed., 1921, 56 pp.
ept. 67, 2d ser., 1933, 11 pp.

After this publication has served your purpose and if you have
no further need for it, please return it to the Bureau of Mines.
The use of this mailing label to do so will be official business,
and no postage stamps will be required

UNITED STATES
DEPARTMENT OF THE INTERIOR
BUREAU OF MINES

PENALTY FOR PRIVATE USE TO AVOID
PAYMENT OF POSTAGE, \$300

OFFICIAL BUSINESS
RETURN PENALTY LABEL
This label can be used only for returning
official publications. The address must not
be changed.

BUREAU OF MINES,
WASHINGTON, D. C.

ASARCO ALV 0006343