

PLAINTIFF'S EXHIBIT

SA-441

ehp

ENVIRONMENTAL
HEALTH PERSPECTIVES

Volume 9, December 1974

RECEIVED

JUN 12

5:03PM

PRINTED

JUN 12

5:03PM

Environmental Health Perspectives
Vol. 9, pp. 313-315, 1974

Tumorigenic Effect of Fibrous Dusts in Experimental Animals

by F. Pott,* F. Huth,* and K. H. Friedrichs*

NOTICE
This material may be
protected by copyright
© 1974 by IRLS Code 1

Fibrous dusts (chrysotile, glass fibers, nemalite, palygorscite, and gypsum) and granular dusts (actinolite, biotite, hematite, pectolite, sanidine, and talcum) were injected intraperitoneally into rats. The fibrous dusts (other than gypsum) resulted in a high incidence of mesothelioma (30 - 67%). Gypsum produced only 5% and granular dusts none at all. It is suggested that the fibrous shape leads to a high multiplication rate of cells and predisposes to tumor formation. Fibroids, in the other hand, does not so predispose. Milled chrysotile with 99.8% fibers than 5 μ m in length are carcinogenic in our experience. The carcinogenicity of glass fibers in our experiments may have significance for occupational situations.

The starting point of our investigations was the question whether the tumorigenic effect of asbestos fibers depends on physicochemical properties of the fiber or the shape of the fibers which are characteristic for all kinds of asbestos. For this purpose chemically different fibrous forms were compared to chemically similar dusts having different forms.

Dusts Tested

Tables 1 and 2 list the dusts tested in the animal experiments with respect to their chemical composition, particle shape, fiber length, and particle size. The fiber length and particle size were estimated by evaluation of electron micrographs.

Experimental Methods

The dusts were injected intraperitoneally in Wistar rats. We could not see any difference between the reaction of peritoneum and pleura. In addition, the injection did not essentially disturb the general status of the animals.

*Medizinisches Institut für Lufthygiene und Silikonforschung, Universität Düsseldorf, Düsseldorf, Germany.

Table 1. Estimation of fiber length of fibrous dust by Electron microscopy.

Dust	Chemical composition	Fiber length	
		<2 μ m, %	<5 μ m, %
Chrysotile A	Mg silicate	78.7	93.9
Chrysotile (milled)	Mg silicate	97.4	99.8
Glass	Na, Ca borosilicates	49.9	72.6
Gypsum	Ca sulfate	85.0	75.0
Nemalite	Mg hydroxide	91.5	96.4
Palygorscite	Mg, Al silicates	87.5	70.0

Table 2. Estimation of particle size of grain dusts, by electron microscopy.

Dust	Chemical composition	Particle size	
		<2 μ m, %	<5 μ m, %
Actinolite	Ca, Mg, Fe silicates	a	a
Biotite	K, Fe, Mg, Al silicates	86.5	96.3
Hematite	Fe oxide (precipitated)	a	a
Hematite	Fe oxide (mineral)	a	a
Pectolite	Ca, Na silicates	26.2	98.4
Sanidine	K, Al silicates	85.6	97.2
Talc	Mg silicate	a	a

a No particle size analysis possible.

The best experimental method would be the inhalation of the dusts, but this method leads to

enormous technical difficulties. The time required to produce tumors following injection of dusts is already long. This time would be prolonged tremendously by an inhalation method, since it takes months to get an effective dose at the vulnerable cell structures. The necessary concentration time of the dusts and the time to produce the tumors may even exceed normal life span.

The test dusts were suspended in saline solution at concentrations up to 25 mg/2 ml. For higher dosages, the injections were repeated once a week. The different test groups consisted 40 rats. Pure saline was injected in 80 control animals. The rats were observed until spontaneous death or sacrifice. All tumors were studied histologically.

Results

The UICC standard asbestos as well as palygorskite, nemalite, and glass fibers induced tumors in the abdominal cavity of 30–67% of the rats. Several dusts chemically closely related to chrysotile (like actinolite, biotite, pectolite, talcum), but of granular or platy shape with only a few fibers did not lead to the development of tumors except for a few cases. Calcium sulfate (gypsum) did show a fibrous shape but dissolved in the animal tissue and induced tumors only in 5% of the animals. Table 3 shows the tumor rate within the different experimental groups. Histologically nearly all the tumors were sarcomatous mesotheliomata.

In current experiments the lowest effective dosage of glass fibers was 2 mg/rat. The glass fibers were uncoated. The average diameter of the glass fibers was about 0.5 μ m. With this dosage the first mesotheliomata occurred at 17 months. It seems of importance that 2 mg of glass fibers induced only slight adhesions around the liver lobes; 10 mg led to slight or medium adhesions and fibrous alterations, especially around the liver and the stomach; 50 mg of fiber dust induced tremendous adhesions on all abdominal organs with strong fibrosis. The fibrotic alterations generally could be differentiated from the numerous tumors. With 100 mg glass fiber only 6.5 months were necessary for tumor induction in the first rat.

Activity of Fibers Within the Tissues

We are confronted with the question: Why do special shapes of particles lead to formation of

Table 3. Tumor rate after intraperitoneal injection of fibrous and granular dusts.

Dust	Doses (i.p.), mg	Time required to produce first tumor in animals of group, days	Tumor rate, %
Chrysotile A	6	343	67.5
Chrysotile A	25	276	65
Chrysotile A	4 × 25	270	37.5
Chrysotile A (milled)	4 × 25	400	30
Glass fibers	4 × 25	197	57.5
Nemalite	4 × 25	249	62.5
Palygorskite	3 × 25	257	65
Gypsum	4 × 25	546	5
Pectolite	4 × 25	569	2.5
Sanidine	4 × 25	743	2.5
Talcum	4 × 25	587	2.5
Actinolite	4 × 25	—	—
Biotite	4 × 25	—	—
Hematite	4 × 25	—	—
(precipitated)			
Hematite	4 × 25	—	—
(mineral)			
NaCl (control)	4 × 2 ml	—	—

tumors? We don't have a complete answer, but, we now make the following speculation. Animal experiments and the histological investigation showed formation of tumors in the mesothelium in an overwhelming degree. Some authors have reported lung tumors following asbestos application. In our experiments subcutaneous injection of chrysotile A in one case only led to tumors in the subcutaneous connective tissue space. In addition, the storage of fibers by the lymph nodes never induced tumors of the lymphatic system. These observations suggest that tissues have different vulnerabilities to fibrous dusts. Epithelial and epithelium-like cells obviously conflict with fibrous dusts in an intensive degree; this intensive contact can be well observed in cell cultures. Beck and Bruch (1) demonstrated by electron microscopy a partial invagination of long asbestos and glass fibers by fibroblasts (L-cells) of mice. Although these cells can not phagocytose the fibers completely they remain dense around the foreign bodies. Such a relation between fiber and cells could support a chronically enhanced reproduction of cells within quickly regenerating epithelial and mesothelial cells. For decades it has been known that in epithelial organs like the liver and the respiratory tract chronically enhanced regeneration can change to abnormal

regene
prolife
We
wheth
cells
irritat
Fin
duced
as sta
Even
30–4
that
degre

The P

The
facto
exper
diam
speci
by th
has
The
and
kinog
plete
brell
shor
duce
only
99.8
also
can
thar
dian
duce
dian
reac
yet