

AR 226-1205

1089P 8EHQ-0103-00373

RECEIVED
FEB 19 2003


3M

Certified Mail

January 27, 2003

8EHQ-20-373

Document Processing Center
EPA East - Room 6428 Attn: Section 8(e)
Office of Pollution Prevention and Toxics
US EPA
1200 Pennsylvania Avenue NW
Washington DC 20460-0001



RE: TSCA 8(E) SUPPLEMENTAL SUBMISSION: Docket No. 8EHQ-998-374
Perfluorooctane Sulfonate (PFOS) CAS# 2795-39-3

Dear Docket Coordinators:

3M has previously informed the EPA (September 14, 1998) of the results of a multi-generation reproductive/developmental study in rats conducted with potassium perfluorooctane sulfonate (PFOS) in which perinatal mortality occurred in the F₁ generation pups at the two highest dose levels (1.6 and 3.2 mg/kg/day). 3M subsequently informed the EPA (March 28, 2001) of the results of a follow-up study that more precisely defined the No-Observed-Effects Level (0.4 mg/kg/day) and the mechanism for the observed toxicity.

In today's submission, 3M has new results to report from a study conducted to better define the dose-response for the perinatal effects previously observed. 3M has received a final report (see enclosure) indicating an effect on mean pup body weight per litter at the lowest maternal treatment level (0.4 mg/kg/day) that was statistically significant ($p < 0.05$) on post-natal days 1-5 when compared to control litters. The effect on mean pup body weight per litter also displayed a dose-response over the range of doses tested (0.4, 0.8, 1.0, 1.2, 1.6, and 2.0 mg/kg/day). The lower 95% confidence interval of the benchmark dose representing a 10% decrease in mean pup weight per litter on post-natal day 1 was 0.88 mg/kg/day maternal dose.

Please contact John Butenhoff (651-733-9162) if you have any questions or if we can provide additional information.

Sincerely,

Larry R. Zobel, MD MPH
Staff Vice President and Medical Director

Contain NO CBI

Enclosure

000001

MR 67014

2003 FEB 19 AM 9:04

RECEIVED
OPT NCIC